

REVIEW ARTICLE

WILEY

Semaglutide for the treatment of overweight and obesity: A review

Natasha Chidekel Bergmann MD¹  | Melanie J. Davies MD^{2,3}  |
Ildiko Lingvay MD⁴  | Filip K. Knop MD^{5,6,7} 

¹Center for Clinical Metabolic Research, Gentofte Hospital, University of Copenhagen, Hellerup, Denmark

²Diabetes Research Centre, University of Leicester, Leicester, UK

³NIHR Leicester Biomedical Research Centre, Leicester, UK

⁴Department of Internal Medicine/Endocrinology and Department of Population and Data Sciences, University of Texas Southwestern Medical Center, Dallas, Texas, USA

⁵Steno Diabetes Center Copenhagen, Herlev, Denmark

⁶Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

⁷Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

Correspondence

Professor Filip K. Knop, MD, PhD, Center for Clinical Metabolic Research, Gentofte Hospital, Gentofte Hospitalsvej 7, 3rd floor, DK-2900 Hellerup, Denmark.
Email: filip.krag.knop.01@regionh.dk

Funding information

Novo Nordisk A/S; Novo Nordisk, Grant/Award Number: GPP3

Abstract

Obesity is a chronic, relapsing disease associated with multiple complications and a substantial morbidity, mortality and health care burden. Pharmacological treatments for obesity provide a valuable adjunct to lifestyle intervention, which often achieves only limited weight loss that is difficult to maintain. The Semaglutide Treatment Effect in People with obesity (STEP) clinical trial programme is evaluating once-weekly subcutaneous semaglutide 2.4 mg (a glucagon-like peptide-1 analogue) in people with overweight or obesity. Across STEP 1, 3, 4 and 8, semaglutide 2.4 mg was associated with mean weight losses of 14.9%–17.4% in individuals with overweight or obesity without type 2 diabetes from baseline to week 68; 69%–79% of participants achieved $\geq 10\%$ weight loss with semaglutide 2.4 mg (vs. 12%–27% with placebo) and 51%–64% achieved $\geq 15\%$ weight loss (vs. 5%–13% with placebo). In STEP 5, mean weight loss was -15.2% with semaglutide 2.4 mg versus -2.6% with placebo from baseline to week 104. In STEP 2 (individuals with overweight or obesity, and type 2 diabetes), mean weight loss was -9.6% with semaglutide 2.4 mg versus -3.4% with placebo from baseline to week 68. Improvements in cardiometabolic risk factors, including high blood pressure, atherogenic lipids and benefits on physical function and quality of life were seen with semaglutide 2.4 mg. The safety profile of semaglutide 2.4 mg was consistent across trials, primarily gastrointestinal adverse events. The magnitude of weight loss reported in the STEP trials offers the potential for clinically relevant improvement for individuals with obesity-related diseases.

KEYWORDS

anti-obesity drug, GLP-1 analogue, obesity therapy, weight management

1 | INTRODUCTION

Obesity is a highly prevalent, chronic, relapsing disease requiring long-term management.^{1–3} The clinical complications of obesity affect almost every organ system, and the impact of obesity on morbidity, mortality and health care costs is substantial.^{4–7} The prevalence of

obesity has risen globally for the past several decades, a trend predicted to continue.^{8,9} Worldwide obesity prevalence is 13%,¹⁰ but many countries have a much higher prevalence; for example, prevalence in adults increased from 31% to 42% between 1999–2000 and 2017–2018 in the USA,¹¹ with an increase from 10% to 40% across most European countries over a 10-year period to 2017.⁸

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2022 The Authors. *Diabetes, Obesity and Metabolism* published by John Wiley & Sons Ltd.

Lifestyle modification is the guideline-recommended foundation of treatment for individuals with overweight or obesity,^{7,12-14} but typically achieves only modest weight loss that is often regained.¹⁵⁻¹⁷ Furthermore, this loss is challenging to maintain because of metabolic adaptation that promotes gradual weight regain.^{13,17} Pharmacological treatments for obesity provide a valuable adjunct to lifestyle interventions, but until recently the available agents only offered moderate weight loss over that achieved with lifestyle intervention.¹⁸ Semaglutide is a potent long-acting glucagon-like peptide-1 (GLP-1) analogue that requires once-weekly administration.¹⁹ The Semaglutide Treatment Effect in People with obesity (STEP) clinical trial development programme is evaluating once-weekly subcutaneous semaglutide 2.4 mg in people with overweight or obesity. Based on data from the STEP trials, once-weekly subcutaneous semaglutide 2.4 mg has been approved in Canada, Europe, the UK and the USA for chronic weight management in adults with overweight (with weight-related comorbidities) or obesity.²⁰⁻²³

The aim of this review is to describe the study designs and clinical outcomes of the published trials from the STEP programme, discuss clinical implications of the data and practicalities of using semaglutide 2.4 mg in individuals with obesity and place the STEP programme into the context of the current and future obesity pharmacotherapy landscapes.

2 | PATHOPHYSIOLOGY OF OBESITY

Obesity is associated with alterations in the quantity, distribution and function of adipose tissue. Most cases result from complex interactions between genetic, metabolic, neuroendocrine, behavioural and environmental factors involved in the regulation of energy balance and fat storage.^{24,25} In addition to these complex interactions, the imbalance in activity level and energy intake ultimately resulting in increased fat storage may be impacted by individual epigenetics and the gut microbiome.²⁶

3 | MANAGEMENT OF OBESITY

Older pharmacological options for chronic weight management, such as orlistat, phentermine-topiramate and naltrexone-bupropion, typically show moderate efficacy (~3%-9% mean weight loss over that achieved with lifestyle intervention alone).^{18,27-31} Liraglutide 3.0 mg once daily administered subcutaneously was the first GLP-1 receptor agonist (GLP-1RA) to be approved for weight management, after demonstrating weight losses of 4%-6% over those achieved with lifestyle intervention alone in clinical trials of 20-56 weeks' duration. In these trials, 46%-76% of participants lost $\geq 5\%$ and 23%-37% lost $\geq 10\%$ of their baseline body weight.^{18,32-35}

Semaglutide is a GLP-1 analogue shown to reduce energy intake, reduce hunger and increase feelings of satiety and fullness.^{19,36} This effect has been shown to arise via GLP-1 receptor activation in the central nervous system, with further indirect modulation of neuronal activity involved in appetite regulation and food intake and preference.³⁷ The

STEP programme was designed to comprehensively explore the efficacy of once-weekly subcutaneous semaglutide 2.4 mg in people with overweight or obesity, with each trial addressing a specific research question. For the six completed trials discussed in this review, these are: STEP 1 (large pivotal study), weight loss³⁸; STEP 2, weight loss in type 2 diabetes³⁹; STEP 3, weight loss in combination with intensive behavioural therapy⁴⁰; STEP 4, effect of continuing versus withdrawing semaglutide on weight-loss maintenance⁴¹; STEP 5, weight maintenance over 2 years⁴²; and STEP 8, head-to-head comparison of semaglutide and liraglutide.⁴³ In addition to these STEP trials, a regional phase IIIa trial, STEP 6, assessed the effect of semaglutide versus placebo for weight management in 401 adults from east Asia (Japan and South Korea) with obesity, with or without type 2 diabetes.⁴⁴

4 | STEP TRIAL DESIGNS

Table 1 compares key aspects of the trial designs, eligibility criteria, participant demographics and baseline characteristics of the global trials.

In STEP 1, the largest trial (N = 1961),³⁸ participants were randomized to receive once-weekly subcutaneous semaglutide 2.4 mg or matching placebo for 68 weeks. At week 68, treatments (including lifestyle intervention) were discontinued and an off-treatment extension followed a representative subset of participants for a further year.

STEP 2³⁹ enrolled individuals with type 2 diabetes [glycated haemoglobin (HbA1c) 7.0%-10.0%], and randomized them to once-weekly subcutaneous semaglutide 2.4 mg, semaglutide 1.0 mg (the approved dose in type 2 diabetes) or placebo for 68 weeks.

STEP 3⁴⁰ was a 68-week trial that compared once-weekly subcutaneous semaglutide 2.4 mg with placebo as an adjunct to intensive behavioural therapy and an initial low-calorie meal-replacement diet for all participants.

STEP 4⁴¹ was a 68-week withdrawal trial, designed to assess the effect on weight change of continuing versus discontinuing once-weekly subcutaneous semaglutide 2.4 mg after an initial 20-week semaglutide run-in period.

STEP 5⁴² is the longest trial of the STEP programme. Participants were randomized to receive once-weekly subcutaneous semaglutide 2.4 mg or matching placebo for 104 weeks.

STEP 6⁴⁴ was a 68-week trial in adults from east Asia with a body mass index (BMI) of ≥ 27.0 kg/m² with ≥ 2 weight-related comorbidities or a BMI of ≥ 35.0 kg/m² with ≥ 1 weight-related comorbidity (one comorbidity had to be either hypertension, dyslipidaemia or, in Japan only, type 2 diabetes). Participants were randomized to semaglutide 2.4 mg, 1.7 mg or placebo.

STEP 8⁴³ was a head-to-head comparison in which participants were randomized to receive once-weekly subcutaneous semaglutide 2.4 mg or matching placebo, or once-daily liraglutide 3.0 mg or matching placebo for 68 weeks.

All trials followed the same dose-escalation regimen. Participants treated with subcutaneous semaglutide 2.4 mg were initiated on a dose of 0.25 mg once weekly and the dose was escalated every 4 weeks to 0.5 mg, 1.0 mg and 1.7 mg, until the target dose of 2.4 mg was reached at week 16. If participants were unable to tolerate the

TABLE 1 STEP trials 1-5 and 8: study designs and participant population baseline demographics and clinical characteristics^{16,38-43}

	STEP 1 ^{16,38}	STEP 2 ^{16,39}	STEP 3 ^{16,40}	STEP 4 ^{16,41}	STEP 5 ^{16,42}	STEP 8 ⁴³
	Weight management	Weight management in type 2 diabetes	Weight management with intensive behavioural therapy	Sustained weight management	Two-year weight management	Semaglutide vs. liraglutide
N	1961	1210	611	902 enrolled; 803 randomized	304	338
Participants	Adults with BMI ≥30, or BMI ≥27 with ≥1 comorbidity, and without diabetes ^a	Adults with BMI ≥27 and type 2 diabetes, with HbA1c 7.0%-10.0% ^b	Adults with BMI ≥30, or BMI ≥27 with ≥1 comorbidity, and without diabetes ^a	Adults with BMI ≥30, or BMI ≥27 with ≥1 comorbidity, and without diabetes ^a	Adults with BMI ≥30, or BMI ≥27 with ≥1 comorbidity, and without diabetes ^a	Adults with BMI ≥30, or BMI ≥27 with ≥1 comorbidity, and without diabetes ^a
Treatment arms and randomization ratios	Semaglutide 2.4 mg vs. placebo 2:1 ratio	Semaglutide 2.4 mg vs. placebo 1:1 ratio	Semaglutide 2.4 mg vs. placebo 2:1 ratio	All receive semaglutide for 20 weeks (dose escalation) Then semaglutide 2.4 mg vs. placebo 2:1 ratio	Semaglutide 2.4 mg vs. placebo 1:1 ratio	Semaglutide 2.4 mg vs. placebo 3.0 mg vs. liraglutide 3:1:3:1 ratio
Duration of study	68 weeks of treatment	68 weeks	68 weeks	68 weeks (randomization into two arms at week 20)	104 weeks	68 weeks
Primary endpoints	% change in body weight ≥5% weight loss	% change in body weight ≥5% weight loss (semaglutide 2.4 mg vs. placebo)	% change in body weight ≥5% weight loss	% change in body weight	% change in body weight ≥5% weight loss	% change in body weight
Confirmatory secondary endpoints (in hierarchical testing order)	≥10% weight loss ≥15% weight loss Change in: Waist circumference Systolic blood pressure Physical functioning (SF-36 and IWQOL-lite-CT)	≥10% weight loss ≥15% weight loss Change in: Waist circumference % change in body weight (semaglutide 2.4 vs. 1.0 mg) HbA1c Systolic blood pressure Physical functioning (SF-36 and IWQOL-lite-CT)	≥10% weight loss ≥15% weight loss Change in: Waist circumference Systolic blood pressure Physical functioning (SF-36)	Change in: Waist circumference Systolic blood pressure Physical functioning (SF-36)	≥10% weight loss ≥15% weight loss Change in: Waist circumference Systolic blood pressure	≥10% weight loss ≥15% weight loss ≥20% weight loss
Background treatment	Lifestyle intervention Reduced-calorie diet (500 kcal/day deficit relative to estimated energy expenditure) and increased physical activity (150 min/week)	Lifestyle intervention Reduced-calorie diet (500 kcal/day deficit relative to estimated energy expenditure) and increased physical activity (150 min/week)	Intensive behavioural therapy Low-calorie diet (1000-1200 kcal/day) as meal replacements for first 8 weeks post-randomization then a hypocaloric diet (1200-1800 kcal/day) of conventional food for remainder of the 68 weeks. Participants were prescribed	Lifestyle intervention Reduced-calorie diet (500 kcal/day deficit relative to estimated energy expenditure) and increased physical activity (150 min/week)	Lifestyle intervention Reduced-calorie diet (500 kcal/day deficit relative to estimated energy expenditure) and increased physical activity (150 min/week)	Lifestyle intervention Reduced-calorie diet (500 kcal/day deficit relative to estimated energy expenditure) and increased physical activity (150 min/week)

TABLE 1 (Continued)

STEP 1 ^{16,38}		STEP 2 ^{16,39}	STEP 3 ^{16,40}	STEP 4 ^{16,41}	STEP 5 ^{16,42}	STEP 8 ⁴³
Weight management		Weight management in type 2 diabetes	Weight management with intensive behavioural therapy	Sustained weight management	Two-year weight management	Semaglutide vs. liraglutide
			100 min/week of physical activity at randomization, increasing by 25 min every 4 weeks, to reach 200 min/week 30 individual intensive behavioural therapy visits with a registered dietitian			
Sex, female, n (%)	1453 (74.1)	616 (50.9)	495 (81.0)	634 (79.0)	236 (77.4)	Sema/lira/placebo: 102 (81.0)/97 (76.4)/66 (77.6)
Age, years: mean (SD)	46 (13)	55 (11)	46 (13)	46 (12)	47 (11)	Sema/lira/placebo: 48 (14)/49 (13)/51 (12)
Race, n (%)						
White	1472 (75.1)	751 (62.1)	465 (76.1)	672 (83.7)	283 (93.1)	Sema/lira/placebo: 94 (74.6)/95 (74.8)/60 (70.6)
Black/African American	111 (5.7)	100 (8.3)	116 (19.0)	104 (13.0)	12 (3.9)	Sema/lira/placebo: 25 (19.8)/20 (15.7)/19 (22.4)
Asian	261 (13.3)	317 (26.2)	11 (1.8)	19 (2.4)	2 (0.7)	Sema/lira/placebo: 4 (3.2)/6 (4.7)/3 (3.5)
Other ^c	117 (6.0)	42 (3.5)	19 (3.1)	8 (1.0)	7 (2.3)	Sema/lira/placebo: 3 (2.4)/6 (4.7)/3 (3.5)
Mean body weight, kg	105.3	99.8	105.8	107.2	106.0	104.5
HbA1c, %; mean (SD)	5.7 (0.32)	8.1 (0.8)	5.7 (0.3)	5.7 (0.3)	5.7 (0.3)	Sema/lira/placebo: 5.5 (0.3)/5.5 (0.3)/5.6 (0.4)
BMI, kg/m ² ; mean (SD)	37.9 (6.7)	35.7 (6.3)	38.0 (6.7)	38.4 (6.9)	38.5 (6.9)	Sema/lira/placebo: 37.0 (7.4)/37.2 (6.4)/38.8 (6.5)
Waist circumference, cm; mean (SD)	114.7 (14.7)	114.6 (14.1)	113.0 (15.5)	115.3 (15.5)	115.7 (14.8)	Sema/lira/placebo: 111.8 (16.3)/113.5 (15.0)/115.4 (15.1)
Blood pressure, mmHg; mean (SD)						

(Continues)

TABLE 1 (Continued)

	STEP 1 ^{16,38}	STEP 2 ^{16,39}	STEP 3 ^{16,40}	STEP 4 ^{16,41}	STEP 5 ^{16,42}	STEP 8 ⁴³
	Weight management	Weight management in type 2 diabetes	Weight management with intensive behavioural therapy	Sustained weight management	Two-year weight management	Semaglutide vs. liraglutide
Systolic	126.5 (14.3)	130.0 (13.5)	124.4 (14.8)	127 (14)	125.5 (14.5)	Sema/lira/placebo: 125 (14)/126 (16)/123 (14)
Diastolic	80.3 (9.6)	79.8 (9.0)	80.5 (9.7)	81 (10)	80.1 (9.4)	Sema/lira/placebo: 81 (9)/81 (10)/79 (9)
Prediabetes ^d	885 (43.6)	NA	305 (49.9)	408 (45.3)	141 (46.4)	Sema/lira/placebo: 43 (34.1)/45 (35.4)/34 (40.0)
Number of comorbidities at screening, n (%) ^e						
0	491 (25.0)	0	148 (24.2)	214 (26.7)	NA	Sema/lira/placebo: 32 (25.4)/25 (19.7)/16 (18.8)
1	524 (26.7)	124 (10.2)	146 (23.9)	238 (29.6)		Sema/lira/placebo: 31 (24.6)/29 (22.8)/17 (20.0)
2	433 (22.1)	219 (18.1)	139 (22.7)	171 (21.3)		Sema/lira/placebo: 25 (19.8)/29 (22.8)/21 (24.7)
3	279 (14.2)	317 (26.2)	100 (16.4)	111 (13.8)		Sema/lira/placebo: 17 (13.5)/24 (18.9)/9 (10.6)
4	139 (7.1)	290 (24.0)	45 (7.4)	53 (6.6)		Sema/lira/placebo: 10 (7.9)/11 (8.7)/9 (10.6)
≥ 5	95 (4.8)	260 (21.5)	33 (5.4)	16 (2.0)		Sema/lira/placebo: 11 (8.7)/9 (7.1)/13 (15.3)

Abbreviations: BMI, body mass index; HbA_{1c}, glycated haemoglobin; IWQOL-Lite-CT, impact of weight on quality of life-lite clinical trials version questionnaire; lira, liraglutide; NA, not available; SD, standard deviation; sema, semaglutide; SF-36, short form-36 version 2 health survey, acute version.

^aKey inclusion criteria for STEP 1, 3-5 and 8 trials were age ≥18 years, BMI ≥30 kg/m² or ≥27 kg/m² with ≥1 weight-related comorbidity (treated or untreated): hypertension, dyslipidaemia, obstructive sleep apnoea or cardiovascular disease, history of at least one self-reported unsuccessful dietary effort to lose weight and no type 1 or type 2 diabetes.

^bKey inclusion criteria for the STEP 2 trial were age ≥18 years, type 2 diabetes diagnosed ≥180 days prior, BMI ≥27 kg/m², HbA_{1c} 7%-10% (53-86 mmol/mol) and treatment with diet and exercise alone or stable treatment with metformin, sulphonylurea, sodium-glucose co-transporter-2 inhibitor, glitazone as single-agent therapy or ≤3 agents for diabetes according to local label.

^cNative American, Alaska Native, Native Hawaiian, other Pacific Islander and other ethnic group. In STEP 1, this category included participants who answered 'not applicable', which is the way race or ethnic group was recorded in France.

^dFor STEP 1, the presence of prediabetes was determined by investigators based on available information (e.g. medical records, concomitant medication and blood glucose variables) and in accordance with American Diabetes Association criteria.

^eInformation collected at screening on comorbidities was based on medical history and included: type 2 diabetes (STEP 2 only), dyslipidaemia, hypertension, coronary artery disease, cerebrovascular disease, obstructive sleep apnoea, impaired glucose metabolism (not included for STEP 2), reproductive system disorders, liver disease, kidney disease, osteoarthritis, gout, thyroid disease (STEP 3 only) and asthma or chronic obstructive pulmonary disease.

2.4 mg dose because of adverse events (AEs), lower maintenance doses were permitted if the participant would otherwise discontinue trial treatment completely. It was recommended that participants made at least one attempt to re-escalate to the recommended target dose.³⁸⁻⁴⁴ Diet and physical activity (lifestyle intervention) were recorded daily in a diary or by use of a smartphone application or other tools, reviewed during counselling sessions.

4.1 | Estimands used to assess treatment efficacy

In line with regulatory guidance, the STEP trials were designed to report results using two estimands.⁴⁵ The treatment policy (primary) estimand assessed effects regardless of treatment discontinuation or rescue medication, while the secondary trial product estimand modelled treatment effects in all randomly assigned participants, assuming they had remained on treatment for the duration of the trials, and without initiation of rescue medication. The data reported in this review are based on the primary treatment policy estimand, since the statistical analyses were performed using this approach. Results using the two estimands were largely consistent.

5 | CLINICAL EVIDENCE FROM THE STEP PROGRAMME

5.1 | Demographics and baseline characteristics

STEP 1, 3, 4, 5 and 8 showed broadly similar participant demographics and baseline characteristics (Table 1).^{38,40-43} In contrast, STEP 2 had a lower proportion of female participants (51% vs. 74%-81% in the other trials), a greater mean age (55 years vs. 46-49 years) and a higher mean HbA1c (8.1% vs. 5.5%-5.7%), which is to be expected for a trial enrolling individuals with type 2 diabetes.³⁹ Mean BMI was lower in STEP 2 than the other trials (35.7 kg/m² vs. 37.5-38.4 kg/m²).³⁸⁻⁴³ The STEP 6 trial population comprised mainly Japanese participants (90%) with the remainder from South Korea. Mean body weight and BMI was lower for this population than in other STEP trials (87.5 kg and 31.9 kg/m², respectively).⁴⁴

5.2 | Efficacy outcomes

Table 2 presents results of the primary and selected key secondary and exploratory endpoints across the trials. All trials met their primary endpoints. Across the 68-week long trials in individuals with overweight or obesity with comorbidities without type 2 diabetes (STEP 1, 3, 4 and 8), semaglutide 2.4 mg was associated with a mean weight loss of 14.9%-17.4% from baseline to week 68 (Table 2).^{38,40,41,43} Furthermore, >84% of participants were receiving the full dose of semaglutide at week 68.

In STEP 1, mean weight loss with semaglutide plus usual lifestyle intervention was 14.9% (vs. 2.4% with placebo), whereas in STEP 3, mean weight loss with semaglutide plus intensive behavioural

therapy was 16.0% (vs. 5.7% with placebo),^{38,40} In STEP 4, the mean decrease in body weight during the 20-week run-in period with semaglutide treatment was 10.6%. Individuals randomized to continue semaglutide lost an additional 7.9% in body weight from weeks 20 to 68, whereas individuals who switched to placebo experienced a mean 6.9% increase.⁴¹ In STEP 8, mean weight loss was greater with semaglutide 2.4 mg than with liraglutide 3.0 mg from baseline to week 68 (15.8% vs. 6.4%)⁴³

In the STEP 1, 3, 4 and 8 trials (the 68-week long trials in participants without diabetes), weight loss of $\geq 5\%$ (a threshold widely accepted as indicating clinically meaningful response to therapy)⁴⁶ was achieved by 86%-89% of participants receiving semaglutide 2.4 mg versus 29%-48% receiving placebo (Table 2).^{38,40,41,43} In these trials, 69%-79% of participants achieved $\geq 10\%$ weight loss with semaglutide 2.4 mg (vs. 12%-27% with placebo), 51%-64% achieved $\geq 15\%$ weight loss (vs. 5%-13% with placebo), and 32%-40% achieved $\geq 20\%$ weight loss (vs. 2%-5% with placebo) (Table 2).^{38,40,41,43}

Among participants with type 2 diabetes in the STEP 2 trial, the reduction in body weight with semaglutide 2.4 mg was 9.6% (vs. 7.0% for semaglutide 1.0 mg and 3.4% for placebo) from baseline to week 68 (Table 2). In STEP 2, 69% of participants achieved $\geq 5\%$ weight loss with semaglutide 2.4 mg (vs. 57% with semaglutide 1.0 mg and 29% with placebo). In addition, participants were more likely to probably lose $\geq 10\%$, $\geq 15\%$ and $\geq 20\%$ body weight with semaglutide 2.4 mg versus placebo (46% vs. 8%, 26% vs. 3% and 13 vs. 2%, respectively) (Table 2).³⁹

In STEP 5, mean weight loss from baseline to week 104 was 15.2% with semaglutide 2.4 mg versus 2.6% with placebo. Participants were more likely to probably lose $\geq 5\%$, $\geq 10\%$ and $\geq 15\%$ and $\geq 20\%$ body weight with semaglutide versus placebo (77% vs. 34%, 62% vs. 13%, 52% vs. 7.0% and 36% vs. 2%, respectively).⁴²

In STEP 6 (east Asian population with or without diabetes), mean weight loss from baseline to week 68 was 13.2% with semaglutide 2.4 mg, 9.6% with semaglutide 1.7 mg and 2.1% with placebo. As in the other STEP trials, a larger proportion of participants achieved $\geq 5\%$, $\geq 10\%$ and $\geq 15\%$ reduction in baseline bodyweight with semaglutide 2.4 mg compared with placebo (83% vs. 21%, 61% vs. 5% and 41% vs. 3%, respectively).⁴⁴

Secondary endpoint results from the STEP trials indicated improvement in cardiometabolic risk factors including waist circumference, blood pressure, lipids and C-reactive protein with semaglutide 2.4 mg, as well as benefits on physical function and quality of life from baseline to week 68 (Table 2).³⁸⁻⁴³ Improvements in body composition with semaglutide 2.4 mg were observed in a subpopulation of participants assessed by dual-energy X-ray absorptiometry in STEP 1,³⁸ and with reductions in visceral fat areas with semaglutide 2.4 mg observed in a subpopulation of participants assessed by computed tomography scan in STEP 6.⁴⁴ Changes in glycaemic status (shift from prediabetes to normoglycaemia; tested while on treatment) were observed in STEP 1 and STEP 6. Further analyses of participants with baseline prediabetes from the STEP 1, 3 and 4 trials showed that treatment with semaglutide resulted in significantly more participants with normoglycaemia at week 68.⁴⁷

TABLE 2 Efficacy results from the STEP 1-5 and 8 trials^a

	STEP 1 ³⁸		STEP 2 ³⁹		STEP 3 ⁴⁰		STEP 4 ⁴¹		STEP 5 ⁴²		STEP 8 ⁴³	
	Weight management		Weight management in type 2 diabetes		Weight management with intensive behavioural therapy management		Sustained weight management		Two-year weight management		Semaglutide vs. liraglutide	
	Semaglutide 2.4 mg (n = 1306)	Placebo (n = 655)	Semaglutide 2.4 mg (n = 404)	Semaglutide 1.0 mg (n = 403)	Placebo (n = 403)	Semaglutide 2.4 mg (n = 407)	Placebo (n = 204)	Semaglutide 2.4 mg (n = 535)	Placebo (n = 268)	Semaglutide 2.4 mg (n = 152)	Semaglutide 2.4 mg (n = 126)	Placebo (n = 85) ^b
% weight change	-14.9	-2.4	-9.6	-7.0	-3.4	-16.0	-5.7	Baseline to week 20: -10.6; week 20-68: -7.9; week 0-68: -17.4	Week 20-68: -15.2 6.9; week 0-68: -5.0	-2.6	-15.8	-1.9
Treatment difference	-12.4		2.4 mg vs. 1.0 mg -2.7			-10.3		-14.8		-12.6	Sema vs. lira -9.4	
% of participants with weight reduction of: ^c												
≥5%	86.4	31.5	68.8	57.1	28.5	86.6	47.6	88.7	47.6	77.1	34.4	29.5
≥10%	69.1	12.0	45.6	28.7	8.2	75.3	27.0	79.0	20.4	61.8	13.3	15.4
≥15%	50.5	4.9	25.8	13.7	3.2	55.8	13.2	63.7	9.2	52.1	7.0	6.4
≥20%	32.0	1.7	13.1	4.7	1.6	35.7	3.7	39.6	4.8	36.1	2.3	2.6
Change from baseline to end of treatment in:												
Waist circumference change, cm	-13.54	-4.13	-9.4	-6.7	-4.5	-14.6	-6.3	-6.4	3.3	-14.4	-5.2	-2.0
Treatment difference	-9.42		2.4 mg vs. 1.0 mg -2.7			-8.3		-9.7		-9.2	Sema vs. lira -6.6	
Systolic blood pressure, mmHg	-6.16	-1.06	-3.9	-2.9	-0.5	-5.6	-1.6	0.5	4.4	-5.7	-1.6	3.2
Treatment difference	-5.10		2.4 mg vs. 1.0 mg -1.0			-3.9		-3.9		-4.2	Sema vs. lira -2.8	
Diastolic blood pressure, mmHg	-2.83	-0.42	-1.6	-0.6	-0.9	-3.0	-0.8	0.3	0.9	-4.4	-0.8	0.7
Treatment difference	-2.41		2.4 mg vs. 1.0 mg -0.9			-2.2		-0.6		-3.7	Sema vs. lira -4.5	
SF-36 physical functioning score	2.21	0.41	2.5	2.4	1.0	2.4	1.6	1.0	-1.5	NA	NA	NA

TABLE 2 (Continued)

	STEP 1 ³⁸		STEP 2 ³⁹		STEP 3 ⁴⁰		STEP 4 ⁴¹		STEP 5 ⁴²		STEP 8 ⁴³	
	Weight management		Weight management in type 2 diabetes		Weight management with intensive behavioural therapy management		Sustained weight management		Two-year weight management		Semaglutide vs. liraglutide	
Treatment difference	1.80		2.4 mg vs. 1.0 mg 0.1 2.4 mg vs. placebo 1.5		0.8		2.5		-		-	
IWQOL-Lite-CT physical function score	14.67	5.25	10.1	8.7	5.3	NA	NA	NA	NA	NA	NA	NA
Treatment difference	9.43		2.4 mg vs. 1.0 mg 1.4 2.4 mg vs. placebo 4.8		-		-		-		-	
Body weight, kg	-15.3	-2.6	-9.7	-6.9	-3.5	-16.8	-6.2	-7.1	-16.1	-3.2	-15.3	-6.8
Treatment difference	-12.7		2.4 mg vs. 1.0 mg -2.7 2.4 mg vs. placebo -6.1		-10.6		-13.2		-12.9		Sema vs. lira -8.5	-1.6
BMI, kg/m ²	-5.54	-0.92	-3.5	-2.5	-1.3	-6.0	-2.2	-2.6	-5.9	-1.6	NA	NA
Treatment difference	-4.61		2.4 mg vs. 1.0 mg -1.0 2.4 mg vs. placebo -2.3		-3.8		-4.7		-12.6		-	
HbA1c, %	-0.45	-0.15	-1.6	-1.5	-0.4	-0.51	-0.27	-0.1	-0.4	-0.1	-0.2	-0.1
Treatment difference	-0.29		2.4 mg vs. 1.0 mg -0.2 2.4 mg vs. placebo -1.2		-0.24		-0.2		-0.3		Sema vs. lira -0.2	0.1
Fasting plasma glucose, mg/dl	-8.35	-0.48	-38.0	-32.2	-1.4 ^c	-6.73	-0.65	-0.8	-7.6	1.7	-8.3	-4.3
Treatment difference	-7.87		2.4 mg vs. 1.0 mg -5.7 2.4 mg vs. placebo -36.6		-6.09		-7.5		-9.3		Sema vs. lira -3.9	3.3
Lipid levels, % change ^d												
Total cholesterol	-3.0	0	-1.0	-2.0	-1.0	-3.8	2.1	5	-3.3	1.4	-7.1	-0.1
Treatment difference	-3.0		2.4 mg vs. 1.0 mg 1.0 ^e 2.4 mg vs. placebo -1.0		-5.8		-6		-4.6		Sema vs. lira -7.0	-3.3
HDL cholesterol	5.0	1.0	7.0	5.0	4.0	6.5	5.0	18	9.6	8.1	-0.3	1.9
Treatment difference	4.0		2.4 mg vs. 1.0 mg 2.0 2.4 mg vs. placebo 3.0		1.5		0		1.3		Sema vs. lira -2.2	-0.9
LDL cholesterol	-3.0	1.0	0	-1.0	0	-4.7	2.6	1	-6.1	-2.7	-6.5	0.9
Treatment difference	-4.0		2.4 mg vs. 1.0 mg 1.0 2.4 mg vs. placebo 0		-7.1		-6		-3.4		Sema vs. lira -7.3	-1.1
VLDL cholesterol	-22.0	-7.0	-21.0	-17.0	-10.0	-22.5	-6.6	-6	-18.9	3.3	-20.7	-10.9
Treatment difference	-16.0		2.4 mg vs. 1.0 mg -5.0 2.4 mg vs. placebo -12.0		-17.0		-18		-21.5		Sema vs. lira -11.0	-4.1
Free fatty acids	-17.0	-7.0	-16.0	-14.0	-1.0	-11.9	4.0	-18	0.3	7.0	-12.6	-8.8
												2.6

(Continues)

TABLE 2 (Continued)

	STEP 1 ³⁸	STEP 2 ³⁹	STEP 3 ⁴⁰	STEP 4 ⁴¹	STEP 5 ⁴²	STEP 8 ⁴³
	Weight management	Weight management in type 2 diabetes	Weight management with intensive behavioural therapy management	Sustained weight management	Two-year weight management	Semaglutide vs. liraglutide
Treatment difference	−11.0	2.4 mg vs. 1.0 mg −3.0 2.4 mg vs. placebo −16.0	−15.3	−5	−6.2	Sema vs. lira −4.2
Triglycerides	−22.0	−22.0	−22.5	−6	−19.0	−20.7
Treatment difference	−16.0	2.4 mg vs. 1.0 mg −6.0 2.4 mg vs. placebo −14.0	−6.5	−18	−21.9	Sema vs. lira −11.0

Abbreviations: BMI, body mass index; HbA_{1c}, glycated haemoglobin; HDL, high-density lipoprotein; IWQOL-Lite-CT, impact of weight on quality of life-lite clinical trials version questionnaire; LDL, low-density lipoprotein; NA, not available; SF-36, short form-36 version 2 health survey, acute version; VLDL, very low-density lipoprotein.

^aResults are estimated data for the treatment policy estimand (effects assessed regardless of treatment discontinuation or rescue intervention); data and treatment differences are reported for the randomized period [i.e. baseline (week 20 in STEP 4) to week 68 for STEP 1, 2, 3, 4 and 8, and from baseline to week 104 for STEP 5].

^bPooled placebo data.

^cObserved percentages of participants who had body weight reductions of ≥5%, ≥10%, ≥15% and ≥20% from baseline to week 68 (week 104 for STEP 5) during the in-trial observation period, based on the number of participants for whom data were available at the week 68 visit (week 104 for STEP 5).

^dFor STEP 1 and 2, lipid parameters were initially analysed on a log scale as estimated ratio of end of treatment value to baseline. For interpretation, these data are expressed as relative percentage change and were calculated using the formula (estimated ratio − 1) × 100.

In STEP 2, mean reduction in HbA_{1c} from baseline to week 68 with semaglutide 2.4 mg was 1.6%, versus 1.5% with semaglutide 1.0 mg and 0.4% with placebo. More patients receiving semaglutide decreased use of concomitant glucose-lowering medications. Furthermore, urine albumin-to-creatinine ratios were improved with semaglutide in STEP 2.³⁹

5.3 | Safety

The safety profile of once-weekly subcutaneous semaglutide 2.4 mg was broadly consistent across the STEP trials. AEs, serious AEs, AEs leading to discontinuation, AEs reported in ≥10% of participants in any trial and safety focus area AEs are detailed in Table 3.³⁸⁻⁴³

AEs reported for semaglutide 2.4 mg were typical of the GLP-1RA class in general and were primarily gastrointestinal (GI) events. Most events were transient, and mild or moderate in severity. No new safety concerns arose from the STEP trials. The occurrence of cholelithiasis was consistent with the known associations between rapid weight loss and increased risk of cholelithiasis, and with previous reports of gallbladder-related disorders with GLP-1RAs.⁴⁸⁻⁵⁰ There were no notable increases in the incidence of acute pancreatitis with semaglutide 2.4 mg. In the STEP 2 trial, participants were excluded if they had uncontrolled and potentially unstable diabetic retinopathy or maculopathy. In terms of AEs, diabetic retinopathy events were reported in 4.0% of patients receiving semaglutide 2.4 mg, 2.7% with semaglutide 1.0 mg and 2.7% with placebo.³⁹

GI AEs generally led to more participants discontinuing treatment in the semaglutide 2.4 mg groups compared with placebo groups (0.8%-4.5% vs. 0%-1.2%, respectively), but overall, few participants discontinued treatment for this reason (Table 3).³⁸⁻⁴³ No new safety signals were observed with longer-term semaglutide 2.4 mg treatment in STEP 5.⁴² In STEP 8, rates of AEs (which were mostly GI-related) were similar with semaglutide 2.4 mg and liraglutide 3.0 mg (95.2% vs. 96.1%). Of note, rates of discontinuation due to AEs were lower with semaglutide than with liraglutide (3.2% vs. 12.6%).⁴³

STEP 2 was the only trial to include a once-weekly semaglutide 1.0 mg treatment arm in addition to the placebo arm. A slightly higher proportion of participants experienced GI AEs in the semaglutide 2.4 mg group versus the semaglutide 1.0 mg group (63.5% vs. 57.5%, respectively, and vs. 34.3% in the placebo group).³⁹ In the STEP 6 trial, which included once-weekly semaglutide 1.7 mg, GI AEs were reported in fewer participants in the semaglutide 2.4 mg group compared with the 1.7 mg group (59% and 64%, respectively).⁴⁴

6 | CLINICAL IMPLICATIONS OF THE STEP PROGRAMME

6.1 | Clinical implications of the trial outcomes for weight management

Across the STEP trials, once-weekly subcutaneous semaglutide 2.4 mg demonstrated mean weight loss of 14.9%-17.4% at 68 weeks in participants without diabetes with the mean baseline weight of

TABLE 3 Number and proportion of participants with adverse events in the STEP 1-5 and 8 trials

	STEP 1 ³⁸				STEP 2 ³⁹				STEP 3 ⁴⁰				STEP 4, ^{41a}				STEP 5 ⁴²				STEP 8 ⁴³			
	Weight management				Weight management in type 2 diabetes				Weight management with intensive behavioural therapy				Sustained weight management				Two-year weight management				Semaglutide vs. liraglutide			
	Semaglutide 2.4 mg (n = 1306)	Placebo (n = 655)	Semaglutide 2.4 mg (n = 403)	Semaglutide 1.0 mg (n = 402)	Semaglutide 2.4 mg (n = 402)	Placebo (n = 402)	Semaglutide 2.4 mg (n = 407)	Placebo (n = 204)	Semaglutide 2.4 mg (n = 535)	Placebo (n = 268)	Semaglutide 2.4 mg (n = 152)	Placebo (n = 152)	Semaglutide 2.4 mg (n = 126)	Placebo (n = 85)										
n (%) of participants with AEs	1171 (89.7)	566 (86.4)	353 (87.6)	329 (81.8)	309 (76.9)	390 (95.8)	196 (96.1)	435 (81.3)	201 (75.0)	146 (96.1)	136 (89.5)	120 (95.2)	122 (96.1)	81 (95.3)										
Any AE	128 (9.8)	42 (6.4)	40 (9.9)	31 (7.7)	37 (9.2)	37 (9.1)	6 (2.9)	41 (7.7)	15 (5.6)	12 (7.9)	18 (11.8)	10 (7.9)	14 (11.0)	6 (7.1)										
Serious AEs	92 (7.0)	20 (3.1)	25 (6.2)	20 (5.0)	14 (3.5)	24 (5.9)	6 (2.9)	13 (2.4)	6 (2.2)	9 (5.9)	7 (4.6)	4 (3.2)	16 (12.6)	3 (3.5)										
AE leading to discontinuation	59 (4.5)	5 (0.8)	17 (4.2)	14 (3.5)	4 (1.0)	14 (3.4)	0	NA	NA	6 (3.9)	1 (0.7)	1 (0.8)	8 (6.3)	1 (1.2)										
GI disorders leading to discontinuation	1 (0.1)	1 (0.2)	1 (0.2)	1 (0.2)	1 (0.2)	0	0	1 (0.2)	1 (0.4)	1 (0.7)	0	0	0	0										
Fatal events																								
GI AEs reported in ≥10% of participants ^c																								
Nausea	577 (44.2)	114 (17.4)	136 (33.7)	129 (32.1)	37 (9.2)	237 (58.2)	45 (22.1)	75 (14.0)	13 (4.9)	81 (53.3)	33 (21.7)	77 (61.1)	75 (59.1)	19 (22.4)										
Diarrhoea	412 (31.5)	104 (15.9)	86 (21.3)	89 (22.1)	48 (11.9)	147 (36.1)	45 (22.1)	77 (14.4)	19 (7.1)	53 (34.9)	36 (23.7)	35 (27.8)	23 (18.1)	22 (25.9)										
Vomiting	324 (24.8)	43 (6.6)	88 (21.8)	54 (13.4)	11 (2.7)	111 (27.3)	22 (10.8)	55 (10.3)	8 (3.0)	46 (30.3)	7 (4.6)	32 (25.4)	26 (20.5)	5 (5.9)										
Constipation	306 (23.4)	62 (9.5)	70 (17.4)	51 (12.7)	22 (5.5)	150 (36.9)	50 (24.5)	62 (11.6)	17 (6.3)	47 (30.9)	17 (11.2)	49 (38.9)	40 (31.5)	20 (23.5)										
Safety focus areas																								
GI disorders	969 (74.2)	314 (47.9)	256 (63.5)	231 (57.5)	138 (34.3)	337 (82.8)	129 (63.2)	224 (41.9)	70 (26.1)	125 (82.2)	82 (53.9)	106 (84.1)	105 (82.7)	47 (55.3)										
Gall-bladder-related disorders	34 (2.6)	8 (1.2)	1 (0.2)	4 (1.0)	3 (0.7)	20 (4.9)	3 (1.5)	15 (2.8)	10 (3.7)	4 (2.6)	2 (1.3)	1 (0.8)	4 (3.1)	1 (1.2)										
Hepatic disorders	31 (2.4)	20 (3.1)	10 (2.5)	10 (2.5)	14 (3.5)	8 (2.0)	4 (2.0)	11 (2.1)	4 (1.5)	3 (2.0)	3 (2.0)	2 (1.6)	1 (0.8)	3 (3.5)										
Acute pancreatitis	3 (0.2)	0	1 (0.2)	0	1 (0.2)	0	0	0	0	0	0	0	1 (0.8)	0										
Cardiovascular disorders	107 (8.2)	75 (11.5)	6 (1.5)	6 (1.5)	5 (1.2)	40 (9.8)	22 (10.8)	26 (4.9)	30 (11.2)	17 (11.2)	32 (21.1)	16 (12.7)	18 (14.2)	9 (10.6)										
Allergic reactions	96 (7.4)	54 (8.2)	26 (6.5)	22 (5.5)	18 (4.5)	35 (8.6)	19 (9.3)	26 (4.9)	11 (4.1)	23 (15.1)	8 (5.3)	9 (7.1)	11 (8.7)	10 (11.8)										
Injection-site reactions	65 (5.0)	44 (6.7)	12 (3.0)	6 (1.5)	10 (2.5)	22 (5.4)	12 (5.9)	14 (2.6)	6 (2.2)	10 (6.6)	15 (9.9)	0	14 (11.0)	5 (5.9)										
Malignant neoplasms	14 (1.1)	7 (1.1)	5 (1.2)	7 (1.7)	8 (2.0)	3 (0.7)	1 (0.5)	6 (1.1)	1 (0.4)	2 (1.3)	4 (2.6)	3 (2.4)	3 (2.4)	1 (1.2)										
Psychiatric disorders	124 (9.5)	83 (12.7)	24 (6.0)	23 (5.7)	15 (3.7)	60 (14.7)	24 (11.8)	46 (8.6)	35 (13.1)	26 (17.1)	25 (16.4)	7 (5.6)	19 (15.0)	9 (10.6)										
Acute renal failure	3 (0.2)	2 (0.3)	4 (1.0)	2 (0.5)	2 (0.5)	0	0	1 (0.2)	1 (0.4)	0	0	1 (0.8)	0	1 (1.2)										
Hypoglycaemia	8 (0.6)	5 (0.8)	23 (5.7)	22 (5.5)	12 (3.0)	2 (0.5)	0	3 (0.6)	3 (1.1)	4 (2.6)	0	0	1 (0.8)	0										

Abbreviations: AE, adverse event; GI, gastrointestinal; NA, not available.

^aData are reported for the randomized period (i.e. weeks 20-68).^bPooled placebo data.^cAE in ≥5% of participants in STEP 4.


Administration	<ul style="list-style-type: none"> • Injections should be on the same day each week • The time of day and injection site can vary each week 										
Dose-escalation schedule	 <table border="1"> <tbody> <tr> <td>Weeks 1–4</td> <td>0.25 mg</td> </tr> <tr> <td>Weeks 5–8</td> <td>0.5 mg</td> </tr> <tr> <td>Weeks 9–12</td> <td>1.0 mg</td> </tr> <tr> <td>Weeks 13–16</td> <td>1.7 mg</td> </tr> <tr> <td>Weeks 17+</td> <td>2.4 mg</td> </tr> </tbody> </table> <p>If 2.4 mg is not tolerated, stay at 1.7 mg for a further 4 weeks and then re-escalate</p>	Weeks 1–4	0.25 mg	Weeks 5–8	0.5 mg	Weeks 9–12	1.0 mg	Weeks 13–16	1.7 mg	Weeks 17+	2.4 mg
Weeks 1–4	0.25 mg										
Weeks 5–8	0.5 mg										
Weeks 9–12	1.0 mg										
Weeks 13–16	1.7 mg										
Weeks 17+	2.4 mg										
Missed doses	<ul style="list-style-type: none"> • If a dose is missed and the next scheduled dose is more than 2 days away, the dose should be administered as soon as possible • If more than 5 days have passed, the dose should be skipped, and dosing resumed on the scheduled day 										
Management of GI AEs	<ul style="list-style-type: none"> • Counsel patients on the potential for GI side effects • Consider slower dose escalation for individuals who have challenging GI symptoms in the first few weeks • Counsel patients on dietary modifications for upper GI side effects; recommend increasing fibre and water intake for constipation and consider stool softeners • Short-term use of the over-the-counter medications for more persistent / severe GI symptoms may be considered • For patients with vomiting, dietary measures, such as smaller volumes of food intake and possibly more frequently, may be helpful, along with maintenance of hydration • Since GI AEs are dose-dependent, a lower GLP-1RA dosage could be considered for those unable to tolerate the recommended maintenance dose 										
Switching from liraglutide or other GLP-1RAs	<ul style="list-style-type: none"> • Consider patient counselling, dose titration, management of GI side effects, and re-evaluation of response 										
Risk of hypoglycaemia	<ul style="list-style-type: none"> • Semaglutide lowers blood glucose and is associated with a low risk for hypoglycaemia • Consider reducing the dose of insulin secretagogue or insulin to reduce the risk of hypoglycaemia 										
Concomitant medications	<ul style="list-style-type: none"> • Semaglutide causes delay of gastric emptying and therefore has the potential to affect absorption of oral medications • Although in clinical pharmacology trials of semaglutide 1.0 mg absorption was not affected, monitoring is advised 										
Contraindications	<ul style="list-style-type: none"> • Personal or family history of medullary thyroid carcinoma or multiple endocrine neoplasia syndrome type 2 										
Special populations	<ul style="list-style-type: none"> • Monitor blood glucose in patients with diabetes • Discontinue prior to pregnancy • Consider the benefits of breast feeding alongside the mother's clinical need • There are no data to on the presence of semaglutide in human milk; animal data suggest possible exposure 										

FIGURE 1 Key considerations for using once-weekly subcutaneous semaglutide 2.4 mg in clinical practice (approved in Canada, Europe, the UK and the USA).^{20–23} Abbreviations: AE, adverse event; GI, gastrointestinal; GLP-1RA, glucagon-like peptide-1 receptor agonist




FIGURE 2 Future landscape of obesity pharmacotherapy. Ongoing trials with semaglutide, and selected ongoing and completed trials with other investigational agents. Studies in closed boxes are completed, while those in open boxes are ongoing. Abbreviations: CVD, cardiovascular disease; CVOT, cardiovascular outcomes trial; HF, heart failure; LTE, long-term extension; OA, osteoarthritis; T2D, type 2 diabetes

AUTHOR CONTRIBUTIONS

Drafting the manuscript (NCB, MJD, IL, FKK); critical revision of the manuscript for important intellectual content (NCB, MJD, IL, FKK); responsibility for content (NCB, MJD, IL, FKK).

ACKNOWLEDGMENTS

The authors thank Paul Barlass, PhD, of Axis, a division of Spirit Medical Communications Group Ltd., and Sarah Stowell, a contract writer working on behalf of Axis who provided medical writing assistance under the direction of the authors, funded by Novo Nordisk A/S, Denmark, in accordance with Good Publication Practice 3 (GPP3) guidelines (www.ismpp.org/gpp3).

CONFLICTS OF INTEREST

NCB declared no conflict of interest. MJD is a consultant, advisory board member and speaker for Boehringer Ingelheim, Eli Lilly, Novo Nordisk and Sanofi-Aventis; advisory board member and speaker for AstraZeneca; advisory board member for Gilead Sciences Ltd, Janssen, Lexicon, Pfizer and Servier; speaker for Mitsubishi Tanabe Pharma Corporation, Napp Pharmaceuticals and Takeda Pharmaceuticals International Inc.; received grants to support investigator and investigator-initiated trials from AstraZeneca, Boehringer Ingelheim, Eli Lilly, Janssen, Novo Nordisk and Sanofi-Aventis. MJD is co-funded by the NIHR Leicester Biomedical Research Centre. IL obtained research funding, advisory/consulting fees, and/or other support from AstraZeneca, Bayer, Boehringer Ingelheim, Gl Dynamics, Intarcia, Intercept, Janssen, Eli Lilly, Mannkind, Merck, Mylan, Novartis, Novo Nordisk, Pfizer, Sanofi, TARGETPharma, Valeritas and Zealand Pharma. FKK served on scientific advisory panels and/or been part of speaker bureaus for, served as a consultant to and/or received research support from Amgen, AstraZeneca, Bayer, Boehringer Ingelheim, Carmot Therapeutics, Eli Lilly, Gubra, Lupin, MedImmune, MSD/Merck, Mundipharma, Norgine, Novo Nordisk, Pharmacosmos, Sanofi, ShouTi, Zealand Pharma and Zucara; minority shareholder in Antag Therapeutics.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/dom.14863>.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Natasha Chidekel Bergmann  <https://orcid.org/0000-0002-6797-8041>

Melanie J. Davies  <https://orcid.org/0000-0002-9987-9371>

Ildiko Lingvay  <https://orcid.org/0000-0001-7006-7401>

Filip K. Knop  <https://orcid.org/0000-0002-2495-5034>

REFERENCES

- Frühbeck G, Busetto L, Dicker D, et al. The ABCD of obesity: an EASO position statement on a diagnostic term with clinical and scientific implications. *Obes Facts*. 2019;12(2):131-136.
- NCD Risk Factor Collaboration (NCD-RisC). Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet*. 2016;387(10026):1377-1396.
- Bray GA, Kim KK, Wilding JPH, World Obesity Federation. Obesity: a chronic relapsing progressive disease process. A position statement of the World Obesity Federation. *Obes Rev*. 2017;18(7):715-723.
- Guh DP, Zhang W, Bansback N, Amarsi Z, Birmingham CL, Anis AH. The incidence of co-morbidities related to obesity and overweight: a systematic review and meta-analysis. *BMC Public Health*. 2009;9:88.
- Pereira-Miranda E, Costa PRF, Queiroz VAO, Pereira-Santos M, Santana MLP. Overweight and obesity associated with higher depression prevalence in adults: a systematic review and meta-analysis. *J Am Coll Nutr*. 2017;36(3):223-233.
- Simon GE, Von Korff M, Saunders K, et al. Association between obesity and psychiatric disorders in the US adult population. *Arch Gen Psychiatry*. 2006;63(7):824-830.
- Garvey WT, Mechanick JL, Brett EM, et al. American Association of Clinical Endocrinologists and American College of Endocrinology comprehensive clinical practice guidelines for medical care of patients with obesity. *Endocr Pract*. 2016;22(Suppl 3):1-203.
- Agha M, Agha R. The rising prevalence of obesity: part A: impact on public health. *Int J Surg Oncol (N Y)*. 2017;2(7):e17.
- Janssen F, Bardoutsos A, Vidra N. Obesity prevalence in the long-term future in 18 European countries and in the USA. *Obes Facts*. 2020;13(5):514-527.
- World Health Organization. Obesity and overweight. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>. Accessed March 31, 2022.
- Hales CM, Carroll MD, Fryar CD, Ogden CL. Prevalence of obesity and severe obesity among adults: United States, 2017-2018. *NCHS Data Brief*. 2020;360:1-8.
- Durrer Schutz D, Busetto L, Dicker D, et al. European practical and patient-centred guidelines for adult obesity management in primary care. *Obes Facts*. 2019;12(1):40-66.
- Jensen MD, Ryan DH, Apovian CM, et al. AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *J Am Coll Cardiol*. 2014;63(25 Pt B):2985-3023.
- Yumuk V, Tsigos C, Fried M, et al. European guidelines for obesity management in adults. *Obes Facts*. 2015;8(6):402-424.
- Wadden TA, Tronieri JS, Butryn ML. Lifestyle modification approaches for the treatment of obesity in adults. *Am Psychol*. 2020;75(2):235-251.
- Kushner RF, Calanna S, Davies M, et al. Semaglutide 2.4 mg for the treatment of obesity: key elements of the STEP trials 1 to 5. *Obesity (Silver Spring)*. 2020;28(6):1050-1061.
- Sumithran P, Prendergast LA, Delbridge E, et al. Long-term persistence of hormonal adaptations to weight loss. *N Engl J Med*. 2011;365(17):1597-1604.
- Singh AK, Singh R. Pharmacotherapy in obesity: a systematic review and meta-analysis of randomized controlled trials of anti-obesity drugs. *Expert Rev Clin Pharmacol*. 2020;13(1):53-64.
- Blundell J, Finlayson G, Axelsen M, et al. Effects of once-weekly semaglutide on appetite, energy intake, control of eating, food preference and body weight in subjects with obesity. *Diabetes Obes Metab*. 2017;19(9):1242-1251.
- Novo Nordisk. WEGOVY (semaglutide) highlights of prescribing information. <https://www.novo-pi.com/wegovy.pdf>. Accessed November 30, 2021.
- Medicines & Healthcare Products Regulatory Agency. Wegovy 2.4 mg, solution for injection in pre-filled pen – summary of product characteristics. <https://mhraproducts4853.blob.core.windows.net/docs/>

- 4b161649ffc66e938bc76f4e45a4d3797af54914. Accessed March 31, 2022.
22. European Medicines Agency. Wegovy – summary of product characteristics. <https://www.ema.europa.eu/en/medicines/human/EPAR/wegovy>. Accessed March 31, 2022.
23. Health Canada. Wegovy – product monograph. <https://health-products.canada.ca/dpd-bdpp/info.do?lang=en&code=101171>. Accessed March 31, 2022.
24. Mechanick JL, Hurley DL, Garvey WT. Adiposity-based chronic disease as a new diagnostic term: the American Association of Clinical Endocrinologists and American College of Endocrinology position statement. *Endocr Pract.* 2017;23(3):372-378.
25. Bray GA, Frühbeck G, Ryan DH, Wilding JP. Management of obesity. *Lancet.* 2016;387(10031):1947-1956.
26. Theilade S, Christensen MB, Vilsbøll T, Knop FK. An overview of obesity mechanisms in humans: endocrine regulation of food intake, eating behaviour and common determinants of body weight. *Diabetes Obes Metab.* 2021;23(Suppl 1):17-35.
27. Broom I, Wilding J, Stott P, Myers N, UK Multimorbidity Study Group. Randomised trial of the effect of orlistat on body weight and cardiovascular disease risk profile in obese patients: UK multimorbidity study. *Int J Clin Pract.* 2002;56(7):494-499.
28. Lindgärde F. The effect of orlistat on body weight and coronary heart disease risk profile in obese patients: the Swedish multimorbidity study. *J Intern Med.* 2000;248(3):245-254.
29. Rössner S, Sjöström L, Noack R, Meinders AE, Nosedá G. Weight loss, weight maintenance, and improved cardiovascular risk factors after 2 years treatment with orlistat for obesity. *Obes Res.* 2000;8(1):49-61.
30. Hollander PA, Elbein SC, Hirsch IB, et al. Role of orlistat in the treatment of obese patients with type 2 diabetes. A 1-year randomized double-blind study. *Diabetes Care.* 1998;21(8):1288-1294.
31. Finer N, James WP, Kopelman PG, Lean ME, Williams G. One-year treatment of obesity: a randomized, double-blind, placebo-controlled, multicentre study of orlistat, a gastrointestinal lipase inhibitor. *Int J Obes Relat Metab Disord.* 2000;24(3):306-313.
32. Astrup A, Rössner S, Van Gaal L, et al. Effects of liraglutide in the treatment of obesity: a randomised, double-blind, placebo-controlled study. *Lancet.* 2009;374(9701):1606-1616.
33. Wadden TA, Hollander P, Klein S, et al. Weight maintenance and additional weight loss with liraglutide after low-calorie-diet-induced weight loss: the SCALE maintenance randomized study. *Int J Obes (Lond).* 2013;37(11):1443-1451.
34. Pi-Sunyer X, Astrup A, Fujioka K, et al. A randomized, controlled trial of 3.0 mg of liraglutide in weight management. *N Engl J Med.* 2015;373(1):11-22.
35. Blackman A, Foster GD, Zammit G, et al. Effect of liraglutide 3.0 mg in individuals with obesity and moderate or severe obstructive sleep apnea: the SCALE sleep apnea randomized clinical trial. *Int J Obes (Lond).* 2016;40(8):1310-1319.
36. Friedrichsen M, Breitschaft A, Tadayon S, Wizert A, Skovgaard D. The effect of semaglutide 2.4 mg once weekly on energy intake, appetite, control of eating, and gastric emptying in adults with obesity. *Diabetes Obes Metab.* 2021;23(3):754-762.
37. Gabery S, Salinas CG, Paulsen SJ, et al. Semaglutide lowers body weight in rodents via distributed neural pathways. *JCI Insight.* 2020;5(6):e133429.
38. Wilding JPH, Batterham RL, Calanna S, et al. Once-weekly semaglutide in adults with overweight or obesity. *N Engl J Med.* 2021;384(11):989-1002.
39. Davies M, Færch L, Jeppesen OK, et al. Semaglutide 2.4 mg once a week in adults with overweight or obesity, and type 2 diabetes (STEP 2): a randomised, double-blind, double-dummy, placebo-controlled, phase 3 trial. *Lancet.* 2021;397(10278):971-984.
40. Wadden TA, Bailey TS, Billings LK, et al. Effect of subcutaneous semaglutide vs placebo as an adjunct to intensive behavioral therapy on body weight in adults with overweight or obesity: the STEP 3 randomized clinical trial. *JAMA.* 2021;325(14):1403-1413.
41. Rubino D, Abrahamsson N, Davies M, et al. Effect of continued weekly subcutaneous semaglutide vs placebo on weight loss maintenance in adults with overweight or obesity: the STEP 4 randomized clinical trial. *JAMA.* 2021;325(14):1414-1425.
42. Garvey WT, Batterham RL, Bhatta M, et al. Two-year effect of semaglutide 2.4 mg once-weekly in adults with overweight or obesity (STEP 5): a randomized, double-blind, placebo controlled, phase 3 trial. *Nat Med.* 2022; in press.
43. Rubino DM, Greenway FL, Khalid U, et al. Effect of weekly subcutaneous semaglutide vs daily liraglutide on body weight in adults with overweight or obesity without diabetes: the STEP 8 randomized clinical trial. *JAMA.* 2022;327(2):138-150.
44. Kadowaki T, Isendahl J, Khalid U, et al. Semaglutide once a week in adults with overweight or obesity, with or without type 2 diabetes in an east Asian population (STEP 6): a randomised, double-blind, double-dummy, placebo-controlled, phase 3a trial. *Lancet Diabetes Endocrinol.* 2022;10(3):193-206.
45. Wharton S, Astrup A, Endahl L, et al. Estimating and reporting treatment effects in clinical trials for weight management: using estimands to interpret effects of intercurrent events and missing data. *Int J Obes (Lond).* 2021;45(5):923-933.
46. Williamson DA, Bray GA, Ryan DH. Is 5% weight loss a satisfactory criterion to define clinically significant weight loss? *Obesity (Silver Spring).* 2015;23(12):2319-2320.
47. Perreault L, Davies M, Frias JP, et al. Changes in glucose metabolism and glycemic status with once-weekly subcutaneous semaglutide 2.4 mg among participants with prediabetes in the STEP program. *Diabetes Care.* 2022;dc211785. doi:10.2337/dc21-1785 Online ahead of print.
48. Stokes CS, Lammert F. Excess body weight and gallstone disease. *Visc Med.* 2021;37(4):254-260.
49. Faillie JL, Yu OH, Yin H, Hillaire-Buys D, Barkun A, Azoulay L. Association of bile duct and gallbladder diseases with the use of incretin-based drugs in patients with type 2 diabetes mellitus. *JAMA Intern Med.* 2016;176(10):1474-1481.
50. Nauck MA, Muus Ghorbani ML, Kreiner E, Saevereid HA, Buse JB, LEADER Publication Committee on behalf of the LEADER Trial Investigators. Effects of liraglutide compared with placebo on events of acute gallbladder or biliary disease in patients with type 2 diabetes at high risk for cardiovascular events in the LEADER randomized trial. *Diabetes Care.* 2019;42(10):1912-1920.
51. Hall KD, Kahan S. Maintenance of lost weight and long-term management of obesity. *Med Clin North Am.* 2018;102(1):183-197.
52. Ryan DH, Yockey SR. Weight loss and improvement in comorbidity: differences at 5%, 10%, 15%, and over. *Curr Obes Rep.* 2017;6(2):187-194.
53. Wing RR, Lang W, Wadden TA, et al. Benefits of modest weight loss in improving cardiovascular risk factors in overweight and obese individuals with type 2 diabetes. *Diabetes Care.* 2011;34(7):1481-1486.
54. Dambha-Miller H, Day AJ, Strelitz J, Irving G, Griffin SJ. Behaviour change, weight loss and remission of type 2 diabetes: a community-based prospective cohort study. *Diabet Med.* 2020;37(4):681-688.
55. Aminian A, Zajichek A, Tu C, et al. How much weight loss is required for cardiovascular benefits? Insights from a metabolic surgery matched-cohort study. *Ann Surg.* 2020;272(4):639-645.
56. Lingvay I, Sumithran P, Cohen RV, le Roux CW. Obesity management as a primary treatment goal for type 2 diabetes: time to reframe the conversation. *Lancet.* 2022;399(10322):394-405.
57. Wadden TA, Neiberg RH, Wing RR, et al. Four-year weight losses in the look AHEAD study: factors associated with long-term success. *Obesity (Silver Spring).* 2011;19(10):1987-1998.
58. Svetkey LP, Stevens VJ, Brantley PJ, et al. Comparison of strategies for sustaining weight loss: the weight loss maintenance randomized controlled trial. *JAMA.* 2008;299(10):1139-1148.

59. Torgerson JS, Hauptman J, Boldrin MN, Sjöström L, Xenical in the prevention of diabetes in obese subjects (XENDOS) study: a randomized study of orlistat as an adjunct to lifestyle changes for the prevention of type 2 diabetes in obese patients. *Diabetes Care*. 2004;27(1):155-161.
60. Davies MJ, Bergenstal R, Bode B, et al. Efficacy of liraglutide for weight loss among patients with type 2 diabetes: the SCALE diabetes randomized clinical trial. *JAMA*. 2015;314(7):687-699.
61. Husain M, Bain SC, Jeppesen OK, et al. Semaglutide (SUSTAIN and PIONEER) reduces cardiovascular events in type 2 diabetes across varying cardiovascular risk. *Diabetes Obes Metab*. 2020;22(3):442-451.
62. Ryan DH, Lingvay I, Colhoun HM, et al. Semaglutide effects on cardiovascular outcomes in people with overweight or obesity (SELECT) rationale and design. *Am Heart J*. 2020;229:61-69.
63. Aroda VR, Ahmann A, Cariou B, et al. Comparative efficacy, safety, and cardiovascular outcomes with once-weekly subcutaneous semaglutide in the treatment of type 2 diabetes: insights from the SUSTAIN 1-7 trials. *Diabetes Metab*. 2019;45(5):409-418.
64. Wharton S, Davies M, Dicker D, et al. Managing the gastrointestinal side effects of GLP-1 receptor agonists in obesity: recommendations for clinical practice. *Postgrad Med*. 2022;134(1):14-19.
65. Almandoz JP, Lingvay I, Morales J, Campos C. Switching between glucagon-like peptide-1 receptor agonists: rationale and practical guidance. *Clin Diabetes*. 2020;38(4):390-402.
66. Jain AB, Ali A, Gorgojo Martínez JJ, et al. Switching between GLP-1 receptor agonists in clinical practice: expert consensus and practical guidance. *Int J Clin Pract*. 2021;75(2):e13731.
67. Lingvay I, Faerch L, Koroleva A, et al. Influence of baseline characteristics on weight loss with semaglutide 2.4 mg in adults with overweight/obesity and type 2 diabetes (STEP 2). Presented at the American Diabetes Association (ADA) virtual meeting, June 25-29, 2021.
68. Donath MY, Bircel R. GLP-1 effects on islets: hormonal, neuronal, or paracrine? *Diabetes Care*. 2013;36(Suppl 2):S145-S148.
69. ClinicalTrials.gov. A research study on how well semaglutide works in adolescents with overweight or obesity (NCT04102189). <https://clinicaltrials.gov/ct2/show/NCT04102189>. Accessed November 30, 2021.
70. ClinicalTrials.gov. Research study to investigate how well semaglutide works in people living with heart failure and obesity (STEP-HFpEF) (NCT04788511). <https://clinicaltrials.gov/ct2/show/NCT04788511>. Accessed November 30, 2021.
71. ClinicalTrials.gov. Research study to look at how well semaglutide works in people living with heart failure, obesity and type 2 diabetes (STEP HFpEF DM) (NCT04916470). <https://clinicaltrials.gov/ct2/show/NCT04916470>. Accessed November 30, 2021.
72. ClinicalTrials.gov. Research study of how well semaglutide works in people living with overweight or obesity (STEP 7) (NCT04251156). <https://clinicaltrials.gov/ct2/show/NCT04251156>. Accessed March 31, 2022.
73. ClinicalTrials.gov. Research study looking at how well semaglutide works in people suffering from obesity and knee osteoarthritis (NCT05064735). <https://clinicaltrials.gov/ct2/show/NCT05064735>. Accessed November 30, 2021.
74. ClinicalTrials.gov. Research study looking at how well semaglutide works in people living with obesity and prediabetes (STEP 10) (NCT05040971). <https://clinicaltrials.gov/ct2/show/NCT05040971>. Accessed November 30, 2021.
75. Coskun T, Sloop KW, Loghin C, et al. LY3298176, a novel dual GIP and GLP-1 receptor agonist for the treatment of type 2 diabetes mellitus: from discovery to clinical proof of concept. *Mol Metab*. 2018;18:3-14.
76. Rosenstock J, Wysham C, Frias JP, et al. Efficacy and safety of a novel dual GIP and GLP-1 receptor agonist tirzepatide in patients with type 2 diabetes (SURPASS-1): a double-blind, randomised, phase 3 trial. *Lancet*. 2021;398(10295):143-155.
77. Frias JP, Davies MJ, Rosenstock J, et al. Tirzepatide versus semaglutide once weekly in patients with type 2 diabetes. *N Engl J Med*. 2021;385(6):503-515.
78. Ludvik B, Giorgino F, Jódar E, et al. Once-weekly tirzepatide versus once-daily insulin degludec as add-on to metformin with or without SGLT2 inhibitors in patients with type 2 diabetes (SURPASS-3): a randomised, open-label, parallel-group, phase 3 trial. *Lancet*. 2021;398(10300):583-598.
79. ClinicalTrials.gov. A study of tirzepatide (LY3298176) in participants with type 2 diabetes who have obesity or are overweight (SURMOUNT-2) (NCT04657003). <https://clinicaltrials.gov/ct2/show/NCT04657003>. Accessed November 30, 2021.
80. ClinicalTrials.gov. A study of tirzepatide (LY3298176) in participants after a lifestyle weight loss program (SURMOUNT-3) (NCT04657016). <https://clinicaltrials.gov/ct2/show/NCT04657016>. Accessed November 30, 2021.
81. ClinicalTrials.gov. A study of tirzepatide (LY3298176) in participants with obesity or overweight for the maintenance of weight loss (SURMOUNT-4) (NCT04660643). <https://clinicaltrials.gov/ct2/show/NCT04660643>. Accessed November 30, 2021.
82. Frias JP, Nauck MA, Van J, et al. Efficacy and safety of LY3298176, a novel dual GIP and GLP-1 receptor agonist, in patients with type 2 diabetes: a randomised, placebo-controlled and active comparator-controlled phase 2 trial. *Lancet*. 2018;392(10160):2180-2193.
83. Jastreboff AM, Aronne LJ, Ahmad NN, et al. Tirzepatide once weekly for the treatment of obesity. *N Engl J Med*. 2022;387:205-216. doi: 10.1056/NEJMoa2206038
84. ClinicalTrials.gov. A study of tirzepatide (LY3298176) once a week versus insulin glargine once a day in participants with type 2 diabetes and increased cardiovascular risk (SURPASS-4) (NCT03730662). <https://clinicaltrials.gov/ct2/show/NCT03730662>. Accessed June 24, 2022.
85. ClinicalTrials.gov. A study of tirzepatide (LY3298176) versus placebo in participants with type 2 diabetes inadequately controlled on insulin glargine with or without metformin (SURPASS-5). (NCT04039503). <https://clinicaltrials.gov/ct2/show/NCT04039503>. Accessed June 24, 2022.
86. Frias JP, Bastyr EJ 3rd, Vignati L, et al. The sustained effects of a dual GIP/GLP-1 receptor agonist, NNC0090-2746, in patients with type 2 diabetes. *Cell Metab*. 2017;26(2):343-352.e2.
87. ClinicalTrials.gov. Research study of a new medicine (NNC9204-1706) in people with overweight or obesity (NCT03661879). <https://clinicaltrials.gov/ct2/show/NCT03661879>. Accessed March 25, 2022.
88. Tillner J, Posch MG, Wagner F, et al. A novel dual glucagon-like peptide and glucagon receptor agonist SAR425899: results of randomised, placebo-controlled first-in-human and first-in-patient trials. *Diabetes Obes Metab*. 2019;21(1):120-128.
89. ClinicalTrials.gov. A study to test whether different doses of BI 456906 are effective in treating adults with type 2 diabetes (NCT04153929). <https://clinicaltrials.gov/ct2/show/NCT04153929>. Accessed November 30, 2021.
90. Kim JK, Kim JY, Lee SM, et al. Potential of a novel long-acting glucagon analog, HM15136, for the treatment of obesity. *Diabetes*. 2019;68(Suppl 1):991.
91. ClinicalTrials.gov. A first-in-human study looking at the safety of ZP8396 and how it works in the body of healthy trial participants (NCT05096598). <https://clinicaltrials.gov/ct2/show/NCT05096598>. Accessed November 30, 2021.
92. Enebo LB, Berthelsen KK, Kankam M, et al. Safety, tolerability, pharmacokinetics, and pharmacodynamics of concomitant administration of multiple doses of cagrilintide with semaglutide 2.4 mg for weight management: a randomised, controlled, phase 1b trial. *Lancet*. 2021;397(10286):1736-1748.

93. ClinicalTrials.gov. A research study to investigate how well NNC0165-1875 in combination with semaglutide works in people with obesity (NCT04969939). <https://clinicaltrials.gov/ct2/show/NCT04969939>. Accessed March 25, 2022.
94. ClinicalTrials.gov. Long-term extension trial of setmelanotide (NCT03651765). <https://clinicaltrials.gov/ct2/show/NCT03651765>. Accessed November 30, 2021.
95. ClinicalTrials.gov. Setmelanotide (RM-493), melanocortin-4 receptor (MC4R) agonist, in Bardet-Biedl Syndrome (BBS) and Alström Syndrome (AS) patients with moderate to severe obesity (NCT03746522). <https://clinicaltrials.gov/ct2/show/NCT03746522>. Accessed November 30, 2021.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Bergmann NC, Davies MJ, Lingvay I, Knop FK. Semaglutide for the treatment of overweight and obesity: A review. *Diabetes Obes Metab*. 2023;25(1):18-35. doi:[10.1111/dom.14863](https://doi.org/10.1111/dom.14863)