

Pseudorandomness (III) & One-Way Functions

601.642/442: Modern Cryptography

Fall 2020

Recap: One-bit stretch PRG \implies Poly-bit stretch PRG

- A PRG with **one-bit stretch** can be used to construct a PRG with arbitrary **polynomial-bit stretch**.

Recap: One-bit stretch PRG \implies Poly-bit stretch PRG

- A PRG with **one-bit stretch** can be used to construct a PRG with arbitrary **polynomial-bit stretch**.

Construction of $G_{poly} : \{0, 1\}^n \rightarrow \{0, 1\}^{\ell(n)}$

Let $G : \{0, 1\}^n \rightarrow \{0, 1\}^{n+1}$ be a one-bit stretch PRG.

$$\begin{aligned} s &= x_0 \\ G(x_0) &= x_1 \| b_1 \\ &\vdots \\ G(x_{\ell(n)-1}) &= x_{\ell(n)} \| b_{\ell(n)} \end{aligned}$$

$$G_{poly}(s) := b_1 \dots b_{\ell(n)}$$

Recap: Pseudorandomness of G_{poly}

- In order to show $\left\{ G_{poly}(s); s \xleftarrow{\$} \{0,1\}^n \right\} \approx_c \left\{ r \xleftarrow{\$} \{0,1\}^{\ell(n)} \right\}$ we considered the following hybrid experiments:

Recap: Pseudorandomness of G_{poly}

- In order to show $\left\{ G_{poly}(s); s \xleftarrow{\$} \{0,1\}^n \right\} \approx_c \left\{ r \xleftarrow{\$} \{0,1\}^{\ell(n)} \right\}$ we considered the following hybrid experiments:

<u>Experiment \mathcal{H}_1</u>	<u>Experiment \mathcal{H}_2</u>	<u>Experiment $\mathcal{H}_{\ell(n)}$</u>
$s = x_0$	$s = x_0$	$s = X_0$
$G(x_0) = x_1 b_1$	$s_1 u_1 = x_1 u_1$	$s_1 u_1 = x_1 u_1$
$G(x_1) = x_2 b_2$	$G(x_1) = x_2 b_2$	$s_2 u_2 = x_2 u_2$
\dots	\dots	\dots
\dots	\dots	\dots
$G(x_{\ell(n)-1}) = x_{\ell(n)} b_{\ell(n)}$	$G(X_{\ell(n)-1}) = x_{\ell(n)} b_{\ell(n)}$	$s_{\ell(n)} u_{\ell(n)} = x_{\ell(n)} u_{\ell(n)}$
Output $G(s) := b_1 b_2 \dots b_{\ell(n)}$	Output $G(s) := u_1 b_2 \dots b_{\ell(n)}$	Output $G(s) := u_1 u_2 \dots u_{\ell(n)}$

Recap: Pseudorandomness of G_{poly}

- In order to show $\left\{ G_{poly}(s); s \xleftarrow{\$} \{0,1\}^n \right\} \approx_c \left\{ r \xleftarrow{\$} \{0,1\}^{\ell(n)} \right\}$ we considered the following hybrid experiments:

<u>Experiment \mathcal{H}_1</u>	<u>Experiment \mathcal{H}_2</u>	<u>Experiment $\mathcal{H}_{\ell(n)}$</u>
$s = x_0$	$s = x_0$	$s = X_0$
$G(x_0) = x_1 b_1$	$s_1 u_1 = x_1 u_1$	$s_1 u_1 = x_1 u_1$
$G(x_1) = x_2 b_2$	$G(x_1) = x_2 b_2$	$s_2 u_2 = x_2 u_2$
\dots	\dots	\dots
\dots	\dots	\dots
$G(x_{\ell(n)-1}) = x_{\ell(n)} b_{\ell(n)}$	$G(X_{\ell(n)-1}) = x_{\ell(n)} b_{\ell(n)}$	$s_{\ell(n)} u_{\ell(n)} = x_{\ell(n)} u_{\ell(n)}$
Output $G(s) := b_1 b_2 \dots b_{\ell(n)}$	Output $G(s) := u_1 b_2 \dots b_{\ell(n)}$	Output $G(s) := u_1 u_2 \dots u_{\ell(n)}$

- And established that $\forall i \in [\ell(n) - 1], \mathcal{H}_i \approx_c \mathcal{H}_{i+1}$.

Contrapositive Point of View

- So far, we have proved statements in the forward direction.

“If G is a one-bit stretch PRG, then G_{poly} is a poly-bit stretch PRG.”

Contrapositive Point of View

- So far, we have proved statements in the forward direction.

“If G is a one-bit stretch PRG, then G_{poly} is a poly-bit stretch PRG.”

- Let's now think about the **contrapositive** of these statements.

*“If G_{poly} is a **not** poly-bit stretch PRG, then G is **not** a one-bit stretch PRG.”*

Contrapositive Point of View

- So far, we have proved statements in the forward direction.
“If G is a one-bit stretch PRG, then G_{poly} is a poly-bit stretch PRG.”
- Let’s now think about the **contrapositive** of these statements.
*“If G_{poly} is a **not** poly-bit stretch PRG, then G is **not** a one-bit stretch PRG.”*
- If G_{poly} is not a PRG, then there exists a n.u. PPT adversary \mathcal{A} who can distinguish between its output on a random input and a uniformly sampled string with some **non-negligible** advantage μ .

Contrapositive Point of View

- So far, we have proved statements in the forward direction.
“If G is a one-bit stretch PRG, then G_{poly} is a poly-bit stretch PRG.”
- Let’s now think about the **contrapositive** of these statements.
*“If G_{poly} is a **not** poly-bit stretch PRG, then G is **not** a one-bit stretch PRG.”*
- If G_{poly} is not a PRG, then there exists a n.u. PPT adversary \mathcal{A} who can distinguish between its output on a random input and a uniformly sampled string with some **non-negligible** advantage μ .

Contrapositive Point of View

- So far, we have proved statements in the forward direction.

“If G is a one-bit stretch PRG, then G_{poly} is a poly-bit stretch PRG.”

- Let’s now think about the **contrapositive** of these statements.

*“If G_{poly} is a **not** poly-bit stretch PRG, then G is **not** a one-bit stretch PRG.”*

- If G_{poly} is not a PRG, then there exists a n.u. PPT adversary \mathcal{A} who can distinguish between its output on a random input and a uniformly sampled string with some **non-negligible** advantage μ .

Can we somehow use \mathcal{A} to also break security of G ?

Contrapositive Point of View

- We just proved security of G_{poly} using a sequence of hybrids.

Contrapositive Point of View

- We just proved security of G_{poly} using a sequence of hybrids.
- If we assume that an adversary \mathcal{A} exists, who has non-negligible advantage in breaking the security of G_{poly} , then at least one of the steps of our previous proof must break down.

Contrapositive Point of View

- We just proved security of G_{poly} using a sequence of hybrids.
- If we assume that an adversary \mathcal{A} exists, who has non-negligible advantage in breaking the security of G_{poly} , then at least one of the steps of our previous proof must break down.
- By hybrid lemma, \mathcal{A} can distinguish between at least 1 pair of consecutive hybrids (say \mathcal{H}_i and \mathcal{H}_{i+1}) with at least $\mu/\ell(n)$ advantage.

Contrapositive Point of View

- We just proved security of G_{poly} using a sequence of hybrids.
- If we assume that an adversary \mathcal{A} exists, who has non-negligible advantage in breaking the security of G_{poly} , then at least one of the steps of our previous proof must break down.
- By hybrid lemma, \mathcal{A} can distinguish between at least 1 pair of consecutive hybrids (say \mathcal{H}_i and \mathcal{H}_{i+1}) with at least $\mu/\ell(n)$ advantage.

Contrapositive Point of View

- We just proved security of G_{poly} using a sequence of hybrids.
- If we assume that an adversary \mathcal{A} exists, who has non-negligible advantage in breaking the security of G_{poly} , then at least one of the steps of our previous proof must break down.
- By hybrid lemma, \mathcal{A} can distinguish between at least 1 pair of consecutive hybrids (say \mathcal{H}_i and \mathcal{H}_{i+1}) with at least $\mu/\ell(n)$ advantage.

Lemma (Alternate way to state Hybrid Lemma)

Let X^1, \dots, X^m be distribution ensembles for $m = \text{poly}(n)$. Suppose there exists a distinguisher/adversary \mathcal{A} that distinguishes between X^1 and X^m with probability μ . Then $\exists i \in [m-1]$, such that \mathcal{A} distinguishes between X^i and X^{i+1} with advantage at least μ/m .

Contrapositive Point of View

- We just proved security of G_{poly} using a sequence of hybrids.
- If we assume that an adversary \mathcal{A} exists, who has non-negligible advantage in breaking the security of G_{poly} , then at least one of the steps of our previous proof must break down.
- By hybrid lemma, \mathcal{A} can distinguish between at least 1 pair of consecutive hybrids (say \mathcal{H}_i and \mathcal{H}_{i+1}) with at least $\mu/\ell(n)$ advantage.
- In our previous proof, we relied on the security of G to argue indistinguishability of each pair of consecutive hybrids.

Contrapositive Point of View

- We just proved security of G_{poly} using a sequence of hybrids.
- If we assume that an adversary \mathcal{A} exists, who has non-negligible advantage in breaking the security of G_{poly} , then at least one of the steps of our previous proof must break down.
- By hybrid lemma, \mathcal{A} can distinguish between at least 1 pair of consecutive hybrids (say \mathcal{H}_i and \mathcal{H}_{i+1}) with at least $\mu/\ell(n)$ advantage.
- In our previous proof, we relied on the security of G to argue indistinguishability of each pair of consecutive hybrids.
- We will now use \mathcal{A} that has non-negligible advantage in distinguishing between \mathcal{H}_i and \mathcal{H}_{i+1} , to construct another adversary \mathcal{B} to break security of G .

Contrapositive Point of View

- We just proved security of G_{poly} using a sequence of hybrids.
- If we assume that an adversary \mathcal{A} exists, who has non-negligible advantage in breaking the security of G_{poly} , then at least one of the steps of our previous proof must break down.
- By hybrid lemma, \mathcal{A} can distinguish between at least 1 pair of consecutive hybrids (say \mathcal{H}_i and \mathcal{H}_{i+1}) with at least $\mu/\ell(n)$ advantage.
- In our previous proof, we relied on the security of G to argue indistinguishability of each pair of consecutive hybrids.
- We will now use \mathcal{A} that has non-negligible advantage in distinguishing between \mathcal{H}_i and \mathcal{H}_{i+1} , to construct another adversary \mathcal{B} to break security of G .
- However, since G is a secure PRG, no such n.u. PPT \mathcal{A} should exist. This will give us a contradiction and imply that our assumption was incorrect. G_{poly} is in fact secure.

Proof via Reduction

- How do we construct \mathcal{B} ?

Proof via Reduction


- How do we construct \mathcal{B} ?
- We consider the game-based definition of PRG.

Proof via Reduction

- How do we construct \mathcal{B} ?
- We consider the game-based definition of PRG.

Proof via Reduction

- How do we construct \mathcal{B} ?
- We consider the game-based definition of PRG.




$$\left| \Pr[b' = 1 | b = 1] - \Pr[b' = 1 | b = 0] \right| \leq \nu(n)$$

Proof by Reduction

- This proof technique is also called **proof by reduction**.

Proof by Reduction

- This proof technique is also called **proof by reduction**.



Proof by Reduction

- If y is pseudorandom, i.e., sampled as $y = G(s)$, then the input to \mathcal{A} is distributed identically to the output of \mathcal{H}_i .

Proof by Reduction

- If y is pseudorandom, i.e., sampled as $y = G(s)$, then the input to \mathcal{A} is distributed identically to the output of \mathcal{H}_i .
- Otherwise, i.e., y is (truly) random, and therefore the input to \mathcal{A} is distributed identically to the output of \mathcal{H}_{i+1} .

Proof by Reduction

- If y is pseudorandom, i.e., sampled as $y = G(s)$, then the input to \mathcal{A} is distributed identically to the output of \mathcal{H}_i .
- Otherwise, i.e., y is (truly) random, and therefore the input to \mathcal{A} is distributed identically to the output of \mathcal{H}_{i+1} .
- Hence, \mathcal{B} has the **same advantage** in distinguishing between the output of G and a pseudorandom string that \mathcal{A} has in distinguishing between \mathcal{H}_i and \mathcal{H}_{i+1} .

Proof by Reduction

- If y is pseudorandom, i.e., sampled as $y = G(s)$, then the input to \mathcal{A} is distributed identically to the output of \mathcal{H}_i .
- Otherwise, i.e., y is (truly) random, and therefore the input to \mathcal{A} is distributed identically to the output of \mathcal{H}_{i+1} .
- Hence, \mathcal{B} has the **same advantage** in distinguishing between the output of G and a pseudorandom string that \mathcal{A} has in distinguishing between \mathcal{H}_i and \mathcal{H}_{i+1} .
- Moreover, since \mathcal{A} is n.u. PPT, so is \mathcal{B} . This is a contradiction!

Proof by Reduction

- If y is pseudorandom, i.e., sampled as $y = G(s)$, then the input to \mathcal{A} is distributed identically to the output of \mathcal{H}_i .
- Otherwise, i.e., y is (truly) random, and therefore the input to \mathcal{A} is distributed identically to the output of \mathcal{H}_{i+1} .
- Hence, \mathcal{B} has the **same advantage** in distinguishing between the output of G and a pseudorandom string that \mathcal{A} has in distinguishing between \mathcal{H}_i and \mathcal{H}_{i+1} .
- Moreover, since \mathcal{A} is n.u. PPT, so is \mathcal{B} . This is a contradiction!
- Hence, $G_{\text{poly}(n)}$ is a PRG.

Proof by Reduction: Key Points

- These are four important things that you must work through for a valid reduction:

Proof by Reduction: Key Points

- These are four important things that you must work through for a valid reduction:
 - ① **Input Mapping:** How to map the input that “outer adversary” \mathcal{B} receives from the challenger to an input to the “internal adversary” \mathcal{A} ?

Proof by Reduction: Key Points

- These are four important things that you must work through for a valid reduction:
 - 1 **Input Mapping:** How to map the input that “outer adversary” \mathcal{B} receives from the challenger to an input to the “internal adversary” \mathcal{A} ?
 - 2 **Input Distribution:** Does the input mapping provide the right distribution of inputs that \mathcal{A} expects?

Proof by Reduction: Key Points


- These are four important things that you must work through for a valid reduction:
 - ① **Input Mapping:** How to map the input that “outer adversary” \mathcal{B} receives from the challenger to an input to the “internal adversary” \mathcal{A} ?
 - ② **Input Distribution:** Does the input mapping provide the right distribution of inputs that \mathcal{A} expects?
 - ③ **Output Mapping:** How do we map the output that \mathcal{A} provides to an output for \mathcal{B} ?

Proof by Reduction: Key Points

- These are four important things that you must work through for a valid reduction:
 - ① **Input Mapping:** How to map the input that “outer adversary” \mathcal{B} receives from the challenger to an input to the “internal adversary” \mathcal{A} ?
 - ② **Input Distribution:** Does the input mapping provide the right distribution of inputs that \mathcal{A} expects?
 - ③ **Output Mapping:** How do we map the output that \mathcal{A} provides to an output for \mathcal{B} ?
 - ④ **Probability:** When we assume existence of \mathcal{A} , we also assume that \mathcal{A} wins with non-negligible advantage. What is the probability/advantage that \mathcal{B} wins, given the mappings above?


One-Way Functions

One Way Functions




- A function is one-way if it “easy to compute,” but “hard to invert”

One Way Functions




- A function is one-way if it “easy to compute,” but “hard to invert”
- Necessary for the existence of most cryptographic primitives (e.g., multi-messgae encryption, digital signatures)

One Way Functions




- A function is one-way if it “easy to compute,” but “hard to invert”
- Necessary for the existence of most cryptographic primitives (e.g., multi-message encryption, digital signatures)
- Also sufficient for some cryptographic primitives (e.g., pseudorandom generators, secret-key encryption, digital signatures).

One Way Functions



- A function is one-way if it “easy to compute,” but “hard to invert”
- Necessary for the existence of most cryptographic primitives (e.g., multi-message encryption, digital signatures)
- Also sufficient for some cryptographic primitives (e.g., pseudorandom generators, secret-key encryption, digital signatures).

One Way Functions



- A function is one-way if it “easy to compute,” but “hard to invert”
- Necessary for the existence of most cryptographic primitives (e.g., multi-message encryption, digital signatures)
- Also sufficient for some cryptographic primitives (e.g., pseudorandom generators, secret-key encryption, digital signatures).

How to define one-way functions?

Defining One Way Functions: Attempt 1

Attempt 1: A function $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ is a one-way function (OWF) if it satisfies the following two conditions:

- **Easy to compute:** there is a polynomial-time algorithm \mathcal{C} s.t.
 $\forall x \in \{0, 1\}^*,$

$$\Pr [\mathcal{C}(x) = f(x)] = 1.$$

Defining One Way Functions: Attempt 1

Attempt 1: A function $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ is a one-way function (OWF) if it satisfies the following two conditions:

- **Easy to compute:** there is a polynomial-time algorithm \mathcal{C} s.t.
 $\forall x \in \{0, 1\}^*,$

$$\Pr [\mathcal{C}(x) = f(x)] = 1.$$

- **Hard to invert:** for every non-uniform PPT adversary \mathcal{A} , for any input length $n \in \mathbb{N}$

Probability of Inversion is Negligible

Defining One Way Functions: Attempt 1

Attempt 1: A function $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ is a one-way function (OWF) if it satisfies the following two conditions:

- **Easy to compute:** there is a polynomial-time algorithm \mathcal{C} s.t.
 $\forall x \in \{0, 1\}^*,$

$$\Pr [\mathcal{C}(x) = f(x)] = 1.$$

- **Hard to invert:** for every non-uniform PPT adversary \mathcal{A} , for any input length $n \in \mathbb{N}$

$$\Pr [\mathcal{A} \text{ inverts } f(x) \text{ for random } x] \leq \textit{negligible}.$$

This is called **average-case** hardness.

One Way Functions: Definition

Definition (One Way Function)

A function $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ is a one-way function (OWF) if it satisfies the following two conditions:

- **Easy to compute:** there is a polynomial-time algorithm \mathcal{C} s.t.
 $\forall x \in \{0, 1\}^*,$

$$\Pr [\mathcal{C}(x) = f(x)] = 1.$$

- **Hard to invert:** there exists a negligible function $\nu : \mathbb{N} \rightarrow \mathbb{R}$ s.t.
for every non-uniform PPT adversary \mathcal{A} and $\forall n \in \mathbb{N}$:

$$\Pr \left[x \stackrel{\$}{\leftarrow} \{0, 1\}^n, x' \leftarrow \mathcal{A}(1^n, f(x)) : f(x') = f(x) \right] \leq \nu(n).$$


- The above definition is also called **strong** one-way functions.

One Way Functions: Game Based Definition

It is also instructive to think of that definition in this game-based form.


One Way Functions: Game Based Definition

It is also instructive to think of that definition in this game-based form.



One Way Functions: Game Based Definition

It is also instructive to think of that definition in this game-based form.



We say that $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ is a one-way function if there exists a negligible function $\nu : \mathbb{N} \rightarrow \mathbb{R}$ s.t. for every n.u. PPT adversary \mathcal{A} and $\forall n \in \mathbb{N}$:

$$\Pr[\mathcal{A} \text{ wins}] \leq \nu(n).$$

Injective OWFs and One Way Permutations (OWP)

- **Injective or 1-1 OWFs:** each image has a unique pre-image:

$$f(x_1) = f(x_2) \implies x_1 = x_2$$

- **One Way Permutations (OWP):** 1-1 OWF with the additional conditional that “each image has a pre-image”

(Equivalently: domain and range are of same size.)

Existence of OWFs

Do OWFs exist?

Do OWFs exist?

- NOT Unconditionally — proving that f is one-way requires proving (at least) $\mathbf{P} \neq \mathbf{NP}$.

Do OWFs exist?

- NOT Unconditionally — proving that f is one-way requires proving (at least) $\mathbf{P} \neq \mathbf{NP}$.
- However, we can construct them ASSUMING that certain problems are hard.

Do OWFs exist?

- NOT Unconditionally — proving that f is one-way requires proving (at least) $\mathbf{P} \neq \mathbf{NP}$.
- However, we can construct them ASSUMING that certain problems are hard.
- Such constructions are sometimes called “candidates” because they are based on an assumption or a conjecture.

Factoring Problem

- Consider the **multiplication** function $f_{\times} : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$:

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

Factoring Problem

- Consider the **multiplication** function $f_{\times} : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$:

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- The first condition helps exclude the trivial factor 1.

Factoring Problem

- Consider the **multiplication** function $f_{\times} : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$:

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- The first condition helps exclude the trivial factor 1.
- Is f_{\times} a OWF?

Factoring Problem

- Consider the **multiplication** function $f_{\times} : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$:

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- The first condition helps exclude the trivial factor 1.
- Is f_{\times} a OWF?
- Clearly not!** With prob. $1/2$, a random number (of any fixed size) is even. I.e., xy is even w/ prob. $\frac{3}{4}$ for random (x, y) .

Factoring Problem

- Consider the **multiplication** function $f_{\times} : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$:

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- The first condition helps exclude the trivial factor 1.
- Is f_{\times} a OWF?
- Clearly not!** With prob. $1/2$, a random number (of any fixed size) is even. I.e., xy is even w/ prob. $\frac{3}{4}$ for random (x, y) .
- Inversion: given number z , output $(2, z/2)$ if z is even and $(0, 0)$ otherwise! (succeeds 75% time)

Factoring Problem (continued)

- Eliminate such trivial small factors.

Factoring Problem (continued)

- Eliminate such trivial small factors.
- Let Π_n be the set of all **prime** numbers $< 2^n$.

Factoring Problem (continued)

- Eliminate such trivial small factors.
- Let Π_n be the set of all **prime** numbers $< 2^n$.
- Choose numbers p and q randomly from Π_n and multiply.

Factoring Problem (continued)

- Eliminate such trivial small factors.
- Let Π_n be the set of all **prime** numbers $< 2^n$.
- Choose numbers p and q randomly from Π_n and multiply.
- This is unlikely to have small trivial factors.
[Factoring Assumption] For every (non-uniform PPT) adversary \mathcal{A} , there exists a negligible function ν such that

$$\Pr \left[p \xleftarrow{\$} \Pi_n; q \xleftarrow{\$} \Pi_n; N = pq : \mathcal{A}(N) \in \{p, q\} \right] \leq \nu(n).$$

Factoring Problem (continued)

- Factoring assumption is a well established conjecture.

Factoring Problem (continued)

- Factoring assumption is a well established conjecture.
- Studied for a long time, with no known polynomial-time attack.

Factoring Problem (continued)

- Factoring assumption is a well established conjecture.
- Studied for a long time, with no known polynomial-time attack.
- Best known algorithms for breaking Factoring Assumption:

$$2^{O(\sqrt{n \log n})} \quad (\text{provable})$$

$$2^{O(\sqrt[3]{n \log^2 n})} \quad (\text{heuristic})$$

Factoring Problem (continued)

- Factoring assumption is a well established conjecture.
- Studied for a long time, with no known polynomial-time attack.
- Best known algorithms for breaking Factoring Assumption:

$$2^{O(\sqrt{n \log n})} \quad (\text{provable})$$

$$2^{O(\sqrt[3]{n \log^2 n})} \quad (\text{heuristic})$$

- Can we construct OWEs from the Factoring Assumption?

Multiplication Function

- Going back to the multiplication function $f_{\times} : \mathbb{N}^2 \rightarrow \mathbb{N}$.

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

Multiplication Function

- Going back to the multiplication function $f_{\times} : \mathbb{N}^2 \rightarrow \mathbb{N}$.

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- **Observation 1:** If randomly chosen x and y happen to be primes, no PPT \mathcal{A} can invert (except with negligible probability). Call it the **GOOD** case.

Multiplication Function

- Going back to the multiplication function $f_{\times} : \mathbb{N}^2 \rightarrow \mathbb{N}$.

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- **Observation 1:** If randomly chosen x and y happen to be primes, no PPT \mathcal{A} can invert (except with negligible probability). Call it the **GOOD** case.
- If GOOD case occurs with probability $> \varepsilon$,
 \Rightarrow every PPT \mathcal{A} must fail to invert f_{\times} with probability at least ε .

Multiplication Function

- Going back to the multiplication function $f_{\times} : \mathbb{N}^2 \rightarrow \mathbb{N}$.

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- **Observation 1:** If randomly chosen x and y happen to be primes, no PPT \mathcal{A} can invert (except with negligible probability). Call it the **GOOD** case.
- If GOOD case occurs with probability $> \varepsilon$,
 \Rightarrow every PPT \mathcal{A} must fail to invert f_{\times} with probability at least ε .
- Now suppose that ε is a **noticeable function** (say e.g. an inverse polynomial, i.e., $\frac{1}{p(\cdot)}$)
 \Rightarrow every \mathcal{A} must fail to invert f_{\times} with **noticeable** probability.

Multiplication Function

- Going back to the multiplication function $f_{\times} : \mathbb{N}^2 \rightarrow \mathbb{N}$.

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- **Observation 1:** If randomly chosen x and y happen to be primes, no PPT \mathcal{A} can invert (except with negligible probability). Call it the **GOOD** case.
- If GOOD case occurs with probability $> \varepsilon$,
 \Rightarrow every PPT \mathcal{A} must fail to invert f_{\times} with probability at least ε .
- Now suppose that ε is a **noticeable function** (say e.g. an inverse polynomial, i.e., $\frac{1}{p(\cdot)}$)
 \Rightarrow every \mathcal{A} must fail to invert f_{\times} with **noticeable** probability.
- This is already useful!

Multiplication Function

- Going back to the multiplication function $f_{\times} : \mathbb{N}^2 \rightarrow \mathbb{N}$.

$$f_{\times}(x, y) = \begin{cases} \perp & \text{if } x = 1 \vee y = 1 \\ x \cdot y & \text{otherwise} \end{cases}$$

- **Observation 1:** If randomly chosen x and y happen to be primes, no PPT \mathcal{A} can invert (except with negligible probability). Call it the **GOOD** case.
- If GOOD case occurs with probability $> \varepsilon$,
 \Rightarrow every PPT \mathcal{A} must fail to invert f_{\times} with probability at least ε .
- Now suppose that ε is a **noticeable function** (say e.g. an inverse polynomial, i.e., $\frac{1}{p(\cdot)}$)
 \Rightarrow every \mathcal{A} must fail to invert f_{\times} with **noticeable** probability.
- This is already useful!
- Usually called a **weak OWF**.

Noticeable Functions

Let us start by formally defining noticeable functions. These are functions that are **at most polynomially small**.

Noticeable Functions

Let us start by formally defining noticeable functions. These are functions that are **at most polynomially small**.

Definition (Noticeable Function)

A function $\nu(n)$ is noticeable if $\exists c, n_0$ such that $\forall n > n_0, \nu(n) \geq \frac{1}{n^c}$.

Noticeable Functions

Let us start by formally defining noticeable functions. These are functions that are **at most polynomially small**.

Definition (Noticeable Function)

A function $\nu(n)$ is noticeable if $\exists c, n_0$ such that $\forall n > n_0, \nu(n) \geq \frac{1}{n^c}$.

Note that a non-negligible function is not necessarily a noticeable function. Example:

$$f(n) = \begin{cases} 1 & \text{if } n \text{ is even} \\ 2^{-n} & \text{if } n \text{ is odd} \end{cases}.$$

This function is non-negligible, but not noticeable. **Why?**