

MA 6.101

Probability and Statistics

**Tejas Bodas**

Assistant Professor, IIT Hyderabad

# Logistics

# Logistics

- ▶ Tutorial has been uploaded.

# Logistics

- ▶ Tutorial has been uploaded.
- ▶ Practice a lot of problems from Bertsekas.

# Logistics

- ▶ Tutorial has been uploaded.
- ▶ Practice a lot of problems from Bertsekas.
- ▶ Quiz 1 has been corrected.

# Logistics

- ▶ Tutorial has been uploaded.
- ▶ Practice a lot of problems from Bertsekas.
- ▶ Quiz 1 has been corrected.
- ▶ Paper distribution tomorrow at 3:30-5pm ?

# Logistics

- ▶ Tutorial has been uploaded.
- ▶ Practice a lot of problems from Bertsekas.
- ▶ Quiz 1 has been corrected.
- ▶ Paper distribution tomorrow at 3:30-5pm ?
- ▶ Venue will be emailed later.

# Function of random variables



# Function of random variables

- ▶ Consider  $Y = |X|$  where  $X$  is the outcome of an experiment where an integer is chosen uniformly from  $-4$  to  $4$ .

# Function of random variables

- ▶ Consider  $Y = |X|$  where  $X$  is the outcome of an experiment where an integer is chosen uniformly from  $-4$  to  $4$ .
- ▶  $p_X(x) = \frac{1}{9}$  for  $x \in \{-4, -3, \dots, 3, 4\}$ .

# Function of random variables

- ▶ Consider  $Y = |X|$  where  $X$  is the outcome of an experiment where an integer is chosen uniformly from  $-4$  to  $4$ .
- ▶  $p_X(x) = \frac{1}{9}$  for  $x \in \{-4, -3, \dots, 3, 4\}$ .
- ▶ What is the range  $\Omega'$  for  $Y$ ?

# Function of random variables

- ▶ Consider  $Y = |X|$  where  $X$  is the outcome of an experiment where an integer is chosen uniformly from  $-4$  to  $4$ .
- ▶  $p_X(x) = \frac{1}{9}$  for  $x \in \{-4, -3, \dots, 3, 4\}$ .
- ▶ What is the range  $\Omega'$  for  $Y$ ?  $\Omega' = \{0, \dots, 4\}$ .

# Function of random variables

- ▶ Consider  $Y = |X|$  where  $X$  is the outcome of an experiment where an integer is chosen uniformly from  $-4$  to  $4$ .
- ▶  $p_X(x) = \frac{1}{9}$  for  $x \in \{-4, -3, \dots, 3, 4\}$ .
- ▶ What is the range  $\Omega'$  for  $Y$ ?  $\Omega' = \{0, \dots, 4\}$ .
- ▶ What is  $p_Y(2)$ ?


# Function of random variables

- ▶ Consider  $Y = |X|$  where  $X$  is the outcome of an experiment where an integer is chosen uniformly from  $-4$  to  $4$ .
- ▶  $p_X(x) = \frac{1}{9}$  for  $x \in \{-4, -3, \dots, 3, 4\}$ .
- ▶ What is the range  $\Omega'$  for  $Y$ ?  $\Omega' = \{0, \dots, 4\}$ .
- ▶ What is  $p_Y(2)$ ?
- ▶  $p_Y(2) = \sum_{\{x: |x|=2\}} p_X(x) = p_X(-2) + p_X(2) = \frac{2}{9}$ .

# Function of random variables

# Function of random variables

►  $p_Y(2) = \sum_{\{x:|x|=2\}} p_X(x) = p_X(-2) + p_X(2) = \frac{2}{9}.$



Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  
$$p_Y(y) = \sum_{\{x:g(x)=y\}} p_X(x).$$



# Function of random variables

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  
$$p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x).$$

Proof:

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  
$$p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x).$$

Proof:

► Then what is  $p_Y(y)$ ?

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  
$$p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x).$$

Proof:

- ▶ Then what is  $p_Y(y)$ ?
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : Y(\omega) = y\}.$

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  
$$p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x).$$

Proof:

- ▶ Then what is  $p_Y(y)$ ?
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : Y(\omega) = y\}.$
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : g(X(\omega)) = y\}.$

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  $p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x)$ .

Proof:

- ▶ Then what is  $p_Y(y)$ ?
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : Y(\omega) = y\}$ .
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : g(X(\omega)) = y\}$ .
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : X(\omega) = g^{-1}(y)\}$ .

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  $p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x)$ .

Proof:

- ▶ Then what is  $p_Y(y)$ ?
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : Y(\omega) = y\}$ .
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : g(X(\omega)) = y\}$ .
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : X(\omega) = g^{-1}(y)\}$ .
- ▶ Is there a problem with the text in red?

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  $p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x)$ .

Proof:

- ▶ Then what is  $p_Y(y)$ ?
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : Y(\omega) = y\}$ .
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : g(X(\omega)) = y\}$ .
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : X(\omega) = g^{-1}(y)\}$ .
- ▶ Is there a problem with the text in red?
- ▶ Is  $g^{-1}(y)$  a value or a set?



# Function of random variables

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  
$$p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x).$$

Proof Continued:

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  
$$p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x).$$

Proof Continued:

- ▶  $g^{-1}(y)$  is a value if  $g(\cdot)$  is one to one.

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  
$$p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x).$$

Proof Continued:

- ▶  $g^{-1}(y)$  is a value if  $g(\cdot)$  is one to one.
- ▶ If  $g(\cdot)$  is many to one, then  $g^{-1}(y) := \{x : g(x) = y\}$ .

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  $p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x)$ .

Proof Continued:

- ▶  $g^{-1}(y)$  is a value if  $g(\cdot)$  is one to one.
- ▶ If  $g(\cdot)$  is many to one, then  $g^{-1}(y) := \{x : g(x) = y\}$ .
- ▶ In that case,  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : X(\omega) \in g^{-1}(y)\}$ .

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  $p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x)$ .

Proof Continued:

- ▶  $g^{-1}(y)$  is a value if  $g(\cdot)$  is one to one.
- ▶ If  $g(\cdot)$  is many to one, then  $g^{-1}(y) := \{x : g(x) = y\}$ .
- ▶ In that case,  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : X(\omega) \in g^{-1}(y)\}$ .
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : X(\omega) \in \{x : g(x) = y\}\}$ .

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  $p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x)$ .

Proof Continued:

- ▶  $g^{-1}(y)$  is a value if  $g(\cdot)$  is one to one.
- ▶ If  $g(\cdot)$  is many to one, then  $g^{-1}(y) := \{x : g(x) = y\}$ .
- ▶ In that case,  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : X(\omega) \in g^{-1}(y)\}$ .
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : X(\omega) \in \{x : g(x) = y\}\}$ .
- ▶ Now  $\mathbb{P}\{\omega \in \Omega : X(\omega) \in B\}$

# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  $p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x)$ .

Proof Continued:

- ▶  $g^{-1}(y)$  is a value if  $g(\cdot)$  is one to one.
- ▶ If  $g(\cdot)$  is many to one, then  $g^{-1}(y) := \{x : g(x) = y\}$ .
- ▶ In that case,  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : X(\omega) \in g^{-1}(y)\}$ .
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : X(\omega) \in \{x : g(x) = y\}\}$ .
- ▶ Now  $\mathbb{P}\{\omega \in \Omega : X(\omega) \in B\} = \sum_{\{x \in B\}} p_X(x)$  for  $B \in \mathcal{F}'$ .



# Function of random variables

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then  $p_Y(y) = \sum_{\{x: g(x)=y\}} p_X(x)$ .

Proof Continued:

- ▶  $g^{-1}(y)$  is a value if  $g(\cdot)$  is one to one.
- ▶ If  $g(\cdot)$  is many to one, then  $g^{-1}(y) := \{x : g(x) = y\}$ .
- ▶ In that case,  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : X(\omega) \in g^{-1}(y)\}$ .
- ▶  $p_Y(y) = \mathbb{P}\{\omega \in \Omega : X(\omega) \in \{x : g(x) = y\}\}$ .
- ▶ Now  $\mathbb{P}\{\omega \in \Omega : X(\omega) \in B\} = \sum_{\{x \in B\}} p_X(x)$  for  $B \in \mathcal{F}'$ .
- ▶ Proof follows after setting  $B = \{x : g(x) = y\}$  □

$$E[g(X)]$$

$$E[g(X)]$$

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then,  
$$E[Y] = \sum_x g(x)p_X(x)$$

Proof

$$E[g(X)]$$

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then,  
 $E[Y] = \sum_x g(x)p_X(x)$

Proof

$$\begin{aligned} E[Y] &= \sum_y yp_Y(y) \\ &= \sum_y \sum_{\{x:g(x)=y\}} g(x)p_X(x) \end{aligned}$$

$$E[g(X)]$$

Suppose  $Y = g(X)$  and  $X$  is discrete with pmf  $p_X(\cdot)$ . Then,  
 $E[Y] = \sum_x g(x)p_X(x)$

Proof

$$\begin{aligned} E[Y] &= \sum_y yp_Y(y) \\ &= \sum_y \sum_{\{x:g(x)=y\}} g(x)p_X(x) \\ &= \sum_x g(x)p_X(x). \end{aligned}$$



# Function of continuous random variables

# Function of continuous random variables

- ▶ Consider  $Y = aX + b$  where  $X$  is a continuous random variable.

# Function of continuous random variables

- ▶ Consider  $Y = aX + b$  where  $X$  is a continuous random variable.
- ▶ What is  $F_Y(y)$  and  $f_Y(y)$ ?



# Function of continuous random variables

- ▶ Consider  $Y = aX + b$  where  $X$  is a continuous random variable.
- ▶ What is  $F_Y(y)$  and  $f_Y(y)$ ?
- ▶  $F_Y(y) = P(Y \leq y) = P(aX + b \leq y)$ .

# Function of continuous random variables

- ▶ Consider  $Y = aX + b$  where  $X$  is a continuous random variable.
- ▶ What is  $F_Y(y)$  and  $f_Y(y)$ ?
- ▶  $F_Y(y) = P(Y \leq y) = P(aX + b \leq y)$ .
- ▶  $F_Y(y) = F_X(\frac{y-b}{a})$  if  $a > 0$

# Function of continuous random variables

- ▶ Consider  $Y = aX + b$  where  $X$  is a continuous random variable.
- ▶ What is  $F_Y(y)$  and  $f_Y(y)$ ?
- ▶  $F_Y(y) = P(Y \leq y) = P(aX + b \leq y)$ .
- ▶  $F_Y(y) = F_X(\frac{y-b}{a})$  if  $a > 0$
- ▶  $f_Y(y) = \frac{dF_Y(y)}{dy} = \frac{1}{a} f_X(\frac{y-b}{a})$  when  $a > 0$

# Function of continuous random variables

- ▶ Consider  $Y = aX + b$  where  $X$  is a continuous random variable.
- ▶ What is  $F_Y(y)$  and  $f_Y(y)$ ?
- ▶  $F_Y(y) = P(Y \leq y) = P(aX + b \leq y)$ .
- ▶  $F_Y(y) = F_X(\frac{y-b}{a})$  if  $a > 0$
- ▶  $f_Y(y) = \frac{dF_Y(y)}{dy} = \frac{1}{a}f_X(\frac{y-b}{a})$  when  $a > 0$
- ▶  $F_Y(y) = 1 - F_X(\frac{y-b}{a})$  if  $a < 0$

# Function of continuous random variables

- ▶ Consider  $Y = aX + b$  where  $X$  is a continuous random variable.
- ▶ What is  $F_Y(y)$  and  $f_Y(y)$ ?
- ▶  $F_Y(y) = P(Y \leq y) = P(aX + b \leq y)$ .
- ▶  $F_Y(y) = F_X(\frac{y-b}{a})$  if  $a > 0$
- ▶  $f_Y(y) = \frac{dF_Y(y)}{dy} = \frac{1}{a} f_X(\frac{y-b}{a})$  when  $a > 0$
- ▶  $F_Y(y) = 1 - F_X(\frac{y-b}{a})$  if  $a < 0$
- ▶  $f_Y(y) = \frac{dF_Y(y)}{dy} = \frac{-1}{a} f_X(\frac{y-b}{a})$  when  $a < 0$

# Function of continuous random variables

- ▶ Consider  $Y = aX + b$  where  $X$  is a continuous random variable.
- ▶ What is  $F_Y(y)$  and  $f_Y(y)$ ?
- ▶  $F_Y(y) = P(Y \leq y) = P(aX + b \leq y)$ .
- ▶  $F_Y(y) = F_X(\frac{y-b}{a})$  if  $a > 0$
- ▶  $f_Y(y) = \frac{dF_Y(y)}{dy} = \frac{1}{a} f_X(\frac{y-b}{a})$  when  $a > 0$
- ▶  $F_Y(y) = 1 - F_X(\frac{y-b}{a})$  if  $a < 0$
- ▶  $f_Y(y) = \frac{dF_Y(y)}{dy} = \frac{-1}{a} f_X(\frac{y-b}{a})$  when  $a < 0$
- ▶ In general,  $f_Y(y) = \frac{1}{|a|} f_X(\frac{y-b}{a})$

# Function of continuous random variables

Consider  $Y = aX + b$  where  $X$  is a continuous random variable. Then  $f_Y(y) = \frac{1}{|a|} f_X\left(\frac{y-b}{a}\right)$ .

# Function of continuous random variables

Consider  $Y = aX + b$  where  $X$  is a continuous random variable. Then  $f_Y(y) = \frac{1}{|a|} f_X\left(\frac{y-b}{a}\right)$ .

- ▶ What if  $Y = g(X)$  where  $g(\cdot)$  is continuous, differentiable and monotone. Any guess?



# Function of continuous random variables

Consider  $Y = aX + b$  where  $X$  is a continuous random variable. Then  $f_Y(y) = \frac{1}{|a|} f_X\left(\frac{y-b}{a}\right)$ .

- ▶ What if  $Y = g(X)$  where  $g(\cdot)$  is continuous, differentiable and monotone. Any guess?
- ▶ Since  $g(\cdot)$  is monotone and continuous it is invertible. Let  $h(\cdot)$  denote the inverse function.

# Function of continuous random variables

Consider  $Y = aX + b$  where  $X$  is a continuous random variable. Then  $f_Y(y) = \frac{1}{|a|} f_X\left(\frac{y-b}{a}\right)$ .

- ▶ What if  $Y = g(X)$  where  $g(\cdot)$  is continuous, differentiable and monotone. Any guess?
- ▶ Since  $g(\cdot)$  is monotone and continuous it is invertible. Let  $h(\cdot)$  denote the inverse function. Then  $h(Y) = X$ .

Consider  $Y = g(X)$  where  $g$  is monotone, continuous, differentiable. Then  $f_Y(y) = \left|\frac{dh}{dy}(y)\right| f_X(h(y))$  where  $h$  is the inverse function of  $g$ .

# Function of continuous random variables

Consider  $Y = g(X)$  where  $g$  is monotone, continuous, differentiable. Then  $f_Y(y) = f_X(h(y)) \left| \frac{dh}{dy}(y) \right|$  where  $h$  is the inverse function of  $g$ .

Proof:

# Function of continuous random variables

Consider  $Y = g(X)$  where  $g$  is monotone, continuous, differentiable. Then  $f_Y(y) = f_X(h(y)) \left| \frac{dh}{dy}(y) \right|$  where  $h$  is the inverse function of  $g$ .

Proof:

- ▶ Since  $g(\cdot)$  is monotone and continuous it is invertible. Let  $h(\cdot)$  denote the inverse function.

# Function of continuous random variables

Consider  $Y = g(X)$  where  $g$  is monotone, continuous, differentiable. Then  $f_Y(y) = f_X(h(y)) \left| \frac{dh}{dy}(y) \right|$  where  $h$  is the inverse function of  $g$ .

Proof:

- ▶ Since  $g(\cdot)$  is monotone and continuous it is invertible. Let  $h(\cdot)$  denote the inverse function. Then  $X = h(Y)$ .

# Function of continuous random variables

Consider  $Y = g(X)$  where  $g$  is monotone, continuous, differentiable. Then  $f_Y(y) = f_X(h(y)) \left| \frac{dh}{dy}(y) \right|$  where  $h$  is the inverse function of  $g$ .

Proof:

- ▶ Since  $g(\cdot)$  is monotone and continuous it is invertible. Let  $h(\cdot)$  denote the inverse function. Then  $X = h(Y)$ .
- ▶  $F_Y(y) = P(g(X) \leq y)$ .

# Function of continuous random variables

Consider  $Y = g(X)$  where  $g$  is monotone, continuous, differentiable. Then  $f_Y(y) = f_X(h(y)) \left| \frac{dh}{dy}(y) \right|$  where  $h$  is the inverse function of  $g$ .

Proof:

- ▶ Since  $g(\cdot)$  is monotone and continuous it is invertible. Let  $h(\cdot)$  denote the inverse function. Then  $X = h(Y)$ .
- ▶  $F_Y(y) = P(g(X) \leq y)$ .
- ▶ Is  $P(g(X) \leq y) = P(X \leq h(y))$  always?

# Function of continuous random variables

Consider  $Y = g(X)$  where  $g$  is monotone, continuous, differentiable. Then  $f_Y(y) = f_X(h(y)) \left| \frac{dh}{dy}(y) \right|$  where  $h$  is the inverse function of  $g$ .

Proof:

- ▶ Since  $g(\cdot)$  is monotone and continuous it is invertible. Let  $h(\cdot)$  denote the inverse function. Then  $X = h(Y)$ .
- ▶  $F_Y(y) = P(g(X) \leq y)$ .
- ▶ Is  $P(g(X) \leq y) = P(X \leq h(y))$  always?
- ▶ Are the two events  $\{g(X) \leq y\}$  and  $X \leq h(y)$  same?



# Function of continuous random variables

Consider  $Y = g(X)$  where  $g$  is monotone, continuous, differentiable. Then  $f_Y(y) = f_X(h(y)) \left| \frac{dh}{dy}(y) \right|$  where  $h$  is the inverse function of  $g$ .



Proof:

- ▶ Since  $g(\cdot)$  is monotone and continuous it is invertible. Let  $h(\cdot)$  denote the inverse function. Then  $X = h(Y)$ .
- ▶  $F_Y(y) = P(g(X) \leq y)$ .
- ▶ Is  $P(g(X) \leq y) = P(X \leq h(y))$  always?
- ▶ Are the two events  $\{g(X) \leq y\}$  and  $X \leq h(y)$  same?
- ▶ If they are same, then the two probabilities are equal.

# Function of continuous random variables



# Function of continuous random variables

- Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?



# Function of continuous random variables

- Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?



- Same when  $g$  is increasing and compliments when  $g$  is decreasing.

# Function of continuous random variables

# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?

# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?
- ▶ Same when  $g$  is increasing and compliments when  $g$  is decreasing.

# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?
- ▶ Same when  $g$  is increasing and compliments when  $g$  is decreasing.
- ▶ CASE 1:  $g(x)$  is non-decreasing



# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?
- ▶ Same when  $g$  is increasing and compliments when  $g$  is decreasing.
- ▶ CASE 1:  $g(x)$  is non-decreasing
- ▶  $F_Y(y) = P(g(X) \leq y) = P(X \leq h(y)) = F_X(h(y))$ .

# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?
- ▶ Same when  $g$  is increasing and compliments when  $g$  is decreasing.
- ▶ CASE 1:  $g(x)$  is non-decreasing
- ▶  $F_Y(y) = P(g(X) \leq y) = P(X \leq h(y)) = F_X(h(y))$ .
- ▶  $f_Y(y) = \frac{d}{dy}(F_X(h(y))) =$

# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?
- ▶ Same when  $g$  is increasing and compliments when  $g$  is decreasing.
- ▶ CASE 1:  $g(x)$  is non-decreasing
- ▶  $F_Y(y) = P(g(X) \leq y) = P(X \leq h(y)) = F_X(h(y))$ .
- ▶  $f_Y(y) = \frac{d}{dy}(F_X(h(y))) = f_X(h(y)) \frac{dh}{dy}(y)$  where  $\frac{dh}{dy}(y) \geq 0$  as  $h$  is also non-decreasing.

# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?
- ▶ Same when  $g$  is increasing and compliments when  $g$  is decreasing.
- ▶ CASE 1:  $g(x)$  is non-decreasing
- ▶  $F_Y(y) = P(g(X) \leq y) = P(X \leq h(y)) = F_X(h(y))$ .
- ▶  $f_Y(y) = \frac{d}{dy}(F_X(h(y))) = f_X(h(y)) \frac{dh}{dy}(y)$  where  $\frac{dh}{dy}(y) \geq 0$  as  $h$  is also non-decreasing.
- ▶ Rewritten therefore as  $f_Y(y) = f_X(h(y)) \left| \frac{dh}{dy}(y) \right|$

# Function of continuous random variables

# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?

# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?
- ▶ Same when  $g$  is increasing and compliments when  $g$  is decreasing.

# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?
- ▶ Same when  $g$  is increasing and compliments when  $g$  is decreasing.
- ▶ CASE 2:  $g(x)$  is non-increasing



# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?
- ▶ Same when  $g$  is increasing and compliments when  $g$  is decreasing.
- ▶ CASE 2:  $g(x)$  is non-increasing
- ▶  $F_Y(y) = P(g(X) \leq y) = P(X > h(y)) = 1 - F_X(h(y)).$

# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?
- ▶ Same when  $g$  is increasing and compliments when  $g$  is decreasing.
- ▶ CASE 2:  $g(x)$  is non-increasing
- ▶  $F_Y(y) = P(g(X) \leq y) = P(X > h(y)) = 1 - F_X(h(y)).$
- ▶  $f_Y(y) = -\frac{d}{dy}(F_X(h(y))) =$

# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?
- ▶ Same when  $g$  is increasing and compliments when  $g$  is decreasing.
- ▶ CASE 2:  $g(x)$  is non-increasing
- ▶  $F_Y(y) = P(g(X) \leq y) = P(X > h(y)) = 1 - F_X(h(y))$ .
- ▶  $f_Y(y) = -\frac{d}{dy}(F_X(h(y))) = -f_X(h(y))\frac{dh}{dy}(y)$  where  $\frac{dh}{dy}(y) \leq 0$  as  $h$  is non-increasing as well.

# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?
- ▶ Same when  $g$  is increasing and compliments when  $g$  is decreasing.
- ▶ CASE 2:  $g(x)$  is non-increasing
- ▶  $F_Y(y) = P(g(X) \leq y) = P(X > h(y)) = 1 - F_X(h(y)).$
- ▶  $f_Y(y) = -\frac{d}{dy}(F_X(h(y))) = -f_X(h(y))\frac{dh}{dy}(y)$  where  $\frac{dh}{dy}(y) \leq 0$  as  $h$  is non-increasing as well.
- ▶ Rewritten therefore as  $f_Y(y) = f_X(h(y))|\frac{dh}{dy}(y)|.$  □

# Function of continuous random variables

- ▶ Are the two events  $\{g(X) \leq y\}$  and  $\{X \leq h(y)\}$  same ?
- ▶ Same when  $g$  is increasing and compliments when  $g$  is decreasing.
- ▶ CASE 2:  $g(x)$  is non-increasing
- ▶  $F_Y(y) = P(g(X) \leq y) = P(X > h(y)) = 1 - F_X(h(y))$ .
- ▶  $f_Y(y) = -\frac{d}{dy}(F_X(h(y))) = -f_X(h(y))\frac{dh}{dy}(y)$  where  $\frac{dh}{dy}(y) \leq 0$  as  $h$  is non-increasing as well.
- ▶ Rewritten therefore as  $f_Y(y) = f_X(h(y))|\frac{dh}{dy}(y)|$ . □

HW: What about the case when  $g$  is not monotone ?

# Multiple random variables

# A running example

# A running example

- ▶ Consider an experiment of tossing a coin and a dice together.



# A running example

- ▶ Consider an experiment of tossing a coin and a dice together.
- ▶  $\Omega = \{0, 1\} \times \{1, 2, 3, 4, 5, 6\}$ .

# A running example

- ▶ Consider an experiment of tossing a coin and a dice together.
- ▶  $\Omega = \{0, 1\} \times \{1, 2, 3, 4, 5, 6\}$ .  $\mathcal{F} = 2^\Omega$ .

# A running example

- ▶ Consider an experiment of tossing a coin and a dice together.
- ▶  $\Omega = \{0, 1\} \times \{1, 2, 3, 4, 5, 6\}$ .  $\mathcal{F} = 2^\Omega$ .  $\mathbb{P}(\omega) = \frac{1}{12}$ .

# A running example

- ▶ Consider an experiment of tossing a coin and a dice together.
- ▶  $\Omega = \{0, 1\} \times \{1, 2, 3, 4, 5, 6\}$ .  $\mathcal{F} = 2^\Omega$ .  $\mathbb{P}(\omega) = \frac{1}{12}$ .
- ▶ Let  $X$  and  $Y$  denote the random variables depicting outcome of a coin and dice respectively.

# A running example

- ▶ Consider an experiment of tossing a coin and a dice together.
- ▶  $\Omega = \{0, 1\} \times \{1, 2, 3, 4, 5, 6\}$ .  $\mathcal{F} = 2^\Omega$ .  $\mathbb{P}(\omega) = \frac{1}{12}$ .
- ▶ Let  $X$  and  $Y$  denote the random variables depicting outcome of a coin and dice respectively.
- ▶ For  $\omega = (1, 5)$  we have  $X(\omega) = 1$  and  $Y(\omega) = 5$ .

# A running example

- ▶ Consider an experiment of tossing a coin and a dice together.
- ▶  $\Omega = \{0, 1\} \times \{1, 2, 3, 4, 5, 6\}$ .  $\mathcal{F} = 2^\Omega$ .  $\mathbb{P}(\omega) = \frac{1}{12}$ .
- ▶ Let  $X$  and  $Y$  denote the random variables depicting outcome of a coin and dice respectively.
- ▶ For  $\omega = (1, 5)$  we have  $X(\omega) = 1$  and  $Y(\omega) = 5$ .
- ▶ We are now interested in the joint PMF  $p_{XY}(x, y)$  and joint CDF  $F_{XY}(x, y)$  of  $X$  and  $Y$  together.

## An example

- ▶ We are now interested in the joint PMF  $p_{XY}(x, y)$  and joint CDF  $F_{XY}(x, y)$  of  $X$  and  $Y$  together.

## An example

- ▶ We are now interested in the joint PMF  $p_{XY}(x, y)$  and joint CDF  $F_{XY}(x, y)$  of  $X$  and  $Y$  together.
- ▶  $p_{XY}(x, y) := \mathbb{P}\{\omega \in \Omega : X(\omega) = x \text{ and } Y(\omega) = y\}.$



## An example

- ▶ We are now interested in the joint PMF  $p_{XY}(x, y)$  and joint CDF  $F_{XY}(x, y)$  of  $X$  and  $Y$  together.
- ▶  $p_{XY}(x, y) := \mathbb{P}\{\omega \in \Omega : X(\omega) = x \text{ and } Y(\omega) = y\}.$
- ▶  $F_{XY}(x, y) := \mathbb{P}\{\omega \in \Omega : X(\omega) \leq x \text{ and } Y(\omega) \leq y\}.$

## An example

- ▶ We are now interested in the joint PMF  $p_{XY}(x, y)$  and joint CDF  $F_{XY}(x, y)$  of  $X$  and  $Y$  together.
- ▶  $p_{XY}(x, y) := \mathbb{P}\{\omega \in \Omega : X(\omega) = x \text{ and } Y(\omega) = y\}$ .
- ▶  $F_{XY}(x, y) := \mathbb{P}\{\omega \in \Omega : X(\omega) \leq x \text{ and } Y(\omega) \leq y\}$ .
- ▶ We can use PMF to calculate  $P((X, Y) \in A)$ .

# An example

- ▶ We are now interested in the joint PMF  $p_{XY}(x, y)$  and joint CDF  $F_{XY}(x, y)$  of  $X$  and  $Y$  together.
- ▶  $p_{XY}(x, y) := \mathbb{P}\{\omega \in \Omega : X(\omega) = x \text{ and } Y(\omega) = y\}$ .
- ▶  $F_{XY}(x, y) := \mathbb{P}\{\omega \in \Omega : X(\omega) \leq x \text{ and } Y(\omega) \leq y\}$ .
- ▶ We can use PMF to calculate  $P((X, Y) \in A)$ .
- ▶  $P((X, Y) \in A) = \mathbb{P}\{\omega \in \Omega : (X(\omega), Y(\omega)) \in A\}$ .

# An example

- ▶ We are now interested in the joint PMF  $p_{XY}(x, y)$  and joint CDF  $F_{XY}(x, y)$  of  $X$  and  $Y$  together.
- ▶  $p_{XY}(x, y) := \mathbb{P}\{\omega \in \Omega : X(\omega) = x \text{ and } Y(\omega) = y\}$ .
- ▶  $F_{XY}(x, y) := \mathbb{P}\{\omega \in \Omega : X(\omega) \leq x \text{ and } Y(\omega) \leq y\}$ .
- ▶ We can use PMF to calculate  $P((X, Y) \in A)$ .
- ▶  $P((X, Y) \in A) = \mathbb{P}\{\omega \in \Omega : (X(\omega), Y(\omega)) \in A\}$ . Therefore  $P((X, Y) \in A) = \sum_{(x,y) \in A} p_{XY}(x, y)$ .

# An example

- ▶ We are now interested in the joint PMF  $p_{XY}(x, y)$  and joint CDF  $F_{XY}(x, y)$  of  $X$  and  $Y$  together.
- ▶  $p_{XY}(x, y) := \mathbb{P}\{\omega \in \Omega : X(\omega) = x \text{ and } Y(\omega) = y\}$ .
- ▶  $F_{XY}(x, y) := \mathbb{P}\{\omega \in \Omega : X(\omega) \leq x \text{ and } Y(\omega) \leq y\}$ .
- ▶ We can use PMF to calculate  $P((X, Y) \in A)$ .
- ▶  $P((X, Y) \in A) = \mathbb{P}\{\omega \in \Omega : (X(\omega), Y(\omega)) \in A\}$ . Therefore  $P((X, Y) \in A) = \sum_{(x,y) \in A} p_{XY}(x, y)$ .
- ▶ Suppose  $A$  is the event that you get a head and the roll is even. What is  $P((X, Y) \in A)$ ?

# Marginals

# Marginals

- ▶ What is  $p_{XY}(1, i)$ ?

# Marginals

- ▶ What is  $p_{XY}(1, i)$ ? ( $= \frac{1}{12}$ ).



# Marginals

- ▶ What is  $p_{XY}(1, i)$ ? ( $= \frac{1}{12}$ ).
- ▶  $\sum_i p_{XY}(1, i) =$

# Marginals

- ▶ What is  $p_{XY}(1, i)$ ? ( $= \frac{1}{12}$ ).
- ▶  $\sum_i p_{XY}(1, i) = \mathbb{P}\{\omega \in \Omega : X(\omega) = 1\}$

# Marginals

- ▶ What is  $p_{XY}(1, i)$ ? ( $= \frac{1}{12}$ ).
- ▶  $\sum_i p_{XY}(1, i) = \mathbb{P}\{\omega \in \Omega : X(\omega) = 1\} = \frac{1}{2}$

# Marginals

- ▶ What is  $p_{XY}(1, i)$ ? ( $= \frac{1}{12}$ ).
- ▶  $\sum_i p_{XY}(1, i) = \mathbb{P}\{\omega \in \Omega : X(\omega) = 1\} = \frac{1}{2} = p_X(x)$ .

# Marginals

- ▶ What is  $p_{XY}(1, i)$ ? ( $= \frac{1}{12}$ ).
- ▶  $\sum_i p_{XY}(1, i) = \mathbb{P}\{\omega \in \Omega : X(\omega) = 1\} = \frac{1}{2} = p_X(x)$ .
- ▶ Similarly,  $p_{XY}(1, i) + p_{XY}(0, i) =$

# Marginals

- ▶ What is  $p_{XY}(1, i)$ ? ( $= \frac{1}{12}$ ).
- ▶  $\sum_i p_{XY}(1, i) = \mathbb{P}\{\omega \in \Omega : X(\omega) = 1\} = \frac{1}{2} = p_X(x)$ .
- ▶ Similarly,  $p_{XY}(1, i) + p_{XY}(0, i) = \frac{1}{6}$

# Marginals

- ▶ What is  $p_{XY}(1, i)$ ? ( $= \frac{1}{12}$ ).
- ▶  $\sum_i p_{XY}(1, i) = \mathbb{P}\{\omega \in \Omega : X(\omega) = 1\} = \frac{1}{2} = p_X(x)$ .
- ▶ Similarly,  $p_{XY}(1, i) + p_{XY}(0, i) = \frac{1}{6} = p_Y(i)$ .

# Marginals

- ▶ What is  $p_{XY}(1, i)$ ? ( $= \frac{1}{12}$ ).
- ▶  $\sum_i p_{XY}(1, i) = \mathbb{P}\{\omega \in \Omega : X(\omega) = 1\} = \frac{1}{2} = p_X(x)$ .
- ▶ Similarly,  $p_{XY}(1, i) + p_{XY}(0, i) = \frac{1}{6} = p_Y(i)$ .

The marginal PMF's  $p_X$  and  $p_Y$  can be obtained from the joint PMF as follows:

$$p_X(x) = \sum_y p_{XY}(x, y) \text{ and } p_Y(y) = \sum_x p_{XY}(x, y).$$

This is true in general, and requires a proof (later).