




A. ASENSIO RAMOS

HE I MULTIPLETS: HAZEL PHYSICS

ChroTel Helium 1083 nm

16 Apr 2015 07:48:20 UTC



N
W



ADVANCED FORWARD MODELING AND INVERSION OF STOKES PROFILES RESULTING FROM THE JOINT ACTION OF THE HANLE AND ZEEMAN EFFECTS

A. ASENSIO RAMOS AND J. TRUJILLO BUENO¹

Instituto de Astrofísica de Canarias, Vía Láctea s/n, E-38205 La Laguna, Tenerife, Spain; aasensio@iac.es, jtb@iac.es

AND

E. LANDI DEGL'INNOCENTI

Università degli Studi di Firenze, Dipartimento di Astronomia e Scienza dello Spazio,
Largo Enrico Fermi 2, I-50125 Florence, Italy; landie@arcetri.astro.it

Received 2007 December 11; accepted 2008 April 15

ABSTRACT

A big challenge in solar and stellar physics in the coming years will be to decipher the magnetism of the solar outer atmosphere (chromosphere and corona) along with its dynamic coupling with the magnetic fields of the underlying photosphere. To this end, it is important to develop rigorous diagnostic tools for the physical interpretation of spectro-polarimetric observations in suitably chosen spectral lines. Here we present a computer program for the synthesis and inversion of Stokes profiles caused by the joint action of atomic level polarization and the Hanle and Zeeman effects in some spectral lines of diagnostic interest, such as those of the He I 10830 Å and 5876 Å (or D₃) multiplets. It is based on the quantum theory of spectral line polarization, which takes into account in a rigorous way all the relevant physical mechanisms and ingredients (optical pumping, atomic level polarization, level crossings and repulsions, Zeeman, Paschen-Back, and Hanle effects). The influence of radiative transfer on the emergent spectral line radiation is taken into account through a suitable slab model. The user can either calculate the emergent intensity and polarization for any given magnetic field vector or infer the dynamical and magnetic properties from the observed Stokes profiles via an efficient inversion algorithm based on global optimization methods. The reliability of the forward modeling and inversion code presented here is demonstrated through several applications, which range from the inference of the magnetic field vector in solar active regions to determining whether or not it is canopy-like in quiet chromospheric regions. This user-friendly diagnostic tool called “HAZEL” (from HAnle and ZEeman Light) is offered to the astrophysical community, with the hope that it will facilitate new advances in solar and stellar physics.

Subject headings: magnetic fields — methods: data analysis — methods: numerical — polarization — radiative transfer — scattering — Sun: chromosphere



Search or jump to...

Pull requests Issues Marketplace Explore

Fork me on GitHub

aasensio / hazel2

Unwatch 3

Star 1

Fork 1

Code

Issues 7

Pull requests 0

Projects 0

Wiki

Insights

Settings

Hazel v2.0: synthesis and inversion of Stokes profiles caused by the joint action of atomic level polarization and the Hanle and Zeeman effects <https://aasensio.github.io/hazel2/index.html>

Edit

Add topics

52 commits

2 branches

0 releases

1 contributor

MIT

Branch: master ▾

New pull request

Create new file

Upload files

Find file

Clone or download ▾

 aasensio	Fixed some issues	Latest commit 330de60 21 hours ago
 docs	Fixed some issues	21 hours ago
 examples	Fixed some issues	21 hours ago
 gui	Fixed a bug appearing with newest gfortran compilers	3 months ago
 hazel	Fixed some issues	21 hours ago
 src	Fixed some issues	21 hours ago
 test	Added badges for coveralls and travis	4 months ago
 .gitignore	Notebooks added for docs. Improvements on inversion	3 months ago
 .travis.yml	Added zarr in travis	3 months ago
 LICENSE	Deployment to pypi	5 months ago
 MANIFEST.in	Added tests to travis	5 months ago
 README.md	Typo on README	3 months ago
 TODO	Fixed issue #12	a day ago
 deploy_testpypi	Deployment to pypi	5 months ago
 setup.cfg	Removed gregor data	5 months ago
 setup.py	Removed matplotlib as requirement	2 months ago

README.md



Hazel v2.0

GitHub [aasensio/hazel2](#) build [passing](#) coverage [33%](#) license [MIT](#) ADS [2008ApJ...683..542A](#) arXiv [0804.2695](#)


HAZEL

- ▶ 5-term atom
- ▶ Simple slab with constant physical properties
- ▶ Line formation is “hidden” on the optical depth
- ▶ Atomic level polarization + Zeeman (Paschen-Back) + Hanle effect
- ▶ Flat spectrum
- ▶ Fixed pumping radiation field
- ▶ Very simple formal solution
- ▶ Serial/MPI version



HAZEL2

- ▶ Invert He I (many lines simultaneously) + photospheric + telluric lines
- ▶ Flexibility on how to define topology:
 $(ph1+ph2) \rightarrow (ch1+ch2) \rightarrow te1$
- ▶ Sparsity (and other) regularization (WIP)
- ▶ 2D regularization (TBD)
- ▶ Python with wrappers to the time consuming parts coded in F90
- ▶ Fast versions using deep learning (TBD)
- ▶ Collaborations are welcome:
<https://github.com/aasensio/hazel2>


PARA-HELIUM







ORTHO-HELIUM







MODEL ATOM




ENERGY SPLITTING




ENERGY SPLITTING



GEOMETRY



GEOMETRY: YOU'LL ALWAYS COME TO THIS SLIDE




STATISTICAL EQUILIBRIUM EQUATIONS

$$\begin{aligned}
\frac{d}{dt} \beta^{LS} \rho_Q^K(J, J') &= -2\pi i \sum_{K'Q'} \sum_{J''J'''} N_{\beta^{LS}}(KQJJ', K'Q'J''J''')^{LS} \rho_{Q'}^{K'}(J'', J''') \\
&+ \sum_{\beta_\ell L_\ell K_\ell Q_\ell J_\ell J'_\ell} \beta_\ell L_\ell S \rho_{Q_\ell}^{K_\ell}(J_\ell, J'_\ell) \mathbb{T}_A(\beta^{LS} K Q J J', \beta_\ell L_\ell S K_\ell Q_\ell J_\ell J'_\ell) \\
&+ \sum_{\beta_u L_u K_u Q_u J_u J'_u} \beta_u L_u S \rho_{Q_u}^{K_u}(J_u, J'_u) \left[\mathbb{T}_E(\beta^{LS} K Q J J', \beta_u L_u S K_u Q_u J_u J'_u) \right. \\
&\quad \left. + \mathbb{T}_S(\beta^{LS} K Q J J', \beta_u L_u S K_u Q_u J_u J'_u) \right] \\
&- \sum_{K'Q'J''J'''} \beta^{LS} \rho_{Q'}^{K'}(J'', J''') \left[\mathbb{R}_A(\beta^{LS} K Q J J' K' Q' J'' J''') \right. \\
&\quad \left. + \mathbb{R}_E(\beta^{LS} K Q J J' K' Q' J'' J''') + \mathbb{R}_S(\beta^{LS} K Q J J' K' Q' J'' J''') \right]
\end{aligned}$$

EMISSION/ABSORPTION

$$\begin{aligned}
\eta_i^A(\nu, \Omega) &= \frac{h\nu}{4\pi} \mathcal{N} \sum_{\beta_\ell L_\ell S \beta_u L_u} (2L_\ell + 1) B(\beta_\ell L_\ell S \rightarrow \beta_u L_u S) \\
&\times \sum_{KQK_\ell Q_\ell} \sqrt{3(2K+1)(2K_\ell+1)} \\
&\times \sum_{j_\ell J_\ell J'_\ell J''_\ell} \sum_{j_u J_u J'_u M_\ell M'_\ell M_u q q'} (-1)^{1+J''_\ell - M_\ell + q'} C_{J_\ell}^{j_\ell}(\beta_\ell L_\ell S, M_\ell) C_{J''_\ell}^{j_\ell}(\beta_\ell L_\ell S, M_\ell) \\
&\times C_{J_u}^{j_u}(\beta_u L_u S, M_u) C_{J'_u}^{j_u}(\beta_u L_u S, M_u) \sqrt{(2J_\ell + 1)(2J'_\ell + 1)(2J_u + 1)(2J'_u + 1)} \\
&\times \begin{pmatrix} J_u & J_l & 1 \\ -M_u & M_l & -q \end{pmatrix} \begin{pmatrix} J'_u & J'_l & 1 \\ -M_u & M'_l & -q' \end{pmatrix} \begin{pmatrix} 1 & 1 & K \\ q & -q' & -Q \end{pmatrix} \\
&\times \begin{pmatrix} J''_\ell & J'_\ell & K_\ell \\ M_\ell & -M'_\ell & -Q_\ell \end{pmatrix} \left\{ \begin{matrix} L_u & L_\ell & 1 \\ J_\ell & J_u & S \end{matrix} \right\} \left\{ \begin{matrix} L_u & L_\ell & 1 \\ J'_\ell & J'_u & S \end{matrix} \right\} \\
&\times \text{Re} \left[T_Q^K(i, \Omega)^{\beta_\ell L_\ell S} \rho_{Q_\ell}^{K_\ell}(J''_\ell, J'_\ell) \Phi(\nu_{\beta_u L_u S j_u, \beta_\ell L_\ell S j_\ell M_\ell} - \nu) \right].
\end{aligned}$$

EMISSION/ABSORPTION



HAZEL: THINGS TO IMPROVE

- ▶ 5-term atom 
- ▶ Simple slab with constant physical properties 
- ▶ Line formation is “hidden” on the optical depth 
- ▶ Atomic level polarization + Zeeman (Paschen-Back) + Hanle effect
- ▶ Flat spectrum 
- ▶ Fixed pumping radiation field 
- ▶ Very simple formal solution 
- ▶ Serial/MPI version 