



Cambridge IGCSE™

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

ADDITIONAL MATHEMATICS

0606/11

Paper 1 Non-calculator

May/June 2026

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen. Do **not** use correction fluid or tape.
- Do **not** write on any bar codes.
- Calculators must **not** be used in this paper.
- You must show all necessary working clearly.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages.

List of formulas

Equation of a circle with centre (a, b) and radius r .

$$(x - a)^2 + (y - b)^2 = r^2$$

Curved surface area, A , of cone of radius r , sloping edge l .

$$A = \pi r l$$

Surface area, A , of sphere of radius r .

$$A = 4\pi r^2$$

Volume, V , of pyramid or cone, base area A , height h .

$$V = \frac{1}{3} Ah$$

Volume, V , of sphere of radius r .

$$V = \frac{4}{3} \pi r^3$$

Quadratic equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial theorem

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series

$$u_n = a + (n - 1)d$$

$$S_n = \frac{1}{2} n(a + l) = \frac{1}{2} n\{2a + (n - 1)d\}$$

Geometric series

$$u_n = ar^{n-1}$$

$$S_n = \frac{a(1 - r^n)}{1 - r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1 - r} \quad (|r| < 1)$$

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulas for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} ab \sin C$$





1 Solve the equation $|5x + 4| = |3x - 2|$.

[3]



2 A circle has equation $(x-3)^2 + (y-4)^2 = 20$.

(a) Write down the coordinates of the centre of the circle.

[1]

(b) Write down the radius of the circle.

[1]

(c) The line AB is a diameter of this circle.
The point A has coordinates $(-1, 6)$.

Find the coordinates of the point B .

[2]

- 3 Show that $(3 + 5 \cot \theta)^2 + (3 - 5 \cot \theta)^2 - 50 \operatorname{cosec}^2 \theta$, where $0 < \theta < \frac{\pi}{2}$, can be simplified to a constant.

[4]

- 4 Given that $\int_0^a \frac{1}{3x+2} dx = \ln 4$, find the value of the constant a .

[5]

- 5 The polynomial p is such that $p(x) = ax^3 + 4x^2 - ax - 28$, where a is a positive constant.

It is given that $p(a) = 0$.

- (a) Find the value of a .

[4]

- (b) Using your value of a , find $p(x)$ as a product of a linear factor and a quadratic factor.

[2]

- (c) Hence show that the equation $p(x) = 0$ has exactly one root.

[2]



- 6 The function f is such that $f(x) = 3x^2 + 8x$ for $x \geq k$, where k is a constant. It is given that f^{-1} exists. k has the least possible value for which f^{-1} exists.

(a) Find the value of k .

[2]

(b) Find the range of f .

[2]

(c) Find an expression for f^{-1} .

[3]



(d) The function g is such that $g(x) = 3^{-x}$ for $x > -6$.

(i) Write down the domain of gf .

[1]

(ii) Solve the equation $gf(x) = 27$.

[4]

7 A curve has equation $y = \frac{4 + e^{2x}}{2 + e^x}$.

(a) Show that, when $\frac{dy}{dx} = 0$, $e^{2x} + 4e^x - 4 = 0$.

[5]

- (b) Hence find the x -coordinate of the stationary point on the curve.
Give your answer in the form $\ln(a\sqrt{b} + c)$, where a , b and c are integers.

[4]

8 (a) Write down the value of $\lg(10000)$.

[1]

(b) Solve the equation $\log_2(\log_3 x) = 2$.

[3]

(c) Solve the equation $\log_x 5 = \log_5 x^4$.

[4]

- 9 (a) A geometric progression has common ratio r and first term a , where $a \neq 0$.
The 4th term is -8 times the 7th term.

(i) Find the value of r .

[2]

- (ii) The sum of the first 5 terms of this progression is $\frac{99}{32}$.

Find the value of a .

[2]

- (b) The first three terms of a different geometric progression are $\cos \theta \sin \theta$, $\cos \theta \sin^3 \theta$ and $\cos \theta \sin^5 \theta$, for $0 < \theta < \frac{\pi}{2}$.

Show that the sum to infinity of this progression is $\tan \theta$.

[4]

10 A curve has equation $y = f(x)$, where $f''(x) = (2x + 10)^{-\frac{1}{2}} - 2x$.

The curve has a stationary point at $\left(3, \frac{64}{3}\right)$.

Find the equation of the curve.

[8]



- 11 A triangle OAB is such that $\vec{OA} = \mathbf{a}$ and $\vec{OB} = \mathbf{b}$.

The point X lies on OA such that $\vec{OA} = 3\vec{OX}$.

The point Y lies on XB such that $\vec{XB} = 5\vec{XY}$.

The point Z lies on OB such that $\vec{OB} = m\vec{OZ}$, where m is a scalar.

$\vec{OX} = n\vec{ZY}$, where n is a scalar.

Use a vector method to find the values of m and n .

[8]

Question 12 is on the back page.





12 Find the value of n such that ${}^{n+3}C_6 = 12 \times {}^{n+2}C_5$.

[3]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in our Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge International Education is the name of our awarding body and a part of Cambridge University Press & Assessment, which is a department of the University of Cambridge.

