Grasp-relevant Representation of Objects in the Human Parietal and Occipito-temporal Cortices

Amirali Soltani Tehrani[1], Kristin Woodard[2], Emalie McMahon[2], Leah Ettensohn[2], Natalia Pallis-Hassani[2], John Ingeholm[2], Leslie Ungerleider[2], Chris Baker[2], Maryam Vaziri-Pashkam[1,2]

[1]Movement and Visual Perception Lab, Department of Psychological & Brain Sciences, University of Delaware.

[2]Laboratory of Brain and Cognition, National Institute of Mental Health

Introduction

- Human visual system is capable of various behaviors toward objects, from recognizing them to performing action on them.
- Multiple studies have shown object representation in both human occipito-temporal and parietal cortices.
- Previous studies suggest distinct roles for object-selective occipito-temporal and parietal regions.

How do these object representations support such diverse behaviors?

Behavioral Tasks

Odd-one-out (subjective similarity)



- 70 objects
- 112 points per pair

Grasp Similarity



- 58 objects
- 16 sensors in three coordinates (48D vector)

RDMs are $\rho = 0.27$ correlated!

fMRI Experiment





- 64 3D-printed objects
- Data collected in four sessions

Object Localizer
Objects > Scrambled

Grasp Localizer
Grasp> Touch





Multiple response profiles were observed in the distinct ROIs



Distinct patterns in the higher-level visual regions in relation to the behaviors



LOT
Inf IPS
AIP
Pre-C
SMG



First principal components create a manifold reflecting phase transition in addition to ROI separation

Conclusion

- **Occipito-temporal and posterior parietal regions represent features relevant for subjective judgements when viewing objects.**
- **Anterior parietal regions jointly represent features relevant for subjective judgements and grasp when planning and performing movements.**
- **Experimental phase and function jointly determine patterns of response in object-selective cortex.**

@aasolteh.bsky.social

@aasolteh