

Unitary Coupled Cluster Theory

Cheng Lab

April 6, 2018

1 Derivation of Polarisation Propogator $G_{pr,rs}$ in the new unitary transformed excitation/deexcitation scheme

The Polarisation Propogator in superoperator representation is given by :

$$\begin{aligned} G_{pq,rs}(\omega) &= \langle \Psi_{gr} | [a_p a_q^\dagger, (\omega \hat{I} - \hat{H})^{-1} a_r^\dagger a_s] | \Psi_{gr} \rangle \\ &= \langle \Psi_{gr} | [q_I, (\omega \hat{I} - \hat{H})^{-1} q_J^\dagger] | \Psi_{gr} \rangle \end{aligned}$$

A Binary Product can be defined as follows :

$$(A|B) = \langle \Psi_{gr} | [A^\dagger, B] | \Psi \rangle$$

Another definition used in this derivation is :

$$(A|\hat{O}|B) \equiv (A|\hat{O}B)$$

Polarisation Propogator can be written in terms of binary product notation as :

$$G(\omega) = (q_i^\dagger | (\omega \hat{I} - \hat{H})^{-1} q_j^\dagger)$$

In our self consistant polarisation propagation theory, the unitary transformed excitation/deexcitation operators are defined as :

$$y_i^\dagger = \exp(\sigma) b_i^\dagger \exp(-\sigma)$$

Using the resolution of identity in G :

$$\begin{aligned} \hat{I} &= \sum_I \{|y_I^\dagger\rangle \langle y_I^\dagger| - |y_I\rangle \langle y_I|\} \\ G(\omega) &= (q_I | y_I^\dagger) (y_I^\dagger | (\omega \hat{I} - \hat{H})^{-1} q_j^\dagger) - (q_I | y_I) (y_I | (\omega \hat{I} - \hat{H})^{-1} q_j^\dagger) \end{aligned}$$

Taking the second half of the expression and expanding through tailors expansion :

$$\begin{aligned}
(y_I | (\omega \hat{I} - \hat{H})^{-1} q_j^\dagger) &= \langle \Psi_{gr} | [y_I, (\omega \hat{I} - \hat{H})^{-1} q_j^\dagger] | \Psi_{gr} \rangle \\
&= \langle \Psi_{gr} | [y_I, \frac{1}{\omega} \left((\hat{I} - \hat{H}) \right)^{-1} q_j^\dagger] | \Psi_{gr} \rangle \\
&= \langle \Psi_{gr} | [y_I, \frac{1}{\omega} \left((\hat{I} + \frac{1}{\omega} \hat{H} + \frac{1}{\omega^2} \hat{H}^2 \dots) \right) q_j^\dagger] | \Psi_{gr} \rangle \\
&= \frac{1}{\omega} \left(\langle \Psi_{gr} | [y_I, (\hat{I} q_j^\dagger)] | \Psi_{gr} \rangle + \langle \Psi_{gr} | [y_I, \frac{1}{\omega} \hat{H} q_j^\dagger] | \Psi_{gr} \rangle + \langle \Psi_{gr} | [y_I, \frac{1}{\omega^2} \hat{H}^2 q_j^\dagger] | \Psi_{gr} \rangle \right) \\
&= \frac{1}{\omega} \left((y_I | (\hat{I} q_j^\dagger)) + (y_I | \frac{1}{\omega} \hat{H} q_j^\dagger) + (y_I | \frac{1}{\omega^2} \hat{H}^2 q_j^\dagger) \right) \\
&= \frac{1}{\omega} \left((y_I | (\hat{I} q_j^\dagger)) + (y_I | \frac{1}{\omega} \hat{H} q_j^\dagger) + (y_I | \frac{1}{\omega^2} \hat{H}^2 q_j^\dagger) \dots \right) \\
&= \frac{1}{\omega} (y_I | (\hat{I} + \frac{1}{\omega} \hat{H} + \frac{1}{\omega^2} \hat{H}^2 \dots) q_j^\dagger) \\
&= \frac{1}{\omega} (y_I | (\hat{I} - \frac{\hat{H}}{\omega})^{-1} q_j^\dagger) \\
&= (y_I | (\omega \hat{I} - \hat{H})^{-1} q_j^\dagger)
\end{aligned}$$

Using this result in the previously derived expression :

$$\begin{aligned}
\hat{I} &= \sum_I \{ |y_I^\dagger\rangle (y_I^\dagger | - |y_I\rangle) (y_I | \} \\
G(\omega) &= (q_I^\dagger | y_I^\dagger) (y_I^\dagger | (\omega \hat{I} - \hat{H})^{-1} q_j^\dagger) - (q_I^\dagger | y_I) (y_I | (\omega \hat{I} - \hat{H})^{-1} q_j^\dagger) \\
&= (q_I^\dagger | y_I^\dagger) (y_I^\dagger | (\omega \hat{I} - \hat{H})^{-1} q_j^\dagger) - (q_I^\dagger | y_I) (y_I | (\omega \hat{I} - \hat{H})^{-1} q_j^\dagger)
\end{aligned}$$

Again using resolution of Identity Operator :

$$\begin{aligned}
\hat{I} &= \sum_I \{ |y_I\rangle (y_I | - |y_I^\dagger\rangle) (y_I^\dagger | \} \\
G(\omega) &= (q_I^\dagger | y_I^\dagger) (y_I^\dagger | (\omega \hat{I} - \hat{H})^{-1} |y_J^\dagger\rangle) (y_J^\dagger | q_j^\dagger) - (q_I^\dagger | y_I) (y_I | (\omega \hat{I} - \hat{H})^{-1} |y_J^\dagger\rangle) (y_J^\dagger | q_j^\dagger) \\
&\quad - (q_I^\dagger | y_I^\dagger) (y_I^\dagger | (\omega \hat{I} - \hat{H})^{-1} |y_J\rangle) (y_J | q_j^\dagger) + (q_I^\dagger | y_I) (y_I | (\omega \hat{I} - \hat{H})^{-1} |y_J\rangle) (y_J | q_j^\dagger)
\end{aligned}$$

1.1