

# Extraction of partonic transverse momentum distributions from semi-inclusive deep inelastic scattering and Drell-Yan data

Alessandro Bacchetta,<sup>1,2,\*</sup> Filippo Delcarro,<sup>1,2,†</sup> Cristian Pisano,<sup>1,2,\*</sup> Marco Radici,<sup>2,‡</sup> and Andrea Signori<sup>3,4,5,§</sup>

<sup>1</sup>*Dipartimento di Fisica, Università di Pavia, via Bassi 6, I-27100 Pavia*

<sup>2</sup>*INFN Sezione di Pavia, via Bassi 6, I-27100 Pavia, Italy*

<sup>3</sup>*Theory Center, Thomas Jefferson National Accelerator Facility,  
12000 Jefferson Avenue, Newport News, VA 23606, USA*

<sup>4</sup>*Department of Physics and Astronomy, VU University Amsterdam,  
De Boelelaan 1081, NL-1081 HV Amsterdam, the Netherlands*

<sup>5</sup>*Nikhef, Science Park 105, NL-1098 XG Amsterdam, the Netherlands*

(Dated: Tuesday 28<sup>th</sup> February, 2017, 16:35)

We present an extraction of unpolarized partonic transverse momentum distributions (TMDs) from a simultaneous fit of available data measured in semi-inclusive deep inelastic scattering and in Drell-Yan processes through the production of photon and  $Z$  bosons. To connect data at different scales, we use TMD evolution at next-to-leading logarithmic accuracy. The analysis is restricted to the low-transverse-momentum region, with no matching to fixed-order calculations at high transverse momentum. We introduce specific choices to deal with TMD evolution at low scales, of the order of 1 GeV<sup>2</sup>. This could be considered as a first attempt at a global fit of TMDs.

PACS numbers: 13.60.Le, 13.87.Fh, 14.20.Dh

## Contents

<b>I. Introduction</b>	2
<b>II. Formalism</b>	2
A. Semi-inclusive DIS	2
B. Drell-Yan processes	4
C. TMDs and their evolution	5
<b>III. Data analysis</b>	8
A. Hermes data	8
B. Compass data	8
C. Low-energy Drell-Yan data	9
D. Z-boson production data	9
E. The replica method	10
<b>IV. Results</b>	12
A. Agreement between data and theory	12
Semi-inclusive DIS	12
Drell-Yan processes	12
B. Transverse momentum dependence at 1 GeV	13
C. Comments on flavor-dependent fits	13
<b>V. Conclusions and outlook</b>	13
<b>Acknowledgments</b>	16
<b>References</b>	18

---

\*Electronic address: [alessandro.bacchetta@unipv.it](mailto:alessandro.bacchetta@unipv.it)

†Electronic address: [filippo.delcarro01@ateneopv.it](mailto:filippo.delcarro01@ateneopv.it)

‡Electronic address: [marco.radici@pv.infn.it](mailto:marco.radici@pv.infn.it)

§Electronic address: [asignori@jlab.org](mailto:asignori@jlab.org)

## I. INTRODUCTION

Parton distribution functions describe the internal structure of the nucleon in terms of its elementary constituents (quarks and gluons). They cannot be easily computed from first principles, because they require the ability to carry out Quantum Chromodynamics (QCD) calculations in a nonperturbative regime. Many experimental observables in hard scattering experiments involving hadrons are related to parton distribution functions (PDFs) and fragmentation functions (FFs), in a way that is specified by factorization theorems (see, e.g., Refs. [1, 2]). These theorems also elucidate the universality properties of PDFs and FFs (i.e., the fact that they are the same in different processes) and their evolution equations (i.e., how they get modified by the change in the hard scale of the process). Availability of measurements of different processes in different experiments makes it possible to test the reliability of factorization theorems and extract PDFs and FFs through so-called global fits. On the other side, the knowledge of PDFs and FFs allows us to make predictions for other hard hadronic processes. These general statements apply equally well to standard collinear PDFs and FFs and to transverse-momentum-dependent parton distribution functions (TMD PDFs) and fragmentation functions (TMD FFs). Collinear PDFs describe the distribution of partons integrated over all components of partonic momentum except the one collinear to the parent hadron; hence, collinear PDFs are functions only of the parton longitudinal momentum fraction  $x$ . TMD PDFs (or TMDs for short) include also the dependence on transverse momentum components  $k_\perp^2$ . They can be interpreted as three-dimensional generalizations of collinear PDFs. Similar arguments apply to collinear FFs and TMD FFs.

There are several differences between collinear and TMD distributions. From the formal point of view, factorization theorems for the two types of functions are qualitatively different, implying also different universality properties and evolution equations [3]. From the experimental point of view, observables related to TMDs require the measurement of some transverse momentum component much smaller than the hard scale of the process [4, 5]. For instance, Deep-Inelastic Scattering (DIS) is characterized by a hard scale represented by the 4-momentum squared of the virtual photon ( $-Q^2$ ). In inclusive DIS this is the only scale of the process, and access is limited to collinear PDFs and FFs. In semi-inclusive DIS (SIDIS) also the transverse momentum of the outgoing hadron ( $P_{hT}$ ) can be measured [6, 7]. If  $P_{hT}^2 \ll Q^2$ , TMD factorization can be applied and the process is sensitive to TMDs [2].

If polarization is taken into account, several TMDs can be introduced [6, 8–11]. Attempts to extract some of them have already been presented in the past [12–19]. In this work, we focus on the simplest ones, i.e., the unpolarized TMD PDF  $f_1^q(x, k_\perp^2)$  and the unpolarized TMD FF  $D_1^{q \rightarrow h}(z, P_{hT}^2)$ , where  $z$  is the fractional energy carried by the detected hadron  $h$ . Despite their simplicity, the phenomenology of these unpolarized TMDs present several challenges [20]: the functional form of TMDs at low partonic transverse momentum, its possible dependence on the parton flavor [21], the implementation of TMD evolution [3, 22], the matching to fixed-order calculations in collinear factorization [23].

We take into consideration three kinds of processes: semi-inclusive DIS, and Drell–Yan processes (DY) with the production of virtual photons and  $Z$  bosons. To date, they represent **almost** [AB:why almost? what else could be used?] all possible processes where experimental information is available for unpolarized TMD extractions. The only important process currently missing is electron-positron annihilation, which is particularly important for the determination of TMD FFs [22]. This work can therefore be considered as the first attempt at a global fit of TMDs.

The paper is organized as follows. In Sec. II, the general formalism for TMDs in SIDIS and DY processes is briefly outlined, including a description of the assumptions and approximations in the phenomenological implementation of TMD evolution equations. In Sec. ??, the criteria for selecting the data analyzed in the fit are summarized and commented. In Sec. IV, the results of our global fit are presented and discussed. In Sec. V, we draw some conclusions.

## II. FORMALISM

Shall we add pictures for the kinematics of SIDIS and DY data ? E.g. see Fig. 1 in [21].

### A. Semi-inclusive DIS

In one-particle **SIDIS**, a lepton  $\ell$  with momentum  $l$  scatters off a hadron target  $N$  with mass  $M$  and momentum  $P$ . In the final state, the scattered lepton momentum  $l'$  is measured together with one hadron  $h$  with mass  $M_h$  and momentum  $P_h$ . The corresponding reaction formula is

$$\ell(l) + N(P) \rightarrow \ell(l') + h(P_h) + X. \quad (1)$$

The space-like momentum transfer is  $q = l - l'$ , with  $Q^2 = -q^2$ . We introduce the usual invariants

$$x = \frac{Q^2}{2P \cdot q}, \quad y = \frac{P \cdot q}{P \cdot l}, \quad z = \frac{P \cdot P_h}{P \cdot q}, \quad \gamma = \frac{2Mx}{Q}. \quad (2)$$

The available data refer to **SIDIS** hadron multiplicities, namely to the differential number of hadrons produced per corresponding inclusive DIS event. In terms of cross sections, we define the multiplicities as

$$m_N^h(x, z, |\mathbf{P}_{hT}|, Q^2) = \frac{d\sigma_N^h/(dx dz d|\mathbf{P}_{hT}| dQ^2)}{d\sigma_{\text{DIS}}/(dx dQ^2)}, \quad (3)$$

where  $d\sigma_N^h$  is the differential cross section for the **SIDIS** process and  $d\sigma_{\text{DIS}}$  is the corresponding inclusive one, and where  $\mathbf{P}_{hT}$  is the component of  $\mathbf{P}_h$  transverse to  $\mathbf{q}$ . In the single-photon-exchange approximation, the multiplicities can be written as ratios of structure functions (see [7] for details):

$$m_N^h(x, z, |\mathbf{P}_{hT}|, Q^2) = \frac{2\pi |\mathbf{P}_{hT}| F_{UU,T}(x, z, \mathbf{P}_{hT}^2, Q^2) + 2\pi \varepsilon |\mathbf{P}_{hT}| F_{UU,L}(x, z, \mathbf{P}_{hT}^2, Q^2)}{F_T(x, Q^2) + \varepsilon F_L(x, Q^2)}, \quad (4)$$

where

$$\varepsilon = \frac{1 - y - \frac{1}{4}\gamma^2 y^2}{1 - y + \frac{1}{2}y^2 + \frac{1}{4}\gamma^2 y^2}, \quad (5)$$

and the structure function  $F_{XY,Z}$  corresponds to a lepton with polarization  $X$  scattering on a target with polarization  $Y$  by exchanging a virtual photon in a polarization state  $Z$ .

The semi-inclusive cross section can be expressed in a factorized form in terms of TMDs only in the kinematical limits  $M^2 \ll Q^2$  and  $\mathbf{P}_T^2 \ll Q^2$ . In these limits, the structure function  $F_{UU,L}$  of Eq. (4) can be neglected [24]. The structure function  $F_L$  in the denominator contains contributions involving powers of the strong coupling constant  $\alpha_S$  at an order that goes beyond the level reached in this analysis; hence, it will be consistently neglected (see also Ref. [21]).

To express the structure functions in terms of TMD PDFs and FFs, we rely on the factorized formula for **SIDIS** [2, 25–32]:

$$\begin{aligned} F_{UU,T}(x, z, \mathbf{P}_{hT}^2, Q^2) &= \sum_a \mathcal{H}_{UU,T}^a(Q^2; \mu^2) \\ &\times \int d\mathbf{k}_\perp d\mathbf{P}_\perp f_1^a(x, \mathbf{k}_\perp^2; \mu^2) D_1^{a \rightarrow h}(z, \mathbf{P}_\perp^2; \mu^2) \delta(z\mathbf{k}_\perp - \mathbf{P}_{hT} + \mathbf{P}_\perp) \\ &+ Y_{UU,T}(Q^2, \mathbf{P}_{hT}^2) + \mathcal{O}(M^2/Q^2). \end{aligned} \quad (6)$$

Here,  $\mathcal{H}_{UU,T}$  is the hard scattering part;  $f_1^a(x, \mathbf{k}_\perp^2; \mu^2)$  is the TMD distribution of unpolarized partons with flavor  $a$  in an unpolarized proton, carrying longitudinal momentum fraction  $x$  and transverse momentum  $\mathbf{k}_\perp$  at the factorization scale  $\mu^2$ , which in the following we choose to be equal to  $Q^2$ . The  $D_1^{a \rightarrow h}(z, \mathbf{P}_\perp^2; \mu^2)$  is the TMD fragmentation function describing the fragmentation of an unpolarized parton with flavor  $a$  into an unpolarized hadron  $h$  carrying longitudinal momentum fraction  $z$  and transverse momentum  $\mathbf{P}_\perp$ . The term  $Y_{UU,T}$  is introduced to ensure a matching to the perturbative fixed-order calculations at higher transverse momenta.

The applicability of Eq. (6) relies on the possibility of neglecting  $M^2/Q^2$  corrections. At large  $Q^2$  this should not pose serious problems. However, fixed-target DIS experiments typically collect a large amount of data at relatively low  $Q^2$  values, which may lead to problems (see, e.g., recent discussions in Refs. [33? ]).

Eq. (6) can be expanded in powers of  $\alpha_S$ . In the present analysis, we will consider only the leading order terms in  $\alpha_S$ , i.e., stop at order  $\alpha_S^0$ . In this case  $\mathcal{H}_{UU,T}(Q^2, \mu^2) \approx 1$  and  $Y_{UU,T} \approx 0$ . However, perturbative corrections include large logarithms  $L \equiv \log(z^2 Q^2 / P_{hT}^2)$ , so that  $\alpha_S L \approx 1$ . In the present analysis, we will take into account all powers of the form  $\alpha_S^n L^{2n} \approx 1$  (Leading Logarithms –LL) and  $\alpha_S^n L^n \approx 1$  (Next-to-Leading Logarithms – NNL).

In these approximations (LO in  $\alpha_S$  and NLL), only the first term in Eq. (6) is relevant (often in the literature this has been called  $W$  term). We expect this term to provide a good description of the structure function only in the region where  $P_{hT}^2/z^2 \ll Q^2$ . It can happen that  $Y_{UU,T}$ , defined in the standard way (see, e.g., Ref. [26]), gives large contributions also in this region, but it is admissible to redefine it in order to avoid this problem [23]. We leave a detailed treatment of the matching to the high  $P_{hT}^2 \approx Q^2$  region to future investigations.

To the purpose of applying TMD evolution equations, need to calculate the Fourier transform of the the part of Eq. (6) involving TMDs. The structure function thus reduces to

$$F_{UU,T}(x, z, \mathbf{P}_{hT}^2, Q^2) \approx \sum_a \int_0^\infty \frac{d\zeta_T}{2\pi} \zeta_T J_0(\zeta_T |\mathbf{P}_{hT}|/z) \tilde{f}_1^a(x, \zeta_T; Q^2) \tilde{D}_1^{a \rightarrow h}(z, \zeta_T; Q^2).$$

**E' questa la formula usata nel codice, o serve dire altro?** where we introduced the Fourier transforms of the TMD PDF and FF according to

$$\tilde{f}_1^a(x, \zeta_T; \mu^2) = \int_0^\infty \frac{d|\mathbf{k}_\perp|}{2\pi} |\mathbf{k}_\perp| J_0(\zeta_T |\mathbf{k}_\perp|) f_1^a(x, \mathbf{k}_\perp^2; \mu^2), \quad (7)$$

$$\tilde{D}_1^{a \rightarrow h}(z, \zeta_T; \mu^2) = \int_0^\infty \frac{d|\mathbf{P}_\perp|}{2\pi z^2} |\mathbf{P}_\perp| J_0(\zeta_T |\mathbf{P}_\perp|/z) D_1^{a \rightarrow h}(z, \mathbf{P}_\perp^2; \mu^2). \quad (8)$$

### B. Drell–Yan processes

In a Drell–Yan process, two hadrons  $A$  and  $B$  with momenta  $P_A$  and  $P_B$  collide at a center-of-mass energy squared  $s = (P_A + P_B)^2$  and produce a virtual photon or a  $Z$  boson plus hadrons. The boson decays into a lepton-antilepton pair. The reaction formula is

$$A(P_A) + B(P_B) \rightarrow [\gamma^*/Z + X \rightarrow] \ell^+(l) + \ell^-(l') + X. \quad (9)$$

The invariant mass of the virtual photon is  $Q^2 = q^2$  with  $q = l + l'$ . We introduce the rapidity of the virtual photon/ $Z$  boson

$$\eta = \frac{1}{2} \log \left( \frac{q^0 + q_z}{q^0 - q_z} \right). \quad (10)$$

where the  $z$  direction is defined along the momentum of hadron  $A$ .

The cross section can be written in terms of structure functions [34, 35]. For our purposes, we need the unpolarized cross section integrated over  $d\Omega$  and over the azimuthal angle of the virtual photon,

$$\frac{d\sigma}{dQ^2 dq_T^2 d\eta} = \sigma_0^{\gamma, Z} \left( F_{UU}^1 + \frac{1}{2} F_{UU}^2 \right). \quad (11)$$

The elementary cross sections are

$$\sigma_0^\gamma = \frac{4\pi\alpha_{\text{em}}^2}{3Q^2 s}, \quad \sigma_0^Z = \frac{\pi^2\alpha_{\text{em}}}{s(\sin^2\theta_W \cos^2\theta_W)} B_R(Z \rightarrow \ell^+\ell^-) \delta(Q^2 - M_Z^2), \quad (12)$$

where  $\theta_W$  is Weinberg's angle,  $M_Z$  is the mass of the  $Z$  boson, and  $B_R$  is the branching ratio for the  $Z$  boson decay in two leptons. We adopted the narrow-width approximation, i.e., we neglect contributions for  $Q^2 \neq M_Z^2$ . We used the values  $\sin^2\theta_W = 0.2313$ ,  $M_Z = 91.18$  GeV, and  $B_R(Z \rightarrow \ell^+\ell^-) = 3.366$ .

[ Secondo me nella prima delle eq.(10) ci vuole  $\pi^2$ . Infatti, partendo da Ref.[14] abbiamo

$$\frac{d\sigma}{d^4q} = \frac{\alpha^2}{sQ^2} \frac{8\pi}{3} \left[ F_{UU}^1 + \frac{1}{2} F_{UU}^2 \right]$$

Ora  $d^4q = dq^+ dq^- dq_x dq_y$ . Lo Jacobiano della trasformazione  $||dQ^2 d\eta/dq^+ dq^-|| = 2$ . Mentre quello per  $||dq_T^2 d\theta_q/dq_x dq_y|| = 2$ . Quindi

$$\frac{d\sigma}{dQ^2 d\eta dq_T^2 d\theta_q} = \frac{1}{4} \frac{d\sigma}{d^4q} = \frac{\alpha^2}{sQ^2} \frac{2\pi}{3} \left[ F_{UU}^1 + \frac{1}{2} F_{UU}^2 \right]$$

L'ulteriore integrazione in  $d\theta_q$  fornisce un  $2\pi$ , quindi

$$\frac{d\sigma}{dQ^2 d\eta dq_T^2} = \frac{\alpha^2}{sQ^2} \frac{4\pi^2}{3} \left[ F_{UU}^1 + \frac{1}{2} F_{UU}^2 \right]$$

Vi torna? ] [AB: I started from Eq. 57 of Ref. [35] and I integrated over the solid angle. The integration over  $\phi$  is trivial and gives a  $2\pi$ . The integration over  $\cos\theta$  gives a factor  $8/3$  in front of  $F_{UU}^1$  and a factor  $4/3$  in front of  $F_{UU}^2$ . The  $F$  factor in the denominator, neglecting hadron masses, is equal to  $2s$  (see first line of p. 3). Finally, there is the factor 2 in the denominator coming from the first Jacobian of Marco. ]

Similarly to the SIDIS case, in the kinematical limit  $q_T^2 \ll Q^2$  and neglecting the hadron masses the structure function  $F_{UU}^2$  can be neglected.

The longitudinal momentum fractions can be written in terms of rapidity in the following way

$$x_A = \frac{Q}{\sqrt{s}} e^\eta, \quad x_B = \frac{Q}{\sqrt{s}} e^{-\eta}. \quad (13)$$

Some experiments use the variable  $x_F$ , which is connected to the other variables by the following relations

$$\eta = \sinh^{-1} \left( \frac{\sqrt{s}}{Q} \frac{x_F}{2} \right), \quad x_A = \sqrt{\frac{Q^2}{s} + \frac{x_F^2}{4}} + \frac{x_F}{2}, \quad x_B = x_A - x_F. \quad (14)$$

The structure function  $F_{UU}^1$  can be written as

$$\begin{aligned} F_{UU}^1(x_A, x_B, \mathbf{q}_T^2, Q^2) &= \sum_a \mathcal{H}_{UU}^{1a}(Q^2; \mu^2) \\ &\times \int d\mathbf{k}_{\perp A} d\mathbf{k}_{\perp B} f_1^a(x_A, \mathbf{k}_{\perp A}^2; \mu^2) f_1^{\bar{a}}(x_B, \mathbf{k}_{\perp B}^2; \mu^2) \delta(\mathbf{k}_{\perp A} - \mathbf{q}_T + \mathbf{k}_{\perp B}) \\ &+ Y_{UU}^1(Q^2, \mathbf{q}_T^2) + \mathcal{O}(M^2/Q^2). \end{aligned} \quad (15)$$

As in the SIDIS case, with the above kinematical limits the  $Y_{UU}$  term and corrections from higher twists of order  $M^2/Q^2$  or higher can be neglected. In our analysis, we consider the hard coefficients only up to leading order in the couplings, i.e.,

$$\mathcal{H}_{UU,\gamma}^{1a}(Q^2; \mu^2) \approx \frac{e_a^2}{N_c}, \quad \mathcal{H}_{UU,Z}^{1a}(Q^2; \mu^2) \approx \frac{V_a^2 + A_a^2}{N_c}, \quad (16)$$

where<sup>1</sup>

$$V_a = I_{3a} - 2e_a \sin \theta_W, \quad A_a = I_{3a}. \quad (17)$$

The structure function can be conveniently expressed as a Fourier transform of the right-handside of Eq. (15) as

$$F_{UU}^1(x_A, x_B, \mathbf{q}_T^2, Q^2) = \sum_a \mathcal{H}_{UU}^{1a} \int_0^\infty \frac{d\zeta_T}{2\pi} \zeta_T J_0(\zeta_T |\mathbf{q}_T|) \tilde{f}_1^a(x_A, \zeta_T; \mu^2) \tilde{f}_1^{\bar{a}}(x_B, \zeta_T; \mu^2). \quad (18)$$

Stessa osservazione che in SIDIS: è questa la formula usata nel codice, o serve dire altro?

### C. TMDs and their evolution

Following the formalism of Refs. [2, 29], the unpolarized TMD distribution and fragmentation functions in configuration space for a parton flavor  $a$  at a certain scale  $\mu^2$  can be written as

$$\tilde{f}_1^a(x, \zeta_T; \mu^2) = \sum_{i=q,\bar{q},g} (C_{a/i} \otimes f_1^i)(x; \mu_b^2) e^{S(\mu_b^2, \mu^2)} e^{g_K(\zeta_T) \ln(\mu^2/Q_0^2)} \tilde{f}_{\text{1NP}}^a(x, \zeta_T), \quad (19)$$

$$\tilde{D}_1^{a \rightarrow h}(z, \zeta_T; \mu^2) = \sum_{i=q,\bar{q},g} (\hat{C}_{a/i} \otimes D_1^{i \rightarrow h})(z; \mu_b^2) e^{S(\mu_b^2, \mu^2)} e^{g_K(\zeta_T) \ln(\mu^2/Q_0^2)} \tilde{D}_{\text{1NP}}^{a \rightarrow h}(z, \zeta_T). \quad (20)$$

---

<sup>1</sup> We remind the reader that the value of weak isospin  $I_3$  is equal to +1 for  $u, c, t$  and  $-1$  for  $d, s, b$ .

The  $C$  and  $\hat{C}$  are perturbatively calculable Wilson coefficients for the TMD distribution and fragmentation functions, respectively. They are convoluted with the corresponding collinear functions according to

$$(C_{a/i} \otimes f_1^i)(x, \bar{b}_*; \mu_b^2) = \int_x^1 \frac{du}{u} C_{a/i}\left(\frac{x}{u}, \alpha_S(\mu_b^2)\right) f_1^i(u; \mu_b^2), \quad (21)$$

$$(\hat{C}_{a/i} \otimes D_1^{i \rightarrow h})(z, \bar{b}_*; \mu_b^2) = \int_z^1 \frac{du}{u} \hat{C}_{a/i}\left(\frac{z}{u}, \bar{b}_*; \mu_b^2\right) D_1^{i \rightarrow h}(u; \mu_b^2). \quad (22)$$

In the present analysis, we consider only the leading-order term in the  $\alpha_S$  expansion, i.e.,

$$C_{a/i}\left(\frac{x}{u}, \alpha_S(\mu_b^2)\right) \approx \delta_{ai} \delta(1 - x/u), \quad \hat{C}_{a/i}\left(\frac{z}{u}, \alpha_S(\mu_b^2)\right) \approx \delta_{ai} \delta(1 - z/u). \quad (23)$$

As a consequence, the expression for the evolved TMD functions reduces to

$$\tilde{f}_1^a(x, \zeta_T; \mu^2) = f_1^a(x; \mu_b^2) e^{S(\mu_b^2, \mu^2)} e^{g_K(\zeta_T) \ln(\mu^2/Q_0^2)} \tilde{f}_{\text{1NP}}^a(x, \zeta_T), \quad (24)$$

$$\tilde{D}_1^{a \rightarrow h}(z, \zeta_T; \mu^2) = D_1^{a \rightarrow h}(z; \mu_b^2) e^{S(\mu_b^2, \mu^2)} e^{g_K(\zeta_T) \ln(\mu^2/Q_0^2)} \tilde{D}_{\text{1NP}}^{a \rightarrow h}(z, \zeta_T). \quad (25)$$

The Sudakov exponent  $S$  can be written as

$$S(\mu_b^2, \mu^2) = - \int_{\mu_b^2}^{\mu^2} \frac{dk_T^2}{k_T^2} \left[ A(\alpha_s(k_T)) \ln\left(\frac{Q^2}{k_T^2}\right) + B(\alpha_s(k_T)) \right], \quad (26)$$

The convolutions are only valid for small  $\zeta_T \ll 1/\Lambda_{\text{QCD}}$ . At larger  $\zeta_T$ , the TMDs need to match the nonperturbative expressions  $\tilde{f}_{\text{1NP}}^a$  and  $\tilde{D}_{\text{1NP}}^{a \rightarrow h}$ , respectively, that must be constrained by fitting experimental data. The evolution of TMDs from the initial scale  $Q_0$  to  $\mu$  is carried out through perturbatively calculable Sudakov factors  $S$  and  $\hat{S}$ , respectively, and through a nonperturbative universal term  $g_K$  at large  $\zeta_T$  that accounts for the radiation of soft gluons emitted by the considered parton.

The matching between small (perturbative) and large (nonperturbative)  $\zeta_T$  is controlled by the  $\mu_b$  scale, which naturally should be proportional to  $1/\zeta_T$ . We choose

$$\mu_b = \frac{2e^{-\gamma_E}}{\bar{b}_*}, \quad (27)$$

where  $\gamma_E$  is the Euler constant and

$$\bar{b}_* \equiv b_{\text{max}} \left( \frac{1 - e^{-\zeta_T^4/b_{\text{max}}^4}}{1 - e^{-\zeta_T^4/b_{\text{min}}^4}} \right)^{1/4}. \quad (28)$$

This variable replaces the simple dependence upon  $\zeta_T$  in the convolutions of Eqs. (21), (22) and in the perturbative Sudakov factors  $S$  and  $\hat{S}$ ; namely, in the perturbative parts of the TMD definitions of Eqs. (24), (25). In fact, at large  $\zeta_T$  these parts are no longer reliable. Therefore, the  $\bar{b}_*$  is chosen to saturate on the maximum value  $b_{\text{max}}$ , as suggested by the CSS formalism [2, 29].<sup>2</sup> On the other hand, at small  $\zeta_T$  the TMD formalism is not reliable and should be matched to the fixed-order collinear calculations. The way the matching is implemented is arbitrary. In any case, the TMD contribution can be arbitrarily modified at small  $\zeta_T$ . In our approach, we choose to saturate  $b_*$  at the minimum value  $b_{\text{min}} \propto 1/Q$ . With the appropriate choices, for  $\zeta_T = 0$  the Sudakov exponent vanishes, as it should [37?]. Our choice partially corresponds to modifying the resummed logarithms as in Ref. [38] and to other similar modifications proposed in the literature [23?]. One advantage of these kind of prescriptions is that by integrating over the impact parameter  $\zeta_T$ , the collinear expression for the cross section, in terms of collinear PDFs, is recovered, at least at leading order [23].

**AB: I disagree with the following statement** In general, both  $b_{\text{max}}$  and  $b_{\text{min}}$  must not be considered as free parameters; rather, they should be regarded as arbitrary scales separating perturbative from nonperturbative regimes [39]. We choose to fix them to the values

$$b_{\text{max}} = 2e^{-\gamma_E} \text{ GeV}^{-1} = 1.123 \text{ GeV}^{-1}, \quad b_{\text{min}} = 2e^{-\gamma_E}/Q. \quad (29)$$

The motivations are the following:

---

<sup>2</sup> We remind that different schemes are possible to deal with the high- $\zeta_T$  region like the so-called “complex- $b$  prescription” [36].

- with the above choices, the scale  $\mu_b$  is constrained between 1 GeV and  $Q$ , so that the collinear PDFs are never computed at a scale lower than 1 GeV and the lower limit of the integrals contained in the definition of the perturbative Sudakov factor can never become larger than the upper limit;
- at  $Q_0 = 1$  GeV,  $b_{\max} = b_{\min}$  and there are no evolution effects; the TMD is simply given by the corresponding collinear function multiplied by a nonperturbative contribution depending on  $k_{\perp}$  (plus possible corrections of order  $\alpha_S$  from the Wilson coefficients).

Following Refs. [40–42], for the nonperturbative Sudakov factor we make the traditional choice  $g_K(\zeta_T) = -g_2 \zeta_T^2/2$  with  $g_2$  a free parameter. Recently, several alternative forms have been proposed [39, 43] including the suggestion of not including such term [44].

We parametrize the intrinsic nonperturbative parts of the TMDs in the following ways

$$\tilde{f}_{\text{1NP}}^a(x, \zeta_T) = e^{-\langle \mathbf{k}_{\perp a}^2 \rangle \frac{\zeta_T^2}{4}} \left( 1 - \frac{\lambda \langle \mathbf{k}_{\perp a}^2 \rangle^2}{1 + \lambda \langle \mathbf{k}_{\perp a}^2 \rangle} \frac{\zeta_T^2}{4} \right), \quad (30)$$

$$\tilde{D}_{\text{1NP}}^{a \rightarrow h}(z, \zeta_T) = \frac{\langle \mathbf{P}_{\perp a \rightarrow h}^2 \rangle e^{-\langle \mathbf{P}_{\perp a \rightarrow h}^2 \rangle \frac{\zeta_T^2}{4}} + \lambda_F \langle \mathbf{P}_{\perp a \rightarrow h}'^2 \rangle \left( 1 - \langle \mathbf{P}_{\perp a \rightarrow h}'^2 \rangle \frac{\zeta_T^2}{4} \right) e^{-\langle \mathbf{P}_{\perp a \rightarrow h}'^2 \rangle \frac{\zeta_T^2}{4}}}{\langle \mathbf{P}_{\perp a \rightarrow h}^2 \rangle + \lambda_F \langle \mathbf{P}_{\perp a \rightarrow h}'^2 \rangle}. \quad (31)$$

After performing the anti-Fourier transform, the  $f_{\text{1NP}}$  and  $D_{\text{1NP}}$  in momentum space correspond to the normalized linear combination of a Gaussian and a weighted Gaussian:

$$f_{\text{1NP}}^a(x, \mathbf{k}_{\perp}) = \frac{1}{\pi} \frac{(1 + \lambda \mathbf{k}_{\perp}^2)}{\langle \mathbf{k}_{\perp a}^2 \rangle + \lambda \langle \mathbf{k}_{\perp a}^2 \rangle^2} e^{-\frac{\mathbf{k}_{\perp}^2}{\langle \mathbf{k}_{\perp a}^2 \rangle}}, \quad (32)$$

$$D_{\text{1NP}}^{a \rightarrow h}(z, \mathbf{P}_{\perp}) = \frac{1}{\pi} \frac{1}{\langle \mathbf{P}_{\perp a \rightarrow h}^2 \rangle + \lambda_F \langle \mathbf{P}_{\perp a \rightarrow h}'^2 \rangle^2} \left( e^{-\frac{\mathbf{P}_{\perp}^2}{\langle \mathbf{P}_{\perp a \rightarrow h}^2 \rangle}} + \lambda_F \mathbf{P}_{\perp}^2 e^{-\frac{\mathbf{P}_{\perp}^2}{\langle \mathbf{P}_{\perp a \rightarrow h}'^2 \rangle}} \right). \quad (33)$$

Based on the analyses of Refs. [21, 22], we consider that the Gaussian width of the TMD distribution depends on the parton flavor  $a$  and on its fractional momentum  $x$  according to

$$\langle \mathbf{k}_{\perp a}^2 \rangle(x) = \langle \hat{\mathbf{k}}_{\perp a}^2 \rangle \frac{(1-x)^{\alpha} x^{\sigma}}{(1-\hat{x})^{\alpha} \hat{x}^{\sigma}}, \quad (34)$$

where  $\alpha$ ,  $\sigma$ , and  $\langle \hat{\mathbf{k}}_{\perp a}^2 \rangle \equiv \langle \mathbf{k}_{\perp a}^2 \rangle(\hat{x})$  with  $\hat{x} = 0.1$ , are free parameters. Similarly, we have

$$\langle \mathbf{P}_{\perp a \rightarrow h}^2 \rangle(z) = \langle \hat{\mathbf{P}}_{\perp a \rightarrow h}^2 \rangle \frac{(z^{\beta} + \delta)(1-z)^{\gamma}}{(\hat{z}^{\beta} + \delta)(1-\hat{z})^{\gamma}}, \quad (35)$$

where  $\beta$ ,  $\gamma$ ,  $\delta$ , and  $\langle \hat{\mathbf{P}}_{\perp a \rightarrow h}^2 \rangle \equiv \langle \mathbf{P}_{\perp a \rightarrow h}^2 \rangle(\hat{z})$  with  $\hat{z} = 0.5$ , are free parameters.

### III. DATA ANALYSIS

One of the main goals of our fit is to test the universality of TMD parton distribution and fragmentation functions among different processes. To achieve this we included measurements taken from SIDIS, Drell-Yan and  $Z$  boson production from different experimental collaborations. In this chapter we describe the data sets considered for each process and the reasons behind the kinematic cuts applied.

Tab. I refers to the data sets for SIDIS off proton target (HERMES experiment) and presents their kinematic ranges. The same holds for Tab. II, Tab. III, Tab. IV for SIDIS off deuteron (HERMES and COMPASS experiments), Drell-Yan events at low energy and  $Z$  boson production respectively. For each kinematic variable in all the considered data sets, we fit the average value in each bin. **Specify how we treat the nuclear targets, like D= p+n incoherent, then DY ...**

#### A. Hermes data

The semi-inclusive DIS data are taken from HERMES [45] and COMPASS [46] experiments. Both HERMES and COMPASS data have been already analyzed in a previous works [21, 47].

HERMES data [45] are grouped in two sets, distinguished by the inclusion or subtraction of the vector meson contribution. In our work we considered only data set where contributions from vector mesons have been subtracted. The collaboration measured the multiplicities for SIDIS in a fixed-target experiment using hydrogen ( $p$ ) and deuteron ( $D$ ) targets and separating charged pions and kaons produced in the final state. The data include 8 different channels for every combination of target and final-state hadron for a total of 2688 points.

These are grouped in bins of  $(x, z, Q^2, P_{hT})$  with the average values of  $(x, Q^2)$  ranging from about  $(0.04, 1.25 \text{ GeV}^2)$  to  $(0.4, 9.2 \text{ GeV}^2)$ . The collinear energy fraction  $z$  (see (2)) ranges in  $0.1 \leq z \leq 0.9$ . The transverse momentum of the detected hadron lies in  $0.1 \text{ GeV} \leq |P_{hT}| \leq 1 \text{ GeV}$ .

#### B. Compass data

Compass collaboration instead extracted multiplicities of charge-separated but unidentified hadrons produced in SIDIS off a deuteron ( $^6\text{LiD}$ ) target [46]. The data are organised in multidimensional bins,  $(x, z, Q^2, P_{hT})$ . The number of data is an order of magnitude higher compared to the HERMES experiment.

The data cover a range in  $(x, Q^2)$  from  $(0.0052, 1.11 \text{ GeV}^2)$  to  $(0.0932, 7.57 \text{ GeV}^2)$  and the interval  $0.2 \leq z \leq 0.8$ . Similarly to HERMES, for COMPASS  $P_{hT}^2 \lesssim 1 \text{ GeV}^2$ .

To avoid issues related to the normalization of Compass multiplicities (see the *erratum* to [46]), we normalize all the data in  $(x, z, Q^2)$  to the value of the first bin in  $P_{hT}^2$ . We define the *normalized* multiplicity as:

$$m_{\text{norm}}(x, z, \mathbf{P}_{hT}^2, Q^2) = \frac{m_N^h(x, z, \mathbf{P}_{hT}^2, Q^2)}{m_N^h(x, z, \text{Min}[\mathbf{P}_{hT}^2], Q^2)}, \quad (36)$$

where the multiplicity  $m_N^h$  is defined in (3). Fitting normalized multiplicities the first data point of each bin is considered as a fixed parameter and excluded from the degrees of freedom of the system.

The application of the TMD formalism to SIDIS crucially depends on the capability of separating the current fragmentation region from the target fragmentation region and from a soft fragmentation region. This has been recently discussed in [33], where the authors point out the possible overlap among the three fragmentation regions when the hard scale  $Q$  is sufficiently low. In this paper we do not explore this effect and we leave it to future studies. As described in Tabs. I and II, we identify the current fragmentation region operating a cut on  $z$  only, namely  $0.2 < z < 0.7$ .<sup>3</sup>

Another requirement for the applicability of TMD factorization is the presence of two separate hard scales in the process. In SIDIS, those are the  $Q^2$  and  $P_{hT}^2$ , which should satisfy the hierarchy  $\Lambda_{\text{QCD}}^2 \ll P_{hT}^2 \ll Q^2$ . In order to satisfy  $\Lambda_{\text{QCD}}^2 \ll Q^2$ , we request  $Q^2 > 1.4 \text{ GeV}^2$ . The second condition is  $P_{hT}^2 \ll Q^2$ , together with the further constraint  $P_{hT}^2/z^2 \ll Q^2$ . To comply with these, we impose  $P_{hT} < \min[0.2 Q, 0.7 Qz] + 0.5 \text{ GeV}$ . The specific values of the coefficients are chosen to maximize the goodness of the fit procedure. All these choices are summarized in Tabs. I and II.

---

<sup>3</sup> The implementation of the collinearity  $R$  criterion proposed in [33] crucially depends on the value of  $\langle k_T^2 \rangle$ , features that requires independent determinations of the quark properties.



### C. Low-energy Drell-Yan data

We analyze Drell-Yan events collected by fixed-target experiments at low-energy. These data set have been considered also in previous works, e.g. [48]. We use data from E288 [49] measured at  $\sqrt{s} = 19.4, 23.8$  and  $27.4$  GeV<sup>2</sup>, denoted with the label “200”, “300” and “400” respectively. We also include data from E605 [50] at  $\sqrt{s} = 38.8$  GeV<sup>2</sup>.

The explored  $Q$  values are higher compared to the SIDIS case, see Tab. III. E288 provides data at fixed rapidity, whereas E605 explores a range of values for  $x_F$  (see (14)). As discussed for SIDIS data, we can apply TMD factorization if  $\Lambda_{\text{QCD}}^2 \ll q_T^2 \ll Q^2$ , where  $q_T$  is the transverse momentum of the intermediate electroweak boson, reconstructed from the kinematics of the final state leptons. To fulfill this condition, we choose  $q_T < 0.2 Q + 0.5$  GeV. Again, the values of the coefficients are chosen to maximize the goodness of the fit procedure.

### D. Z-boson production data

In order to reach higher  $Q$  and  $q_T$  values, we also consider  $Z$  boson production in collider experiments at Tevatron. We analyze data from CDF and D0, collected during Tevatron Run I [51, 52] at  $\sqrt{s} = 1.8$  TeV and Run II [53, 54] at  $\sqrt{s} = 1.96$  TeV. The invariant mass distribution peaks at the  $Z$ -pole,  $Q = M_Z$ , while the transverse momentum of the exchanged  $Z$  ranges in  $0 < q_T < 20$  GeV. We use the same kinematic condition applied to Drell-yan events:  $q_T < 0.2 Q + 0.5$  GeV = 18.7 GeV, since  $Q$  is fixed to  $M_Z$ .

The observable is  $d\sigma/dq_T$ , apart from the case of D0 Run II, for which the published data refer to  $1/\sigma \times d\sigma/dq_T$ . In order to work with the same observable in all the cases considered, we multiply the D0-Run II data by the total cross section of the process  $\sigma_{exp} = 255.8 \pm 16$  pb [54]. In this case, we add in quadrature the uncertainties of the total cross section and of the published data.

We normalize our functional form with factors listed in Tab. IV. These are the same normalization factors used by the authors in [48] to fit  $Z$  boson production and differ from the experimental ones.

	HERMES $p \rightarrow \pi^+$	HERMES $p \rightarrow \pi^-$	HERMES $p \rightarrow K^+$	HERMES $p \rightarrow K^-$
Reference	[45]			
Cuts	$Q^2 > 1.4 \text{ GeV}^2$ $0.2 < z < 0.7$ $P_{hT} < \text{Min}[0.2 Q, 0.7 Qz] + 0.5 \text{ GeV}$			
Points	190	190	189	187
Max. $Q^2$	9.2 GeV <sup>2</sup>			
$x$ range	$0.06 < x < 0.4$			

TABLE I: Semi-inclusive DIS proton-target data (Hermes experiment).

	HERMES $D \rightarrow \pi^+$	HERMES $D \rightarrow \pi^-$	HERMES $D \rightarrow K^+$	HERMES $D \rightarrow K^-$	COMPASS $D \rightarrow h^+$	COMPASS $D \rightarrow h^-$
Reference	[45]				[46]	
Cuts	$Q^2 > 1.4 \text{ GeV}^2$ $0.2 < z < 0.7$ $P_{hT} < \text{Min}[0.2 Q, 0.7 Qz] + 0.5 \text{ GeV}$					
Points	190	190	189	189	3125	3127
Max. $Q^2$	9.2 GeV <sup>2</sup>				10 GeV <sup>2</sup>	
$x$ range	0.06 < $x$ < 0.4				0.006 < $x$ < 0.12	
Notes					Observable: $m_{\text{norm}}(x, z, \mathbf{P}_{hT}^2, Q^2)$ , eq. (36)	

TABLE II: Semi-inclusive DIS deuteron-target data (Hermes and Compass experiments).

	E288 200	E288 300	E288 400	E605
Reference	[49]	[49]	[49]	[50]
Cuts	$q_T < 0.2 Q + 0.5 \text{ GeV}$			
Points	45	45	78	35
$\sqrt{s}$	19.4 GeV	23.8 GeV	27.4 GeV	38.8 GeV
$Q$ range	4-9 GeV	4-9 GeV	5-9, 11-14 GeV	7-9, 10.5-18 GeV
Kin. var.	$y=0.4$	$y=0.21$	$y=0.03$	$-0.1 < x_F < 0.2$

TABLE III: Low energy Drell-Yan data collected by the E288 and E605 experiments at Tevatron, with different center-of-mass energies.

	CDF Run I	D0 Run I	CDF Run II	D0 Run II
Reference	[51]	[52]	[53]	[54]
Cuts	$q_T < 0.2 Q + 0.5 \text{ GeV} = 18.7 \text{ GeV}$			
Points	31	14	37	8
$\sqrt{s}$	1.8 TeV	1.8 TeV	1.96 TeV	1.96 TeV
Normalization	1.114	0.992	1.049	1.048

TABLE IV:  $Z$  boson production data collected by the CDF and D0 experiments at Tevatron, with different center-of-mass energies.

### E. The replica method

I edited a bit the text, but some overlap remains with the text in [21].

In this section we describe the replica method and we give a definition of the  $\chi^2$  function minimized by the fit procedure. The fit and the error analysis were carried out using a similar Monte Carlo approach as in Ref. [21, 55, 56], and taking inspiration from the work of the Neural-Network PDF (NNPDF) collaboration (see, e.g., [57–59]). The approach consists in creating  $\mathcal{M}$  replicas of the data points. In each replica (denoted by the index  $r$ ), each data point  $i$  is shifted by a Gaussian noise with the same variance as the measurement. Each replica, therefore, represents a possible outcome of an independent experimental measurement, which we denote by  $m_{N,r}^h(x, z, \mathbf{P}_{hT}^2, Q^2)$ . The number of replicas is chosen so that the mean and standard deviation of the set of replicas accurately reproduces the original data points. We see that 200 replicas are sufficient for the purpose.

A minimization procedure is applied to each replica separately, by minimizing the following error function: <sup>4</sup>

$$E_r^2(\{p\}) = \sum_i \frac{\left(m_{N,r}^h(x_i, z_i, \mathbf{P}_{hT}^2, Q_i^2) - m_{N,\text{theo}}^h(x_i, z_i, \mathbf{P}_{hT}^2; \{p\})\right)^2}{\left(\Delta m_{N,\text{stat}}^h + \Delta m_{N,\text{sys}}^h\right)(x_i, z_i, \mathbf{P}_{hT}^2, Q_i^2) + \left(\Delta m_{N,\text{theo}}^h(x_i, z_i, \mathbf{P}_{hT}^2)\right)^2}. \quad (37)$$

The sum runs over the  $i$  experimental points, including all species of targets  $N$  and final-state hadrons  $h$ . In each  $z$  bin for each replica the values of the collinear fragmentation functions  $D_1^{a \rightarrow h}$  are independently modified with a Gaussian noise with standard deviation equal to the theoretical error  $\Delta D_1^{a \rightarrow h}$ . In this work we rely on different parametrizations for  $D_1^{a \rightarrow h}$ : for pions we use the DSEHS analysis [60] at NLO in  $\alpha_s$ ; for kaons we use the DSS parametrization [61] at LO in pQCD. The uncertainties  $\Delta D_1^{a \rightarrow h}$  are estimated from the plots in [62]; they represents the only source of uncertainty in  $\Delta m_{N,\text{theo}}^h$ . Statistical and systematical experimental uncertainties  $\Delta m_{N,\text{stat}}^h$  and  $\Delta m_{N,\text{sys}}^h$  are taken from the experimental collaborations. We do not take into account the covariance among different kinematic bins.

MINUIT minimizes the error function in (37) calculating its gradient with respect to the vector of parameters  $\{p\}$ . The final outcome is a set of  $\mathcal{M}$  different vectors of best-fit parameters,  $\{p_{0r}\}$ ,  $r = 1, \dots, \mathcal{M}$ , with which we can calculate any observable, its mean, and its standard deviation. The distribution of these values needs not to be

<sup>4</sup> Note that the error for each replica is taken to be equal to the error on the original data points. This is consistent with the fact that the variance of the  $\mathcal{M}$  replicas should reproduce the variance of the original data points.

necessarily Gaussian. In fact, in this case the  $1\sigma$  confidence interval is different from the 68% interval. The latter can simply be computed for each experimental point by rejecting the largest and the lowest 16% of the  $\mathcal{M}$  values.

Although the minimization is performed on the function defined in (37), the agreement of the  $\mathcal{M}$  replicas with the original data is better expressed in terms of a  $\chi^2$  function defined as in (37) but with the replacement  $m_{N,r}^h \rightarrow m_N^h$ , i.e., with respect to the original data set. If the model is able to give a good description of the data, the distribution of the  $\mathcal{M}$  values of  $\chi^2/\text{d.o.f.}$  should be peaked around one.

## IV. RESULTS

AS: Check if we are using the last version available of the plots - Filippo ?

In the following we detail the results of a fit to the data sets described in Sec. III with a a flavor-independent configuration for the transverse momentum dependence of unpolarized TMDs In Tab. V we present the total  $\chi^2$ . As usual, the number of degrees of freedom (d.o.f.) is given by the number of data points analyzed reduced by the number of free parameters in the error function. The overall quality of the fit is good, with a global  $\chi^2/\text{d.o.f.} = 1.55 \pm 0.05$ . Uncertainties are computed as the 68% confidence level (C.L.) from the replica methodology. The C.L. is not necessarily Gaussian.

### A. Agreement between data and theory

The partition of the global  $\chi^2$  among SIDIS off proton, SIDIS off deuteron, Drell-Yan and  $Z$  production events is given in Tab. VI, VII, VIII, IX respectively.

#### *Semi-inclusive DIS*

For SIDIS at HERMES off a proton, events with a kaon in the final state have in general a lower  $\chi^2$ . This is due to the large uncertainties for the kaon multiplicities. On the contrary, the major contribution to the  $\chi^2$  comes from events with a  $\pi^+$  in the final state. In [21, 63] a poor agreement between experiment and theory (which relies on the DSS parametrization [61] for collinear FFs) at the level of the collinear multiplicities affected the quality of the fit, especially for  $\pi^\pm$ . Instead, in this work we use a newer parametrization of the collinear FFs (DSEHS [60]), based on a fit which includes HERMES collinear pion multiplicities. This significantly improves the agreement at the collinear level with respect to [21, 63]. The poor  $\chi^2$  for  $\pi^\pm$  production off a proton at HERMES is mainly due to a bad agreement in the TMD multipliities at low  $z$  values (see the first two blocks from the top in Fig. 1). For kaon production off the proton at HERMES the agreement at low  $z$  is better than for the pions (see the first two blocks from the top in Fig. 2), which, combined with larger experimental uncertainties, results in lower  $\chi^2$ .

For SIDIS at HERMES off a deuteron, the situation is slightly different with respect to the proton case. For pion production the  $\chi^2$  is lower with respect to the scattering off a proton because the experimental uncertainties for  $D \rightarrow \pi^\pm$  are slightly larger than for  $p \rightarrow \pi^\pm$  (compare the first two blocks from the top with the last two ones in Fig. 1). On the contrary, for kaon production the  $\chi^2$  is higher with respect to the scattering off a proton because the experimental uncertainties for  $D \rightarrow K^\pm$  are slightly smaller than for  $p \rightarrow K^\pm$  (compare the first two blocks from the top with the last two ones in Fig. 2). SIDIS at COMPASS involve scattering off deuteron only,  $D \rightarrow h^\pm$ , and we identify  $h \equiv \pi$ .

#### *Drell-Yan processes*

Points	Parameters	$\chi^2$	$\chi^2/\text{d.o.f.}$
8059	11	$12629 \pm 363$	$1.55 \pm 0.05$

TABLE V: Total number of points analyzed, number of free parameters, and  $\chi^2$  values.

	HERMES $p \rightarrow \pi^+$	HERMES $p \rightarrow \pi^-$	HERMES $p \rightarrow K^+$	HERMES $p \rightarrow K^-$
Points	190	190	189	187
$\chi^2/\text{points}$	4.83	2.47	0.91	0.82

TABLE VI: Number of points analyzed and  $\chi^2$  values for SIDIS off a proton target.

Description of Hermes data (version from Dropbox, Feb. 9<sup>th</sup>). Legend for  $z$  values needs to be added too.

	HERMES $D \rightarrow \pi^+$	HERMES $D \rightarrow \pi^-$	HERMES $D \rightarrow K^+$	HERMES $D \rightarrow K^-$	COMPASS $D \rightarrow h^+$	COMPASS $D \rightarrow h^-$
Points	190	190	189	189	3125	3127
$\chi^2/\text{points}$	3.46	2.00	1.31	2.54	1.11	1.61

TABLE VII: Number of points analyzed and  $\chi^2$  values for SIDIS off a deuteron target.

	E288 200	E288 300	E288 400	E605
Points	45	45	78	35
$\chi^2/\text{points}$	0.99	0.84	0.32	1.12

TABLE VIII: Number of points analyzed and  $\chi^2$  values for fixed-target Drell-Yan experiments at low energy.

	CDF Run I	D0 Run I	CDF Run II	D0 Run II
Points	31	14	37	8
$\chi^2/\text{points}$	1.36	1.11	2.00	1.73

TABLE IX: Number of points analyzed and  $\chi^2$  values for  $Z$  boson production at Tevatron.

Description of Compass data (version from Dropbox, Feb. 9<sup>th</sup>). Description of low energy Drell-Yan data (version from Dropbox, Feb. 9<sup>th</sup>). Legends need to be added too. Fix the y-axis label. Description of Z-boson production data (version from Dropbox, Feb. 9<sup>th</sup>). Legends need to be added too. Fix the y-axis label.

### B. Transverse momentum dependence at 1 GeV

Tab. X summarizes the values of the nonperturbative parameters  $b_{\min}$  and  $b_{\max}$  (which delimit the range in  $\zeta_T$  where transverse momentum resummation is computed perturbatively) and  $g_2$ , which quantifies the amount of soft gluons radiated. Tab. XI collects the best-fit values for the parametrization of the nonperturbative part of the TMDs (Eqs. (30) and (31)); central values and standard deviations are based on the replica methodology (see Sec. III E), using 68% confidence levels.

We stress here that a prescription involving both  $b_{\min}$  and  $b_{\max}$  is crucial in order to fit SIDIS data with TMD evolution.

Average square transverse momenta and their kinematic dependence. (version from Dropbox, Feb. 9<sup>th</sup>). Full sets of replicas and 68% confidence level bands compared to the results from other fits. Fix some of the labels on the vertical axis.

	$b_{\max}$ [GeV <sup>-1</sup> ] (fixed)	$b_{\min}$ [GeV <sup>-1</sup> ] (fixed)	$g_2$ [GeV <sup>2</sup> ]
All replicas	$2e^{-\gamma_E}/\text{GeV}$	$2e^{-\gamma_E}/Q$	$0.13 \pm 0.01$
Replica 105	$2e^{-\gamma_E}/\text{GeV}$	$2e^{-\gamma_E}/Q$	0.128

TABLE X: Values of parameters common to TMD PDFs and FFs

### C. Comments on flavor-dependent fits

## V. CONCLUSIONS AND OUTLOOK




FIG. 1: Hermes multiplicities for production of pions off a proton and a deuteron for different  $\langle x \rangle$ ,  $\langle z \rangle$ , and  $\langle Q^2 \rangle$  bins as a function of the transverse momentum of the detected hadron  $P_{hT}$ .




FIG. 2: Hermes multiplicities for production of kaons off a proton and a deuteron for different  $\langle x \rangle$ ,  $\langle z \rangle$ , and  $\langle Q^2 \rangle$  bins as a function of the transverse momentum of the detected hadron  $P_{hT}$ .




FIG. 3: Compass multiplicities for production of negative hadrons (pions) off a deuteron for different  $\langle x \rangle$ ,  $\langle z \rangle$ , and  $\langle Q^2 \rangle$  bins as a function of the transverse momentum of the detected hadron  $P_{hT}$ . Multiplicities are normalized to the first bin in  $P_{hT}$  for each  $\langle z \rangle$  value. For clarity, each  $\langle z \rangle$  bin has been shifted by an offset indicated in the legend.

### Acknowledgments

This work is supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No. 647981, 3DSPIN). AS acknowledges support from U.S. Department of Energy contract DE-AC05-06OR23177, under which Jefferson Science Associates, LLC, manages and operates






FIG. 4: Compass multiplicities for production of positive hadrons (pions) off a deuteron for different  $\langle x \rangle$ ,  $\langle z \rangle$ , and  $\langle Q^2 \rangle$  bins as a function of the transverse momentum of the detected hadron  $P_{hT}^2$ . Multiplicities are normalized to the first bin in  $P_{hT}$  for each  $\langle z \rangle$  value. For clarity, each  $\langle z \rangle$  bin has been shifted by an offset indicated in the legend.

Jefferson Lab. The work of AS has been funded partly also by the program of the Stichting voor Fundamenteel Onderzoek der Materie (FOM), which is financially supported by the Nederlandse Organisatie voor Wetenschappelijk




FIG. 5: Drell-Yan differential cross section for different experiments and different values of  $\sqrt{s}$  and for different  $\langle Q \rangle$  bins. For clarity, each  $\langle Q \rangle$  bin has been shifted by an offset indicated in the legend.

TMD PDFs	$\langle \hat{k}_\perp^2 \rangle$ [GeV <sup>2</sup> ]	$\alpha$	$\sigma$		$\lambda$ [GeV <sup>-2</sup> ]	
All replicas	$0.28 \pm 0.06$	$2.95 \pm 0.05$	$0.17 \pm 0.02$		$0.86 \pm 0.78$	
Replica 105	0.285	2.98	0.173		0.39	
TMD FFs	$\langle \hat{P}_\perp^2 \rangle$ [GeV <sup>2</sup> ]	$\beta$	$\delta$	$\gamma$	$\lambda_F$ [GeV <sup>-2</sup> ]	$\langle \hat{P}_\perp^2 \rangle$ [GeV <sup>2</sup> ]
All replicas	$0.21 \pm 0.02$	$1.65 \pm 0.49$	$2.28 \pm 0.46$	$0.14 \pm 0.07$	$5.50 \pm 1.23$	$0.13 \pm 0.01$
Replica 105	0.212	2.10	2.52	0.094	5.29	0.135

TABLE XI: 68% confidence intervals of best-fit parameters for TMD PDFs and FFs

Onderzoek (NWO).

- 
- [1] J. C. Collins, D. E. Soper, and G. Sterman, Adv. Ser. Direct. High Energy Phys. **5**, 1 (1989), hep-ph/0409313.
  - [2] J. Collins, *Foundations of Perturbative QCD*, Cambridge Monographs on Particle Physics, Nuclear Physics and Cosmology (Cambridge University Press, 2011), ISBN 9780521855334, URL <http://books.google.it/books?id=0xGi1KW9vykC>.
  - [3] T. C. Rogers, Eur. Phys. J. **A52**, 153 (2016), 1509.04766.
  - [4] A. Bacchetta, Eur. Phys. J. **A52**, 163 (2016).
  - [5] M. Radici, AIP Conf. Proc. **1735**, 020003 (2016).
  - [6] P. J. Mulders and R. D. Tangerman, Nucl. Phys. **B461**, 197 (1996), erratum-ibid. **B484** (1997) 538, hep-ph/9510301.
  - [7] A. Bacchetta, M. Diehl, K. Goeke, A. Metz, P. J. Mulders, and M. Schlegel, JHEP **02**, 093 (2007), hep-ph/0611265.
  - [8] D. Boer and P. J. Mulders, Phys. Rev. **D57**, 5780 (1998), hep-ph/9711485.
  - [9] A. Bacchetta and P. J. Mulders, Phys. Rev. **D62**, 114004 (2000), hep-ph/0007120.
  - [10] P. J. Mulders and J. Rodrigues, Phys. Rev. **D63**, 094021 (2001), hep-ph/0009343.
  - [11] D. Boer, S. Cotogno, T. van Daal, P. J. Mulders, A. Signori, and Y.-J. Zhou, JHEP **10**, 013 (2016), 1607.01654.




FIG. 6: Cross section differential with respect to the transverse momentum  $q_T$  of a  $Z$  boson produced from  $p\bar{p}$  collisions at Tevatron. The four panels refer to different experiments (CDF and D0) with two different values for the center-of-mass energy ( $\sqrt{s} = 1.8$  TeV and  $\sqrt{s} = 1.96$  TeV).




FIG. 7: write the caption here (a) and another here (b).

- [12] A. Bacchetta and M. Radici, Phys.Rev.Lett. **107**, 212001 (2011), 1107.5755.
- [13] M. Anselmino, M. Boglione, and S. Melis, Phys.Rev. **D86**, 014028 (2012), 1204.1239.
- [14] M. G. Echevarria, A. Idilbi, Z.-B. Kang, and I. Vitev, Phys. Rev. **D89**, 074013 (2014), 1401.5078.
- [15] Z. Lu and I. Schmidt, Phys. Rev. **D81**, 034023 (2010), 0912.2031.
- [16] V. Barone, M. Boglione, J. O. Gonzalez Hernandez, and S. Melis, Phys. Rev. **D91**, 074019 (2015), 1502.04214.
- [17] C. Lefky and A. Prokudin, Phys. Rev. **D91**, 034010 (2015), 1411.0580.
- [18] M. Anselmino, M. Boglione, U. D'Alesio, S. Melis, F. Murgia, et al., Phys.Rev. **D87**, 094019 (2013), 1303.3822.
- [19] Z.-B. Kang, A. Prokudin, P. Sun, and F. Yuan, Phys. Rev. **D93**, 014009 (2016), 1505.05589.
- [20] A. Signori, Ph.D. thesis, Vrije U., Amsterdam (2016), URL <http://inspirehep.net/record/1493030/files/Thesis-2016-Signori.pdf>.




FIG. 8: write the caption here (a) and another here (b).

- [21] A. Signori, A. Bacchetta, M. Radici, and G. Schnell, JHEP **1311**, 194 (2013), 1309.3507.
- [22] A. Bacchetta, M. G. Echevarria, P. J. G. Mulders, M. Radici, and A. Signori, JHEP **11**, 076 (2015), 1508.00402.
- [23] J. Collins, L. Gamberg, A. Prokudin, T. C. Rogers, N. Sato, and B. Wang, Phys. Rev. **D94**, 034014 (2016), 1605.00671.
- [24] A. Bacchetta, D. Boer, M. Diehl, and P. J. Mulders, JHEP **08**, 023 (2008), 0803.0227.
- [25] J. C. Collins and D. E. Soper, Nucl. Phys. **B193**, 381 (1981).
- [26] J. C. Collins, D. E. Soper, and G. Sterman, Nucl. Phys. **B250**, 199 (1985).
- [27] X. Ji and F. Yuan, Phys. Lett. **B543**, 66 (2002), hep-ph/0206057.
- [28] X. Ji, J.-P. Ma, and F. Yuan, Phys. Rev. **D71**, 034005 (2005), hep-ph/0404183.
- [29] S. Aybat and T. C. Rogers, Phys. Rev. **D83**, 114042 (2011), 1101.5057.
- [30] M. G. Echevarria, A. Idilbi, and I. Scimemi, JHEP **1207**, 002 (2012), 1111.4996.
- [31] M. G. Echevarria, A. Idilbi, A. Schfer, and I. Scimemi, Eur.Phys.J. **C73**, 2636 (2013), 1208.1281.
- [32] J. C. Collins and T. C. Rogers, Phys.Rev. **D87**, 034018 (2013), 1210.2100.
- [33] M. Boglione, J. Collins, L. Gamberg, J. O. Gonzalez-Hernandez, T. C. Rogers, and N. Sato, Phys. Lett. **B766**, 245 (2017), 1611.10329.
- [34] D. Boer and W. Vogelsang, Phys. Rev. **D74**, 014004 (2006), hep-ph/0604177.
- [35] S. Arnold, A. Metz, and M. Schlegel, Phys. Rev. **D79**, 034005 (2009), 0809.2262.
- [36] E. Laenen, G. F. Sterman, and W. Vogelsang, Phys. Rev. Lett. **84**, 4296 (2000), hep-ph/0002078.
- [37] G. Parisi and R. Petronzio, Nucl.Phys. **B154**, 427 (1979).
- [38] G. Bozzi, S. Catani, G. Ferrera, D. de Florian, and M. Grazzini, Phys. Lett. **B696**, 207 (2011), 1007.2351.
- [39] J. Collins and T. Rogers, Phys. Rev. **D91**, 074020 (2015), 1412.3820.
- [40] P. M. Nadolsky, D. R. Stump, and C. P. Yuan, Phys. Rev. **D61**, 014003 (2000), hep-ph/9906280.
- [41] F. Landry, R. Brock, P. M. Nadolsky, and C. P. Yuan, Phys. Rev. **D67**, 073016 (2003), hep-ph/0212159.
- [42] A. V. Konychev and P. M. Nadolsky, Phys. Lett. **B633**, 710 (2006), hep-ph/0506225.
- [43] C. A. Aidala, B. Field, L. P. Gamberg, and T. C. Rogers, Phys. Rev. **D89**, 094002 (2014), 1401.2654.
- [44] U. D'Alesio, M. G. Echevarria, S. Melis, and I. Scimemi, JHEP **11**, 098 (2014), 1407.3311.
- [45] A. Airapetian et al. (HERMES Collaboration), Phys. Rev. **D87**, 074029 (2013), 1212.5407.
- [46] C. Adolph et al. (COMPASS), Eur. Phys. J. **C73**, 2531 (2013), [Erratum: Eur. Phys. J.C75,no.2,94(2015)], 1305.7317.
- [47] M. Anselmino, M. Boglione, J. O. G. H., S. Melis, and A. Prokudin (2013), 1312.6261.
- [48] U. D'Alesio, M. G. Echevarria, S. Melis, and I. Scimemi, JHEP **11**, 098 (2014), 1407.3311.
- [49] A. S. Ito et al., Phys. Rev. **D23**, 604 (1981).
- [50] G. Moreno et al., Phys. Rev. **D43**, 2815 (1991).
- [51] T. Affolder et al. (CDF), Phys. Rev. Lett. **84**, 845 (2000), hep-ex/0001021.
- [52] B. Abbott et al. (D0), Phys. Rev. **D61**, 032004 (2000), hep-ex/9907009.
- [53] T. Aaltonen et al. (CDF), Phys. Rev. **D86**, 052010 (2012), 1207.7138.
- [54] V. M. Abazov et al. (D0), Phys. Rev. Lett. **100**, 102002 (2008), 0712.0803.
- [55] A. Bacchetta, A. Courtoy, and M. Radici, JHEP **1303**, 119 (2013), 1212.3568.
- [56] M. Radici, A. Courtoy, A. Bacchetta, and M. Guagnelli, JHEP **05**, 123 (2015), 1503.03495.
- [57] S. Forte, L. Garrido, J. I. Latorre, and A. Piccione, JHEP **0205**, 062 (2002), hep-ph/0204232.
- [58] R. D. Ball et al. (NNPDF Collaboration), Nucl. Phys. **B809**, 1 (2009), 0808.1231.
- [59] R. D. Ball et al., Nucl. Phys. **B838**, 136 (2010), 1002.4407.
- [60] D. de Florian, R. Sassot, M. Epele, R. J. Hernandez-Pinto, and M. Stratmann, Phys. Rev. **D91**, 014035 (2015), 1410.6027.
- [61] D. de Florian, R. Sassot, and M. Stratmann, Phys. Rev. **D75**, 114010 (2007), hep-ph/0703242.
- [62] M. Epele, R. Llubaroff, R. Sassot, and M. Stratmann, Phys. Rev. **D86**, 074028 (2012), 1209.3240.
- [63] A. Signori, A. Bacchetta, and M. Radici, Int. J. Mod. Phys. Conf. Ser. **25**, 1460020 (2014), 1309.5929.