



اللهم انصر اهل غزة وثبت اقدامهم

فهي كالغيث

Quick Review
Keep Calm & Pass Exam

عدنا

بسم الله لنبدء سعادة جديدة

الملف تحديث دوري لإضافة أفكار جديدة وتعديل أي أخطاء

الملف ومجربات الكرام لوجه الله خاصة

1. **A 77 year old man was diagnosed with a right renal tumor size 3.5 cm. he underwent a radical right nephrectomy. Which of following post operative cares is of high priority?**
 - A. Take body temperature
 - B. Measure hourly urine output**
 - C. Provide sips of clear liquid
 - D. Turn client from side to side

2. **A newborn baby is evaluated on the Apgar score immediately after his after his. The score measures the quality of new born pulse rate, reflexes, after the birth?**
 - A. It helps to plan the treatment for congenital diseases
 - B. It serves as permanent record for the newborn babies
 - C. It provides the bases for comparison, a few minutes later**
 - D. It helps identifying the abnormalities related to muscular

3. **A full term male infant was born at 39 weeks gestation and weight 4300 grams by cesarean section 35year old mother with GDM. the mother did not follow the antenatal care schedule until delivery. HR 110 RR 55 TEM 36.8 . which of the following the infant is more liable to immediately after?**
 - A. Hypoinsulinemia
 - B. Hypoglycemia**
 - C. Hyperglycemia
 - D. Hypotonia

4. **A 44 year old obese patient was subject for a surgery called bariatric gastric bypass to specific. The nurse and the physician are giving information about the procedure to the patient. Which of the following statement is the best describing the patient understanding?**
 - A. Same effect with liposuction
 - B. An opening will placed in my medication
 - C. The surgery will reduce the size of my stomach**
 - D. Easiest way to lose weight no need to exercise after surgery

5. **A nurse is caring for a 55 years patient with abdominal pain. On assessment the nurse noticed lower abdominal distention on palpation and was concerned about urinary retention. Which of the following clinical indicators supports the nurses concerns?**
 - A. Hyperthermia
 - B. Urine output of 35 ml/hour
 - C. Imbalanced electrolytes level
 - D. Voiding small amounts of urine at a time**

6. **Nurse is preparing to administer oxygen 4 LPM via nasal cannula complaining of shortness of breath. Which of the following actions will protect the mucosal layer of the respiratory tract?**
 - A. Check for bubbling in the humidifier**
 - B. Change the position of patient frequently
 - C. Stop oxygen after oxygen saturation reaches 98%
 - D. Give oxygen as long as needed as there is no side effect

7. A 66 year old woman with a history of unstable angina and hypertension presents to the emergency department with a dull chest pain that she describes as similar to heartburn. The pain radiates down the left arm. She had taken sublingual nitroglycerin tablets without any relief. An electrocardiograph is performed and shows evaluated S-T segments. The nurse will administer a thrombolytic medication by intravenous infusion. Which of the following factors places this candidate at high risk for bleeding?
- A. Unstable angina
 - B. Hypertension**
 - C. Age
 - D. Elevated S-Segment
8. The nurse is monitoring a client in the Medical- Surgical Unit who is recovering overnight following a femoral angioplasty and observes that his leg has become pale and cool to the touch. After paging the health care provider for an emergency evaluation, the nurse receives an angry telephone call asking why this department always calls at 2 AM with ridiculous questions. Which of the following is the most appropriate response?
- A. "how would you feel if it was your father not getting the medical care he needed"
 - B. "I am worried that this man may lose his leg unless something is done immediately"**
 - C. "your behaviour at this time is extremely unprofessional, and I will be making a formal report to my supervisor"
 - D. I am sorry, but it is hospital policy to report any post-operative complications to the health care provider on call. May I ask what's has been bothering you about this department's pages
9. A patient with long-standing diabetes mellitus type1 (is scheduled for surgical amputation of 4 gangrenous toes on the right foot. Which surgical intervention would this be classified as?
- A. Palliative**
 - B. Curative
 - C. Reconstructive
 - D. Diagnostic
10. Which of the following the priority arrangement for postoperative patient assessment?
- A. Cardiovascular, Respiration, Surgery site, Neurological signs.**
 - B. Respiration, Cardiovascular, Neurological signs, Surgery site
 - C. Neurological signs, Cardiovascular, Respiration, Surgery site
 - D. Surgery site, Respiration, Cardiovascular, Neurological signs
11. A 26 years old married woman is admitted in the plastic surgery for the correction of burn strictures and skin grafting on her neck and face under general anesthesia. While discussing the treatment with her, the plastic surgeon explained that she will have a series of surgeries but she needs to be on family planning until the treatment is completed. The patient asked the nurse whether she will be normal again. What initial assessment is required?
- A. Detailed history and physical examination
 - B. Patient's acceptance for the treatment plan
 - C. Need for psychological support to reduce anxiety**
 - D. Family's involvement and consent for her treatment
12. What is the most appropriate blood test for patient who is going to surgery?
- A. WBC
 - B. Albumin
 - C. Prothrombin time**

- D. Haemoglobin
13. A 38-year-old patient is about to have lumbar disk surgery during preoperative care, the nurse instructs the patient including the family members how to do log rolling to change patient position. One of the family members asks why they must do such action in turning the patient postoperative. Which of the following is the nurse's best response?
- A. Facilitate good circulation
B. Avoid movement
C. **Prevent post-operative bedsores**
D. Makes changing of patient position easier
14. A 46-year-old patient is in the male Urology Ward after the surgical removal of the stone from his left kidney through percutaneous nephrolithotomy under general anesthesia. He has dull aching pain in left lumbar region. His nephrostomy bag is attached through a tube in his left kidney for a few days (see image). What findings should alert the nurse to?
- A. Abdominal discomfort
B. Patient very exhausted
C. Presence of blood and stone gravels in urine
D. **Urine output less than 30 ml/hour**
15. Surgery for pyloric stenosis movement?
- A- Pylorotomy
B- Pylorectomy
C- Pylorostomy
D- **Pyloromyotomy**
16. The nurse is assessing a child (an infant) with pyloric stenosis. Which of the following is likely to note?
- A. Diarrhea
B. **Projectile vomiting**
C. Swallowing difficulties
D. Currant jelly like stool
17. A 5-year-old child was seen to the Emergency Department with abdominal pain, palpable sausage-shaped mass, and intussusception is suspected. Which of the following is the best diagnostic evaluation to?
- A. X-ray
B. Endoscopy
C. Rectal biopsy
D. **Ultrasonography**
18. A male patient admitted to the Medical Unit was diagnosed with deep venous thrombosis complaining of pain on both legs. Which of the following nursing diagnosis most likely describes the problem?
- A. Risk for injury
B. Fluid volume excess
C. Electrolyte imbalance
D. **Impaired tissue perfusion**
19. The nurse stated to a psychiatric patient "I will stay here with you". Which of the following therapeutic communication techniques that the nurse used?
- A. Accepting
B. Exploring

C. Offering self

D. Reflecting

20. **Psychiatric patient with depression. The doctor order antidepressant with low side effect. Which of the following is the most appropriate type of antidepressant?**

- A. Anti depressant nor Adrenaline
- B. Monoamine oxidase inhibitors (MAOIs)

C. Selective serotonin reuptake inhibitors (SSRIs)

21. **The nurse manager received complaints from some nursing to improper work distribution and discrimination How should the nurse manager handle the situation?**

A. Investigate the complaints

- B. Individual counselling
- C. Identify list of problems
- D. Continue to observe

22. **Patient have depression, physician advice to procedure ECT. What is this procedure?**

- A. Involves a brief electrical stimulation of the brain while the patient is under anaesthesia**
- B. A Medical procedure that involves passing a mild electric current through your heart.

23. **A patient with symptoms including tingling in the fingers this question will be the result of any deficiency?**

- A. Sodium
- B. Potassium
- C. Magnesium

D. Calcium

24. **The child had a cleft lip repaired and is currently in elbow restraints. What should the nurse do?**

- A. Remove the restrain every 2 hours
- B. Remove the restrain when the parents are with the child**
- C. Put mittens

25. **The doctor ordered a blood sample from the patient. A nurse wants to insert an IV line for the patient after selecting a vein for IV-line placement. What is the next step they should take?**

- A. Wipe the vein with alcohol swap
- B. Apply tourniquet above vein
- C. Flatten the palpation/ palpate the vein**
- D. Needle insertion

26. **While inserting an NGT (Nasogastric Tube) for a patient before abdominal surgery, which of the following should prompt the insertion to be stopped?**

- A. Saliva
- B. Discomfort
- C. Tearing eye
- D. Inability speaking**

27. **A 10 years old child had an appendectomy yesterday. Which pain scale should the nurse use to assess child pain?**

- A. Numeric

B. Faces

- C. FLACC
- D. Cries.

28. **What is the best type of evidence-based practice for teaching diabetic patients?**

- A. Randomize control study
- B. Comparison of diabetic teaching types

C. Systematic review of randomize control

- D. Study about diabetic teaching type.

29. **A teenager girl cannot administer insulin to herself. What should a nurse do for the patient?**

- A. I make her come back to the hospital every few weeks

B. I teach the patient

- C. I teach her parents

30. **A woman came to the clinic with her husband for long-term family planning and contraception. What advice should the physician give them?**

A. Copper IUD

- B. Hormonal IUD
- C. Combined

31. **Pancreatic cancer is life threatening disease, there is low chance of survive if diagnosed in end stage , only 1% can survive for 2 year , what is best describe of mortality rate?**

- A. Incidence rate higher than mortality rate

B. mortality rate higher than incidence rate

- C. incidence rate not related to mortality rate
- D. Mortality rate and incidence rate are approximately the same

32. **From where is human chorionic gonadotropin (HCG) secreted in early pregnancy?**

A. Embryo

- B. Corpus luteum
- C. Ovary
- D. Pituitary gland

33. **What does the blastocyst secrete?**

A. HCG

- B. Estrogen
- C. ADH
- D. .Progesterone

34. **A patient 4 years old needs to receive a doctor-prescribed medication at a dosage of 10mg/kg/day for one week. The medication is available in a concentration of 125mg/5ml. The patient's body weight is 15 kg. How many ml per day should be given to the patient?**

- A. 2 ml QID

B. 3 ml Bid

- C. 5 ml OD
- D. 6 ml PRN

35. **The vegetarian people that not eat red meat. Which of deficiency is high risk?**

- A. Vitamin D

- B. Vitamin C
- C. Folic Acid
- D. Riboflavin**

36. A nurse receive pt one day post operative she expected to give antibiotic But she didn't find any antibiotic for the pt written in medication record after reviewing Doctor order she found antibiotic in order and the another nurse not give it. And the pt missed 3 doses. What is the appropriate measure the nurse do to prevent harmful from the nurse malpractice ?
- A. Ask Physician to make another prescription
 - B. Complain incident report**
 - C. Write the rational in pt record
37. A 36-weeks pregnant woman is experiencing irregular contractions and has a closed cervix. What would be the appropriate advice during contractions?
- A. Drinking warm milk
 - B. Encourage in regular walking**
 - C. Monitoring fetal heart movement
38. The nurse is preparing to transfer a 55-year-old patient on a the CT scan unit. The patient is too heavy for the nurse to transport, nurse went to get an assistive device to transfer the patient. Which of the following transfer device is the most appropriate for the nurse to use?
- A. Board
 - B. Handle
 - C. Trapeze
 - D. Mechanical lift**
39. How can In Vitro Fertilization Happen?
- A. A woman egg cells are combined with man sperms inside uterus and fertilized egg and implant in the uterus
 - B. A woman's egg cells are combined with a man's sperm cells outside the uterus. The fertilized egg is the implanted in the woman's uterus**
40. Death rate of disease in proportion to those who are at high risk to this disease?
- A. Specific Mortality
 - B. Case Fatality**
 - C. Specific Morbidity
41. What is the complication of thalassemia?
- A. Hepatomegaly**
 - B. Bleeding
 - C. Heart Failure
42. A 5-year-old has been diagnosed with Wilms tumor. The doctor discussed the condition and available treatment options with the parents and provided them with an appointment to make a decision. What is the description of this action?
- A. Inspiring
 - B. Empower**
 - C. Enabling
 - D. Encouraging

43. A nurse is teaching a 13- years- old patient how to give an insulin injection. What is the description of the nurse's action?
- A. Inspiring
 - B. **Empower**
 - C. Enabling
 - D. Encouraging
44. Hyperemesis gravidarum is a common occurrence for pregnant women, and there are home remedies that have been proven to relieve nausea and vomiting. Which one of the following is one of them?
- A. Cucumber
 - B. **Ginger**
 - C. Coffee
 - D. Cinnamon
45. What is the diet for a patient after cholecystectomy?
- A. **Grapes and gelatin**
 - B. Cola and macaroni
 - C. Cheese cake and yogurt
46. The infant with congenital diaphragmatic hernia, who has been admitted to the NICU, what intervention should be done first?
- A. **Mechanical ventilation and intubation**
 - B. Feeding via NGT (nasogastric tube).
47. The older woman, who hasn't been doing BSE (Breast Self Examination) for 7 years, came to the clinic. What advice should the nurse give her regarding performing BSE? How frequently should she perform it?
- A. **Monthly**
 - B. Annually
 - C. Quarterly
 - D. Biannual
48. The patient is bedridden and not allowed to walk. How can the nurse measure the weight of a bedridden patient?
- A. **Hoist scale**
 - B. -Sitting scale
 - C. Don't do weight measure
 - D. Standing scale
49. 25-year-old woman has a family history of breast cancer. The nurse reviews the procedure for breast self-examination (BSE) and tells her that the best time for a woman to perform a breast self-examination is:
- A. a few days before her period.
 - B. during her menstrual period.
 - C. on the last day of menstrual flow.
 - D. **3- 7days after the beginning of her period**
50. The nurse is instructing a female client how to do breast self exam. Which of the following is the best time to perform this exam?

- A. After ovulation
 - B. After period**
 - C. Two weeks after period
 - D. Three days before period
51. A patient with renal calculi is undergoing a percutaneous nephrostomy and asks the nurse to explain the procedure. What should the nurse's answer be?
- A. Administering a chemical agent to dissolve the stone.
 - B. Using laser waves
 - C. Incision in renal pelvic to remove stone
 - D. Using an incision in the skin with a scope and forceps to remove stones**
52. A 36 weeks pregnant woman went into labor and said that she uses cocaine. Who should the nurse report it to?
- A. Obstetric doctor**
 - B. Hospital police
 - C. Not tell anyone.
 - D. Husband or legal guardian
53. A newborn presented with mild fever cough, poor feeding, sneezing, crying and appears restless. What advice did the nurse give to the mother?
- A. Amoxicillin for 3 days
 - B. Antiviral for a week
 - C. Give antipyretic for mild fever and discomfort**
 - D. Cough depressant
54. The doctor is conducting an assessment while the patient is uncomfortable and asks, What should the nurse do?
- A. Cover the unassessed area**
 - B. Stop the assessment and write a report
 - C. Ignore the patient
 - D. Complete the assessment
55. Lanugo hair found on?
- A. Back**
 - B. holder
 - C. forehead
56. A patient has a low-grade fever and is shivering What is appropriate action?
- A. Analgesic
 - B. Antipyretic**
 - C. Antibiotic
57. From where is human chorionic gonadotropin (HCG) secreted in early pregnancy?
- A. Embryo**
 - B. Corpus luteum
 - C. Ovary
 - D. Pituitary gland

58. What does the blastocyst secrete?
A. **HCG**
B. Estrogen
C. ADH
D. Progesterone
59. Every patient complain from chest pain came to ER the doctors order Troponin level, This considering as?
A. Overuse
B. **Misuse**
C. Negligence
60. Patient with measles (rubella) is in airborne precaution nurse should - be were ?
A. Gown
B. Gloves
C. **Face mask**
D. Face Shilded
61. Researcher nurse give all participants ID band and matching them with personal information?
A. A-Justice
B. B-Beneficence
C. C-Autonomy
D. **Confidentially**
62. What is the first principle that empowers nurses' decision-making?
A. Justice
B. **Autonomy**
C. Beneficence
D. Fidelity
63. What is the first ethical principle should the nurse attention when make NCP?
A. Justice
B. Non-maleficence
C. Beneficence
D. Autonomy
64. A nurse forced a patient who wished to take a one-day postoperative rest to take a walk. Which ethical principle did the nurse violate?
A. Justice
B. Beneficence
C. **Autonomy**
D. Nonmaleficence
65. A patient with symptoms including tingling in the fingers this question will be the result of any deficiency?
A. Sodium
B. Potassium
C. Magnesium
D. **Calcium**

66. The child had a cleft lip repaired and is currently in elbow restraints. What should the nurse do?
- A. Remove the restrain every 2 hours
 - B. Remove the restrain when the parents are with the child**
 - C. Put mittens.
67. The doctor ordered a blood sample from the patient. A nurse wants to insert an IV line for the patient after selecting a vein for IV-line placement. What is the next step they should take?
- A. Wipe the vein with alcohol swap
 - B. Apply tourniquet above vein
 - C. Flatten the palpation/ palpate the vein**
 - D. Needle insertion
68. What is the best type of evidence-based practice for teaching diabetic patients?
- A. Randomize control study
 - B. Comparison of diabetic teaching types
 - C. Systematic review of randomize control**
 - D. Study about diabetic teaching type.
69. A teenager girl cannot administer insulin to herself. What should a nurse do for the patient?
- A. I make her come back to the hospital every few weeks
 - B. I teach the patient**
 - C. I teach her parents
70. A 10-year-old girl presents to the Emergency Room (ER) with pain. On assessment, you noticed that when you palpate the left lower quadrant of the child's abdomen, the child feels pain in the right quadrant. Which of the following is the name of this signs?
- A. Rebound tenderness
 - B. McBurney sign
 - C. Roving's sign**
 - D. Obdurat signs
71. Ahmed has as argument with his boss and yells at his kids when he gets home. Which defence mechanism is Ahmed displaying?
- A. Displacement**
 - B. Reaction formation
 - C. Rationalization
 - D. Repression
72. In the year 1930, children had developed immunity against measles without being vaccinated, and as a result, the infection rate among children decreased. What is type of immunity?
- A. Artificial
 - B. Passive
 - C. Herd
 - D. Innate**
73. The patient with Parkinsonism is taking an antibiotic. The doctor changed the dose of his medication. The next day, the patient doesn't move and feels freezing. What can be expected for the patient?
- A. Acute akinesia**
 - B. Neuroleptic malignance syndrome

74. Before administering NGT feeding, what should the nurse do?
- Aspirate 30 cc to check for residual feeding from the last time
 - Check the consistency of the food
 - Ensure the NGT is correctly placed**
75. A nurse performing nursing care plan for a neonate after a birth, which intervention has the highest nursing priority neonate?
- Obtained a dextrostix
 - Give the initial bath
 - Give the vitamin k injection
 - Cover the neonate head with a cap**
76. A primigravida client at 25 weeks gestation visits the clinic and tells the nurse that her lower back aches when she arrives home from work. The nurse should suggest that the client perform?
- Tailor sitting**
 - Leg lifting
 - Shoulder circling
 - Squatting exercises
77. A nurse checked medication. What is the medication nor refill new prescription?
- Diazepam
 - Digoxin
 - Ibuprofen**
 - Tetracycline.
78. Neonate after one day of delivery diagnosed with tracheoesophageal fistula / atresia. What should the nurse observe for neonate?
- continuous crying**
 - Projectile Vomiting
79. Patient came with burn dorsal left arm and dorsal of both leg and his weight 62 kg total intravenous fluid required ?
- 5580**
 - 6000
 - 6500
 - 7000
80. What is the medical abbreviation used for AT bedtime?
- HS**
 - Prn
 - Stat
81. Patient came to ER after accident the vital signs all stable and conscious but expected has increased the pressure (ICP). Which of the following first indicate ICP?
- Change in consciousness (loss of consciousness)**

Handwritten calculation for Parkland formula:

$$TBSA = 9 + 9 + 4.5 = 22.5$$

$$\text{Parkland Formula} = 4ml \times TBSA \times \text{weight}$$

$$= 4 \times 22.5 \times 62 = 5580$$

- B. Unreactive pupil
C. Bradycardia
D. Depress respiratory
82. A 2-year-old child is admitted to pediatric ward. The mother cannot stay the child and she will visit him in the weakened only. Which of the following nursing action indicates an understanding of the emotional needs this child?
A. Give her a warm bath to calm down
B. Allow the child to suck on a pacifier
C. Ask the parents to bring the child favorite toy
D. Tell he parents that frequently visiting is unnecessary
83. A nurse is explaining to the nursing students working on the antepartum unit how to assess edema. Which edema assessment score indicates edema of the lower extremities, face, hands, and sacral area?
A. +1
B. +2
C. +3
D. +4
84. A 36-weeks pregnant woman is experiencing irregular contractions and has a closed cervix. What would be the appropriate advice during contractions?
A. Drinking warm milk
B. Encourage in regular walking
C. Monitoring fetal heart movement
85. Which month does the government provide vaccinations for hepatitis B to children?
A. 0,2,4
B. 0,3,6
C. 0,2,3
86. 3-week-old newborn is diagnosed with Hirschsprung disease Which of the following preoperative care should included in the plan of care
A. Restricting oral intake to clear fluids
B. Administering a tap water enema
C. Inserting a gastrostomy tube
D. Using povidone-iodine to prepare the perineum
87. A researcher want to conduct a research for measles vaccine at 1 year of age then will remeasure again at age of 6 years in the relation to healthy child. What is the type of research?
A. Longitudinal/correlation.
B. Longitudinal/ Cross sectional
C. Cross sectional/correlation
88. Female patient has been advised that laboratory tests confirm genital warts and recurrent many times. The nurse should teach the patient that a Papanicolaou test (Pap smear) is recommended?
A. Once Every 6 months
B. Once Every year
C. Once Every 3 years.

- ❖ For patients with genital warts or a history of recurrent HPV infections, annual Pap smears are recommended. This is because they are at higher risk for cervical dysplasia and cervical cancer.
- ❖ Routine screening for average-risk women typically starts every 3 years (age 21–29), but:
- ❖ High-risk patients (e.g., HPV-positive or recurrent genital warts) need more frequent monitoring, usually annually.

89. A newborn having meconium aspiration. What are the appropriate diagnostic evaluation tests?

- A. Chest-Xray
- B. Blood gas
- C. Ultrasound

90. A nurse is assigned to care for a group of patient in the medication expected to review the medical records of these patient. What patient is at risk for excess fluid volume?

- A. Patient with ileostomy
- B. Patient taking a loop diuretic
- C. Patient with chronic renal failure
- D. Patient hooked to gastrointestinal

91. A nurse is caring for a 72-year-old man patient , is unsteady. The patient requests the nurse to help nurse asked the patient to wait few minutes device to transfer him. Which of the following transfer devices is the nurse to use?

- A. Belt
- B. Board
- C. Handle
- D. Mechanical lift

92. A 40 year old man was admitted to a male medical department with pneumonia. A nurse in the department explained and obtained informed consent for a chest CT scan procedure with contrast. with of the following ethical principles underpinning informed consent?

- A. Autonomy
- B. Non maleficence
- C. Beneficence
- D. Respect

93. A neonatal nurse performs Apgar assessment at 1 minute of birth to evaluate the physical condition of the newborn and immediate need for resuscitation. At 1 minute, Apgar score is 7. At 5 minutes Apgar score is to the progression of scores suggests

- A. A healthy newborn
- B. The need for supplement oxygen
- C. A genetic defect
- D. The infant is becoming stable

94. Which of the following classifications of placenta previa is applicable when the placental edge is 5 cm away from the internal cervical OS?

- A. Total
- B. Partial
- C. Marginal
- D. Complete

95. Gestational diabetes is high blood sugar (diabetes) that starts or is found during pregnancy. The best time for glucose screening tests during pregnancy is:
- 7 – 12 weeks
 - 17 – 18 weeks
 - 24 – 28 weeks**
 - 20 – 35 weeks
96. Assessment of never 9,10 IX , X tools?
- Tuning Fork
 - Pin lights and tongue depressor never XV for Glossopharyngeal , X for Vagus**
97. The antenatal clinic nurse was assessing a 32yaers old gravida 2, para1 pregnant mother on the fundal height at 36 weeks. What is the expected fundus position?
- Umbilicus
 - Xiphoid process**
 - Symphysis pubis
98. A psychiatric patient received an overdose of benzodiazepine. The doctor prescribed an antidote. Which adverse effect of flumazenil should you check before administration?
- Anxiety
 - Seizure**
 - Shivering
 - Chest pain
99. A nurse is caring for an immobile and ventilated patient in the intensive care unit. The patient is at high risk for pressure ulcers. Which intervention should the nurse prioritize to prevent pressure ulcers in this patient?
- Keep the patient's skin dry
 - Keep the head of the bed at 40°
 - Massage the reddened prominent bones
 - Keep the head of the bed at 30° while in a side-lying position**
100. Patient changed his mind after informed research onset. What principle is linked to this patient right?
- Justice
 - Competence
 - Autonomy**
 - Beneficence
101. The Elderly patient, 80 years old, underwent a nurse assessment of activities to measure their ability to perform self-care tasks such as bathing and eating. What type of assessment is the nurse conducting?
- Psychosocial
 - Economic
 - Functional**
 - Psychological
102. A 40 years old female is diagnosed with chlamydia with no signs and symptoms?



A. Subclinical stage

- B. Clinical stage
- C. Susceptibility



103. What Anthropometric measurements are included?

- A. Weight and Wrist
- B. BMI, Wrist and Arm

C. BMI, Wrist and Skinfold thickness



104. A 33-year-old woman has come to the outpatient clinic for treatment of a vaginal infection. Physical assessment reveals yellowish excessive, thin offensive and frothy discharge. Which of the following is the most likely diagnosis?

- A. Candidiasis
- B. Trichomoniasis**
- C. Bacterial vaginosis
- D. Chlamydia

105. 8 month-old child diagnosed with bronchiolitis due to respiratory ncytial virus (RSV) is admitted to the paediatric hospital. The nurse should initiate which of the following isolation precaution?

- A. Contact**
- B. Airborne
- C. Standard
- D. Droplet

106. The nurse begins a morning shift with the following see the doctor in the waiting room (see table)?

- A. 1.A first time mother who delivered two days ago and is having Difficulties breast feeding .
- B. A60 year-old man who has a history of deep vein thrombosis and is taking daily heparin .
- C. A76 year-old women who was hospitalized three days Previously with pneumonia .
- D. A 56-year-old man who has used all his diuretic pills Coughing up sputum .

Which patient should be seen first?

- A. One
- B. Two
- C. Three
- D. Four**

107. A woman patient is admitted for abdominal pain. She complain generalized plan nausea vomiting and constipation rebound tenderness and abdominal rigidity the past hour her localized on right side BP 130/68 HR 92 RR 18 TEM 38.6 What should the nurse suspect?

- A. Intestinal obstruction
- B. Influenza
- C. Appendicitis**
- D. Pyloric stenosis

108. A patient 4 years old needs to receive a doctor-prescribed medication at a dosage of 150 mg for one week. The medication is available in a concentration of 125mg/5ml. How many ml per day should be given to the patient?

A. 3 ml BID

- B. 3 ml QID
- C. 6 ml PRN

109. **Hyperemesis gravidarum is a common occurrence for pregnant women, and there are home remedies that have been proven to relieve nausea and vomiting. Which one of the following is one of them?**
- A. Cucumber
 - B. Ginger**
 - C. Coffee
 - D. Cinnamon
110. **What is the diet for a patient after cholecystectomy?**
- A. Grapes and gelatin**
 - B. Cola and macaroni
 - C. Cheese cake and yogurt
111. **The patient reported having pain and taking morphine, but the pain increased, leading the patient to request an increase in the dose. What is the patient experiencing?**
- A. Drug addiction
 - B. Drug tolerance**
112. **A nurse in the emergency department receives a report that a man has arrived, screaming and yelling that his wife is bleeding in the car. What is the nurse's first action?**
- A. Provide emotional support and reassurance to the husband / Calm the patient
 - B. Triage the wife**
 - C. Notify the attending physician
 - D. Gather a team to prepare instruments for examination
113. **Where is the best place to locate systematic reviews? Systemic reviews, which found located?**
- A. PubMed
 - B. Cochrane library**
 - C. MEDLIN
 - D. CINAHL
114. **The Wassermann test should be done before maternity for the early identification of which of the following?**
- A. Hepatitis C
 - B. HIV
 - C. Syphilis**
 - D. Toxicology
115. **incidence rate for disease still at some rate for 10 year but now prevalence more increase? Disease prevalence higher in 5y old children more than 10 year old?**
- A. Incident rate higher in younger children
 - B. Incident rate higher in older children
 - C. Fatality rate increase in older children
 - D. Duration longer**

116. You have just admitted a patient with bacterial meningitis to the medical-surgical unit. The patient complains of a severe headache with photophobia and has a temperature of 102.60 F orally. Which type of isolation should be suitable for the patient?
- A. Air borne
 - B. Contact
 - C. Droplet
117. A postpartum woman came to the clinic with her baby. The nurse assessed the umbilical cord. What are the signs of cord infection?
- A. Reddened base
 - B. Bluish white
 - C. Black and dry
 - D. Dry
118. Mother of nine children, three of them with congenital anomalies down syndrome; she is a primary school graduate, with low first status. She is not using any method of family planning. According primary health care nurse has referred her counselling. Which of the following phases of home visit accomplishes intervention?
- A. Initial
 - B. Closing
 - C. Action
 - D. Terminal
119. The patient weight 67 kg. He is receiving drug 30mg/kg/24hr. How many grams should the nurse give for 24 hr?
- A. 5 g
 - B. 1.5 g
 - C. 1 g
 - D. 2 g
120. Researcher nurse give all participants ID band and matching them with personal information?
- A. A-Justice
 - B. B-Beneficence
 - C. C-Autonomy
 - D. Confidentially
121. Hyperemesis gravidarum is a common occurrence for pregnant women, and there are home remedies that have been proven to relieve nausea and vomiting. Which one of the following is one of them?
- A. Cucumber
 - B. Ginger
 - C. Coffee
 - D. Cinnamon
122. What is the diet for a patient after cholecystectomy?
- A. Grapes and gelatin
 - B. Cola and macaroni
 - C. Cheese cake and yogurt

123. Patient with colostomy we transfer the patient
A. Belt abdomen
 B. Belt chest
 C. Restrain
124. A 4 month old infant returned immediately from OR room post cleft lip repair which of the following nursing intervention should be considered?
A. Apply elbow restrain
 B. Apply suction when needed
 C. Measure temperature
 D. Put infant in prone position
125. A 22-year-old gravida 2 para 1 with gestational age 38 week admitted to the hospital. The chief complaint is decreased the fetal non-stress test revealed decreased variability and fetal movement. The next morning as part of the antenatal the nurse checks the fetal heart rate by Doppler Sonicaid decreased the fetal heart rate to less than 100 /min. which of the following action the nurse should do first?
 A. Reassure the mother that the FHR is Ok
B. Notify immediately the physician or midwife
 C. Reposition the patient to left lateral position
 D. Ask the mother about the pattern of fetal movement
126. Mother of nine children, three of them with congenital anomalies Down syndrome; she is a primary school graduate, with low status. She is not using any method of family planning. According primary health care nurse referred her for counseling Which of the following types of home visits that the community her nurse should conduct for t client?
 A. Systematic routine
B. Selective
 C. Follow up
 D. Field trip
127. 6month-old boy with hydrocephalus is admitted to the paediatric surgical Ward for ventriculoperitoneal Shunt (VPS) insertion. Which of the following findings should be of the most concern when assessing the child postoperative?
 A. Sunken fontanelle and irritability
 B. decreased head circumference
C. poor feeding and pupillary change
 D. headache and excessive sleepiness
128. A 65-year-old women visited the gynaecological outpatient history reveals that she had 3 pregnancies, one abortion gestational age, had 2 normal deliveries. She smokes 20 Her complaint is that she wets herself when she cough embarrassing for her. Which of the following can be considered as risk factors pelvic floor muscles?
A. Chronic coughing
 B. Diabetes mellitus
 C. Excessive spot
 D. Sedentary life style

129. Nurse plan community program to these who experienced a disease and assess the recovery or rehabilitation or disability . which level of prevention?
- A. Primary
 - B. Secondary
 - C. Tertiary
130. What is an example of secondary level of prevention?
- A. A-How to be smoking cessation .
 - B. B-Given education on HIV to high school students
 - C. Nurse Combine to medication
131. What is an example of primary level of prevention?
- A. B-Given education on HIV to high school students
132. A 29 year old woman had been diagnosed with a 3 cm ovarian cyst. Which of the following is the appropriate step in management?
- A. Cyst aspiration
 - B. Hormonal therapy
 - C. Cyst removal by laparoscopy
 - D. Examination after next menstruation
133. 133- Nurse is preparing to insert a nasogastric tube into a client. The nurse places the client in which position for insertion?
- A. Right side
 - B. Low Fowler's
 - C. High Fowler's
 - D. Supine with the head flat
134. On orientation day, a nurse educator said to a new nurse" our hospital provides the highest level of specialized healthcare in an integrated educational and research setting" which of the following aspects of the hospital's strategic plan is indicated by this statement?
- A. Vision
 - B. Policy
 - C. Values
 - D. Mission
135. Float nurse from medicine floor. Came to telemetry unite Which patient she receives?
- A. MI patient in heparin infusion
 - B. Hypertensive patient on Lasix.
 - C. Atrial Fibrational with anticoagulant drug.
136. Cause of death in Saudia Arabia?
- A. Ischemic heart disease and road accident
137. Diphtheria which type of isolation?
- A. Droplet
 - B. Airborne

138. Which of the following statement is the best description for permanent impairment?
- A. A-Problems in body function and loss
 - B. B-Difficulties to experience life situations
 - C. C-Difficulties in executing activities
 - D. D-Loss of interaction with community
139. A nurse is taking care of a patient who underwent abdominal surgery three days ago. The patient has not been able to breathe deeply and uses to get out of bed since the surgery due to pain. Also, the patient complains of shortness of breath, and the lung sounds are diminished on auscultation. Blood pressure 120/70 mmHg Heart rate 75 /mi Respiratory rate 22 /mi Temperature 36.4 °C Oxygen saturation 89% Which of the following conditions should the nurse suspect?
- A. Sepsis
 - B. Atelectasis
 - C. Congestive heart failure
 - D. Emphysema
140. An autistic child makes no eye contact, unresponsive to continuously spins, twist and head bang. Which of the following is the priority nursing diagnosis?
- A. Risk of injury related to head banging
 - B. Impaired verbal communication related to physical
 - C. Personal identity disorder related to poor ego develop
 - D. Social isolation related to unresponsiveness toward
141. Stages of grief after death The five stages,
- A. Denial, Anger, Bargaining, Depression and Acceptance (DABDA)
142. What is convenient contraceptive for mother who breastfeeding?
- A. high estrogen
 - B. progestin only "mini" pill
 - C. combination ...
143. A patient returns to a clinic for a follow-up visit and is diagnosed as positive for Human Immuno deficiency Virus (HIV). The patient expresses fear related to lack of finances, fear of social avoidance, and hopelessness. Which of the following nursing action should
- A. Referral to a physician infectious disease specialist
 - B. Referral to a community-based HIV clinic
 - C. Referral to the local public health department
 - D. Recommendation to disclose the diagnosis to family
144. A 2-months old infant was brought to the health center for immunization, during assessment the infant temperature registered at 38.1°C. Which is the best course of action that you will take?
- A. Go on with the infant immunization
 - B. Give paracetamol and wait for his fever to subside
 - C. Refer the infant to the physician for further assessment
 - D. Advise the infant mother to bring him back for immunization when he is well
145. Mr X attended in outpatient clinic with symptoms of shortness of breath, diarrhea and severe respiratory distress Which of the following is the best diagnosis of Mrs. A case?
- A. Corona virus

- B. Swine Flue
- C. Zika virus
- D. Hepatitis

146. There is a flu that started in Hong Kong and spread to other areas

- A. Epidemic**
- B. Endemic
- C. Pandemic
- D. Sporadic

147. Obesity spread in Saudi Arabia is?

- A. Epidemic**
- B. Pandemic
- C. Endemic
- D. Sporadic

148. Increase BMI spread in Saudi Arabia is?

- A. Epidemic**
- B. Pandemic
- C. Endemic
- D. Sporadic

149. Unexpected increase case in geographic area consider as?

- A. Endemic
- B. Epidemic**
- C. Sporadic
- D. Pandemic

150. Influenza in HONG Kong and killed millions of people consider as ?

- A. Epidemic**
- B. Pandemic
- C. Endemic
- D. Sporadic

151. One question ask about Malaria transmission occurs in large areas of Central and South America, Africa, Asia?

- A. Epidemic
- B. Endemic**
- C. Pandemic

152. 152- A patients mania, delusion and hallucinations what is the most likely nursing diagnosis?

- A. Impaired thought
- B. Risk of direct violence
- C. Disturbances of sensory perception**

153. A patient will go to rectal surgery, doctor ask the nurse to bath him for the surgery which kind of bath the nurse will do?

- A. Bed bath
- B. Sitz bath**

- C. Partial bath
- D. With Cleansed

154. Which of the following is not a life threatening cause of chest pain?
- A. Pulmonary embolism
 - B. M I (myocardial infarction)
 - C. Acute coronary syndrome
 - D. Esophageal rupture
155. Patient came in ER when the nurse checked the temperature found febrile 40 . Which of the following refill should be?
- A. ICU
 - B. CCU
 - C. Medical
 - D. Communicable disease pavilion
156. What is 4th degree perineal tear?
- A. Perineal and external anal sphincter
 - B. Perineal anal sphincter mucosa
157. The doctor is conducting an assessment while the patient is uncomfortable and asks, What should the nurse do?
- A. Cover the unassessed area
 - B. Stop the assessment and write a report
 - C. Ignore the patient
 - D. Complete the assessment
158. Lanugo hair found on?
- A. Back
 - B. holder
 - C. forehead
159. The child presented with a low-grade fever, is sneezing and appears restless, What instructions would you give to the parents?
- A. Penicillin for 3 days
 - B. Antivirus 10 days
 - C. Antipyretic
 - D. Cough depressant
160. A patient has a low- grade fever and is shivering. What is appropriate action?
- A. Analgesic
 - B. Antipyretic
 - C. Antibiotic
161. A newborn presented with mild fever cough, poor feeding, sneezing, crying and appears restless. What advice did the nurse give to the mother?
- A. Amoxicillin for 3 days
 - B. Antiviral for a week
 - C. Give antipyretic for mild fever and discomfort

D. Cough depressant

162. Paediatric patient with low temperature what should the mother do?

A. Give an antipyretic

B. Administer an antibiotic course of 3 days

C. Administer an antibiotic course of 5 days

163. A 2-months old infant was brought to the health center for immunization, during assessment the infant temperature registered at 38.1C. Which is the best course of action that you will take?

A. Go on with the infant immunization

B. Give paracetamol and wait for his fever to subside

C. Refer the infant to the physician for further assessment

D. Advise the infant mother to bring him back for immunization when he is well

164. A patient with diabetes has high blood sugar. The nurse informed the doctor over the phone that he was busy in an operation, and then the doctor asked the nurse to administer orders for NPH insulin over the phone. What is the reason for the nurse to administer the medication?

A. Emergency situation

B. Physician is busy

C. Trust between nurse and doctor

D. Phone order , the same witness

165. What are the most common causes of maternal mortality?

A. Puerperal sepsis

B. Haemorrhage

C. Pre-eclampsia

D. Diabetes mellitus

166. Research about medical errors?

A. Grounded

B. phenomenology

C. Ethnography

D. Case studies

167. German measles incubation period?

A. 1 week

B. 2 weeks

C. 5 days

After rash 5 days

168. A woman first time do mammogram Pap smear testing. Which of the following is the goal of the nursing practice here?

A. Health promotion

B. Health protection

C. Health prevention

169. A 36 weeks pregnant woman went into labour and said that she uses cocaine. Who should the nurse report it to?

A. Obstetric doctor

- B. Hospital police
- C. Not tell anyone.
- D. Husband or legal guardian

170. A female patient came to the ER two weeks after postpartum and presented with haemorrhage. What type of postpartum haemorrhage is the nurse expecting?
A. Late postpartum haemorrhage
B. Early postpartum haemorrhage
C. Atonic
D. Traumatic
171. A pregnant woman in labour has a cervix dilated to 7cm. What is the frequency of contractions?
A. 3-4 minutes
B. 2-3 minutes
C. 1-2 minutes
172. When is abortion allowed in Saudi Arabia?
A. There is a risk to the mother
B. B There is a risk to the baby
173. 173- There is a difference between invasive and non-invasive fetal assessments. What non-invasive assessment does not harm the mother and child?
A. Amniocentesis
B. Cordocentesis
C. Non stress test
174. 174- A psychiatric patient received an overdose of benzodiazepine. The doctor prescribed an antidote. Which adverse effect of flumazenil should you check before administration?
A. Anxiety
B. Seizure
C. Shivering
D. Chest pain
175. A patient came with neck stiffness. He can't straighten his leg. What is this called?
A. Kernig's sign
B. Brudzinski sign
C. Seizure activity sign
176. A woman is diagnosed with gonorrhoea, and she is worried about her diagnosis. Which of the following is the appropriate description of the disease?
A. Partner test should not be present in
B. Asymptomatic
C. Sexual inactivity
177. A Patient with second degree AV block medication (atropine ,adenosine, diltiazem) After confirmation safety in patient arrest what second action?
A. Check responsiveness
B. Shout for help
C. Start chest compressions

D. Activate EMR

178. **15-years old female needs urgent surgery. The informed consent should taken from which person?**
A. Patient
B. Physician
C. Nurse
D. Legal guardian
179. **What is the complication of thalassemia?**
A. Hepatomegaly
B. Bleeding
C. Heart Failure
180. **A 5-year-old has been diagnosed with Wilms tumor. The doctor discussed the condition and available treatment options with the parents and provided them with an appointment to make a decision. What is the description of this action?**
A. Inspiring
B. Empower
C. Enabling
D. Encouraging
181. **A nurse is teaching a 13- years- old patient how to give an insulin injection. What is the description of the nurse's action?**
A. Inspiring
B. Empower
C. Enabling
D. Encouraging
182. **The child had a cleft lip repaired and is currently in elbow restraints. What should the nurse do?**
A. Remove the restrain every 2 hours
B. Remove the restrain when the parents are with the child
C. Put mittens.
183. **10 years old boy diagnosis appendicitis has serve pain, the tool of pain scale to be use is?**
A. Face pain rating
B. Flacc rating scale
184. **What is the best type of evidence-based practice for teaching diabetic patients?**
A. Randomize control study
B. Comparison of diabetic teaching types
C. Systematic review of randomize control
D. Study about diabetic teaching type.
185. **A teenager girl cannot administer insulin to herself. What should a nurse do for the patient?**
A. I make her come back to the hospital every few weeks
B. I teach the patient
C. I teach her parents

186. The nurse assess low of consciousness for patient after surgery, she asked patient about name and he give her correct answer then she asked about place, time but patient cannot answer, what should nurse do?
- A. Ask patient again later
 - B. Notify the surgeon**
187. A nurse is caring for a patient with diabetes who has a blood glucose level of 30mg/dl after a meal. What is the nurse's priority action?
- A. Measure the blood glucose level again after 30 minutes
 - B. Notify the physician immediately**
 - C. Ask another nurse (or head nurse) to measure the blood glucose level
188. Doctor inserts for patient subclavian catheter (CVL) in right subclavian vein sudden patient developed tachycardia destitution no back flow blood from CVL. What is the expected complication?
- A. Pneumothorax**
 - B. Pulmonary embolism
 - C. Hemothorax
189. Doctor, after inserting a subclavian catheter for the patient, blood flow is observed from the catheter, and the patient is experiencing respiratory restlessness. What is the expected problem with the patient?
- A. Right side failure
 - B. Pneumonia
 - C. Pulmonary embolism**
190. Doctor after inserting a subclavian catheter for the patient, blood flow is observed from the catheter, and the patient is experiencing respiratory restlessness. What is the expected problem with the patient?
- A. Right side failure
 - B. Pneumonia
 - C. Pulmonary embolism
 - D. Pneumothorax**
191. Pregnant women in active labour, how many times the midwife did vaginal examination?
- A. With each contraction
 - B. As much as pot
 - C. Every 4 hours
 - D. Limited vaginal examination**
192. The nurse research study to women eat more salt high risk for hypertension other group low risk. What is the type of study?
- A. Cohort study**
 - B. Randomized control trial
 - C. Case report
193. A patient 4 years old needs to receive a doctor-prescribed medication at a dosage of 10mg/kg/day for one week. The medication is available in a concentration of 125mg/5ml. The patient's body weight is 15 kg. How many ml per day should be given to the patient?
- A. 2 ml QID

B. 3 ml Bid

C. 5 ml OD

D. 6 ml PRN

194. What is the side effect of oxytocin ?

A. Water intoxication

B. Kidney toxicity

C. Diuretics

D. Fluid excess

195. The nurse research study to women eat more salt high risk for hypertension other group low risk. What is the type of study?

A. Cohort study

B. Randomized control trial

C. Case report

196. A teenager girl cannot administer insulin to herself. What should a nurse do for the patient?

A. I make her come back to the hospital every few weeks

B. I teach the patient

C. I teach her parents

197. A 30-year-old man was brought to the hospital by ambulance sitter falling from a height of 10 meters. He was mechanically ventilated on after a Glasgow Coma Score showed a level of six. The nurse is observe for any changes in perceptual, sensory or cognitive Which of the following is an expected patient's response at this time?

A. Slowly obeys commands

B. Exhibits no motor response

C. Reacts towards painful stimuli

D. Uses incomprehensible words

198. A nurse explains to a patient that an endoscopy tube will be inserted down the throat during a gastroscopy which will cause gagging, but it must be swallowed. What type of communication the endoscopy nurse has used here?

A. Assertive

B. Interactive

C. Formative

D. Informative

199. 209- A 45-year-old of man was admitted to the Surgical Ward after removed of pancreatic cyst. The registered nurse checked the postoperative orders Ondansetron (Zofran) PRN was prescribed for the patient. Which of the following complaints from the patient will require the registered nurse to serve the Ondansetron after 12 hours of post - operative period?

A. Wound pain

B. Nausea and vomiting

C. Congestion of flame

D. difficulty in passing urine

200. What is the transmission method of H1N1?

A. Blood

B. Body fluid

C. Saliva

201. Which of the following medications should the nurse give first?

A. Vancomycin 200 mg TID

B. Cefazolin 250 IV BID

B. Furosemide 40 gm STAT

C. Paracetamol 500 PRN

202. Most common cause of hospital acquired infection

A. Environment

B. Blood stream

C. Person to person

203. Post operative position of appendectomy

A. High fowler

B. Right lateral

C. Left lateral

D. Prone

204. Most reliable sign of dehydration

A. Keton in urine

B. Urine gravitate

C. Capillary refill tthan 3 second

205. Patient is psychotic with psychotic drug, with history of mental disorder, the mouth is dry why?

A. Patient is malnourished

B. Due to specific food pt take

C. Due to psychotic medication

206. A patient 4 years old needs to receive a doctor-prescribed medication at a dosage of 12mg/kg/day. The medication is available in a concentration of 125mg/5ml. The patient's body weight is 25kg. How many ml per day should be given to the patient?

A. 4 ml TID

B. 3 ml BID

207. A patient 4 years old needs to receive a doctor-prescribed medication at a dosage of 150 mg for one week. The medication is available in a concentration of 125mg/5ml. How many ml per day should be given to the patient?

A. 3 ml BID

B. 3 ml QID

C. 6 ml PRN

208. Patient after liver biopsy position on right sided. Why should the nurse keep patient on operation site?

A. Enhance blood circulation

B. Prevent bleeding

C. Prevent trapped fluid

209. A 2-day-old new-born is admitted to the nursery. While the nurse is administrating oral feeding, the milk returns through the child's nose and mouth and the infant become cyanotic. Which of the following condition the new-born should have?
- Anorectal malformation
 - Tracheoesophageal fistula**
 - Cleft lip and palate
 - Cardiac condition
210. Which of the following statement indicate nursing action during the first hour after delivery of the placenta?
- Monitor of mothers hemoglobin
 - Assess maternal vital signs every 15 minutes**
 - Ensure that the mother mobilize and empty her bladder
 - Administer 10 units of oxytocin via IV line to ensure uterus is well contracted
211. When the nurse assessed the fundus of a multiparous mother who delivered 2 hours ago, she found the following Level: 2 cm above the umbilicus. Position: deviated to the right. Consistency: Not well contracted what is the next nursing action after massaging the fundus until it becomes firm?
- Assess vital signs
 - Increase V fluids
 - Evacuate the bladder**
 - Ask the mother to res
212. A 29-year-old woman is receiving an intravenous infusion of 5% dextrose in 0.45% sodium chloride. the intravenous line had been started one day prior and now the patient complains of tenderness at the site. On examination, the site is pink coloured, swollen and tender to the touch. Which side effect of intravenous infusion problem is most likely?
- Infiltration
 - Phlebitis**
 - Catheter sepsis
 - Air embolism
213. A patient fell in the bathroom and his left leg was fractured, in order to communicate information about the patient to the next shift. Which of the following documentation should be used by the nurse at the end of the shift?
- Kardex record**
 - Assignment record
 - Shift report
 - Incident report
214. Surgeon instructs a nurse to serve as a witness to an elderly patient's informed consent for surgery. During the explanations to the patient, it becomes clear that the patient is confused and does not understand the procedure, but reluctantly sign the consent form. The nurse should:
- Sign the form as a witness, making a nation that the patient did not appear to understand
 - Not sign the form as a witness and notify the nurse supervisor**
 - Not sign the form and answer the patient's questions after the surgeon leaves he room
215. What is the main limitation of democratic style of leadership?
- it ignores the incompetent personnel

- B. cannot be used in emergency situations**
C. decreases collaboration among the team
D. may end up in inappropriate decision making
216. Nurse is given health education to the parent how to reduce febrile avulsion in the child at home. Which of the following medication in the safest intervention?
A. Analgesic
B. Antipyretic
C. Antibiotic
D. Antiemetic
217. A 40 years old female is diagnosed with chlamydia with no signs and symptoms ?
A. Subclinical stage
B. Clinical stage
C. Susceptibility
218. After thyroidectomy which of the following juice should be offered?
A. Apple juice
B. Tomato juice
C. Orange juice
D. Strawberry
219. A nurse is taking care of a woman who is in active labour. She is multipara at 38 weeks gestation. Her cervix is 5 cm dilated at -2 station with ruptured membranes. On the fetal heart rate tracing, the nurse observes a fetal heart rate baseline of 150 beats per minute with Deceleration in fetal heart rate to as low as 100 beats per minute. Deceleration in heart rate occurs both during and between contractions. With each deceleration heart rate quickly returns to baseline after a few seconds. The nurse has anticipated which of the following?
A. Client will probably deliver quickly
B. Client may need rapid intervention for a prolapsed cord
C. Client has a normal fetal heart rate tracing for a multipara in active labour
D. Client has a fetal heart rate tracing consistent with poor uteroplacental blood flow
220. A 56- year old man with history of COPD is complaining of chest, shortness of breath , lethargy, fever and productive cough. Upon examination, crackles could be heard in the lower lobes BP 11./70 HR 111 RR 18 TEM 37.4 Oxygen Sat 90%
A. Prone
B. Supine
C. High fowler
D. Trendelenburg
221. The nurse has been set priorities for health problem. She is starting to management of time board. Which of the following step of nursing process?
A. Planning
B. Diagnosis
C. Intervention
D. Assessment

222. The child presented with a low-grade fever ,is sneezing and appears restless, What instructions would you give to the parents?
- A. Penicillin for 3 days
 - B. Antivirus 10 days
 - C. Antipyretic**
 - D. Cough depressant
223. The nurse assess low of consciousness for patient after surgery, she asked patient about name and he give her correct answer then she asked about place, time but patient cannot answer, what should nurse do?
- A. Ask patient again later
 - B. Notify the surgeon**
224. 10 years old boy diagnosis appendicitis has serve pain, the tool of pain scale to be use is?
- A. Face pain rating**
 - B. Flacc rating scale
225. Patients with hip or femoral fractures. What should be monitored during the early post-traumatic period?
- A. Fever
 - B. Hypertension
 - C. Acute pain**
 - D. Hematuria
226. patient with chest pain, which radiates to the shoulder, sings of a myocardial infarction (MI). What should be the nurse's first intervention for this patient?
- A. Reduce Anxiety
 - B. Give oxygen**
 - C. Give nitroglycerin
 - D. Give aspirin
227. What is a common finding in most children with congenital heart disease?
- A. Mental retardation
 - B. Delayed physical growth
 - C. Clubbing of fingers
 - D. Cyanosis**
228. A patient is schedule for operation from out patient clinic one day ago came to surgical ward and admitted. Who is responsible to informed patient about surgery?
- A. Nurse
 - B. Head Nurse
 - C. Surgeon**
 - D. Anaesthesia
229. A patient 4 years old needs to receive a doctor-prescribed medication at a dosage of 10mg/kg/day for one week. The medication is available in a concentration of 125mg/5ml. The patient's body weight is 15 kg. How many ml per day should be given to the patient?
- A. 2 ml QID
 - B. 3 ml Bid**

- C. 5 ml OD
- D. 6 ml PRN

230. Term neonate meconium aspiration which of the following initial action by the nurse?
A. Give oxygen
B. Suction mouth and nose
231. A nurse is teaching a 13- years- old patient how to give an insulin injection. What is the description of the nurse's action?
A. Inspiring
B. Empower
C. Enabling
D. Encouraging
232. The child with dehydration and projectile vomiting He diagnosed with hypertrophic pyloric stenosis what's the best for feeding?
A. NGT
B. rehydrate, and continue breast-feeding
C. Rehydrate, and NGT feeding
233. A 3 days old newborn who is diagnosed with Hirschsprung disease presents with failure to pass meconium within 24-48 hours of life. What is the diagnostic test for Hirschsprung disease?
A. Chest X-ray
B. Ultrasound
C. MRI
D. Rectal Biopsy
234. German measles incubation period?
A. 1 week
B. 2 weeks
C. 5 days
5 days after rashes
235. There is a disease that started in a town, then it moved to two cities, spread to other areas, and killed more millions of people. What would this be considered as?
A. Endemic
B. Epidemic
C. Sporadic
D. Pandemic
236. The recommended action for pregnant women to take folic acid is to prevent which specific congenital disabilities?
A. Cleft lip and palate
B. Neural tube defects
237. Scrub nurse experienced wound in her hand. She came to ER and care is done. The next day she has duty. What should she do?
A. Report to head nurse
B. change role with circulating nurse

- C. Wear double gloves in right hand
- D. Wash five minutes before gloving

238. Patient with large liver tumour. The doctor decides to him cryosurgery. What does this type of surgery mean?
- A. **A-Freez/ cold therapy**
 - B. B-Heat therapy
 - C. C-hepatomegaly
239. Cryosurgery was performed on a patient with cancer tumors this surgery.
- A. 1-Freezes the tumors or abnormal or diseased tissue using cold liquid nitrogen organ gas
240. The physician order for a client cap Amoxycillin 500mg BID for 3 days. The available stock is 250 -mg per capsule . How many capsules will you give for 3 days ?
- A. 2 Capsules / 3 days
 - B. **12 Capsules / 3 days**
 - C. 4 Capsules / 3 days
 - D. 6 Capsules / 3 days
241. Hazard factory workers exposed to accidents and lesions and complain from backache and other problems it is called ?
- A. **physical**
 - B. Psychological
 - C. Mechanical
242. A nurse started a blood transfusion for the patient. Then the patient experienced complications due to receiving the wrong blood type. What is this behaviour by the nurse called?
- A. Lack of attention.
 - B. Defect (impaired) documentation.
 - C. **Failure to follow policy.**
 - D. Weak communication.
243. Pancreatic cancer is life threatening disease, there is low chance of survive if diagnosed in end stag , only 1% can survive for 2 year , what is best describe of mortality rate?
- A. Incidence rate higher than mortality rate
 - B. **mortality rate higher than incidence rate**
 - C. incidence rate not related to mortality rate
 - D. Mortality rate and incidence rate are approximately the same
244. What is the side effect of oxytocin ?
- A. **Water intoxication**
 - B. Kidney toxicity
 - C. Diuretics
 - D. Fluid excess
245. A toddler is seen in the Emergency Room with a history of high temperature for 5 days. He had watery eyes and a cough for the last 3days, and then he developed a rash all over the body which started in his face and spread down to his body. The mother did not remember the vaccination schedule of her son. For how long the child should be isolated?

فليطمئن قلبك ولتهدء روحك فالناس للناس..... و مازالت الدنيا بخير

- A. **5 days**
- B. 10 days
- C. 15 days
- D. 20 days

يحيى
GHAITH

Positioning

- Craniotomy → **semi fowler**
- Hip surgery → **keep leg in abduction**
- Cast → **elevate extremity to prevent edema**
- Asthma → **sitting position**
- Icp → **elevate the head of the bed**
- Distended neck vein → **elevate 30 degree**
- Enema → **sims position**
- Post cardiac cath → **semi fowler**
- Spinal pefida → **prone position**
- Seizure → **side lying**
- Shock → **supine with elevated leg**
- Nasogastric tube → **high fowler**
- Nasogastric tube irrigation and tube feeding → **HOB elevated 30-45**
- Central venous catheter → **trendelenburg position**
- Endoscopy → **left lateral position**
- Cleft palate → **supine upright position**
- Cleft palate → **elbow restrain and avoid prone**
- Post tonsillectomy → **side lying/lateral/prone**
- Teratology of fallout → **knee chest position**
- During liver biopsy → **position client supine with the right side of the upper abdomen exposed.**

فليطمئن قلبك ولتهده روحك فالناس للناس..... و مازالت الدنيا بخير

After liver biopsy 🖱️ **right lateral, side lying, position**

..... غيث 🥰.

After. hemorrhoidectomy 🖱️ **lateral, side lying, position, Jackknife position**

perineal or vaginal procedures 🖱️ **lithotomy position**

Central line CVP 🖱️ **trendelenburg position**

The nurse would increase the comfort of the patient with appendicitis by 🖱️ **Flexing the patient's right knee**

Post thyroidectomy position 🖱️ **Place the client in the semi-Fowler's to Fowler's position to reduce swelling and edema in the neck area.**

Mastectomy 🖱️ **with arm on affected side elevated to allow lymph drainage**



Cranial Nerves

Nerve I Olfactory Nerve	smell	Bottle containing something that smell
Nerve II Optic Nerve	Vision	Use visual acuity chart. (Snellen chart
Nerve III Oculomotor Nerve	Eye function , pupil dilation.	Have patient blink their eyes
Nerve IV Oculomotor Nerve	Eye movement	Have patient look down
Nerve V Trigeminal Nerve	Sensory of the face, chewing	pinprick & brushing teeth (Clench jaw. Test for feeling by lightly touching forehead, cheek and jaw.
Nerve VI Abducens Nerve	Eye movement	Have patient look to each side
Nerve VII facial Nerve	Facial expression, wrinkle forehead, taste anterior tongue	Cotton , hammer , sweats and salts
Nerve VIII Vestibulocochlear Nerve	Auditory acuity, balance and postural responses	Rinne test (nap fingers beside patient ear, have them stand on one foot)
Nerve IX Glossopharyngeal Nerve	Taste and swallowing	Tongue depressor and pen light
Nerve X Vagus Nerve	Gag reflex	Tongue depressor
Nerve XI Accessory Nerve	Muscles of neck	
Nerve XII Hypoglossal Nerve	Tongue movement.	stick tongue out

Acid Base Balance

Rule of the B's.. If the pH & the bicarb are both in the same direction = metabolic If they are in different directions = respiratory

RESPIRATORY OR METABOLIC !

PH $\uparrow$ HCO₃ $\uparrow$ or PH $\downarrow$ HCO₃ $\downarrow$ it is **Metabolic**

PH $\uparrow$ HCO₃ $\downarrow$ or PH $\downarrow$ HCO₃ $\uparrow$ it is **Respiratory**

Respiratory نفس الاتجاه تكون Metabolic عكس الاتجاه تكون

pH = 7.35-7.45 acidosis/alkalosis HCO₃
(bicarb) = 22-26 (2+2+2 = 6)
CO₂ = 45-35

ACIDOSIS OR ALKALOSIS !

PH > 7.45 **Alkalosis**

PH < 7.35 **Acidosis**

Compensated or uncompensated !

Un compensated if PCO₂ or HCO₃ are **Normal**

Partially compensated if Nothing is Normal

Compensated if PH is Normal

ex:

PH: 7.30 $\downarrow$

bicarb: 20 = $\downarrow$ = metabolic acidosis

ex:

pH: 7.58 = $\uparrow$

bicarb: 32 = $\uparrow$ = metabolic alkalosis

ex:

pH: 7.22 = $\downarrow$

bicarb: 30 = $\uparrow$ = respiratory acidosis

Vomiting	Metabolic Alkalosis
Nasogastric tube suction	Metabolic Alkalosis
Diarrhoea	Metabolic Acidosis
Ileostomy	Metabolic Acidosis
Dehydration	Metabolic Acidosis
CPR	Respiratory Acidosis
Drowning	Respiratory Acidosis
Cushing syndrome	Metabolic Alkalosis

246. A 12- year- old boy was brought to the Emergency respiratory arrest due to drowning. Cardiac resuscitation what is the major complication that might happen if treated after drowning quickly?
- A. Sepsis
B. Alkalosis
C. Acidosis
D. Hypothermia
247. An ICU nurse reviews the chart of a 47-year-old man patient mechanical ventilator for a long time. Arterial blood gas result see lab results).
- | Test | Result | Normal Values |
|----------------------|--------|---------------|
| ABG HCO ₃ | 24 | 22-28 mmd/L |
| ABG PCO ₃ | 10.66 | 4.7-6.0 KPa |
| PH | 7.16 | 7.36-7.45 |
| ABG PO ₂ | 6.13 | 10.6-14.2 KPa |
| SA O ₂ | 81 | 95-100 % |
- What condition the patient is experiencing presently?
- A. Metabolic acidosis
B. Metabolic alkalosis
C. Respiratory acidosis
D. Respiratory alkalosis
248. Gastric suction can cause :
- A. Metabolic acidosis
B. Respiratory acidosis
C. Metabolic alkalosis
D. Respiratory alkalosis.
249. ABG reading was low PH ,HIGH PCO₂ , NORMAL Hco₃ what the interpretation ?
- A. Compensated respiratory acidosis
B. Uncompensated respiratory acidosis
C. Metabolic acidosis
D. Metabolic alkalosis
250. Patient with ABG PH 7.33 وHCO₃ 30 , PCO₂ 50
- A. Compensate respiratory
B. Compensate metabolic
C. Uncompensated respiratory
D. Uncompensated metabolic

251. The nurse assesses a client with an ileostomy for possible development of which of the following acid-base imbalances?
- A. Respiratory acidosis
 - B. Metabolic acidosis**
 - C. Metabolic alkalosis
 - D. Respiratory alkalosis
252. The nurse is assigned to care for the a patient with Cushing syndrome on adrenal corticoid hormones syndrome on adrenal corticoid hormones Ph 7.2 Which of the following condition should nurse expect to patient
- A. Respiratory alkalosis
 - B. Respiratory acidosis
 - C. Metabolic alkalosis
 - D. Metabolic acidosis**
253. A nurse reviewed a chart of a 42 year-old man whose ABG analysis report is shown PH 7.20 PCO2 35 HCO3 20. Which of the following I the most likely interpretation of the findings in the report?
- A. Respiratory alkalosis
 - B. Metabolic acidosis
 - C. Respiratory acidosis
 - D. Metabolic acidosis**
254. A 20 year old woman, a case of panic attacks, comes to the emergency department. An arterial blood gas analysis is done PH 7.53 7.35-7.45 HCO3 22 22-26 PCO2 27 35-45 what is most likely?
- A. Metabolic acidosis
 - B. Metabolic alkalosis
 - C. Respiratory acidosis
 - D. Respiratory alkalosis**

Intravenous solution


ISOTONIC	HYPOTONIC	HYPERTONIC
• 0.9% NaCl • Lactated Ringer • Ringers' Solution • 5% Dextrose in water	• 0.45% NaCl • 0.33% NaCl • 0.2 % NaCl • 2.5% Dextrose water	• 3% NaCl • 5% NaCl • 3%NaCl or 5% NaCl +D/W • >5% D/W example, D10W

255. A client was admitted in the Emergency Room due to mild metabolic acidosis associated with dehydration and potassium the doctor administered Hartmann's (lactated Ringer's) intravenous fluid and electrolyte replacement. Which of the following elements of the lactated Ringer's solution highest value?
- Calcium
 - Sodium**
 - Potassium
 - Magnesium
256. Ringer Lactate consider as which type of IV solution?
- Hypotonic
 - Hypertonic
 - Isotonic**
 - Hyper alimentation
257. physician orders an intravenous fluid of D5NS at 100cc/hr. This is an example of which of the solution?
- hyper alimentation
 - hypertonic**
 - hypotonic
 - isotonic
258. 13-year-old patient is admitted for diarrhoea and vomiting. He looks pale and lethargic. A nurse is preparing to give IV hypotonic solution.
- | | |
|------------------|-------------|
| Blood pressure | 110/70 mmHg |
| Heart rate | 76 /min |
| Respiratory rate | 18 /min |
| Temperature | 36.1°C |
- Which IV solution is most appropriate?
- 0.9% saline
 - Lactated ringers
 - 10% dextrose in water
 - 0.45% sodium chloride**

CPR Questions

259. **CPR DEPTH** ONE THIRD OF CHEST = **1.5 INCH = 4CM** IN INFANTS , IN ADULT **2 INCH = 5CM**

260. CPR site



261. CPR Technique IS :

- A. **30:2 120**
- B. 15:2 100

262. You are performing CPR on an infant when a second rescuer appears. What is the next step in management?

- A. Immediately transport the patient
- B. Wait until exhausted, then switch
- C. **Have the second rescuer help with CPR, to minimize fatigue**
- D. Have the second rescuer begin ventilations; ratio 30:2

263. What should be your first concern at the scene where a person has been seriously burned?

- A. **Checking the scene for safety.**
- B. Checking the victims breathing and pulse.
- C. Calling your local emergency phone number.
- D. Cooling the burned area.

264. While performing cardiopulmonary resuscitation (CPR) on a 5-year-old child, the nurse palpates for a pulse. Which of the following sites is best for checking the pulse during CPR in a 5-year-old child?

- A. Femoral artery
- B. **Carotid artery.**
- C. Radial artery
- D. Brachial artery

265. After activating the emergency call system, what should be the next immediate action?

- A. Initiate ventricular pacing
- B. Administer a bolus of lidocaine as prescribed
- C. **Defibrillate the patient**
- D. Open the patient's airway



266. A nursing instructor teaches a group of students about basic life support. The instructor asks a student to identify the most appropriate location to assess the pulse of an infant under 1 year of age. Which of the following if stated by the student, would indicate that the student understands the appropriate procedure:

- A. Carotid
- B. Popliteal
- C. Radial
- D. **Brachial**

Chest Tube questions

267. A nurse is transferring a patient with chest tube the X-ray department for chest X-ray. Which location should the nurse place chest tube
- A. Directly on the stretcher in an upright
 - B. On the side of stretcher next to patient
 - C. Hanged on IV pole that is attached to the
 - D. Attached to the stretcher and hanged**
268. 69 year-old man is admitted to the intensive care unit following cardiac surgery. Two hours after admission, the nurse performs a routine assessment and notes the patient's chest tube drainage is 200 milliliters and a dark red color. He has had 60 milliliters output from the indwelling urinary catheter
- Blood pressure 138/68 mmhg
Heart rate 76/min
Respiratory rate 16/min
Body temperature 37.0C oral
Oxygen saturation 94% 6L/min nasal cannula
Which finding should be reported to the doctor?
- A. Dark red chest tube drainage
 - B. Urinary output
 - C. Oxygen saturation**
 - D. Chest tube output volume
269. 40 year-old male patient has a chest tube and properly fixed following cardiac surgery. The chest tube is patent and functioning. When the nurse removes the dressing the tube falls out of the chest. Which is the most appropriate initial intervention the nurse should take?
- A. Reinsert the chest tube and notify the surgeon
 - B. Apply an occlusive dressing to the insertion site**
 - C. Place the open end of the tube in 20 cm of water
 - D. Administer O2 at 10L/min via non-rebreather mask
270. While attempting to get of the bed a patient accidentally disconnects the chest tube from the pleura evac drainage system which of the following actions should the nurse take first
- A. Insert the end of the chest tube in a container of sterile solution**
 - B. Raise the end chest tube above the level of insertion of the chest tube
 - C. Clamp the chest tube near pleura evac drainage system
 - D. Apply pressure dressing to chest tube insertion site
271. The nurse is caring for a client who has just had a chest tube attached to a water seal drainage system. To ensure that the system is functioning effectively the nurse should:
- A. Observe for intermittent bubbling in the water seal chamber**
 - B. Flush the chest tubes with 30-60 ml of NSS every 4-6 hours
 - C. Maintain the client in an extreme lateral position
 - D. Strip the chest tubes in the direction of the client
272. A client chest tube is connected to a chest tube drainage system with a water seal . The nurse noted that the water seal c is fluctuating with each breath that client takes . The fluctuation mwans that
- A. There is an obstruction in the chest tube
 - B. The client is developing emphysema
 - C. The chest tube system is functioning properly**

D. There is leak in the chest tube system

273. 20-Your patient has a chest tube. Your assessing the water seal chamber and you note).11 that the water moves up as the patient inhales and then moves down when the patient ?exhales. What may be causing this to happen

A. This is normal and expected

B. The chest tube has a leak

C. The left chest tube is occluded

D. The water seal suction should be increased 2-5 mmHg

274. 21-You are assisting a MD with the removal of a chest tube. What activity may the MD have ?the patient perform while the chest tube is being removed

A. Valsalva maneuver.

B. Leopold Maneuver

C. Chest Physiotherapy

D. Huff Cough Technique

275. A 67 year-old man is admitted to the Post-anesthesia Recovery unit following chest surgery. The patient has a right chest tube that is attached to low suction. Three hours after admission to the unit, the nurse observes the drainage output from the chest tube is 300 milliliters. What is the most appropriate initial intervention?

A. Notify the doctor

B. Reduce IV infusion rate

C. Strip tube with roller device

D. Re-position in left lateral decubitus

276. The nurse is caring for a client who has had a chest tube inserted and connected to water seal drainage. The nurse determines the drainage system is functioning correctly when which of the following is observed:

A. Continuous bubbling in the water seal chamber

B. Fluctuation in the water seal chamber

C. Suction tubing attached to a wall unit

D. Vesicular breath sounds throughout the lung fields

Infection Control

277. Which of the following is an example of epidemic point source?

- A. Tuberculosis
- B. Public health agency
- C. Contaminated water source**
- D. Communicable disease pavilion

Airborne	Droplet	Contact
TB epidemic disease	Streptococcus	RSV
Corona virus	Meningitis	MRSA
measles	Diphtheria	Staphylococcus
chicken pox	Rubella	Ebola virus
Small pox	influenza	

Tuberculosis

- ❖ TB epidemic disease
- ❖ TB. Wear **N95 mask**
- ❖ TB. Test **Mantoux test / PPD TEST**
- ❖ TB TEST WITHIN **48-72 Hours**
- ❖ **Isoniazid** for TB side effect numbness , hepatitis symptoms , stomach upset ,rashes
- ❖ Tb accurate test **sputum culture** and to be done **early morning**

Meningitis. ...

- ❖ Streptococcal pneumonia
- ❖ meningococcal vaccine. Hib
- ❖ Neck rigidity
- ❖ Droplet
- ❖ Cloudy CSF
- ❖ Lumber puncture
- ❖ Surgical mask. No N95 mask

278. A 32 years old man develops chronic productive cough. He has not been feeling hungry and has lost three kilograms body weight in the past three weeks. On examination of his lung fields, there was diminished breath sound and widespread crackles. An early morning sputum culture was sent to the lab.

- A. Droplet
- B. Contact

C. Airborne

Rationale TB chest XRAY



279. 8 year-old man is diagnosed with tuberculosis (TB) and negative pressure room. Which of the following should wear a facemask?

- A. Patient's health care providers
- B. All people who enter the patient's room**
- C. Person has close contact with the patient
- D. Family members who are at risk for infection

280. A nurse working in medical unit is preparing to with droplet precaution measures in place. The following personal protective equipment; eyewear. What is the correct sequence for putting the equipment on?
- A. Face Mask, Gown, Eyewear, and Gloves
 - B. Gown, Face Mask, Eyewear, and Gloves**
 - C. Eyewear, Goggles, Face Mask, and Gown
 - D. Gloves, Gown, Face Mask, and Eyewear
281. A nurse working in medical unit is going out (removing) with droplet precaution measures in place. The following personal protective equipment; eyewear. What is the correct sequence for putting the equipment off?
- A. Face Mask, Gown, Eyewear, and Gloves
 - B. Gown, Face Mask, Eyewear, and Gloves
 - C. Eyewear, Goggles, Face Mask, and Gown
 - D. Gloves, Eyewear, Gown, and Face Mask**
282. Mr X attended in outpatient clinic with symptoms of shortness of breath, diarrhea and severe respiratory distress. Which of the following is the best diagnosis of Mrs. A case?
- A. Corona virus**
 - B. Swine Flu
 - C. Zika virus
 - D. Hepatitis
283. A nurse receives a telephone call from the admission office of the hospital and is told that a patient with streptococcal meningitis will be admitted to the Medical Unit. The nurse is planning to apply infection control measures for the patient. Which type of isolation precaution the nurse must observe?
- A. Droplet precautions**
 - B. Contact precautions
 - C. Airborne precautions
 - D. Standard precautions
284. A 35 year old patient was admitted to a medical ward with confirmed diagnosis of meningococcal infection. Which of the following infection control precautions the nurse should implement?
- A. Droplet precaution**
 - B. Contact precaution
 - C. Airborne precaution
 - D. Standard precaution
285. A 16 month old child is hospitalized in the intensive care unit with multi resistant sepsis. On the 3rd day. She had explosive diarrhea. A stool sample was sent to the laboratory for C. difficile investigation. Which of the following transmission based precaution is most appropriate?
- A. Combination airborne and droplet
 - B. Contact**
 - C. Droplet
 - D. Airborne
286. Which of the following is the most effective infection control precaution to prevent nosocomial infection?
- A. Hand washing before and after patient contact**
 - B. Wearing gloves and mask for direct patient care
 - C. Isolation precaution

D. Broad spectrum antibiotic

287. Mantoux test done for a patient result was 6.5cm it consider?

- A. Latent
- B. Suspected**
- C. Active

288. When planning discharge teaching for a patient hospitalized for treatment of the 3rd burns over 30% of the body, a nurse knows it is most important to include instructions regarding the loss of large amounts of serum occurring with burns and the resulting loss of immune function. Which of the following instructions should be include?

- A. Wash hands frequently each day**
- B. Wear supplemental oxygen at night
- C. Wear masks while in public spaces
- D. Take a multiple vitamin tablet night

289. An infection control nurse notices purulent exudates , redness and tenderness on the surgical wound site for few post –operative patients in a surgical unit , She discussed with the ward nurse and emphasized that wound infection after the surgery can be prevented. Which of the following is the best possible action to minimize the incidence of wound infection ?

- A. Perform assessment of pain on the wound site
- B. Wash hand before and after each patient activity**
- C. Encourage adequate intake and early ambulation

290. A patient in surgical was transferred to isolation room after the wound swab confirmed to have methicillin Resistant staphylococcus Atreus MRSA. Which of the following measures should the nurse take to prevent infection in the ward?

- A. Clean the would three times a day
- B. Discard all soiled dressing into waste bag
- C. Instruct the patient to wash hands regularly
- D. Wear gloves and gown on every entry into the room**

291. Medical asepsis requires which of the following hand washing techniques?

- A. Use hot water to ensure that pathogens are killed
- B. Use circular motion washing from clean to dirty areas
- C. Rinse soap off keeping hands and forearms lower than downs
- D. Hands shall be held higher than elbows and scrub**

292. A 45-year-old patient admitted with pulmonary tuberculosis. The unit nurse placed the patient in an isolation room with negative air pressure and prepared all the personal protective equipment at the entrance of the room. What type of precaution measure has the nurse activated?

- A. Contact
- B. Droplet
- C. Airborne**
- D. Standard

293. To reduce the risk of transmitting methicillin-resistant staphylococcus auras (MRSA) from an infectious wound, which of the following precautions should be implemented?

- A. Airborne
- B. Contact**
- C. Droplet

D. Reverse isolation

294. A 45 year old patient is admitted with pulmonary tuberculosis. The unit nurse placed the patient in isolation room with negative air pressure and prepared all PPE at the entrance of the room. Which of the following PPE is the most important for the nurse when caring for this patient?
- A. Hair cover
 - B. Sterile gloves
 - C. N95 respirator**
 - D. Protective goggle
295. 8 month-old child diagnosed with bronchiolitis due to respiratory ncytial virus (RSV) is admitted to the pediatric hospital. The nurse should initiate which of the following isolation precaution?
- A. Contact**
 - B. Airborne
 - C. Standard
 - D. Droplet
296. A 5-year-old child is seen in the primary care clinic with mild fever, headache, and malaise for about 2 days and today he has a rash filled with fluids. which of the following is the best suggested diagnosis?
- A. Chicken pox**
 - B. German measles
 - C. Measles
 - D. Scarlet fever
297. A eight year-old boy present to the pediatric clinic with his mother who complains that the child has developed a skin rash. There is a clear drainage from the nose and he appears tired. The rash first appeared on the face and then spread over the trunk and is now beginning to appear over the hands. Some lesions show open ulcerations and other have crusted over. There are scratch marks over the skin. Which finding would indicate the child is ready to return to school
- A. Normal body temperature
 - B. Dry and scabbed lesions**
 - C. Nares of free of drainage
 - D. Absence of rash over trunk
298. A patient with measles (rubella) is on airborne precautions. Which of the following precaution techniques would be essential to implement for non-immune persons entering the room?
- A. Gloves
 - B. Gowns
 - C. Masks**
 - D. Face shields
299. A home health nurse has entered a home to complete an admission assessment on a patient who has a methicillin-resistant Staphylococcus aureus (MRSA) urinary tract infection. The patient will receive intravenous anti-infective via a peripherally inserted central catheter (PICC) for 3 weeks. Which of the following actions should the nurse take FIRST?
- A. Shake the patient's hand
 - B. Place the nursing supply began a clean, dry surface.
 - C. Obtain the patient's written consent for home health care
 - D. Perform hand hygiene per the agency protocol**

300. You have just admitted a patient with bacterial meningitis to the medical-surgical unit. The patient complains of a severe headache with photophobia and has a temperature of 102.60 F orally. Which type of isolation should be suitable for the patient?
- A. Air borne
 - B. Contact
 - C. Droplet**
301. In order to reduce the risk of disease transmission from a patient with diphtheria, which of the following standard precautions would the nurse implement?
- A. Airborne
 - B. Contact
 - C. Droplet**
 - D. Ventilatory
302. Patient with a colostomy complains of itching of the peris ternal skin. On assessment, the skin is covered in a red rash with white patches visible. What is the most likely cause of this condition?
- A. Not changing the pouch regularly
 - B. Candidiasis**
 - C. Consuming acid-producing foods
 - D. Dehydration
303. A 33-year-old woman has come to the outpatient clinic for treatment of a vaginal infection. Physical assessment reveals yellowish excessive, thin offensive and frothy discharge. Which of the following is the most likely diagnosis?
- A. Candidiasis
 - B. Trichomoniasis**
 - C. Bacterial vaginosis
 - D. Chlamydia
304. The nurse is assessing a 65-year-old patient, who reports the fatigue, weight loss, night sweats, and a productive cough with thick sputum The nurse should immediately initiate isolation precautions for which of the following?
- A. Influenza
 - B. Pertussis
 - C. Bacterial pneumonia
 - D. Pulmonary tuberculosis**
305. Which of the following diagnostic tests is definitive for TB?
- A. Chest x-ray
 - B. Mantoux test
 - C. Sputum culture**
 - D. Tuberculin test
306. A patient presented with high fever, headache, vomiting and neck stiffness for the past 3 days, which of the following is the first diagnostic intervention for this patient:
- A. Urine and stool analysis
 - B. lumber puncture with CSF aspiration**
 - C. Complete blood count

D. Chest and abdomen x-ray

307. When a patient is admitted with acute influenza, what type of isolation is MOST appropriate?

- A. Reverse isolation
- B. Contact isolation
- C. strict isolation

D. Respiratory isolation

308. What precautions are necessary when caring for a patient with Hepatitis A?

- A. Gowning before entering the room
- B. Wearing gloves for direct care**
- C. Wearing a mask at all times
- D. Placing the patient in a private room

309. The nurse is assigned to care for the a patient with Ebola virus disease. Which of the following is the most common mode of transmission for Ebola virus?

- A. Vector
- B. Airborne
- C. Direct contact**
- D. Common vehicle

310. A nurse is caring a patient diagnosed with pulmonary tuberculosis and she has to wear N95 mask when she enters to the parent room. Based on the cycle of infection, which of the following is the reason to use the mask?

- A. Break the transmission chain at the portal of entry.**
- B. Break the transmission chain at the portal exit
- C. Kill the agent exits in the patients room.
- D. Eliminate the reservoir of the bacteria

311. After receiving the initial treatment of TB , a 27 years old woman in the chest disease ward was given discharge instruction on dietary management , medication, regimen, hygiene care and follow up visits. What is the instructions need to be emphasized the most?

- A. Increase protein diet to promote healing
- B. More rest and relaxation to restore therapy
- C. Maintain daily exercise schedule to improve health
- D. Dispose sputum as guided to avoid spread of disease**

312. A patient is admitted to the emergency room with an unknown infection disorder. Which of the following is the most important measure a nurse implement to reduce the risks of transmitting these microorganisms other?

- A. Hand hygiene**
- B. Wearing gloves
- C. Wearing a mask
- D. Wearing a gown

313. A nurse is assigned to care for a patient with small that needs to be placed on airborne precaution. Which of the following item should be used before entering the patient room?

- A. Gloves and gown
- B. Gloves and goggles
- C. Gloves and mask
- D. Gloves , gown N 95**

314. Which of the following is the primary level of prevention?
- A. Detect and treat existing disease
 - B. Prevent illness or injury occurring**
 - C. Reduce the extent and severity of health problem
 - D. Minimize disability and restore to prevent function
315. A Nurse in surgical unit prepares for a wound dressing. While she is preparing. She sneeze over the dressing trolley without wearing a facemask. Which of the following is the most appropriate action should the nurse take?
- A. set up a new sterile field**
 - B. put on a mask and continue working
 - C. continue working the most sterile field
 - D. Replace the equipment she thinks is contaminated
316. Which of the following is the most effective infection control precaution to prevent nosocomial infection?
- A. Hand washing before and after patient contact**
 - B. Wearing gloves and mask for direct patient care
 - C. Isolation precaution
 - D. Broad spectrum antibiotic
317. A 16 month old child is hospital in the intensive care unit with multi resistant sepsis. On the 3rd day. She had explosive diarrhea. A stool sample was sent to the laboratory for C. difficile investigation. Which of the following transmission based precaution is most appropriate?
- A. Combination airborne and droplet
 - B. Contact**
 - C. Droplet
 - D. Airborne
318. When planning discharge teaching for a patient hospitalized for treatment of the 3rd burns over 30% of the body, a nurse knows it is most important to include instructions regarding the loss of large amounts of serum occurring with burns and the resulting loss of immune function. Which of the following instructions should be include?
- A. Wash hands frequently each day**
 - B. Wear supplemental oxygen at night
 - C. Wear masks while in public spaces
 - D. Take a multiple vitamin tablet night
319. A 20 year old woman is hospitalized with a strong and uncontrollable cough and has difficulty breathing while coughing. A doctor writes an order for the patient to be transported from the medical surgical department to the radiology department for an xray examination. The nurse prepares to transfer the patient and considers standard precautions and additional transmission precautions. Who of the following would be required to wear a mask?
- A. Radiology staff
 - B. Nurse and patient**
 - C. Patient
 - D. Nurse

Thyroidectomy

320. A patient returned to the Surgical Unit from the thyroidectomy. The nurse observed that the arousable. Blood pressure 90/60 mmHg Heart rate 108 /min What immediate action should the nurse take?
- A. Recheck pulse and blood pressure
 - B. Administer intravenous fluids as ordered
 - C. Place client in modified Trendelenburg's
 - D. **Assess the back of neck surgical dressing for bleeding**
321. Complication of thyroidectomy
- A. Distension
 - B. **Bleeding**
 - C. Vocal cord injury
322. progressive enlargement of a multi-nodular go tracheal compression including pain at the site of the ear and jaw, difficulty swallowing, change of pf breath by compressing the oesophagus. There is in otherwise the patient is at risk. What is the preferred treatment?
- A. Iodine treatment
 - B. Thyroid hormone treatment
 - C. Radioactive iodine treatment
 - D. **Surgical resection of abnormal thyroid**
323. A Post thyroidectomy patient transferred to ward and developed tetany the nurse suspected that patient has ?
- A. Hypokalemia
 - B. **Hypocalcemia**
 - C. Hyponatremia
324. A nurse Wayne is aware that a positive Chvostek's sign indicate
- A. **Hypocalcemia**
 - B. Hyponatremia
 - C. Hypokalemia
 - D. Hypermagnesemia
325. **Trousseau's sign** A sign of hypocalcemia. Carpal spasm can be elicited by compressing the brachial artery with a blood pressure cuff for 3 minutes.



326. **Chvostek's sign** A sign of hypocalcemia. A spasm of the facial muscles elicited by tapping the facial nerve just anterior to the ear

Chvostek's sign: tapping of facial nerve branches
→ twitching of facial muscles ←



327. Patient is being admitted to the Recovery Room following a thyroidectomy. The back of the neck wound is covered with dressing. During the first 15 minutes, the patient started working and having diarrhea . A general assessment is performed with special attention given for the high risk for haemorrhage . Where would bleeding most likely occur?
- Stool
 - Vomitus
 - Dressing
 - Back of neck**
328. A 29 year-old man is in the Surgical Ward on his first post-operative thyroidectomy. He appears drowsy but the he is able to respiration by nodding head. He is developing mild restlessness. What is the initial recommended goal of care?
- Monitor vital signs of thyroid storm**
 - Assess for bilateral vocal fold mobility
 - Monitor for swelling on the neck
 - Monitor for vocal cord paralysis
329. A nuclear plant experienced a leakage, and all involved workers were brought to the Emergency Department for treatment . the medical team prescribed potassium iodide for the workers to block radioactive iodine to be absorbed by an organ in the body. Which of the following organs is the most sensitive to radioactive iodine?
- Brain
 - Lungs
 - Kidney
 - Thyroid**
330. The nurse care for a 60 year old woman who history hypertension, hypothyroidism and elevated cholesterol levels. She takes tablets daily for each of the health problem. The doctor orders a routine dual- x-ray absorptiometry test that shows decrease bone density. Which medication most likely contributed the test result?
- Statins
 - Anti-hypertensive
 - Synthetic thyroid hormones**
 - Cholesterol absorption inhibitors
331. A 36 years old man ha undergone a subtotal thyroidectomy ago. he Is suspected to develop tetany after the surgery. Which of the following symptoms best indicates tetanus?
- Tingling in the fingers**
 - Pain in hands and feet
 - Tension on the suture lines
 - Bleeding on the back of the dressing

332. The nurse is assessing 50 year old woman whose thyroid enlarged. A blood sample was collected and an analysis confirm diagnosis.


TSH 0.12 normal 0.4-6.5 Free T4 normal 50-140. Thyroxin T4 normal 4.5-11.2

- A. **Thyroidectomy**
- B. Incision and drainage
- C. polythyroidectomy
- D. adrenalectomy

Tonsillectomy

333. 3-year-old child with an elevated body temperature is administered oral aspirin. The nurse records the body temperature of the child two hours and The American Academy of pediatrics suggests that removal of the tonsils under certain conditions. Which of the following meets these conditions?

- A. times viral tonsillitis per year
- B. Infrequent snoring and nasal quality
- C. Three times bacterial tonsillitis per year
- D. **Tonsillitis accompanied by adenoid inflammation**



334. 5year-old child is postoperative after tonsillectomy. The nurse should ask the parents to give the child which of food after discharge from the hospital?

- A. Meat and rice
- B. Hot dog and potato chips
- C. **Mashed potatoes and soup**
- D. Cucumbers and tomato salad

335. The nurse is receiving a child postoperative tonsillectomy. Which of the following nursing assessment is suitable for the postop care?

- A. Encourage the child to cough spontaneously
- B. **Observe for subtle signs of haemorrhage**
- C. Place the child in the prone position
- D. Suction the mouth to clear the airway Because of violent behaviour

336. The nurse is receiving a child postoperative tonsillectomy. Which of the following nursing action is suitable for the postoperative care?

- A. Encourage the child to cough spontaneously
- B. Observe for subtle signs of hemorrhage
- C. **Place the child in the prone position**
- D. Suction the mouth to clear the airway Because of violence

337. After tonsillectomy, a child begins to vomit bright red blood .the Initial nursing action is to?

- A. Notify the physician
- B. **Turn the child to the side**
- C. Maintain an NPO status
- D. Administer the prescribed antiemetic

338. A client presents with Hypocalcemia, hyperphosphatemia, muscle cramps, and positive Trousseau's sign. What diagnosis does this support?

- A. Diabetes insipidus
- B. Conn's syndrome

C. Hypoparathyroidism

D. Acromegaly

339. The nurse cares for a client who has undergone a tonsillectomy. The nurse is most concerned about which post-operative finding?

A. Lack of appetite

B. Throat pain

C. Frequent swallowing

D. Nausea

Appendicitis

340. 18-year-old man college student was rushed faint at the school. He complained of severe quadrant. Upon palpation, he jerks even with sample was obtained. What is the most likely diagnosis of this patient?

A. Appendicitis

B. Liver Cirrhosis

C. Kidney stones

D. Duodenal ulcer

341. - year-old child was admitted with suspected appendicitis. A nurse was evaluating the child's condition and the mother stated that the child did have his bowel movement for the past two days and requested for natives. What is the risk of giving laxative to patient with appendicitis?

A. Pain

B. Fever

C. Rupture

D. Diarrhea

342. A woman patient is admitted for abdominal pain. She complain generalized plan nausea vomiting and constipation rebound tenderness and abdominal rigidity the past hour her localized on right side

BP 130/68 HR 92 RR 18 TEM 38.6 What should the nurse suspect

A. Intestinal obstruction

B. Influenza

C. Appendicitis

D. Pyloric stenosis

343. A man is to be discharged from the General appendectomy. The precautionary measures, plans are discussed with him. What is the most important desired outcome after discharge

A. Remain free of post-surgical complications

B. Report fever, redness or drainage from the wound site

C. Use pain management techniques appropriate

D. Resume gradual activities and avoid weight

344. A 17 year-old arrived to the Emergency Room complaining abdominal pain on right lower quadrant. Pain was rated as 9 numeric scale with positive rebound tenderness over the pain

Blood pressure Heart rate Respiratory rate Température

120/70 mmHg 95 /min

20 /min 39.2

Which of the following interventions has the highest priority?

A. Keep NPO

- B. Secure an IV access
- C. Prepare for ultrasound
- D. Prepare for abdominal surgery

345. The nurse would increase the comfort of the patient with appendicitis by:

- A. Having the patient lie prone
- B. Flexing the patient's right knee**
- C. Sitting the patient upright in a chair
- D. Turning the patient onto his or her left side

346. A 10-year-old girl presents to the Emergency Room (ER) with pain. On assessment, you noticed that when you palpate the right lower quadrant of the child's abdomen, the child feels pain in the right quadrant.

Which of the following is the name of this sign?

- A. Rebound tenderness**
- B. McBurney sign
- C. Roving's sign
- D. Obdurate sign

347. A 10-year-old girl presents to the Emergency Room (ER) with pain. On assessment, you noticed that when you palpate the left lower quadrant of the child's abdomen, the child feels pain in the right quadrant.

Which of the following is the name of this sign?

- A. Rebound tenderness
- B. McBurney sign
- C. Roving's sign**
- D. Obdurate sign

Medication and Drug Calculation

348. Treatment of nephritic syndrome is **corticosteroids**
349. **Isoniazid** for TB side effect numbness , hepatitis symptoms , stomach upset , rashes
350. A 66 year old woman with a history of unstable angina and hypertension present to the emergency department with a dull chest pain that she describes as similar to heartburn. The pain radiates down the left arm. She had taken sublingual nitro-glycerine tablets with any relief any electrocardiograph is perform and shows elevated S T segments. Which medication is most likely to given to dissolve the thrombus?
- A. Heparin
 - B. Warfarin
 - C. Streptokinase**
 - D. Aspirin
351. A 62 year man with a history of intracerebral bleeding three months ago, was referred primary health care following acute starch symptoms. Brain CT scan is normal and the patient is receiving oxygen by nasal cannula at 4LPM BP 185/105 HR 82 RR 18 TEM 36.6 SO2 93%
- Which of the following medication would be ordered first?
- A. RtPA
 - B. Aspirin**
 - C. Dopamine
 - D. Nicardipine
- Rational :** Probable acute ischemic stroke; consider fibrinolytic therapy. If the CT scan shows no sign of hemorrhage, it is probable that the patient experienced an ischemic stroke and is a candidate for fibrinolytic therapy. Check for fibrinolytic exclusions such as significant head trauma or stroke in the previous 3 months, history of intracranial hemorrhage, elevated blood pressure, active internal bleeding, or a blood glucose concentration less than 50 milligrams per deciliter. Then repeat the neurologic exam. Consult neurologist or neurosurgeon. If the CT scan indicates hemorrhage, consult neurologists and neurosurgeons, and begin the stroke or hemorrhage pathway. Administer aspirin. If the patient is not a candidate for fibrinolytic therapy, administer aspirin and begin the stroke or hemorrhage pathwa
352. A nurse checks the medication chart of a patient, which has the following order, nitroglycerin Tab sublingual. Which of the following medication administration routes should the nurse use?
- A. In the nose
 - B. In the eyelid
 - C. Under the skin
 - D. Under the tongue**
353. A 28 year man admitted to orthopedic ward complaining of throbbing pain in casted leg. Which of the following nursing intervention should be taken first?
- A. Remove the cast
 - B. Notify a doctor
 - C. Assess pedal pulse**
 - D. Administer PRN medication

354. A 62 year old woman presents to the clinic with a primary complaint of fatigue. An assess shows a smooth and reddened tongue with a loss of papillae and pallid mucous membranes. She complaints of a sense of constant fullness of the stomach, with a decreased appetite and two three loose bowel movement per day. A neurological assessment shows numbness in the feet and lower legs. The patient type of test would the most likely?
- Schilling
 - Erythropoietin
 - Folic acid levels
 - Vitamin B12 levels**
355. A mother of a patient who is on antipsychotic drug asked a doctor about some more detail about the drug. The doctor told the mother that her son is taking an atypical antipsychotic drug. Which of the following is an examples of this type of drug?
- Thioridazine
 - Clozapine**
 - Chlorpromazine
 - Haloperidol
356. A nurse is preparing scheduled medications due at 6 pm. If a doctor orders paracetamol tab 1g QID, and it was supplied from the pharmacy in 250 mg tablets. Which of the following is the most appropriate nursing actions?
- Ask the pharmacy to provide 1g tablets
 - Call the doctor to recheck the dosage
 - Give the patient four 250 mg tablets**
 - Hold the medication and document in nursing notes
357. A patient with a history of atrial fibrillation has an order for 25 milligrams of drug (X), the available supply/dose is 0.25 milligrams. Which of the following doses in correct?
- Dispense two tablets to the patient
 - Administer one tablet twice daily
 - Dispense one half tablet to the patient
 - Return tablets to the pharmacy and re-order**
358. A patient with a history of atrial fibrillation has an order for 0.25 milligrams of drug (X), the available supply/dose is 0.125 milligrams. Which of the following doses in correct?
- Dispense two tablets to the patient**
 - Administer one tablet twice daily
 - Dispense one half tablet to the patient
 - Return tablets to the pharmacy and re-order
359. 13-year-old patient is admitted for diarrhea and vomiting. He looks pale and lethargic. A nurse is preparing to give IV hypotonic solution.
- | | | | |
|------------------|-------------|-------------|---------|
| Blood pressure | 110/70 mmHg | Heart rate | 76 /min |
| Respiratory rate | 18 /min | Temperature | 36.1°C |
- Which IV solution is most appropriate?
- 0.9% saline
 - Lactated ringers
 - 10% dextrose in water
 - 0.45% sodium chloride**

360. A nurse is assigned to care for a group of patient in the medion expected to review the medical records of these patient. What patient is at risk for excess fluid volume?
- A. Patient with ileostomy
 - B. Patient taking a loop diuretic
 - C. Patient with chronic renal failure**
 - D. Patient hooked to gastrointestinal
361. physician orders an intravenous fluid of D5NS at 100cc/hr. This is an example of which of the solution?
- A. hyper alimentation
 - B. hypertonic**
 - C. hypotonic
 - D. isotonic
362. When a patient was first diagnosed with schizophrenia, one of his family members asked the nurse about the possible causes. The nurse said that one reason is that he may have had an excess secretion of a neurotransmitter. Which of the following neurotransmitters?
- A. serotonin
 - B. dopamine**
 - C. glutamate
363. The nurse administered a dose of morphine sulfate as prescribed to a patient who is in the post anesthesia care unit (PACU). The patient appears to be resting comfortably, the respiratory rate is 8 and the O2saturation is 21 oxygen via cannula is 86%. The nurse should IMMEDIATELY administer:
- A. Flumazenil(Romazicon)
 - B. Medazolam(verSED)
 - C. Naloxone (Narcan)**
 - D. Ondansetron (Zofran)
364. Digoxin not given to patient who has which of the following ?
- A. tachycardia
 - B. hypertension
 - C. tachypnea
 - D. bradycardia**
365. Atropine is indicated for increase
- A. Heart rate**
 - B. Respiratory rate
366. Morphine side effect ?
- A. tachycardia
 - B. hypertension
 - C. tachypnea
 - D. bradypnea**
367. patient with ongoing magnesium sulfate the nurse should be alert to ?
- A. temperature
 - B. respiratory rate**
 - C. heart rate
368. Soldier was brought to triage area after being exposed to chemical weapons. Signs and symptoms of nerve gas exposure were noticed. A nurse prepares for medical management. Which medication should the nurse prepare for the patient?

- A. **Atropine**
- B. Adrenaline
- C. Sodium nitrate
- D. Sodium thiosulphate

369. Nitroglycine administration route :

- A. Intramuscular
- B. **Sublingual**
- C. Subcutaneous
- D. Oral

370. A nurse checks the medication chart of a part, which has the following order; Nitroglycerin tab Buccal at stat. Which of the following medication administration routes should the nurse use?

- A. In the ear
- B. Intravenous
- C. Under the skin
- D. **Between cheek and gum**

371. A nurse is caring for a patient who had Coronary Artery bypass Graft Surgery (CABG) four hours ago. The nurse notices that the patient has increased confusion and is restless. The patient reports nausea, weakness and paresthesia in the extremities (see lab results)

Normal Values

Result Test

134-146 mmol/L

145 Sodium

3.5-5.2 mmol/L 6.8 Potassium

2.15-2.62 mmol/L 2.50 Calcium

Which of the following is the best medication?

- A. Naloxone (Narcan)
- B. Hydralazine (Apresoline)
- C. Potassium chloride (KCl)
- D. **Sodium polystyrene sulfonate (Kayexalate)**

372. Which of the following statement by the nurse about the clomid as an ovulation inducing drug?

- A. Given for the first 15 days in each cycle
- B. Maximum dose is 50 mg daily for a month
- C. It increases the risk of birth defects
- D. **It increase the risk of multiple pregnancies**

373. Clomiphene citrate (Clomid) is prescribed for a 32-year-old infertility treatment. The nurse should understand that this medication is used for following actions?

- A. **induce ovulation**
- B. Decrease prolactin level
- C. Reduce endometriosis
- D. Stimulate the release of Follicle-Stimulating Hormone

374. A nurse is caring for a client with bipolar disorder, who is receiving Lithium carbonate. Before Administration of the next dose, the client complains of nausea and vomiting and the nurse finds that the client's lithium blood level 1.5mEq/L. Which of the following actions is considered Apriority?

- A. call the client's physician immediately
- B. **withhold the next dose.**

- C. Administer IV fluids
- D. Repeat the blood lithium level testing

375. A patient was on a regular dose of lithium carbonate. The nurse noticed he has hand tremor, polyuria, diarrhea and vomiting. What immediate action should be taken by the nurse?
- A. Diuretics
 - B. Withholding lithium**
 - C. Calling the psychiatrist
 - D. Monitoring serum lithium level
376. A patient was on a course of lithium carbonate drug. During the nurse found that he complained from nystagmus visual hallucination, and oliguria Which of the following drug related complications best symptoms?
- A. Overdose
 - B. Mild toxicity
 - C. Severe toxicity**
 - D. Moderate toxicity
377. 49-year-old women presented to the Emergency Department complaint of severe chest pain. The ECG showed that the patient myocardial infarction. The doctor ordered the nurse to give the 800 mg of aspirin. What is the primary indication of aspirin in this case?
- A. Breaks down the thrombus
 - B. Decreases the formation of platelet plugs**
 - C. Inhibits the conversion of prothrombine to
 - D. Interferes with vitamin k to maintain
378. which of the following condition is a Contraindication for a woman oral Contraceptives?
- A. Dysmenorrhea
 - B. Menorrhagia
 - C. Thrombophlebitis**
 - D. Toxic shock syndrome
379. A patient was on a course of lithium carbonate drug. During the nurse found that he complained from nystagmus visual hallucination, and oliguria Which of the following drug related complications best symptoms?
- A. Overdose
 - B. Mild toxicity
 - C. Severe toxicity**
 - D. Moderate toxicity
380. A 45-year-old distressed and restless patient in the Psychiatric Ward was unable to sleep during the nights for the last two days. He was ordered sleep medication which was to be administered at 10 pm. At the time medicine administration, the patient was found asleep in bed Which of the following action should the nurse take regarding the dictation?
- A. Discard medicine and cancel the order
 - B. Leave it at the bed side for the patient
 - C. Wake him up and administer
 - D. Hold, record and report**

381. 13- nurse is given health education to the parent how to reduce febrile avulsion in the child at home. Which of the following medication in the safest intervention?
- Analgesic
 - Antipyretic**
 - Antibiotic
 - Antiemetic
382. Doctor order to give Ofloxacin Otic route the nurse understand that medication will be given :
- Nasal
 - Eye
 - Ear**
 - Sublingual.
383. A home care patient with chronic Obstructive reports an upset stomach. The patient is taking and triamcinolone acetate (azmacort). Which of the following counselling should be given
- Theo-dur on an empty stomach
 - Theo-dur and azmacort at the same time
 - Theo-dur and azmacort 12 hours apart
 - Theo-dur with milk or crackers**
384. Which of the following vitamins is necessary for wound healing process?
- Vitamin A
 - Vitamin C**
 - Vitamin D
 - Vitamin K
385. Which of the following vitamin supplements can decrease the incidence of Neural tube defects such as anencephaly and spina bifida new-borns ?
- Vitamin A
 - Riboflavin
 - Folic Acid**
 - Vitamin K
386. A 78-year-old woman who lives in a long-term care facility has been having repeated episodes of urinary tract infections. She is prescribed phenazopyridine one gram to be taken by mouth four times per before meals and at bedtime. The nurse advises the patient that she need to remove milk from the diet while taking the medication. What is the primary purpose for this dietary advice?
- To prevent mal absorption of medication
 - Decrease risk of gastrointestinal upset
 - Reduces effectiveness of medication
 - To make the urine acidic**
387. A nurse received the serum digoxin level result for the patient the day and notes that the result is 2.6 ng/mL (see lab result)

Test	result	normal value
Digoxin (men)	2.6	0.8-2ng/mL

Which of following nursing actions is the most important?

A. Notify the physician

- B. Check previous vital signs of patient
- C. Record normal value on nursing note
- D. Administer scheduled dose of medication

388. A patient is being admitted in medical unit and has orders for fluid restriction of 1400ml / 24 hours. He is receiving IV medications in 50ml solutions QID. How much fluid should the caring nurse allocate for cr intake in 24 hours?

- A. 400 ml
- B. 800 ml
- C. 1000 ml
- D. 1200 ml**

389. A nurse is preparing an order of Amydramine 30 mg PO BD. The available check is 15 mg /5 ml. What is the correct dosage of the medication to be administered?

- A. 5 ML
- B. 10 ML**
- C. 15 ML
- D. 20 ML

390. A nurse is preparing a medication order of 100 mg IV stat. The available hydrocortisone is 200mg per 5 ml distilled water for injection. Which of the following is the correct administered?

- A. 4.5 ML
- B. 3.5 ML
- C. 2.5 ML**
- D. 1.5 ML

391. A nurse is preparing to administer an intravenous infusion of 2000 ml of Ringer's lactate over 12-hours. The administration set has a chamber that delivers 15 drops per ml. How many drops per minute should be administered?

- A. 14
- B. 24
- C. 42**
- D. 56

392. Doctor order to give medication x 300mg and available is 900mg/6ml how many ml the nurse will give

- A. 1.2
- B. 2**
- C. 4
- D. 6

393. A 56-year-old man was admitted with complaint working for three days. The nurse is preparing to administer infusion of saline 1000 ml over six hours. What is the hourly infusion rate that the nurse needs to infusion therapy in mL/hour?

- A. 155
- B. 167**
- C. 190
- D. 217

394. Doctor Order: Heparin 10000 units OD S/C Standard Solution 40000 units/ml How many ml would you administer?
A. 0,25.ml
395. A Physician has ordered clindamycin phosphate 300 mg to be administered intravenously the available stock was labelled . clindamycin phosphate 900mg mg in 6 ml what is the correct dose to be administered by the nurse ?
A. 2 ml
B. 4 ml
C. 6 ml
D. 8 ml
396. A nurse check the medication chart of a patient, which has the following order; Paracetamol 500mg P.O. QID. How many times a day should the nurse administer the medication?
A. Once
B. Twice
C. Three times
D. Four times
397. The following syringe contains 2 gm of morphine sulphate, how many grams are there in 1 ml:
A. 0.5 gm
B. 1 gm
C. 2 gm
D. 4 gm
398. The following syringe contains morphine sulphate in a concentration of 2 mg/cc, how much total morphine this syringe contains:
A. 0.5 mg
B. 1 mg
C. 2mg
D. 4 mg
399. A nurse is assigned to care for a patient in medical ward who has a physician order of 1500ml of normal saline 0.9% to be administered over 12 hour period. Which of the following is the infusion rate?
A. 75 ml/hr
B. 100ml/hr
C. 125ml/hr
D. 150ml/hr
400. Doctor order to give 75ml per hour over 12hours of normal saline how many ml total should be given?
A. 450ml
B. 600ml
C. 900 ml
D. 1000 ml
401. A 37year old female prevented to the Emergency room with complain of symptomatic bradycardia appropriate nursing intervention include establish intervention access what is the possible drug to be given in the IV solution ?

- A. Digoxin
- B. Atropine**
- C. Anticoagulant
- D. Calcium channel blocker

402. A 25 year old woman present to the emergency room with decrease reflexes, hypoventilation, hypotension, and fuced pupils, a family member who is accompanying the patient has bottle of diazepam which the label states was recently the family member also indicates that the patient has a depression. What intervention should the nurse expect to administer?
- A. Naloxone
 - B. Active charcoal**
 - C. Tap water enema
 - D. Magnesium sulfate to reduce the risk seizure
403. Nurse gives health education to patient In outpatient department regarding a medication . she told him "avoid working outdoors in a hot weather, excessive sweating ad during l water daily Which of the following is most likely medication?
- A. Diazepam
 - B. Haloperidol
 - C. Lithium carbonate**
 - D. Monoamine oxidase inhibitor
404. During CPR of an admitted patient following the cardiac arrest. Two Dc shocks and one dose of epinephrine was given. A third dose was delivered followed by anti arrhythmic medication . what medication is the most likely to be given next?
- A. Adenosine
 - B. Lidocaine 2 mg
 - C. Epinephrine 1 mg**
 - D. Sodium bicarbonate
405. A nurse is giving discharge planning instructions to the parents of a one year old child acute otitis media which of the following discharge instruction takes priority?
- A. Administrate antibiotics as prescribed**
 - B. Administer influenza vaccination
 - C. Breast feeding as long as possible
 - D. Continue using of pacifier
406. A 4 years old girl, was playing outside, she came to her mom crying and holding her right upper arm, she went to the hospital with swelling over the upper arm, pain and itching, the appropriate management is:
- A. Maintain patent airway
 - B. Administer s/c Epinephrine**
 - C. Prepare for intubation
407. An older adult client with renal failure comes to the emergency department with a report of nausea and vomiting. The client's heart rate is 45 beats/min. The nurse is most concerned about which medication that the client takes
- A. Nitroglycerin
 - B. Digoxin**
 - C. Doxorubicin
 - D. Furosemide

408. The nurse is caring for a 60-year-old in the Medical-Surgical Unit. On entering the room the nurse finds the patient alert, oriented and. The skin color appears healthy and pink. The nurse observes electrocardiogram and notices absent P-waves. Which initial action is the most important?
- A. Defibrillate
 - B. Cardio version
 - C. Pacemaker preparation
 - D. Intravenous amiodarone**
409. During health education for a patient on Monoamine oxidase inhibitor (MAOI) the nurse instruct him to avoid certain foods that can interact with the MAOI. Food containing with of the following contents should be avoided?
- A. Alcohol
 - B. Caffeine
 - C. Tyramine**
 - D. Folic acid
410. After teaching a 54 year old patient with angina on how to take nitroglycerin sublingual PRN. Which of the following statements reflect the patient understanding?
- A. I have to take this medication once i need it only**
 - B. It is ok to take one tablet daily to prevent the heart attack
 - C. I can take two tablets together at once if the attack is
 - D. This medication will regulate my heart beats and I will be
411. A 45 year old client who diagnosed with brain was schedule for craniotomy. It is important to preventing the developing of cerebral edema after surgery. What medication would the nurse expect to prescribed the client?
- A. Steroids**
 - B. Diuretics
 - C. Ant convulsions
 - D. Antihypertensive
412. A 71 year-old woman who resides in a long-term nursing home fell while walking down stairs. The attending nurse arrives to find the patient sitting motionless on the stairs. She is alert and oriented but wishes to rest. While she rests, the nurse reviews the chart and notes that her medication regimen includes metformin, loratadine, warfarin and diclofenac. Which medication is most likely to increase the patient's risk of injury?
- A. Metformin
 - B. Loratadine**
 - C. Warfarin
 - D. Diclofenac
413. ECG showed S-T elevation to confirm diagnosis of MI the nurse expect to do?
- A. Troponin level**
 - B. Cardiac catheterization
 - C. ECO

Leadership and administration

414. On orientation day, a nurse educator said to a new nurse "our hospital provides the highest level of specialized healthcare in an integrated educational and research setting" which of the following aspects of the hospital's strategic plan is indicated by this statement?
- A. Vision
 - B. Policy
 - C. Values
 - D. Mission**
415. A head nurse of an intensive care unit wrote a report about the needs of her department and gave it to the nursing supervisor. Which of the following types of communication is this?
- A. Upward**
 - B. Horizontal
 - C. Diagonal
 - D. Downward
416. A 45 year old patient has a colostomy bag attached as a result of benign mass removal, while conducting the discharge teaching session, the nurse assessed the patient's wife understanding of the teaching. Which of the following steps of nursing process is exercised here?
- A. Implementation
 - B. Planning
 - C. Assessment
 - D. Evaluation**
417. Which of the following is correct nursing process order?
- A. Diagnosis, assess, plan, evaluate, implement
 - B. Assess, diagnosis, plan, implement, evaluate**
 - C. Evaluate, assess, plan, implement, diagnosis.
 - D. Plan, assess, diagnosis, evaluate, implement
418. A nurse calls a physician regarding a change in a patient's condition. A physician gives orders over the telephone for stat arterial blood gases. Which of the following is the best intervention to be implemented?
- A. Give order stat to health unit coordinator to input in computer
 - B. Write down order and read it back to physician**
 - C. Write down order for ABGs immediately
 - D. Call respiratory therapist stat to draw the ABGs
419. During Cardiopulmonary Resuscitation (CPR) for a 75-year-old man in the Emergency Department, the doctor introduced himself as a leader for the CPR. What is the most appropriate leadership style for this situation?
- A. Autocratic**
 - B. Democratic

- C. Bureaucratic
- D. Laissez-faire

420. A nurse manager is open minded, listens to the team, understands others, makes changes to improve unit operations and procedures. What type of leadership style is this?
- A. Situational
 - B. Democratic**
 - C. Compassionate
 - D. Transformational
421. Leadership using theories which type of leadership?
- A. Autocratic**
 - B. Democratic
 - C. Transnormal
422. Nurse supervising on assistant nurse and delegate assignment to him and say "diving up patient's assignment" what type of leadership
- A. Autocratic
 - B. Laissez-faire**
 - C. situational
 - D. transformational
423. A nurse manager includes staff in decision-making process what unit structure is used?
- A. Divisional
 - B. Centralized
 - C. Functional
 - D. Decentralized**
424. What is the main limitation of democratic style of leadership?
- A. it ignores the incompetent personnel
 - B. cannot be used in emergency situations**
 - C. decreases collaboration among the team
 - D. may end up in inappropriate decision making
425. A nurse manger is preparing and writing a plan for dealing disasters (code red). Which procedure is the top priority for the nurse manager the plan?
- A. Infection control
 - B. Staff orientation**
 - C. Patient education
 - D. Patient relocation
426. Nurse manager was not happy about low results of patient's survey. Which of the following is the first step for nurse manager?
- A. Schedule meeting with staff
 - B. Start changes to improve
 - C. Review evaluation details**
 - D. Report to director
427. Post-operative patient who underwent an abdominal procedure requests a pain medication from the nurse and rates the pain at a level nine. There a standing order for narcotic administration. When the nurse opera the narcotic box and performs a count, the number of pills remaining in the box is different than the number of pills recorded on the sheet. What is the most appropriate initial nursing action?
- A. Notify the nursing supervisor**
 - B. Write the finding on the narcotic sheet

- C. Administer the patient's requested medication
- D. Identify the last nurse who used the narcotic box

428. An 81-year-old bed ridden patient in the Geriatric Ward was pyretic through his nasogastric tube. After an hour, the patient deep sleep, his breathing pattern changed and he did not rep calling his name. Which of the following should be the immediate nursing intervention?
- A. Call the physician to examine the patient
 - B. Check for the correct dosage given
 - C. Check vitals and inform physician**
 - D. Try to wake up the patient
429. The nurses in the Male Medical Unit took a signature on the from a patient who needs an abdominal CT with contrast. What is the best nursing practice?
- A. Consent should be taken after the procedure
 - B. Verbal consent is acceptable for this procedure
 - C. Consent should be taken from the patient's relatives
 - D. Consent should be taken according to the organization's**
430. In the hospital digital dashboard, what types of data displayed?
- A. Staffing
 - B. Financial
 - C. Performance**
 - D. Knowledge-based
431. A newly nursing director assigned to a hospital. That is the first should he do?
- A. Evaluates the staff**
 - B. Change the head nurses
 - C. Change the roles
 - D. Nothing to do
432. Surgeon instructs a nurse to serve as a witness to an elderly patient's informed consent for surgery. During the explanations to the patient, it becomes clear that the patient is confused and does not understand the procedure, but reluctantly sign the consent form. The nurse should:
- A. Sign the form as a witness, making a nation that the patient did not appear to understand
 - B. Not sign the form as a witness and notify the nurse supervisor**
 - C. Not sign the form and answer the patient's questions after the surgeon leaves he room
433. A community nurse assigned to work in a Makkah, one of the most multicultural cities in Saudi Arab has to interact with multicultural population every day What is the most important step the nurse should do before with the clients?
- A. Asking the client about his/her cultural background
 - B. Conducting an appropriate culturological assessment**
 - C. Ensuring that the client has his/her a privacy
 - D. Looking at the client's file to take history
434. The head nurse meets with staff nurses to discuss ways to improve communication among shifts Which of the following statement best exemplifies the final stage of conflict management?
- A. "We need to clearly define the nature of the conflict"
 - B. " I will evaluate the outcomes of the strategies used monthly**

- C. " Let us create a time line for the implementation of our strategies"
- D. " I have to force you to follow the rules to resolve the issue"

435. The nursing director wants to evaluate the quality of nursing care at the in-patient areas. The management team will evaluate on quarterly basis the documentation and the relationship between the patient's length of stay and the quality of care which of the following is the most important data source to identify the quality of care?
- A. Patient's complaints and time taken to resolve them
 - B. Patient's satisfaction level at the time of discharge**
 - C. Details of nursing notes for patient's progress
436. The nurse manager received complaints from some nursing to improper work distribution and discrimination How should the nurse manager handle the situation?
- A. Investigate the complaints**
 - B. Individual counselling
 - C. Identify list of problems
 - D. Continue to observe
437. Which of the following is the most appropriate action for a head nurse starting at a new hospital?
- A. Make immediate change at the unit
 - B. Plan and coordinate new strategies**
 - C. Assess unit activities for at least three months
 - D. Ask about the previous head nurse managerial style
438. After teaching the deep breathing and coughing who is undergoing a surgery, the nurse asked Demonstration and then helped him in correcting what part of therapeutic communication is used
- A. Evaluation**
 - B. Intervention
 - C. Identification
 - D. Demonstration
439. the intensive care unit nurse manager plans to delegates a nurse. What is indicated for a successful delegation?
- A. Nurse Manager supervises nurse.**
 - B. Nurse has authority to change task.
 - C. Nurse Manager checks task sometimes.
 - D. Nurse Manager asks another nurse to check task.
440. nurse accidentally dropped a medication ampoule, informed the charge nurse, and completed an incident report form. The charge nurse arranges medication replacement. Which of the following is the immediate nursing action required?
- A. Revise protocol for medication related incidences
 - B. Allocate senior nurses to medication assignments
 - C. Provide missed medication dosage to patient first**
 - D. Incident reporting must be given the priority
441. Nurse manager first days in new hospital will..?
- A. introduce plan and strategy to top manager in hospital
 - B. introduce himself to staff**
 - C. observation for 3 month before begin work

442. After accessing patients' medical records, which behavior nurse shows that patients confidentiality has been breached?
- A. Reviews patients medical record
 - B. Read patients care plan
 - C. Disclosing patients information**
443. During the assessment phase of a preoperative interview, the patient reports feeling nervous. The patient conveys to the nurse that a parent died in surgery due to malignant hyperthermia .to whom would this information be most important ?
- A. PACU NURSE
 - B. SCRUB NURSE
 - C. ANAESTHESIA TEAM**
 - D. CHARGE NURSE
444. a charge nurse in the nurse was noted poor staffing schedule in one of the shift. The schedule caused problems among staff members in the unit. The women the charge nurse should bring the problem to?
- A. Supervisor**
 - B. Chief nurse
 - C. Follow charge nurse
 - D. Grievance committee
445. nurse responsible for narcotic medication loses on ampule morphine. She reports the problem to nurse manager. Which of the following is the first thing she should say according to the SBAR approach?
- A. I have lost one ampule of morphine**
 - B. We should ask the nurse about the morphine
 - C. I think I left the lid of narcotics box open
 - D. We should open an investigation into problem
446. A nurse intervention a patient recently admitted to long term care facility to obtain information on the patient health perception. The nurse encourage the patient to elaborate this change. Which of the following type of question would be the most effective in this situation?
- A. Analytical
 - B. Focused
 - C. Closed
 - D. Open ended**
447. A nurse enters the room of a patient named Ahmed Saeed to administered medication but the patient not wear his identification bracelet. Which of the following is the most appropriate nursing action?
- A. Ask the patient are you Ahmed Saeed?
 - B. Ask the patient what is your name**
 - C. Ask the patient hat mediation do you take
448. Research data to be collected from children on nursing unit. What type of consent is the most appropriate ?
- A. Informed written consent
 - B. Consent signed by parents**
 - C. Consent from physician

D. Hospital administration consent

449. Nurse manager prepares unit clinical operational plan What is top priority in the plan?
- A. Infection control
 - B. Staff orientation
 - C. Quality projects
 - D. Safe patient care**
450. .What is a characteristic of democratic leadership style?
- A. More changes
 - B. Structured work
 - C. Staff participation**
 - D. No leader decisions
451. Which of the following vaccines in highly recommended to be taken by pilgrims before Haji season?
- A. Meningococcal meningitis**
 - B. Tuberculosis
 - C. Hepatitis A
 - D. Polio myelitis
452. Which of the following is the main reason the makes nurses concerned about adolescents health status?
- A. Take risky behaviors**
 - B. Consider themselves as adult
 - C. Have more health issues
 - D. Transitional period to adulthood
453. A patient fell in the bathroom and his left leg was fractured, in order to communicate information about the patient to the next shift. Which of the following documentation should be used by the nurse at the nurse at the end of the shift?
- A. Kardex record**
 - B. Assignment record
 - C. Shift report
 - D. Incident report
454. A nurse is showing stress and anxiety due to long duty hours. Which of the following should the nurse manager do ?
- A. give day off
 - B. give light tasks
 - C. offer vacation
 - D. talk about coping**

Delegation

455. The nursing assistant with 20 years of experience approaches a recently graduated nurse who recently passed the licensing examination. The nursing assistant states "the only difference between you and me is the size of our pay checks". Which of the following is the most appropriate response for the newly graduated nurse?
- A. assert a hierarchical position
 - B. emphasize the additional education received
 - C. explain the legal difference in the scope of practice

D. focus on the need to work together for quality client care

The licensed practical nurse (LPN) with 20 years of experience approaches the new graduate registered nurse (RN) during orientation. The LPN states, "The only difference between you and me is the size of our paychecks." What would be the best response for the new graduate RN to make initially?

☐ 1. Assert being the manager of the client team (C)

☐ 2. Emphasize the additional education received (C)

☐ 3. Explain the legal difference in the scope of practice (19)

☒ 4. Focus on the need to work together for quality client care (C)

Correct Answered correctly

Explanation

Team building involves recognizing that everyone has personal strengths and specific skill sets that together can be used to provide quality client care. The new graduate should recognize the contributions of the LPN and give respect to the LPN rather than initially be confrontational. Emphasizing common goals, such as safe, quality client care, is usually more effective than debating personnel qualifications.

456. A 14- nurse manager assigns tasks according to clinical competencies of the nurses. Which of the following is the management function?
- A. Delegating**
- B. Evaluating
- C. Planning
- D. Controlling
457. The head nurse of a Coronary Care Unit delegated the staff a senior nurse in that unit What initial step must the head nurse implement before?
- A. Check the hospital policies for delegating tasks**
- B. Explain the task to the senior nurse
- C. Negotiate with the senior nurse
- D. Take the signature of the senior nurse
458. What must be known on legal points of delegation when process to a new nurse?
- A. Evaluation of performance of delegate by clients
- B. Actual time it takes to complete the task by delegat
- C. Institution definition of the job description**
- D. Number of times that the delegate has previously task
459. Which of the following patient is the most appropriate to delegate to a new nurse?
- A. Patient on a strict bed rest**
- B. Patient for discharge teaching
- C. Patient scheduled for an operation
- D. Patient who is hooked to mechanical ventilation
460. A head nurse of a coronary care unit delegated staff scheduling to a senior nurse in that unit. Which of the following steps must the head nurse implement before delegating tasks?
- A. Negotiate with the senior
- B. Take signature of senior
- C. Explain task to senior
- D. Make sure hospital policies for delegating tasks**
461. Which of the following is most appropriate to delegate assistance
- A. Insertion of an oral airway

- B. Changing postoperative dressing
- C. Initial interview on a newly admitted patient
- D. Repositioning a patient to side-lying position**

462. Which of the following patients the nurse should see first?

- A. Patient complaining of muscle aches and fever
- B. Patient scheduled for electrocardiography
- C. Patient newly diagnosed with hypertension
- D. Patient with complaint of chest pain**

463. A patient with carpal tunnel syndrome (CTS) is being prepare surgery . which of the following appropriate to delegate to a assistant?

- A. Assess wrist and hand of patient for discoloration
- B. Assist patient with self care activities such as bathing**
- C. Check digits of the hand for painful tingling sensation
- D. Initiate application of a splint for immobilization during the

464. Nurse manger float RN nurse from surgical ward to CCU which patient should be assigned to this nurse?


- A. Patient just transferred after coronary angiography
- B. Patient postoperative after open heart surgery
- C. Patient on I. V lasix have congestive heart failure**
- D. Patient has ventricular tachycardia and connected to cardiac monitor

465. You are the RN charge nurse on the medical surgical unit, and you are in charge of delegating assignments for the shift. Which of the following would be the most appropriate to delegate to the LPN/LVN?

- A. A 25 year old patient who requires IV antibiotic therapy, and needs a PICC line dressing change before administering the antibiotic.
- B. A 72 year old patient who requires Lasix 20mg IV push
- C. A 50 year old patient who is 3 days post op and requires a simple dressing change**
- D. A 45 year old patient who is a newly diagnosed diabetic and requires discharge teaching

466. Nurse manger pullout nurse from medical ward to CCU which PT should be assigned to this nurse

- A. patient just transferred from cardiac catheterization**
- B. Patient just received with unstable angina on heparin infusion
- C. Patient need discharge education about coronary stenting
- D. Patient on lasix iv have acute left ventricular failure



Scope of practice	
RN	LPN/LVN
▪ Clinical assessment ▪ Initial client education ▪ Discharge education ▪ Clinical judgment ▪ Initiating blood transfusion	▪ Monitoring RN findings ▪ Reinforcing education ▪ Routine procedures (eg, catheterization) ▪ Most medication administrations ▪ Ostomy care ▪ Tube patency & enteral feeding ▪ Specific assessments*

Woman and Child

467. Ovulation occurs 14 days before the onset of the next menstrual cycle
468. A woman informed a nurse that she was never vaccinated against rubella. Which of the following is the best nursing advice?
- A. No need for her to be distress, rubella is not harmful to the fetus
 - B. The vaccine can be administered any time during her pregnancy
 - C. She can get pregnancy any time after receiving the vaccine
 - D. She should be vaccinated after delivery of the baby she get discharge**
469. A 29 year old woman had been diagnosed with a 3 cm ovarian cyst. Which of the following is the appropriate step in management?
- A. Cyst aspiration
 - B. Hormonal therapy
 - C. Cyst removal by laparoscopy
 - D. Examination after next menstruation**
470. A diabetic mother delivered a full term neonate by Caesarean section infant is admitted to the neonatal intensive care unit for observation. This infant is at risk of which of the following complication?
- A. Pneumothorax atelectasis
 - B. Hyperglycemia
 - C. Atelectasis
 - D. Hypoglycemia**
471. The midwife was assessing a 36 year old gravida 4 para 2 mother. The patient was in labour for 10 hour and had extraction. Two saturated pads were fully soaked with blood with hours been admitted in the post natal ward. Which of the following is the appropriate nursing diagnosis?
- A. Anxiety related to blood loss
 - B. Fatigue related to lack of oral intake
 - C. Activity intolerance due to discomfort
 - D. Fluid volume deficit due to uterine atony**
472. Which dietary intake should be initiated in pregnant woman?
- A. Yogurt
 - B. Soft cheese
 - C. Processed cheese**
 - D. Pasteurized milk
473. A 33 old woman presented to the ER with general weakness. The laboratory investigation indicated VIT D deficiency. Which of the following nutrient should be recommended as a good source of vitamin?
- A. Rice
 - B. Green tea
 - C. Orange juice
 - D. Fish liver oils**

474. The midwife was caring for a 30 year old gravida 3 para 3 postpartum normal delivery mother. After three hours, the patient was restless, her skin was cool, clammy and feeling thirsty. What will be midwife's initial action?
- A. Notify the doctor
 - B. Check the vital sign**
 - C. Give patient a drink
 - D. Start fundal massage
475. Which of the following nursing responsibilities should be done immediately following administration of lumbar epidural anesthesia to a woman in labor?
- A. Reposition from side to side
 - B. Administer oxygen
 - C. Assess for maternal hypotension**
476. A nurse is assessing the uterus of a G5P4 patient immediately after delivery. The nurse notes the fundus is not contracted. Which of the following is the most appropriate immediate action should be taken?
- A. Massage the fundus**
 - B. Assess the bladder
 - C. Elevate the mother's legs
 - D. Encourage the mother to void
477. A nurse is assisting during a normal vaginal delivery on a 22 year diabetic patient. The head was delivered without any complication head suddenly retracts against the perineum prompting the physician to immediately ask for the nurse assistance with this dystocia. Which of the following will be the nurse appropriate action to impacted shoulders of the infant?
- A. Fracture the infants clavicle.
 - B. Prepare patient for immediate cesarean section
 - C. Apply fundal pressure to displace anterior shoulder
 - D. Perform supra pubic pressure to release anterior shoulder**
478. At labor room, a nurse assessed the condition of the patient and gathered the following data. Cervical dilatation 2-5 minutes lasting 40-60 seconds increasing bloody show leg discomfort with heaviness what is the significance of the data?
- A. Patient on the first stage of labor**
 - B. Patient on the third stage of labor
 - C. Patient on the second stage of labor
 - D. Patient is experience a prolonged labor
479. A primigravida mother is having her baby through normal vaginal delivery. The baby is completely delivered but the mother is still experiencing the uterine contractions.
- A. Beginning of the third stage of labour**
 - B. Indication of increase in vaginal bleeding
 - C. Need for reducing the rate of intravenous oxytocin
 - D. Uterine contraction will gradually reduce then stop.
480. A mother is the outpatient Clinic for her first post natal visit on the 15th day her normal vaginal delivery. Her physical examination reveals a stable condition, breasts are soft and her sanitary napkin has bright coloured rubra. Which of the following needs further evaluation?
- A. Amount and frequency of breast feeding
 - B. Hydration level and bleeding breast feeding
 - C. Activity, exercise and resting periods
 - D. Uterine size and position**

481. A midwife is discussing the birth spacing measures with a mother whose first baby boy is a 1 year old and was born with spina bifida. The midwife has explained the possible causes of condition of the baby and the measures is necessary if the mother is planned for a next pregnancy?
- Increase iron and calcium supplements
 - Multivitamin and folic acid intake**
 - Genetic screening
 - Immunization before and during pregnancy
482. A 32 weeks pregnant patient present for her follow up appointment in the antenatal clinic. She complains of experiencing frequent heartburn. Which of the following is the most appropriate advice to the patient?
- Drink plain water between meals
 - Raise head of the bed**
 - Eat favorite foods
 - Lie down for one hour after taking food
483. A midwife visits a mother four weeks after delivery. The mother is breastfeeding her baby but she requests the midwife to suggest alternative formula milk as she has to return back to her job and her baby will stay in a day care center. Which of the following teaching plans is suitable for the mother?
- Supplementary medications with bottle feeding
 - Combined schedule of the breastfeed and top feed
 - Hygiene practices with bottle feeding
 - How to express and save milk**
484. A women in labor is progressing well. She has been diagnosed with a large fibroid in the fundus. What should the nurse observe her for after delivery?
- Thrombophlebitis
 - Postpartum depression
 - Postpartum hemorrhage**
 - Loss of bladder tone during puerperium
485. Which of the following assessment is contraindicated while providing intrapartum nursing care for a patient with HELLP syndrome?
- Heart sounds
 - Blood studies
 - Leopold's maneuvers**
 - Deep tendon reflexes
486. For which of the following issues should the nurse observe the movie closely during the 4TH stage of labor?
- Uterine irritability
 - Signs of infection
 - Signs of bleeding**
 - Unwillingness to breastfeed

in the event of abduction.

Care of the Mother

Nursing care of the mother during the fourth stage of labor focuses on observing for hemorrhage and relieving discomfort (Table 13-3).

Observe for Hemorrhage. Important assessments related to hemorrhage are the woman's vital signs, uterine fundus, bladder, lochia, and perineal and labial areas (see Chapter 17).

Vital Signs. Assess the woman's temperature when fourth-stage care begins. Blood pressure, pulse, and respirations should be assessed every 15 minutes during the first hour. A rising pulse rate is an early sign of excessive blood loss because the heart pumps faster to

487. A 32year old woman has just been told that she is pregnant she states ' As much as I love my children , I had had hoped we would not have any more " this statement reflect of which of the following related to pregnancy
- A. Anger
 - B. Denial
 - C. Guilt
 - D. Ambivalence**
488. After nurse assessed a newborn she reported that the baby has syndactyly the student nurse ask the nurse in charge what is the syndactyly ?
Which of the following is the best nursing response
- A. Fistula.
 - B. Abnormal big head
 - C. Extra finger or toes
 - D. Finger or toes wholly of partly united**
489. A newborn is diagnosed with Ventricular Septal Defect (VSD) Which of the following information should the nurse give to the newborn's mother?
- A. Cyanosis will occur most of the time during sleeping
 - B. Breast feeding is not recommended for your child
 - C. The defect might close spontaneously after 6 months**
 - D. Blood pressure is different on the child's arm and the leg
490. A couple asked the nurse which of the first investigation they should do for infertility?
Which of the following should be the proper a nurse answer?
- A. Hysterosalpingogram
 - B. Serum progesterone
 - C. Semen analysis**
 - D. Endometrial biopsy
491. A nurse is assessing a 5 month old infant who has admitted to the pediatric ward with coarctation of aorta
Which of the following is the most common assessment findings ?
- A. Cyanosis and clubbing hands
 - B. Bounding pulse and hypotonicity
 - C. Cyanosis occur frequently during and after feeding
 - D. Blood pressure is different on the arm the leg**
492. A 36 years old woman is in her 18 weeks pregnancy came to antenatal clinic and assessed by the nurse. She has three previous abortion during the first trimesters prior to pregnancy. How the nurse will document the status of the mother?
- A. Parity
 - B. Nullipara**
 - C. Multipara
 - D. Primipar
493. During vaginal examination the nurse palpated the posterior fontanel to be at the right side and upper quadrant of the maternal pelvis ?
- A. ROP
 - B. LOP.
 - C. ROA**

D. LOA

494. A 38 years old gravida 5 para 3 post natal mother delivered via caesarean section. On the fourth post operative day the mid wife noted human's sign was positive . human's sign is result by pain which of the following leg site?
- Calf**
 - Foot
 - Heel
 - Thigh
495. Which of the following vaccines are safe to administer to pregnant women?
- Measles
 - Tetanus**
 - Rubella
 - Varicella
496. Which of the following is independent function of primary health care nurse?
- Complete history and nursing assessment**
 - Referral for diagnostic evolution
 - Medical examination
 - Psychotherapy referral
497. Which of the following statement by the nurse describes the effect of clomid an ovulation inducing drug?
- Given for the first 15 days in each cycle
 - Maximum dose is 50mg daily for a month
 - Increase the risk of birth defects
 - It increases the risk of multiple pregnancy**
498. A 50 year old woman is attending the outpatient clinic for annual check up. Which of following should be reported if inspected during breast examination?
- Everted nipples
 - Symmetrical visible
 - Dimple in the left breast**
 - Right breast is slightly larger than left breast
499. A pregnant woman is 36 weeks pregnant was admitted to antenatal ward for observation after being involved in a car accident the refuses when the nurse tried to touch her abdomen to perform examination saying it is painful. She also present mild vaginal bleeding . Which of the following is the most appropriate diagnosis
- Placenta previa
 - Tubal pregnancy
 - Abruptio placenta**
 - Inevitable abortion
500. Mother of nine children, three of them with congenital anomalies and one down syndrome, she is a primary school graduate, with low financial status, she is not using any method of family planning . according, the primary health are nurse has referred her for counseling. Which of the following is a model concerned with disability as from of social injustice due to stigma, or discrimination
- Health belief
 - Biomedical

C. Sociopolitical

D. Economic

501. A primiparous mother is calling the clinic to report that her baby often regurgitates small amount of breast milk which of the following is appropriate nursing advice the mother?
- A. Elevate the head of the bed**
- B. Feed the baby when he is too hungry
- C. Attend to the clinic with the baby
- D. Put the baby in prone position after feeding
502. A 10 month old infant is admitted to the surgical ward with hydrocephalus. Which of the following indicate increased intracranial pressure?
- A. Bulging**
- B. Decrease blood pressure
- C. Rapid, shallow breathing
- D. Increase body temperature
503. A multigravida and 37 weeks pregnant mother presents to the antenatal Clinic. Her condition is stable and her baby position and heart rate are normal. Which of the following discussion is more appropriate with the mother?
- A. Observer until the 40 weeks complete**
- B. Family planning and birth spacing
- C. Arrange for caesarean section
- D. Possible of induced labor.
504. A 26 years old patient in the gynecological ward complained of pain and swelling in her episiotomy stitches on her second post-natal day. On examination the localized swelling redness foul smell and pussy discharge were identified at the stitches site after examination the nurse checked vital signs BP 116/28 HR 132 RR 28 TEM 39.8. which of the following should the nurse do prior to the patient examination by the gynecologist?
- A. Arrange dressing instrument
- B. Maintain patient privacy**
- C. Arrange stitch removal kit
- D. Collect detailed history
505. A 28-year-old primigravida who is pregnant at 16 weeks of attended at the emergency department because of dark brown discharge. investigation showed a decline in pregnancy test for which of the following is the most likely type of the spontaneous abortion?
- A. Missed**
- B. Threatened
- C. Incomplete
- D. Inevitable
506. A nurse is educating a primigravida woman who is pregnant at 30 weeks on breast feeding. Which of the following statement by the woman indicates that she needs additional teaching?
- A. Breast milk can stored at room temperature
- B. Breast feeding should be based on baby demand
- C. Baby can be held in different ways during feeding
- D. Breast feeding helps the uterus to return to pregnancy size**

507. Which of the following vaccines should be given to a 9 months old ?
- A. Hepatitis
 - B. Varicella & measles
 - C. Oral polio and bacillus cellmate Guerin
 - D. Measles and meningo conjugate quadrivalent (MCV4)**
508. nurse is instructing a female client how to do breast self-exam. Which of the followings is the best time to perform this exam?
- A. After ovulation
 - B. After the period**
 - C. Two weeks after period
 - D. Three days before period
509. A nurse is explaining pudendal block anesthesia to primigravida women who is in active labor. Which of the following relief areas identified by the woman would indicate that teaching was effective?
- A. Back
 - B. Perineum**
 - C. Fundus
 - D. Abdomen
510. Which of the following classifications of placenta previa is applicable when the placental edge is 5 cm away from the internal cervical OS?
- A. Total
 - B. Partial
 - C. Marginal**
 - D. Complete
511. A postdate pregnant woman is admitted for the induction of labour. Her fetal heart rate and vital signs are within normal range her intravenous line is maintained and she is to be started on low doses of labour inducing medication. Which of the following medication the mother is likely to receive intravenously ?
- A. Oxytocin**
 - B. Cervidil
 - C. Cytotec
 - D. Cytoxin
512. Which of the following patients are at risk for cord prolapse?
- A. Fetus that remains at high station**
 - B. Mother with oligohydramnios
 - C. Presenting part at station +
 - D. Intact membranes
513. While taking care of the a patient with a spinal cord injury, the patient suddenly complains of pounding headache upon assessment the patient was found to have diaphoresis , drop in heart and elevated blood pressure , autonomic dysreflexia is suspected and the head on the bed is elevated . which of the of the following is the most appropriate to important immediately?
- A. Notify the physician

B. Assess bladder for distension

C. Continue to monitor for next hour

514. vital statistics and control of communicable disease serve many function. Which of the following is the basic function for health?
- A. Public**
B. Mental
C. Education
D. Maternal & child
515. A 6 months old infant mother decided to wean her child. Which of the following is the best principle of weaning process?
- A. Start the weaning process by 8 month of life
B. Gradually replace one breast session at a time
C. Discontinues the nighttime feeding first
D. Allow the child to take a bottle of milk or juice bed
E.
516. A midwife is conducting a health a health education session to the primigravid mothers. The session is about antepartum care to ensure the health of the mother their babies. Which of the following statement made by mother suggest their appropriate understandings of anti-partum periods ?
- A. Beginning of labor till the baby birth
B. Diagnosis of pregnancy till the baby birth
C. Last three months of pregnancy
D. Forty days after the child is born
517. A 41 weeks pregnant was admitted to labor and delivery unit for induction with oxytocin infusion 3 hour later her contraction are 5 to 6 in 10 minutes and strong which of the following is he best nursing action?
- A. Gives an analgesic
B. Stop the oxytocin & inform the doctor
C. Gives psychological support
D. Change position to left lateral
518. A 26 year old patient in the gynecology ward complained of pain and swelling in episiotomy stitches on her second post natal day. On examination, the localized swelling, redness, foul smell and pussy discharge were identified at the stitches site after examination, the nurse checked her vital signs
BP 116/58 RR 28 HR 132 TEM 39.8
Which of the following should the nurse do prior patient examination by the gynecologist?
- A. Arrange dressing instruments
B. Maintain patient privacy
C. Arrange stitch removal kit
D. Collect detailed history
519. A woman delivered her baby boy 3 hours ago with caput succedaneum. Which of the following the nurse expects to find when she examines the newborn baby caput?
- A. Soft, fluctuant mass filed with fluid**
B. Bilateral mass on both biparietal boons
C. A mass with clear edges that end at the suture lines
D. A hard mass that is filled with blood
520. A 25 year woman was admitted to medical ward for anorexia chemotherapy the nursing diagnosis was imbalanced nutrition less than body requirements related to dysfunctional eating pattern which of the following is the best way to evaluate the outcome of nursing care of this patient?

A. Record daily weight

- B. Monitor vital signs accurately
- C. Schedule meals with family members
- D. Offer small portions of favorites

521. **Total ante natal visits ?**

10 -12 visit (every 4 weeks in the first 28 weeks , every 2 weeks in 32-36 weeks , every week from 36weeks - 40weeks)

522. **What is the antenatal assessment schedule for women the period of conception and 28 weeks of gestation?**

- A. Once a week
- B. Every 2 week
- C. Every 3 weeks
- D. Every 4 weeks**

523. **A nurse is instructing a 20years old nulligravida woman about side effect of oral contraceptive. Which of the following if stated by the woman indicates for further education?**

- A. Nausea
- B. Headache
- C. Weight gain
- D. Ovarian cancer**

524. **A 33 week pregnant mother is on her routine medications that include iron supplement , folic acid , multivitamins and calcium supplement , despite all medication her hemoglobin has not been increase since last two month, she is experiencing more fatigue and lethargy since past few weeks**

HB 80 HCT 0.22

Which of the following intervention is the most desired?

- A. Add vitamin c to increase drug absorption**
- B. Start injectable multivitamins as per regimen
- C. Advise increase intake of organ meat and fortified food
- D. Advise increased food intake by frequent small meals and snacks

525. **A primigravida attend the antenatal clinic for her routine visit the nurse performed an abdominal palpitation and found the fundus to be midway between the symphysis pubis and the umbilicus what are the weeks of gestation according to this findings ?**

- A. 8 Weeks
- B. 16 weeks**
- C. 24 weeks
- D. 28 weeks

526. **What is the meaning of the acronym A of REEDA which is used to assess the perineum after birth?**

- A. The status of edges of the perineal wound it should be closed**
- B. Presence of discharge from the perineal wound
- C. The presence of bruising of the perineal area
- D. Excessive swelling of the perineal area

527. **A pregnant woman has visited the antenatal clinic her last menstrual date was on 10th August this year. Her menstrual cycle was regular at 28 day. What is the expected date of delivery?**

- A. 17th may next year**
- B. 20st may next year
- C. 23rd may next year
- D. 25th may next year

528. A 30 years old woman , gravid 3 para 2 in her 20 weeks pregnancy was complaining of increase leucorrhea during past weeks the nurse reassured the mother and explained the reason of this secretions ?
- A. Vaginal infection
 - B. Expansion of uterus
 - C. Stimulation of cervix
 - D. Production of estrogen**
529. Lochia red sometimes happens during postpartum to bleed heavily with Foul smell for the first three to ten postpartum days, the nurse should expect that may to
- A. Lochia pink
 - B. Sign of abnormal hemoglobin
 - C. Infection**
 - D. Bleeding
530. Which of the following assessment findings indicates laceration of the canal in the fourth stage of labor?
- A. Red-brown lochia
 - B. Firm contracted uterus
 - C. Fundus is palpated at the level of the umbilicus
 - D. More than 1 saturated perineal pad per hour**
531. Which of the following must be checked during the third day of post-partum period by the nurse?
- A. The flow of milk
 - B. Uterine prolapse
 - C. Exclusive breast feeding**
 - D. Complete involution of uterus
532. A child with sever bronchial asthma is started on prednisone. Which of the following is a side effect of prednisone that might be exhibited by the child?
- A. Anorexia**
 - B. Weight loss
 - C. Anemia and fever
 - D. Neurologic symptoms
533. A woman was rushed to the emergency room because if vaginal bleeding mild cramps, tenderness over the uterus, and closed cervix. After the assessment, it was suspected that the patient may be lead to inevitable abortion. What is the possible management of this patient?
- A. bed rest
 - B. surgical management
 - C. induction of oxytocin
 - D. dilatation and curettage**
534. A 28 weeks pregnant mother is in the antenatal clinic with the complaint of vaginal irritation itching and thick white secretion . she is diagnosed as having vaginal candidiasis she had been having the same infection before the pregnancy for which she was prescribed the vaginal cream she is using the same cream for the past few days but her symptoms are not relieved what need to be emphasized to mother
- A. change clothes daily use sanitary napkins
 - B. Avoid self treatment and seek doctor advise**
 - C. Use medicated bubble baths and reduce activates
 - D. Was with warm water every two hours

535. Which of the following nursing responsibilities should be done immediately following administration of lumbar epidural anesthesia to a woman in labour?
- reposition from side to side
 - administer oxygen
 - increase IV fluid as indicated
 - assess for maternal hypotension**
536. the antenatal clinic, a 9-month pregnant woman notifies the nurse that e is concerned that she gained 15Kg during this pregnancy and she is and needs to start a diet program. which of the following is the best nursing response?
- This weight gain is normal during pregnancy**
 - You can reduce the carbohydrates in your diet
 - Wait till you have the baby and start a diet program
 - This weight is lower than you should gain during pregnancy
537. Intrauterine growth curves were used to classify a 32-week-old preterm new-born. Birth weight and gestational age shows the infant's growth rate falls below the 10th percentile. What is the priority nursing diagnosis for a new-born with small for gestational age?
- Risk for injury related to impaired gluconeogenesis
 - Risk for impaired gas exchange related to meconium aspiration
 - Risk for ineffective thermoregulation related to lack of subcutaneous fat**
 - Risk for altered nutrition less than body requirement related to increased metabolic needs
538. A nurse is leading an educational session on the correct use of oral contraceptives. One of the attendees ask the nurse what to do if she missed taking an oral contraceptive for one day? Which of the following should be the nurse advice to her?
- Continue as usual with no back up contraception
 - Take an active pill immediately and take the next pill at the usual time**
 - Take two pills as soon as possible and then one pill daily at the usual time
 - Use back up contraception such as a condom for the next 7 days
539. Which of the following represents secondary infertility?
- Male infertility
 - A couple that does not conceive
 - Infertility that occurs after previous pregnancy**
 - Infertility lasts for more than 3 years
540. A women patient with a body mass index of 35 is admitted to ward. Which of the following nursing diagnosis, related to Imbalance the most appropriate :
- Activity intolerance
 - Less than body requirement
 - More than body requirement**
 - Deficient knowledge on normal nutrition
541. A nurse is providing health education for a primigravida patient who has gestational diabetes mellitus. Which of the following statements made by the patient indicates the need for additional education?
- "I should not do exercises"**
 - "I should follow the prescribed diet"
 - "+I should monitor my blood glucose"

D. "I should report any sign of infection"

542. Gestational diabetes is high blood sugar (diabetes) that starts or is found during pregnancy. The best time for glucose screening tests during pregnancy is:

- A. 7 – 12 weeks
- B. 17 – 18 weeks
- C. 24 – 28 weeks**
- D. 20 – 35 weeks

543. laboring women desires to participate in her cesarean and have pain control. Which of the following methods would satisfy the women's needs

- A. Epidural block**
- B. Pudendal block
- C. Meperidine injection
- D. General anesthesia

544. which of the following statement described the latent phase?

- A. First 3 cm of cervical dilatation**
- B. From onset of labour to full dilatation
- C. When the cervix is 100% effaced
- D. Time of progress from 4 cm to 7 cm

545. A nurse is performing an assessment of a women who is delivery Which assessment finding would indicates a need to physician?

- A. Hemoglobin of 11.0 g/dl
- B. White blood cell count of 12,000
- C. Fetal heart rate of 180 beats per minute**
- D. Maternal pulse rate of 85 beats per minute

Fetal heart rate of 180 beats per minute

A normal fetal heart rate is 120-160 beats per minute. A count of 180 beats per minute could indicate fetal distress and would warrant physician notification. By full term, a normal maternal hemoglobin range is 11-13 g/dL as a

546. Amniotic fluid importance . surrounds ,cushion ,protect baby . allow the fetus to move freely , maintain body temperature to the fetus , and contain urine from the fetus cushion

547. A 34-weeks-pregnant mother experiences a sudden gush o from her vagina and mild uterine contractions. She informs about her condition and requests if she could wait until the delivery. Which of the following is the best desired response for report to the hospital?

- A. Intravenous fluids and medicines need to be administer
- B. Observation is necessary to identify premature labor
- C. Pain and fluid flow both need to be controlled
- D. Fetal heart sound monitoring is necessary**

548. During A Vaginal Delivery of Woman 38 Years Old the Nurse Should Consider the Risk of Which of The Following

- A. Acute bleeding and coma
- B. Low potassium
- C. Brain Injury
- D. Fetal heart rate**

549. Pregnant woman 34 weeks has hypertension this problem is identified as?

- A. Hypertension
- B. pregnancy induced DM
- C. PIH (Pregnancy induced hypertension)**

D. placenta Previa

550. A nurse is explaining to the nursing students working on the antepartum unit how to assess edema. Which edema assessment score indicates edema of the lower extremities, face, hands, and sacral area?

- A. +1
- B. +2
- C. +3**
- D. +4

551. 38weeks-pregnant woman complains that she has been craving to eat non food substance What is the term that describes this pregnant mother's condition?

- A. Pica**
- B. Bulimia
- C. Anorexia
- D. Binge eating

552. pregnant client is making her first Antepartum visit. She has a two year old son born at 40 weeks. a 5 year old daughter born at 38 weeks and 7 year old twin daughters born at 35 weeks. She had a spontaneous abortion 3 years ago at 10 weeks. Using the GTPAL format the nurse should identify that the client is:

- A. G4 T3 P2 A1 L4
- B. G5 T2 P2 A1 L4
- C. G5 T2 P1 A1 L4**
- D. G4 T3 P1 A1 L4

553. multiparous woman is admitted to postpartum ward after vaginal delivery. Assessment showed, lochia: steady trickle of bright red blood and ndus: firm.

Blood pressure 110/70 mmHg
/min
Respiratory rate
Temperature 37.5

Heart rate 80
20 /min

Which of the following is the most likely diagnosis?

- A. Endometritis
- B. Uterine atony
- C. Vulvar hematoma
- D. laceration of the genital tract**

554. A 7-year-old insulin dependent diabetic mother has delivered normally in 38 gestational weeks. The nurse was assessing the insulin requirement this mother after delivery. What is the insulin requirement for this patient?

- A. Higher than before pregnancy
- B. No changes in insulin requirement
- C. Lower than when she was pregnant**
- D. Slightly increased than before deliver

555. Mother of nine children, three of them with congenital anomalies and one down syndrome; she is a primary school graduate, with low financial status. She is not using any method of family planning. Accordingly, the primary health care nurse has referred her for counseling.

Which of the following is a barrier facing the team responsible for providing health services to handicapped individuals?

- A. Sensory limitations
- B. Rigid rules

C. Developmental disabilities

D. Deafness & hearing limitations

556. Mother of nine children, three of them with congenital anomalies and one down syndrome; she is primary school graduate, with low financial status. She is not using any method of family planning. So, the primary health care nurse has referred her for counseling. Which of the following is the best health education method that can be used?
- A. community organization
B. individual counseling
C. group discussion
D. health class
557. multiparous patient on day 1 postpartum is asking the nurse to send her baby to the nursery so she can sleep. What is the most likely phase of psychological adaptation?
- A. Taking-in**
B. Letting-go
C. Taking-go
D. Letting-in
558. A 65-year-old women visited the gynecological outpatient history reveals that she had 3 pregnancies, one abortion gestational age, had 2 normal deliveries. She smokes 20 Her complaint is that she wets herself when she cough embarrassing for her? Which of the following can be considered as risk factors pelvic floor muscles?
- A. Chronic coughing**
B. Diabetes mellitus
C. Excessive spot
D. Sedentary life style
559. At is the recommended weight gain during pregnancy of a woman with MI of < 18.5?
- A. 12.5 -18 Kg**
B. 11.5 -16 Kg
C. 7 -11.5 Kg
D. 5- 9 Kg
560. A 25-year-old mother gravid 2 para q came for a routine check Antenatal Clinic. The nurse assessed fetal heart rate for the 38 pregnant mother. What is the expected normal fetal heart rate per minute?
- A. 90
B. 100
C. 140
D. 170
561. A woman has polycystic ovary syndrome which is 3cm in size what should the nurse do?
- A. A repeat the ultrasound after menstruation**
B. Remove it by laparoscopy.
C. Give the medication
D. None
562. Nancy, a primipara who is due to deliver next week, calls and tells the nurse she has been having contractions every four minutes for an hour. Before asking any further questions, the nurse confirms that Nancy knows how to correctly time contractions, which is
- A. from the end of one contraction to the beginning of the next.

B. from the beginning of one contraction to the end of the next.

C. from the end of one contraction to the end of the next.

D. from the beginning of one contraction to the beginning of the next

563. A 22-year-old gravida 2 para 1 with gestational age 38 week admitted to the hospital. The chief complaint is decreased the fetal non-stress test revealed decreased variability and fetal movement. The next morning as part of the antenatal the nurse checks the fetal heart rate by Doppler Sonicaid decreased the fetal heart rate to less than 100 /min. which of the following action the nurse should do first?

A. Reassure the mother that the FHR is Ok

B. Notify immediately the physician or midwife

C. Reposition the patient to left lateral position

D. Ask the mother about the pattern of fetal movement

564. Pregnant patient admission to triage with abdominal pain on assessment the FHR found 70 p/m. what is the first action for the nurse?

A. Call the doctor

B. Report the finding

C. Put the patient in left lateral position

565. 25 weeks-pregnant, primary gravid woman is on her first antenatal visit, completing physical examination and history, the midwife found out her husband has sickle cell anemia minor. What should be the most appropriate plan of care?

A. Genetic counselling

B. Identify severity of disease

C. Discuss the chances of transferring

D. Amniocentesis to identify genetic abnormality

566. Which of the following statements describe puerperal infection ?

A. presence of fever of 38 or higher from 2 days to 10 days

B. presence of fever of 40

567. When would the nurse schedule a woman for a glucose tolerance test?

A. 6th -10th week of pregnancy

B. 12th -16th week of pregnancy

C. 24th -28th week of pregnancy

D. 32th -36th week of pregnancy

568. A new-born has a diagnosed Developmental Dysplasia of the Hip (DDH) and is using a Pavlik Harness as treatment. Which of the following mechanical factors is associated with DDH?

A. Intrauterine breech position

B. Caesarean section

C. Small infant size

D. Single fetus

569. The relative of a 25-year-old post-partum patient stopped her from taking bath until the 40 days provided with the food by the family that was of saturated fat and a liter of reduced full asking, she said the diet was good for the milk Which of the following should be the prioritized patient's understanding about?

A. A healthy and balanced diet is beneficial

- B. Breast feed is not affected by mother's diet
- C. Maintaining personal hygiene is most**
- D. Rest and relaxation is necessary to regain

570. When is the ideal time to administer analgesia to a women in labour?

- A. A soon as she requests analgesia
- B. When labour is well established
- C. When the women enters into transition phase
- D. when the women progress from latent to active phase**

571. A patient height is 183 cm and weight is 63.5 kg BMI 19. What is the patient weight classified is ?

- A. Underweight
- B. Normal**
- C. Overweight
- D. Obese

572. Which of the following would the nurse included in her discharge education for a postpartum patient regarding signs and symptoms that should be reported immediately?

- A. Lochia rubra persists**
- B. Nipples become red
- C. Lochia decreases in amount
- D. After pains increase with breast feeding

573. Every how many minutes to check fetal pulse with second stage of labor Every ?

- A. 5min**
- B. 10min
- C. 15min
- D. 20min

574. When the nurse assessed the fundus of a multiparous mother who delivered 2 hours ago, she found the following Level: 2 cm above the umbilicus. Position: deviated to the right. Consistency: Not well contracted what is the next nursing action after massaging the fundus until it becomes firm?

- A. Assess vital signs
- B. Increase V fluids
- C. Evacuate the bladder**
- D. Ask the mother to res

575. A 32 years old multigravida woman presents to the outpatient clinic complaining of dysmenorrhea and menorrhagia. She had been diagnosed with uterine fibroids and blood studies has been ordered for her. Which of the following results should the nurse report?

- A. hematocrit, 37%
- B. hemoglobin, 9 g/dl**
- C. white blood cell count, 10,000 cells/mm³
- D. platelets count, 300,000 platelets/ microliter

Mastitis question

Lactational mastitis (infection and inflammation of breast tissue) may result from inadequate milk duct drainage or poor breastfeeding technique. **Bacteria** from the infant's nasopharynx or mother's skin can enter the nipple, especially if it is damaged, and multiply in **stagnant milk**. Manifestations include fever, muscle aches, and breast pain and inflammation (eg, warmth, redness, edema).

Staphylococcus aureus is the most common causative organism and requires **antibiotic treatment** (eg, dicloxacillin, cephalixin). In addition, the nurse should encourage the client to:

- **Continue breastfeeding** frequently (ie, every 2-3 hr) to ensure adequate milk drainage.
- Ensure proper breastfeeding technique (eg, alternate newborn **feeding positions**, **proper latch**).
- Apply **warm compresses** and massage the breast to facilitate complete emptying. Cool compresses can also be used between breastfeeding as needed for comfort.
- Ensure adequate rest, nutrition, and hydration.
- Relieve pain and inflammation with **analgesics** compatible with breastfeeding (eg, acetaminophen, ibuprofen).
- **Wash hands** before and after feeding.

غيث

Breast engorgement is often painful. The management of engorgement varies based on the client's breastfeeding status; for clients who choose **not to breastfeed**, treatment focuses on managing symptoms while promoting **reduced milk production**. Comfort measures include:

- Applying **ice packs to both breasts for 15-20 minutes** every 3-4 hours to reduce blood flow and swelling.
- Applying **chilled, fresh cabbage leaves** to both breasts, replacing with fresh leaves after they wilt. The mechanism of action is unclear but may be related to the cool temperature or to phytoestrogens from the leaves.
- Taking an **anti-inflammatory analgesic** (eg, ibuprofen) as directed to reduce pain.
- Maintaining firm breast support (eg, **supportive bra**, breast binder) until milk flow is diminished.

غيث

Lactational mastitis

Erythema,
induration
& swelling

Lobules
Clogged, inflamed
milk ducts
Clogged nipple pores



غيث

Proper breastfeeding latch

Lips sealed completely
around most of areola
Lower lip turned
outward against
breast

Nose close
to breast

Chin touching
breast

Breastfeeding positions




Cradle hold



Cross cradle hold



Football hold



Lying down


576. **Mastitis** caused by **Streptococcus Aurous**
577. A nurse is providing instructions to a mother who has been diagnosed with mastitis. Which of the following statements if made by the mother indicates a need for further teaching?
- A. I need to take antibiotics. And I should begin to feel better in 24-48 hours.
 - B. I can use analgesics to assist in alleviating some of the discomfort.
 - C. I need to wear a supportive bra to relieve the discomfort.
 - D. "I need to stop breastfeeding until this condition resolves."**
578. A primiparous mother who has mastitis is asking the nurse Feeding Which of the following would be the proper nursing responthe organism of mastitis?
- A. It will not affect the newborn**
 - B. It will be deactivated by salivation
 - C. It will not be expressed in breast milk
 - D. It will be killed by immunoglobulins in breast milk
579. 41- A nurse is giving health education for a mother who has mastitis. Which of the following if stated by the mother about what she needs to do, indicate that additional education is needed?
- A. Take antibiotics
 - B. Use analgesics
 - C. Wear a supportive bra
 - D. Stop breast-feeding**
580. A 23-year-old vaginal delivery primigravida mother was discomfort due to breast engorgement on the second post The mother complained of pain on the breast site and the able to suck the milk. Which of the following will relief the mother's discomfort?
- A. Breast binder
 - B. Well-fitting brassiere
 - C. Encourage breast feeding**
 - D. Lactation suppressing medication
581. The nurse was planning care for a 25-year-old primigravida post-partum mother who had engorgement due to poor feeding technique. the left breast appeared red and swollen and was diagnosed as Which of the following is the best education for the mother
- A. Avoid wearing brassiere
 - B. Begin suckling on the right breast
 - C. Stop pumping milk from the left breast
 - D. Take antibiotics till the soreness subside**
582. After 3 days of breast feeding a post partial patient reports nipple soreness. To relieve her discomfort the nurse should suggest that she:
- A. lubricate her nipples with expressed milk before feeding.**
 - B. dry her nipples with soft towel after feeding.
 - C. Apply warm compresses to her nipples just before feeding.
 - D. Apply soap directly to her nipples, and then rinse

 Measures that help relieve nipple soreness in a breast-feeding patient include lubrication the nipples with a few drops of expressed milk before feedings, applying ice  compresses just before feeding, letting the nipples air dry after feedings, and avoiding the use of soap on the nipples.

583. The nurse is teaching a newly delivered mother about breast feeding which of the following statements indicates that the mother need further teaching?
- A. I will empty my breast and use an alternate breast at
 - B. I will take a daily shower and clean the nipples with
 - C. I will let my nipples dry after feeding to prevent nipple tenderness
 - D. I will always wear a supportive wireless bra
584. Mastitis is an infection of the breast that occurs most often 2-4 days after childbirth. Which of the following is considered first line treatment of mastitis?
- A. Drainage of breast abscess
 - B. Antibiotic therapy and cessation of breast feeding
 - C. Antibiotic therapy and continuation of breast feeding
 - D. Advice mother to stop breastfeeding until infection is clean
585. A 30-year-old primigravida postnatal mother come to the clinic on the She complained of feeling shivering, redness, swelling and pain in her right Breast. Blood pressure 90/62 mmHg Heart rate 84/min Respiratory rate 18/min Temperature 38.8 °C Which is the most common organism that causes mastitis?
- A. Pseudomonas
 - B. Escherichia coli
 - C. Group B streptococcus
 - D. Staphylococcus aureus
586. Immediately after delivery of the placenta the mother starts bleeding profusely from the birth canal. Which of the following is the nurse's immediate action
- A. Administers oxygen by face mask
 - B. Monitors vital signs continuously
 - C. Increase the rate of intravenous infusion
 - D. Rubs the fundus to stimulate contraction
587. postpartum psychological phases sequences?.
- A. Taking in , taking hold ,letting go
 - B. Taking in , taking hold , letting hold
 - C. Taking hold, taking in , letting hold
 - D. Letting go, taking hold , taking in
588. Which of the following statement indicate nursing action during the first hour after delivery of the placenta?
- A. Monitor of mothers hemoglobin
 - B. Assess maternal vital signs every 15 minutes
 - C. Ensure that the mother mobilize and empty her bladder
 - D. Administer 10 units of oxytocin via IV line to ensure uterus is well contracted.

589. The nurse is teaching a 32-week pregnant women how to distinguish between pre-labor (false) contractions and true labor contractions. Which statement about pre-labor contraction is accurate?
- They are regular and increase gradually
 - They are felt in the abdomen**
 - They start at the back and radiate to the abdomen
 - They become more intense during walking
590. Azithromycin is prescribed for an adolescent female who has pneumonia and recurrent chlamydia . What information is most important for the nurse to provide the
- Use two forms of contraception while taking this drug**
 - Have partners screened for human immunodeficiency
 - Decrease intake of high-fat foods, caffeine, and alcohol
 - Report a sudden onset arthralgia to the healthcare provider
591. A mother, who is planned for the labour induction, is started on intravenous medication. She is in the first stage of her labour and is having regular and increasingly stronger uterine contractions. Her cervix is 1 cm dilated for the past few hours; both the mother and the baby are being monitored. Which of the following signs should alert the midwife?
- Baby's head not engaged
 - Decreasing heart rate of the baby**
 - Mother's blood pressure 110/60 mmHg
 - Mother's perspiration and increased thirst
592. A nurse is teaching a 26-year-old primigravida who are 33 weeks pregnant on the how to use a kick chart. Which of the following statements will indicate that she understand the nurse's teaching?
- Fetal movements must be counted three times per day
 - Fetal movements are felt best when the women is on her right side
 - Fetal movements is a reassuring sign which indicates that the fetus healthy**
 - The kick chart is used to record fetal movement for the first time during pregnancy
593. Which test should be performed to screen for cervical neoplasia during antenatal assessment?
- Papanicolaou (PAP)**
 - Vaginal rectal culture
 - Rapid plasma regain test (PPR)
 - Venereal disease research laboratory test (VDRL)
594. A postpartum women who was admitted for 24 hours, community nurse visited her after 2 days of discharge. Which of these are abnormal findings?
- Frequent urination
 - Lochia serosa
 - Uterus below symphysis pubis**
 - Breast full of milk .
595. Second day post delivery which blood color will be the lochia
- Locia rubra**
 - Lochia serosa
 - Lochia alpa

Answer
D. Use two forms of contraception while taking this drug.
من جروب
اجعل من جروب
Rationale
Antibiotics, especially broad spectrum drugs like azithromycin, decrease the effectiveness of oral contraceptives and some spermicides, so the adolescent should be encouraged to use at least two forms of contraception to prevent pregnancy. (A, B, and C) provide important information but do not address the possibility of becoming



596. A 30 year old married woman has dilatation and curettage as a therapeutic abortion. A nurse was preparing for discharge instruction. Which of the following should the nurse include in discharge instruction?
- A. Taking high protein diet
 - B. Use tampons during swimming
 - C. Avoid sexual intercourse for two weeks**
 - D. Continue on bedrest for two weeks at home
597. While the nurse is performing postpartum assessment for a primiparous mother delivered 2 hours ago, she found the following:
- Fundus
level: Midway between symphysis pubis and umbilicus
Position: At the mid line
Consistency: Contracted
Lochia: Constantly
- Which of the following is the most likely diagnosis?
- A. full bladder
 - B. Perineal hematoma
 - C. Birth canal laceration**
 - D. Retained placental part
598. What the purpose of a late vaginal_ rectal culture during ante natal screening ?
- A. To screen for group B streptococci**
 - B. To detect a Chlamydia infection
 - C. To screen for gonorrhea
 - D. To screen for syphilis
599. Which of the following hormones are responsible for implantation of ovum inside Uterine wall??
- A. Progesterone and estrogen**
 - B. LH and. FSH Hormones
 - C. Oxytocin
600. Which hormone is responsible for uterine wall and breast enlargement during pregnancy??
- A. Progestron
 - B. Estrogen**
 - C. HCG
601. A woman at 24 weeks gestational age has fever body ache and has coughing last 5 days she is sent to hospital with admission prescriptions for H1N1influenza . which prescription has the highest priority ?
- A. Assign private room**
 - B. Vital signs every 4 hours
 - C. Obtain specimens for culture
 - D. Ringer lactate IV
602. A woman was diagnosed gestational trophoblastics disease what is the lab investigation was done to diagnose condition ?
- A. Cervical pap smear
 - B. Serum HCG levels**
 - C. Serum estrogen level
 - D. Plasma thyroxin level

603. A community Nursing nurse is planning to conduct prenatal teaching and community assembly for pregnant adolescents. Which teaching strategy would be most effective?
- Offering open sessions for pregnant adolescents and anyone else who wants to attend
 - Designing posters that girls can view individually in community Nursing centre
 - Preparing group class sessions for teaching pregnant adolescents together**
 - Conducting one to one teaching sessions for both mothers and daughters
604. Uterus size after delivery is ?
- 900 grams**
 - 1000 grams
 - 1400grams
 - 1600 grams
605. A pregnant women G1 P0 vaginal delivery observed in the second postpartum day that the perineal pad saturated with bright red lochia rubra what is the priority nursing intervention?
- Massage fundus
 - Obtain vital signs
 - Inform physician
 - Inquire about time of pervious saturated perineal pads**
606. A 32 weeks pregnant patient present for her follow up appointment in the antenatal clinic. She complains of experiencing frequent heartburn. Which of the following is the most appropriate advice to the patient?
- Drink plain water between meals
 - Raise head of the bed**
 - Eat favorite foods
 - Lie down for one hour after taking food
607. A midwife visits a mother four weeks after delivery. The mother is breastfeeding her baby but she requests the midwife to suggest alternative formula milk as she has to return back to her job and her baby will stay in a day care center. Which of the following teaching plans is suitable for the mother?
- Supplementary medications with bottle feeding
 - Combined schedule of the breastfeed and top feed
 - Hygiene practices with bottle feeding
 - How to express and save milk**
608. A multipara mother complained of small vulva with swelling following vaginal delivery of a baby weight 3.8 Kg What is the initial nursing action should the nurse advise the mother to perform?
- Apply ice pack**
 - Maintain bed rest
 - Administer analgesics
 - Encourage fluid intake
609. intrauterine growth curves were used to classify a 32-week-old preterm new-born. Birth weight and gestational age shows the infant's growth rate falls below the 10th percentile. What is the priority nursing diagnosis for a new-born with small for gestational age?
- Risk for injury related to impaired gluconeogenesis
 - Risk for impaired gas exchange related to meconium aspiration
 - Risk for ineffective thermoregulation related to lack of subcutaneous fat**
 - Risk for altered nutrition less than body requirement related to increased metabolic needs

610. A primigravida mother is assisted out of the bed a few normal vaginal delivery. She is taken to the bathroom cleaning herself and to pass urine. She has difficulty in commode seat and is having no urge to urinate. Which of the following intervention is the most desired?
- Teach kegel's exercises
 - Give warm sits bath first
 - Pour warm water over vulva**
 - Identify possible perineal tears
611. The nurse is assessing a young-adult pregnant client with no allergies who has tested positive for gonorrhea. Which of the following medications should the nurse expect to be part of the treatment plan?
- Tetracycline
 - Ciprofloxacin
 - Azithromycin
 - Ceftriaxone**
612. A pregnant mother at early pregnancy was admitted in Emergency Room with leakage of amniotic fluid, vaginal bleeding and lower dominal cramping pain. What is the possible ?diagnosis should the nurse suspect
- Missed
 - Inevitable**
 - Incomplete
 - Threatened
613. A nurse in the postnatal ward is assigned for a multiparous patient has just delivered a healthy newborn. When should the nurse plan to take the patient vital signs?
- Every hour for the first 2 hours
 - Every 30 minutes during the first hour
 - Every 15 minutes during the first hour**
 - Every 5 minutes for the first 30 minutes
614. which valuable information can be obtained from perform abdominal palpation (lepoids maneuvers) during the later pregnancy?
- Weight of the fetus
 - Gestational age of the fetus
 - Number of fetus in current pregnancy
 - Location and presentation of the fetus**
615. A nurse who is caring for a woman in labor, prepares to auscultate fetal heart rate by using Doppler ultrasound device. How does the nurse determine that the fatel heart sounds are correctly?
- Notify if the heart rate is greater than 140 BPM
 - Placing the diaphragm of the Doppler on the woman abdomen
 - Palpating the maternal radial pulse while listening to the rate**
 - Performing Leopold's manoeuvres first to determine the fetal heart
616. A client who is gravida 1, para 0 is admitted in labor. Her cervix is 100% effaced, and she is dilated to 3 cm. Her fetus is at +1 station. The nurse is aware that the fetus' head is:
- Not yet engaged
 - Entering the pelvic inlet
 - Below the ischial spines**
 - Visible at the vaginal opening

to identify a distended or displaced uterus. The uterine fundus is felt above the symphysis pubis with a lateral displacement of the uterus. The nurse also notes an increase in the amount of lochia because the uterus is unable to contract effectively. The bladder is placed, bulges above the symphysis pubis, and feels "boggy" on palpation. Patients experiencing urinary retention because of absence of the urge to void can be helped by assisted early ambulation to the toilet and other measures such as running the water from the lavatory faucet. If ambulation is not possible, the nurse can pour warm water over the vulva and perineal area to help relax the urethral sphincter. Because of the risk of urinary infection associated with urinary stasis, catheterization may be necessary if the patient is unable to void. Constipation commonly occurs because of slowed peristalsis associated with pregnancy hormones and childbirth anesthesia. In addition, perineal discomfort, fear of suture separation at the episiotomy site, and incisional pain (after a cesarean birth) may contribute to decreased frequency in bowel movements. To prevent constipation, nurses should encourage patients to consume foods high in fiber and fluids. Adequate fluid intake also helps to decrease the risk of urinary infection. The nurse should instruct the patient to use a toilet brush to gently clean the perineal area and use of a perineal bottle to gently irrigate the area. The nurse should also use of a perineal bottle to gently irrigate the area. The nurse should also use of a perineal bottle to gently irrigate the area.

617. Less than 24 hours after postpartum calls a nurse in the Mother/Infant Unit. She reports that she has very heavy bright-red bleeding. being discharged, a mother who is newly What would be the best advice to give her?
- A. "Don't worry about it; it is normal."
 - B. "Call your doctor and ask what to do."
 - C. **"Lie down, massage lower abdomen. If it does not work then come to the hospital immediately."**
 - D. "Lie down for about an hour. Then check your bleeding, if it is still as heavy, call me back."
618. A nurse was educating a group of woman on prevention of infection. The nurse asked each woman to state one preventive of vaginal infection. Which woman needs more education?
- A. First woman "keep vaginal area clean & dry"
 - B. Second woman "wear cotton under wear"
 - C. Third woman "wipe from front to back after urination or
 - D. **Fourth woman "Do vaginal douche twice a day "**
619. 30 years old women absent her menstruation for 5 months and her menstrual cycle come every 28 days, she controlled her diet and do heavy exercise this women condition?
- A. Pregnancy
 - B. Primary amenorrhea
 - C. **Secondary amenorrhea**
620. **Molar pregnancy:** is an abnormal form of pregnancy in which a non-viable fertilized egg implants in the uterus and will fail to come to term.
621. Molar pregnancy sings **Rapid increase of uterine growth**
622. A 28-year-old pregnant woman at 9 weeks presents to the o with vaginal bleeding. During assessment, the nurse found height is 12cm. Which of the following is the most likely diagnosis?
- A. Placenta previa
 - B. Abruptio placenta
 - C. Ectopic pregnancy
 - D. **Hydatidiform mole**
623. During a prenatal examination. The nurse draws blood from a young Rh negative client and explain that an indirect Coombs test will be performed to predict whether the fetus is at risk for:
- A. **acute hemolytic disease**
 - B. respiratory distress syndrome
 - C. Protein metabolic deficiency.
 - D. pathologic hyperbilirubinemia
624. Which of the following statements will indicate that she understand the nurse's teaching?
A nurse is teaching a 26-year-old primigravida who are 33 weeks pregnant on the how to use a kick chart.
- A. Fetal movements must be counted three times per day
 - B. Fetal movements are felt best when the women is on her right side
 - C. **Fetal movements is a reassuring sign which indicates that the fetus healthy**
 - D. The kick chart is used to record fetal movement for the first time during pregnancy

625. The antenatal clinic nurse was assessing a 32yaers old gravida 2, para1 pregnant mother on the fundal height at 36 weeks. What is the expected fundus position?
- A. Umbilicus
 - B. Xiphoid process**
 - C. Symphysis pubis
626. A woman is at 30 weeks gestational age admitted to antenatal with premature rupture of membrane . the nurse administered Dexamethasone to her according doctor prescription. She ask what is the Drug for. Which of the following the best answer ?
- A. To promote fetal lung maturation**
 - B. Prevention of chorioamnionitis
 - C. To increase uteroplacental exchange
 - D. Treatment of fetal respiratory distress
627. A 25 years old patient with history of amenorrhea for two month was admitted for hydatiform mole investigation. Which signs and symptoms would the nurse observe ?
- A. Hypotension
 - B. Hyperglycemia
 - C. Rapid uterine growth**
 - D. Painful uterine contraction
628. A women was discharged from gynecological ward after gestational trophoblastic disease (molar pregnancy) Which of the following is the best advice to give her?
- A. Never to fall pregnant again
 - B. To request the doctor to sterilize her
 - C. To consider having her uterus removed
 - D. to avoid falling pregnant for at least one year**
629. Pregnant woman signs and symptoms of hypoglycemia
- A. Blurred vision**
 - B. Dry mouth
 - C. Poly urea
 - D. Flushing
630. postpartum mother is to be discharged on the second day of her forceps delivery. She had sutures on her vaginal and perineum tears. She is breastfeeding her baby and eating the special food provided by her family. what discharge teaching needs more emphasis?
- A. Diet management and exercise plan
 - B. Newborn care and vaccination records
 - C. Hygiene practices and alert signs to report**
 - D. Family planning and child growth monitoring
631. When performing a postpartum assessment on a women the presence of clots in the lochia. The nurse examines the that they are larger than 1 cm, Which of the following nursing actions is most appropriate?
- A. Document the findings
 - B. Notify the physician**
 - C. Reassess the client in 2 hours
 - D. Encourage increased intake of fluids

632. A pregnant woman of 15 week gestation age and Rh –ve has an abortion admitted in the ward what should the nurse do for this woman ?
- Administer Rheum within 72hour**
 - Do not give Rheum since it is not used with abortion
 - Do not give Rheum since the pregnancy is more than 12
 - Make certain she received Rheum on her first clinic visit
633. Lochia red fleshy odor mean :
- Normal**
 - Infection
 - Bleeding
634. The nurse was educating a postpartum woman during discharge about importance of breast feeding. Which of the following if said by the women, indicates the need for further education?
- Breast milk is nutritionally balanced
 - Breast milk reduces the risk of infection
 - Breast feeding promotes mother-child bonding
 - Breast feeding prevents pregnancy**
635. Multiparous mother is attending at the outpatient clinic 2 weeks after er delivery for follow up. While the nurse is assessing the mother, she would not palpate the fundus. Which of the following is the most appropriate nursing action?
- Document normal finding**
 - Massage the fundus to be firm
 - Assess lochia amount and color
 - Admit the mother to the hospital
636. A women breastfeed her infant one or two hours and her infant cries most of the time and she feels pain in her breast. Which of the following instructions are appropriate for the nurse to give the mother:
- Regulate breast feeding every 3 hours**
 - That's normal feeding problem
 - Shift to bottle feeding
 - Start weaning your baby
637. A 17-years-old mother presented to the primary health after delivery. She is suffering from fatigue, anemia, fever vaginal discharge (see lab results)
- 81/50 mmHg Blood pressure
 98 /min Heart rate
 26 /min Respiratory rate
 39.6C Temperature
 Result Test
 4.6 RBC 4.7-6.1 × 10¹²/L (Male) Normal Values 4.2-5.4 ×10¹² /L (Female)
 88 Hb 130-170 g/L (Male) 120-160 g/L (Female)
 2.50 Calcium 2.15-2.62 mmol/L
- Which of the following is considered as the main maternal postpartum haemorrhage complication ?
- Death**
 - Candidacies
 - Cervical cancer
 - Uterine prolapsed

638. A pregnant woman she come to gynocological word she have vaginal bleeding and no FHS and cervex not dialted they diagnosis invetable abortion the treatment for this ledy :
- surgical treatment
 - induction of labor(syntocinon)
 - dialitation and curtage**
639. A gravid 8 para 8 women has just delivered a 4.5Kg infant a pregnancy. Which of the following is a possible complication?
- Postpartum depression
 - Maternal hypoglycemia
 - Postpartum hemorrhage**
 - Pregnancy-induced hypertension
640. A primigravida woman who is pregnant at 30 weeks gestation told the nurse that she is worried that anything happens to her baby. Which of the following should be the proper nurse's response?
- Ask the woman not to worry
 - Ask the woman to express her concerns**
 - Attract the woman's attention to other issue
 - Reassure the woman about the baby condition
641. Which of the following should be included in the nursing care to a woman during 2nd stage of labor?
- Shave the perineum
 - Administer enema to the woman
 - Careful evaluation of prenatal history
 - Watch breathing, bear down with each contortion**
642. Puerperal sepsis symptoms is ?
- A. Temperature of 38c or higher 2 – 10days postpartum**
643. A 17-year-old mother presented to the primary health center ten after delivery. She is suffering from fatigue, anemia, fever and vaginal discharge (see lab results)
- | | |
|------------------|---|
| Blood pressure | 80/50 mmHg |
| Heart rate | 112 /min |
| Respiratory rate | 35 /min |
| Temperature | 39.6°C |
| Test | Result Normal Values |
| RBC | 4 4.7-6.1 × 10 ¹² /L (male) 4.2-5.4 × 10 ¹² /L (female) |
| Hb | 90 130-170 g/L 120-160 g/L (female) |
| HCT | 0.29 0.42-0.52 (male) 0.37-0.48 (female) |
| WBC | 12.8 4.5-10.5 × 10 ⁹ /L |
- Which of the following is the best diagnosis of health problem in this case?
- Severe urinary track infiction
 - Vesico-vaginal fistula
 - Puerperal sepsis**
 - Post-partum haemorrhage
644. What does the relationship of the site of the presenting part to the location on maternal pelvis refer to ?
- Fetal lie
 - Fetal position**
 - Fetal presentation
 - Fetal attitude

645. A 20 weeks pregnant , primary gravid woman visits the antenatal has sickle cell anemia trait and worried this disease transmitted to her baby which of the following should be initial intervention
- Plan for the fetal genetic screening
 - Educate mother that her disease is inactive
 - Discuss the chances of genetic disease in the fetus
 - D. Gather data about the other family members having the disease**
646. woman who is 32 weeks gestation. Her weight was 66 kg last month and today it is 78 kg. Which of the following is the best nursing action?
- Assess the size of her fetus
 - Give health education good nutrition.
 - Advise her to exercise and lose some weight.
 - D. Check her blood pressure and test her urine for protein.**
647. A nurse is caring is caring for a day 1 postpartum patient assessment revealed in addition to red lochia with fleshy odor
- Blood pressure 110/70 Temperature 38 Respiratory rate 20 Heart rate 90
- Which of the following would be the proper nursing interpretation of these findings ?
- A. Normal**
 - Infection
 - Dehydration
 - Hemorrhage
648. A primigravid client who is at 14 weeks gestation has been diagnosed with hyperemesis gravidarum. The nurse explains to the client that the condition is related to high levels of ?
- Testosterone
 - B. Estrogen**
 - Aldosterone
 - Progesteron
649. Upon reviewing the pregnant client's blood test results, the nurse that traces of mercury are present even after the nurse had healthy dietary modifications. Which action of the client does the nurse discuss to reduce risk client?
- Client has five soaked walnuts every day
 - B. Client consumes king mackerel very often**
 - Client eats one medium bowl of flax seeds daily
 - client has cooked soybean seeds as an evening snack
650. A nurse was assessing a newborn delivery by a mother infected with Neisseria gonorrhea. What is the complication that can occur to the newborn?
- Deafness
 - B. Blindness**
 - Mental retardants
 - Hyperbilirubinemia
651. A full term mother presents in the antenatal clinic with mild lower abdominal contraction and show watery discharge for her vagina. She told the midwife that she may be starting her child birth. She is admitted in the labor and delivery unit, support well and is lying down on the bed.
- What knowledge guide is necessary as first intervention ?
- A. Assessment of cervix**
 - Medication and induction
 - Reassurance and support
 - Mode of delivery

652. A pregnant woman visits the Outpatient clinic complaining of excessive vaginal secretion. Which of the following is the appropriate nursing assessment?
- A. Fetal heart rate
 - B. Fundal height
 - C. Signs of infection of labor**
 - D. Fetal presentations and position
653. A pregnant mother with sickle cell anemia is in labor. What should the nurse to ensure the safety of the infant?
- A. Control pain
 - B. Administer oxygen**
 - C. Monitor fetal heart rate
 - D. Monitor maternal vital signs
654. A nurse is preparing to administer RhoGam to Rh-negative women who has delivered a Rh-positive newborn . Which of the following is prevented by this intervention ?
- A. Maternal illness
 - B. Neonatal illness
 - C. Production of antibodies**
 - D. Re- occurrence of Rh positive baby in next mother
655. A 34 years old woman was diagnosed with breast cancer and underwent surgery. She is currently receiving monthly chemotherapy she telephone the clinic and notifies the nurse that she developed a sore throat. Which of the following foods would be the most appropriate to recommended?
- A. Fresh fruits and vegetables**
 - B. Seafood
 - C. Dried fruits nuts
 - D. pasteurized cheese
656. Both parent have sickle anemia disease ?
- A. 100%.**
 - B. 50%.
 - C. 75%.
 - D. 25%
657. Both parent have sickle anemia trait the percentage for the child to have the disease ?
- A. 100%.
 - B. 50%.
 - C. 75%.
 - D. 25%**
658. A 7 year old child is assessed with a urinary tract infection BP 95/55 Tem 36.4 . which of the following is the most appropriate nursing action?
- A. Notify the physician immediately
 - B. Notify the nurse manager immediately
 - C. Document the findings in the nursing note**
 - D. Put the physician

659. A nurse is caring for child who is post tonsillectomy and adenoidectomy. The nurse should the nurse plan to assess which of the following complication?
- A. Pulmonary hypertension
 - B. Hemorrhage**
 - C. Hearing loss
 - D. Orthopnea
660. In the pediatric ward, a newly registered nurse observed that most of the patients have respiratory problems. Which of the following condition needs priority of care?
- A. Pneumonia
 - B. Epiglottitis**
661. An infant was admitted with patent Ductus arteriosus. Indomethacin 0.2mg/kg with prostaglandin inhibitor via intravenous route was prescribed for him. Which of the following is the function of indomethacin?
- A. Assist closure of ductus arteriosus**
 - B. Prevent infection
 - C. Improve tissue perfusion
 - D. Preserve hormone level
662. A 10 year old child admitted to the pediatric ward with rheumatic fever. Which assessment data should the nurse consider when obtain child's history to be most significant?
- A. Family history of congenital heart disease
 - B. Recent episode of streptococcal tonsillitis**
 - C. Lack of interest in certain types of food
 - D. Projectile vomiting and diarrhea for 2 days
663. A nurse is planning to give a 5 year old child an intramuscular injection. Which of the following is the best nursing intervention to minimize fear?
- A. Show a cartoon video injection attention
 - B. Allow child to give injection to a doll**
 - C. Observe another child receiving injection
 - D. Get parents to reassure child
664. A full-term newborn admitted to the neonatal intensive care unit with diagnosis of tracheoesophageal fistula. Which of the following nursing measure to prevent pulmonary complication and improve good prognosis of the newborn condition?
- A. encourage exclusive breast feeding
 - B. insert nasogastric tube for gastric decompression
 - C. intermittent suction by double-lumen catheter**
 - D. prepare for the insertion of tracheostomy
665. A four year old boy is admitted with diabetes insipidus. The nurse is assessing him. Which of the following finding findings should the nurse anticipated.?
- A. Bradycardia
 - B. Low urine output
 - C. Excessive thirst**
 - D. High systolic blood pressure

666. A midwife is discussing the birth spacing measures with a mother whose first baby boy is a 1 year old and was born with spina bifida. The midwife has explained the possible causes of condition of the baby and the measures is necessary if the mother is planned for a next pregnancy?
- A. Increase iron and calcium supplements
 - B. Multivitamin and folic acid intake**
 - C. Genetic screening
 - D. Immunization before and during pregnancy
667. A newborn admitted to the NICU with tracheoesophageal fistula (TOF). which of the following nursing intervention should be included?
- A. Elevated the head for feedings
 - B. Elevating the head but keep the child NPO**
 - C. Insert a nasogastric tube for feeding
 - D. Encourage the mother to breast feed
668. You have learned that in babies and children with developmental dysplasia of the hip (DDH), the hip joint has not formed normally. Which of the following is the most common form of DDH?
- A. acetabular dysplasia
 - B. dislocation
 - C. Preluxion
 - D. Subluxion**
669. Childlern with cerebral palsy **30 -50%** have cognitive impairment
670. A 6 years old child diagnosed with Sickel cell anemia is admitted with sever pain in upper and lower extremities. Which of the following priority should the nurse concern?
- A. hydration and pain management**
 - B. nutrition and restricted hydration
 - C. Nutrition and manage infection
 - D. Manage pain and infection
671. A 2 years old girl is hospitalized with a diagnosis of bronchiolitis. She is receiving oxygen therapy. Which of the following would be the best indicator the effectiveness of the of oxygen therapy?
- A. Oxygen saturation 84%
 - B. Oxygen saturation 90%**
 - C. Respiration rate 12bpm
 - D. Respiration rate 25bpm
672. A nurse is caring a one year old child with congestive heart failure . which nursing goals appropriate?
- A. Relieving pain
 - B. Improving myocardial efficiency**
 - C. Maintaining thermoregulation
 - D. Promoting oral feeding
673. A nurse was assessing a newborn undergone circumcision what is the immediate risk for this newborn the nurse need to observe ?
- A. Bleeding**
 - B. Fever
 - C. Infection
 - D. Pain

674. A 30-year-old pregnant lady in her 33 weeks pregnancy. When the nurse assesses the health condition of the lady, she provides nursing care and health teaching. Which of the following types of home visiting is the best to be conducted?
- Selective
 - Follow up**
 - Field trip
 - systemtic
675. Which of the following circumstances is most likely to cause uterine atony and lead to postpartum hemorrhage?
- endometriosis
 - urine retention**
 - cervical and vaginal tears
 - hypertension
676. Mother of nine children, three of them with congenital anomalies down syndrome; she is a primary school graduate, with low first status. She is not using any method of family planning. According primary health care nurse has refered her counselling. Which of the following phases of home visit accomplishes intervention?
- Initial
 - Closing
 - Action**
 - Terminal
- Rationale .. intervention phase
677. Mother of nine children, three of them with congenital anomalies Down syndrome; she is a primary school graduate, with low status. She is not using any method of family planning. According primary health care nurse referred her for counseling Which of the following types of home visits that the community her nurse should conduct for t client?
- Systematic routine
 - Selective**
 - Follow up
 - Field trip
- Rationale** Selective Home Visiting Programs. These programs focus on individuals or subgroups of the population whose risk of developing a disorder are higher than average, as evidenced by biological, psychological or social risk factors. These families need more sustained or intensive support.
678. Mother of nine children, three of them with congenital anomalies and down syndrome: she is a primary school graduate, with low financial status. She is not using any method of family planning Accordingly, the primary health care nurse has referred her for counseling. Which of the following terms describe any restriction or lack of ability to perform an activity?
- Impairment
 - Abnormality
 - Handicapped
 - Disability**

679. Mother of nine children, three of them with congenital anomalies Down syndrome; she is a primary school graduate, with low status. She is not using any method of family planning. So, the health care nurse has referred her for counseling. Which of the following application the counselor can help Mrs. regarding family planning?

A. Prevention level
B. Gather model
 C. Group teaching
 D. Rejection of Mrs. M expression

Rationale

When you counsel a new client in your village about family planning, you should follow a step-by-step process. **GATHER** is an acronym that will help you remember the 6 basic steps for family planning counselling:-

G Greet the client respectfully.
A Ask them about their family planning needs.
T Tell them about different contraceptive options and methods.
H Help them to make decisions about choices of methods.
E Explain and demonstrate how to use the methods.
R Return/refer; schedule and carry out a return visit and follow up.

When you counsel a new client in your village about family planning, you should follow a step-by-step process. **GATHER** is an acronym that will help you remember the 6 basic steps for family planning counselling (see box below). It is important to know that not every new client in your village needs all the steps — you need to use the GATHER approach sensitively so that it is appropriate to each client's need. Within your community you may need to give more attention to one step than another.

Family planning counselling — the GATHER approach

G Greet the client respectfully.

A Ask them about their family planning needs.

T Tell them about different contraceptive options and methods.

H Help them to make decisions about choices of methods.

E Explain and demonstrate how to use the methods.

R Return/refer; schedule and carry out a return visit and follow up.

It is important to give more emphasis to the points under each counselling step.



680. Mother of nine children, three of them with congenital anomalies and one down syndrome; she is a primary school graduate, with low financial status. She is not using any method of family planning. Accordingly, the primary health care nurse has referred her for counseling. Which of the following is a barrier facing the team responsible for providing health services to handicapped individuals?

A. Sensory limitations
 B. Rigid rules
C. Developmental disabilities
 D. Deafness & hearing limitations

681. A mother of nine children, three of them with congenital anomalies and one down syndrome; she is a primary school graduate, with low financial status. She is not using any method of family planning. So the primary care nurse has referred her for counselling. Which of the following must be focused on by the community health care nurse to provide an effective health education?

A. Educate regardless realistic objectives
B. Use clear and concise language
 C. Use scientific terms during explanation
 D. Explain the negative consequences in the family

682. Mother of nine children, three of them with congenital anomalies and one down syndrome, she is a primary school graduate, with low financial status, she is not using any method of family planning. Accordingly, the primary health care nurse has referred her for counseling. Which of the following is a model concerned with disability as from of social injustice due to stigma, or discrimination

A. Health belief
 B. Biomedical
C. Sociopolitical
 D. Economic

683. A pregnant mother is in the active labour. Her cervix is fully effaced dilated with crowing of the fetal head. The mother is exhausted, perspiring, and is having increasingly painful and severe uterine contractions. Which of the following intervention is best? desired by the assisting midwife

A. exert pressure on abdomen to aid in child birth
 B. increase the rate of intravenous medication
 C. wipe face with wet towel and hold hands
D. instruct mother to push h

Neonates

684. A 15 month old child is admitted for hypospadias repair. After surgery, which of the following instruction should be given to the parents?

- A. Limit activity for 2 weeks
- B. Avoid apply ointment or powder
- C. Give a child a diet that is high in protein
- D. Isolate the child from other children with the infection

use the double diapering technique to protect the stent, don not bathe the child in the tub until the stent or catheter is removed, force fluids, antibiotics, signs of infection, urine will be blood tinged for several days, restrict the infant or toddler from activities, avoid holding infant straddled on the hip, limit activity for 2 weeks

685. A two hour old newborn baby is given to the mother to start on breastfeeding. The mother is reluctant as she thinks is too early for the baby. The mother finally agrees to breast feed the baby after the midwife has discussed the advantages of early of breast feeding. Which of the following statement made by the mother suggests her correct understanding?

- A. Improve natural immunity
- B. Strengthens sucking reflex
- C. Stimulates gut movement

686. A nurse is preparing a teaching session for elementary school students about the safety measures of hazards. Which of the following is the most appropriate topics for safety measure for children at this age?

- A. Sport activities
- B. Riding bicycle
- C. Swimming
- D. Driving

687. Establishes baby's hydrationA 11 month old boy with cleft lip & palate was discharged from the hospital after surgery. The nurse provided the parents with teaching. Which of the following statement by the parents indicate appropriate understanding of the teaching instruction?

- A. I should avoid that my infant's cry's as much as possible
- B. I will use a spoon to feed my child after the surgery
- C. I will put my child in his abdomen after feeding
- D. I will use lotion to ease my child's lip irritation

688. A newborn born by elective caesarian section under general anesthesia ter 28 weeks of pregnancy. His weight is 850 gm, and he is in (20) th percentile in intrauterine growth chart. He is admitted to Neonatal tensive Care Unit. Which of the following is the classification of this newborn according to stational age and birth weight using intrauterine growth chart?

- A. He is appropriate for gestational age
- B. He is extremely low birth weight
- C. He is small for gestational age
- D. He is very low birth weight.

689. A baby born at 38 weeks of gestation with birth weight 1800 gram. which of the following is the classification of this infant?

- A. Low birth weight
- B. Very low birth weight
- C. Appropriate for gestational age
- D. Small for gestational age

690. A baby born at 24 weeks of gestation with birth weight 1000 gram and at 90th . which of the following is the classification of this infant?

- A. Low birth weight
- B. Very low birth weight
- C. Appropriate for gestational age
- D. Large for gestational age

691. A baby born at 29 weeks of gestation with birth weight 950 gram. which of the following is the classification of this infant?

- A. Low birth weight
- B. Very low birth weight
- C. Appropriate for gestational age
- D. Small for gestational age

692. Which of the following would be the proper interpretation by the nurse for a chromosomal analysis showing 46, XY?

- A. Normal male
- B. Abnormal male
- C. Normal female
- D. Abnormal female

693. Which of the following would be the proper interpretation by the nurse for a chromosomal analysis showing 46, XX?

- A. Normal male
- B. Abnormal male
- C. Normal female
- D. Abnormal female

694. To measure fatal heat rate fetus daily activity through ?

- A. Doppler sound
- B. Weighting measures
- C. Non-stress test
- D. Daily fetus kicks

عندنا ثلاث مصطلحات. في المواليد. / حديثي الولادة
الأول
من حيث عدد اسابيع الحمل
Gestation Age
Term
Or
Preterm
Or post term
Term 38-40 weeks
Preterm 30-36
Extremely preterm less than 30
المصطلح الثاني
على الوزن عند الولادة
Birthweight
Normal weight 2.5-3.5
Low birthweight less than 2.5kg
Very Low birthweight less than 1.5kg
Extreme Low birthweight less than 1 kg
تدخل ع المصطلح الثالث
علاقة الوزن بالعمر الجنيني
relation between
birthweight and Gestational age
وهنستخدم فيها الرسم البياني او المنحني
Large for GA more than 90%
بيكون ع الرسمه واقع ف المنطقه البيضاء
Appropriate fo GA
وده بيكون واقع ف الوسط ف المنطقه الصفراء ويكون ما
10%. To 90% بين
Small for GA
وبيكون واقع ف المنطقه %10

695. Immediately following the birth of a full-term newborn, which of the following is the priority nursing diagnosis for this newborn?
- A. airway clearance related to nasal & oral secretions**
 - ineffective thermoregulation related to environmental factors
 - risk for imbalance fluid volume related t weak sucking reflex
 - risk for injury related to immature defense mechanisms
696. A full term newborn is admitted to NICU with a diagnosis of meningocele. Which of the following admission assessment is needed?
- Specific gravity of urine
 - B. Head circumference**
 - Weight and length
 - Palpation of the abdomen
697. A nurse is caring for a full term newborn who was delivered five minutes ago. The infant's Apgar score was 8 at one minutes and five minutes. Which of the following has the highest priority ?
- Assessing the infant's red reflex
 - B. Preventing heat loss from the infant**
 - Maintaining the infant in the supine position
 - Administering humidified oxygen to the infant
698. A30week gestational preterm admitted to NICU 2hours ago the neonate starts to have grunting ,nasal flaring which of the following the nurse recognize regarding signs and symptoms?
- A. Neonate has RDS**
 - It is normally in the first 24 hours of birth
 - This is not significant unless become cyanosis
 - Neonate has hypoglycaemia
699. A 30 weeks of gestational age preterm is admitted to the neonatal intensive care unit 2 hours . the neonate starts to have grunting tachypnea, and nasal flaring. Which of the following is responsible factors for diagnosis of the premature ?
- Bronchial spasm
 - Immature bronchioles
 - C. Lack of surfactant**
 - Pulmonary over load
700. A 7 years old child is admitted to pediatric medical. Ward his urine analysis result revealed massive proteinuria. T he child was diagnosed with Nephrotic Syndrome. Which of the following medication can prevent the relapses of the child condition?
- Antibiotics
 - Corticosteroids
 - Diuretics
 - D. Immunosuppressive**
701. An infant was admitted with patent Ductus arteriosus. Indomethacin 0.2mg/kg with prostaglandin inhibitor via intravenous route was prescribed for him. Which of the following is the function of indomethacin?
- A. Assist closure of ductus arteriosus**
 - Prevent infection
 - Improve tissue perfusion

D. Preserve hormone level

702. A newborn is diagnosed with Ventricular Septal Defect (VSD) Which of the following information should the nurse give to the newborn's mother?

- A. Cyanosis will occur most of the time during sleeping
- B. Breast feeding is not recommended for your child
- C. The defect might close spontaneously after 6 months**
- D. Blood pressure is different on the child's arm and the leg

703. A 10 years old child had an appendectomy yesterday. Which pain scale should the nurse use to assess child pain?

- A. Numeric
- B. Faces**
- C. FLACC
- D. Cries

704. Infant mortality is a critical indicator of children health status. Which of the following statements probably describes the " infant mortality " term?

- A. Deaths of infant during the first year of the life per 100 live births
- B. Death of infants during the first year of life per 1000 live births**
- C. Death of infants during the first year of life per 10000 live births
- D. Death of infants during the first year of life per 10000 live births

705. A child with rheumatic fever is having sever pain , tenderness and swelling in his left joint what should the nurse do to alleviate his pain?

- A. Apply heat on the affected joint
- B. Put the joint in flexion position
- C. Apply cold compresses to the joints**
- D. Ask the child to walk for 15 minutes

706. 9 YEAR old child is admitted to the Postoperative Care Unit following in heart transplant. The nurse administers cyclosporine by intravenous infusion. Fifteen minutes later the child has difficult breathing, his skin feels cold and clammy and he appears restless. Which of the following is the most appropriate initial nursing action?

- A. Ensure airway patency
- B. Administer oxygen therapy
- C. Discontinue intravenous infusion**
- D. Administer intramuscular epinephrine

707. A nurse in the Neonatal Intensive Care Unit is caring for premature newborn, is diagnosed with Respiratory Distress (RDS) and the doctor ordered administering surfactant Surfactant should be given by which of the following routes?

- A. Intravenous
- B. Subcutaneous
- C. Intramuscular
- D. Endtracheal**

708. A 7 year old child is assessed with a urinary tract infection BP 99/55 TEMP 36.4 . which of the following is the most appropriate nursing action?

- A. Notify the physician immediately
- B. Notify the nurse manager immediately
- C. Document the findings in the nurse note**

D. Put the patient in Trendelenburg position

709. In an education session with pregnant mothers, the midwife explains the purpose of amniotic fluid sac or the water bag. After the completion, the midwife is revising the concepts by asking questions. Which of the following statement made by the mothers is most appropriate regarding the fluid bag.
- It provides the baby with immunity against diseases
 - It provides important nutrition and fluid to the baby
 - It protects and maintains baby's temperature**
 - It protects the baby from harmful chemicals in mother's blood
710. A 23-year-old gravid 1 ,para 0 mother is presented to the Antenatal clinic. The health educator educates group of mothers in the waiting area. Which information regarding fetal circulation the educator should stress on?
- One umbilical artery and one umbilical vein
 - One umbilical artery and two umbilical veins
 - Two umbilical arteries and one umbilical vein**
 - Two umbilical arteries and two umbilical veins
711. 37 weeks of gestation neonate is admitted to the Neonatal Intensive Care Unit immediate after delivery because the mother has fever in the last 2 days before delivery. Which of the following is most expected finding for neonatal sepsis?
- decreased urinary output
 - poor feeding
 - stable body weight
 - sudden hyperthermia**
712. Which of organism can cause neonate to develop septicemia including respiratory distress apnea and hypotension within 12 hours of birth
- Escherichia coli
 - Cytomegalovirus
 - varicella zoster virus
 - Group B streptococcus**
713. while a nurse is assessing an infant born 11 hours ago caesarean section, she auscultated moist lung sounds. Which of the following is the most likely interpretation?
- Abnormal finding
 - Normal finding**
 - Pneumothorax
 - Surfactant aspiration
714. **Lanugo hair** is fine hair in preterm baby less than 30weeks
715. **Vernix caseosa**, also known as vernix, is the waxy or cheese-like white substance found coating the skin of newborn human babies. It is produced by dedicated cells and is thought to have some protective roles during fetal development and for a few hours after birth.
716. An 11 years old child has been diagnosed with type 1 diabetes mellitus . which of the following education should the nurse explain about the exercise?
- Extra insulin is require during exercise
 - Extra snacks are needed before exercise**
 - Exercise will increase blood glucose



D. Exercise should be restricted

717. newborn has small, whitish, pinpoint spots over the nose, which the nurse knows are caused by retained sebaceous secretions. When charting this observation, the nurse identifies it as:

A. **Milia**
B. Lanugo
C. Whiteheads
D. Mongolian spots



718. A 25 years old primipara is admitted for labor . the infant is delivered by forceps because of breech presentation and full body assessment shows alarge blue macular marking over left buttocks which o the following the most likely cause ?

A. Echymosis
B. Nervus flames
C. Telangiectasia nevi
D. **Mongolian spots**



719. A nurse was observing the stool color for a newborn on the first day after delivery. What is the expected color of stool for this newborn?

A. Brown
B. Light green
C. Light brown
D. **Brownish green**

720. An infant with developmental dysplasia of the hip is discharging with a Pavlik harness. Which of the following instructions should the nurse give to the infant's parents

A. Put the baby's diaper over the harness
B. Fasten the straps for at least 2 hours every day
C. Press both legs together before applying the harness
D. **Soft, thin clothing should be worn under the harness**

721. A neonatal is admitted to the NICU with a meningomyelocele. HR130 .. RR 28..... TEM 36.7 which of the following action the nurse should perform to prevent infection of the meningomyelocele sac ?

A. Wash the sac with betadine every shift
B. Expose the defect to the room air
C. Apply antibiotic cream every 24 hours
D. **Cover he sac with moist sterile saline dressing**

722. A baby born at 38 weeks of gestational with birth weight 1800gram. Which of the following is the classification of this infant?

A. Low birth weight
B. Very low birth weight
C. Appropriate for gestational age
D. **Small for gestational age**

723. 24 weeks of gestation neonate is admitted to the Neonatal Intensive Unit immediately after delivery with respirator distress syndrome de 1.

heart rate 140 /min respiratory rate 77 /min Temperature 36.5

Which of the following method of feeding is recommended to promote with of this premature?

A. **Enteral feeding of breast milk**
B. Enteral feeding of premature formula
C. Oral breast feeding

- D. Oral premature formula
724. A newborn is delivered by the midwife. The umbilical cord is cut safely by following necessary aseptic techniques. Further newborn is to be taken. Which of the following intervention is the most desired?
- A. Assess sucking response
 - B. Increase mother child bonding
 - C. Assess and record APGAR score
 - D. **Keep dry and maintain thermoregulation**
725. A 2-year-old child is admitted to pediatric ward. The mother cannot stay the child and she will visit him in the weakened only. Which of the following nursing action indicates an understanding of the emotional needs this child?
- A. Give her a warm bath to calm down
 - B. Allow the child to suck on a pacifier
 - C. **Ask the parents to bring the child favorite toy**
 - D. Tell the parents that frequently visiting is unnecessary
726. When assess newborn weight in scale the nurse must avoid heat loss by;
- A. Radiation
 - B. Evaporation
 - C. **Conduction**
 - D. Convection
727. A nurse is caring for a newborn in Well Born Nursery she wraps the baby with blanket and ensures the nursery temperature is suitable for the babies. What type of heat loss is the nurse preventing?
- A. Radiation
 - B. Conduction
 - C. **Convection**
 - D. Evaporation
728. A mother is asking the nurse how sickle cell could be detected. Which of the following is a test detected sickle cell disease of traits?
- A. **Hemoglobin electrophoresis**
 - B. Bone marrow aspiration
 - C. Complete blood count
 - D. Free erythrocyte protoporphyrin
729. An 11-year-old child has been diagnosed with diabetes insipidus?
- A. **Posterior pituitary**
 - B. Adrenal medulla
 - C. Anterior pituitary
 - D. Adrenal cortex
730. A term baby boy has been diagnosed with Down syndrome. Physical examination revealed flattened nose, low set ears, upward slanting eyes, single palmar crease. Which of the following is the most common congenital anomaly associated with this disease?
- A. Developmental dysplasia of hip (DDH)
 - B. **Congenital heart disease**
 - C. Hypospadias
 - D. Pyloric stenosis

731. A neonate is admitted to the neonatal care unit with a meningomyelocele HR 130 RR 28 TEM 36.7 which of the following actions should the nurse perform to prevent infection of the meningomyelocele sac?
- A. Wash the sac with betadine every shift
 - B. Expose the defect the room air
 - C. Apply antibiotic cream every 24 hours
 - D. Cover the sac with sterile dressing**
732. If full term infant weight 3 kg at birth, approximately should the infant weight be at 12 months old ?
- A. 7 kg
 - B. 9 kg**
 - C. 11kg
 - D. 13kg
733. A nurse is performing physical examination on the new born she notes that the baby has cephalhetoma this baby is risk to develop which of the following?
- A. Sudden death
 - B. Pathological jaundice**
 - C. Infected umbilical cord
 - D. Increased intracranial pressure
734. A4 days old baby diagnosed with physiological jaundice . his father is distressed and wants to know why he have this condition . what the nurse should the nurse tell the about the most prominent physiological jaundice
- A. Immature hepatic function**
 - B. Decrease milk intake
 - C. Rh incompatibility
 - D. Red blood cell enzyme defects
735. A newborn with hyperbilirubinemia was started on phototherapy What will be the nurse's instruction regarding feeding?
- A. Feed glucose drinks
 - B. Breastfeed two hourly**
 - C. Bottle feed till the bilirubin level reduce
 - D. Breastfeed alternatively with bottle feeds
736. Which action a nurse needs to include when caring for a newborn receiving phototherapy?
- A. Expose all surfaces
 - B. Prevent stimulation
 - C. Cover the eyes with shield**
 - D. Change position every four hourly
737. A baby born at 35-week was admitted in neonatal intensive 27hours ago, physical examination revealed yellow discoloration sclera and mucus membrane. The result of bilirubin level every 170mol. The infant was diagnosed with neonatal jaundice physician order to start single phototherapy. Which of following should the nurse consider as a priority during phototherapy of this newborn?
- A. ensure proper fitting of eye covering (patches)**
 - B. monitor bilirubin levels every 48 hours
 - C. feed the infant formula every 4 to 5 hours
 - D. avoid removing the infant from phototherapy

738. A nurse prepares to administer a vitamin K injection to a full term the mother wants to know the importance of the injection Which of the following is the best nurse response to the mother
- A. needed for blood clotting to prevent haemorrhage
 - B. accelerate the growth and development of infants
 - C. help in maintain healthy gut and passage of meconium
 - D. protect the infant from developing sever respiratory distress
739. A nurse is caring for child who is post tonsillectomy and adenoidectomy the nurse should plan to assess which of the following complication?
- A. Pulmonary hypertension
 - B. Hemorrhage
 - C. Hearing loss
 - D. Orthopnea
740. Term baby boy is admitted to Neonatal Intensive Care Unit. Physical examination revealed flattened nose, low set ears, upward slanting eyes, glabral crease. Which of the following is the possible diagnosis of the new-born?
- A. Cushing syndrome
 - B. Down syndrome
 - C. Intra-Uterine Growth Retardation
 - D. Congenital hypothyroidism
741. A nurse performing nursing care plan for a neonate after a birth, which intervention has the highest nursing priority neonate?
- A. Obtain a dextrostix
 - B. give the initial bath
 - C. Give the vitamin k injection
 - D. cover the neonate head with a cap
742. During physical assessment of a male infant genitalia, the nurse found that one of the testes are enlarged. Which of the following could be the reason for swollen testes?
- A. Chordee
 - B. Cryptorchidism
 - C. Hydrocele
 - D. Hypospadias
743. A newborn is diagnosed with hypospadias. When teaching the parents of this child, the nurse should tell them to avoid which of the following before the hypospadias repair?
- A. circumcision
 - B. drinking acidic juices
 - C. urinary catheterization
 - D. riding a bicycle
744. A 6-year-old child is diagnosed with sickle cell disease is admitted with severe pain in upper and lower extremities. What type of crisis that the child experiences?
- A. Sequestration
 - B. Aplastic Crisis
 - C. Vasoocclusive
 - D. Hyperhemolytic

745. Mother of a sickle cell anemia child is asking why her child can't go hiking with his friends. Which of the following complications hiking can lead to ?
- A. Enhance iron absorption
 - B. Decrease oxygen consumption
 - C. Inhibit hemoglobin production
 - D. Precipitate vaso-occlusive crisis**
746. The nurse admits a client who is in sickle cell crisis. The nurse should prepare for which intervention as a priority in the management of the client?
- A. Pain management with an opioid
 - B. Intravenous fluid therapy
 - C. Oxygen administration**
 - D. Blood transfusion
747. When performing a new-born assessment, the nurse should measure the vital signs in the following sequence:
- A. Pulse, respirations, temperature
 - B. Temperature, pulse, respirations
 - C. Respirations, temperature, pulse
 - D. Respirations, pulse, temperature**
748. An infant arrives in the Emergency Department not breathing and does not have a pulse. When starting CPR, which of the following arteries is appropriate to perform the pulse checks?
- A. carotid
 - B. Radial
 - C. Brachial**
 - D. temporal
749. A nurse is assessing a 2 days old full-term male neon circumcision. She observed that the circumcised area is re a large amount of fresh blood.
- | | |
|------------------|----------|
| Heart rate | 110 /min |
| Respiratory rate | 40 /min |
| Temperature | 36.6°C |
- Which of the following action should the nurse take?
- A. Apply antibiotic ointment on the affected area
 - B. Give the infant another injection of vitamin K
 - C. Clean the area with betadine to prevent infection
 - D. Apply gentle pressure with a sterile gauze**
750. A maternity nurse is performing a newborn assessment thirty minutes after delivery of a baby who did not receive any prenatal care and has an unknown gestational age. The skin is extensively leathery, cracked and dry and there is an absence of lanugo and vernix. How many weeks 'gestation is this neonate?
- A. <30
 - B. 30-35
 - C. 36-40
 - D. >40**

751. A nurse is caring for a female newborn who born with an imperforate anus. When assessing the newborns urine, she should notify the doctor immediately if the newborn's urine contains which of the following?
- A. meconium
 - B. sugar
 - C. albumin
 - D. crystals
752. A newborn has diagnosed Development Dysplasia of hip DDH and is using a Pavlik Harness as treatment .
- A. Multiple
 - B. Acetabular dysplasia
 - C. Subluxation
 - D. Dislocation
753. Which of the Following Is the Most Commend Site to Obtain a Capillary blood sugar sample from neonate?
- A. Ear lobe
 - B. fingertip
 - C. Heel
 - D. Abdomen
754. A 6 years old preschool returned from the operating room after tonsillectomy. A nurse received the patient from recovery room. Which position the nurse place the child post tonsillectomy?
- A. Supine
 - B. Side lying
 - C. Low fowler
 - D. High fowler
755. female teenager was diagnosed with sickle cell anemia inform a nurse school arranged a hiking activity in nearby mountains. She shows excitement while informing the nurse about her new water bottle for this trip. What is the appropriate initial response by the nurse?
- A. "what do your parents think?"
 - B. "you realize that this type of activity might cause you a pain crisis"
 - C. "you need to get approval from your doctor before attending this activity"
 - D. "it is great that you are hydrating yourself in preparation of this activity"
756. A child with thalassemia was given desferoxamine (Desferal);which of the following should alert the nurse to notify the physician?
- A. Decreased hearing
 - B. Hypertension
 - C. Red urine
 - D. Vomiting
757. A client is diagnosed as having secondary Cushing's syndrome. The nurse knows that the client has most likely been taking which medication:
- A. Estrogen
 - B. Penicillin
 - C. Lovastatin
 - D. Prednisone

758. A nurse is planning to give a 5 year old child an intramuscular injection. Which of the following is the best nursing intervention to minimize fear?
- A. Show a cartoon video injection attention
 - B. **Allow child to give injection to a doll**
 - C. Observe another child receiving injection
 - D. Get parents to reassure child
759. A client who is breastfeeding her newborn requests assistance from the lactation nurse. Which reflex does the nurse explain in order to assist with latching on?
- A. Extrusion reflex
 - B. **Rooting reflex**
 - C. Swallowing reflex
 - D. Tonic neck reflex
760. A healthy baby is born normally via vaginal delivery and when transferred to newborn until the nurse administered vitamin K intramuscularly. Which sits recommended for vitamin K injection ?
- A. Biceps
 - B. Deltoid
 - C. **Vastus lateralis**
 - D. Gluteus maximus
761. The nurse determines that a client understands the purpose of a vitamin K injection for her new-born if the client states that V K is administered for which purpose
- A. New born lack vitamin
 - B. New born have low blood levels
 - C. **New born lack intestinal bacteria**
 - D. New born cannot produce vitamin k in the liver
762. A nurse is administering a shot of vitamin K to 30 days old infant . which of the following target area is the most appropriate
- A. Gluteus maximus
 - B. Gluteus minimus
 - C. **Vastus lateralis**
 - D. Vastus medialis
763. While a nurse is assessing the head of a newborn at the first hour after delivery , she observed a soft edema over the vertex which crosses the suture line. Which of the following would be the proper nurses interpretation?
- A. Cephalohematoma
 - B. Hydrocephaly
 - C. Large head
 - D. **Caput succedaneum**
764. A mother asked the nurse that while she was changing the diaper for her female newborn, she noticed a brick red stain on it. What is the best response by the nurse?
- A. It is a sign of low iron excretion
 - B. **It is expected in female newborn**
 - C. It is due to medication given to the mother
 - D. it due to medication given to the newborn

765. A nurse is assessing a 4-month-old formula fed infant. The parent reported that the infant was irritable, crying excessively, not sleeping well, and vomiting, gastro esophageal reflux is expected. What nursing intervention should the nurse expect to teach the parent ?
- A. Place in an infant seat after eating**
B. Give frequent feedings
C. Position the child in a swing
D. Thin formula with water
766. The normal systolic Bp for new Born:-
- A. 40-60
B. 60-80
C. 80-100
D. 100-120
767. Neonatal mortality rate :
- A. 1:100
B. 1:1000
C. 1:100000
D. 1:1000000
768. New born stomach capacity
- A. 6ml**
B. 12ml
C. 28ml.

Cleft Palate And Cleft Lip

Age for surgery repair

769. Cleft lip **3 – 6 month** after surgery consideration elbow restrain after surgery the infant should not be allow to cry and the infant not allow to breast feed by sucking . never to put the baby prone position after cleft lip repair surgery
770. Cleft palate **12 – 18 month** **position** after surgery **prone position** ,
771. Consideration after cleft lip and cleft palate is :
- ❖ Burping the child frequently every 15 minutes
 - ❖ To prevent cleft lip and cleft palate the pregnant woman u should advice:
 - ❖ Folic acide 400 iu
 - ❖ Increase green vegetables & Citrus intake
 - ❖ Feeding up right setting position

Questions

772. A 32year old gravid 1 and para, is now planning to become pregnant within the next year. The patient herself had born with born with neural tube defects, which were surgically repaired in early childhood. The nurse recommends that the patient begin taking tablet of folic acid. How many micrograms would be most appropriate for this patient?
- A. **400**
B. 600
C. 1800
D. 4000
773. A 2-month-old infant with cleft lip is seen in the primary health care to get the regular vaccine of 2 months. The mother asked proper time for the corrective cleft lip surgery of her infant. Which of following is the best nurse response?
- A. No specific age for repair of cleft lip
B. It is too late, repair should be done immediate after delivery
C. **The age of 2 months old is the time for repair**
D. The proper time for repair after the age of one year,
774. A nurse is assigned to care for a child after a cleft palate repair which of the following types of restraints is very effective for child?
- A. Mummy restraint
B. **Elbow restraint**
C. Wrist restraint
D. Mitt restraint
775. A 4 month old infant returned immediately from OR room post cleft lip repair which of the following nursing intervention should be considered?
- A. **Apply elbow restrain**
B. Apply suction when needed
C. Measure temperature
D. Put infant in prone position

776. The nurse have been teaching a new mother how to feed was born with a cleft lip and palate before surgical repair of Which of the following action from the mother indicate teaching has been successful?
- A. **burping the baby frequently**
 - B. Prevent the infant from crying
 - C. Placing the baby flat during feeding
 - D. Keep the infant prone following feedings
777. Mother came to the Outpatient Department with an infant having cleft and palate. The infant was underweight, so the nurse has to consider Teaching the proper way of feeding the child in the treatment plan. Which of the following is the proper way of feeding
- A. Use a non-squeezable bottle during feeding
 - B. **Feed infant in an upright, sitting position**
 - C. Enlarge nipple holes of bottle to allow more milk to pass through
 - D. Feed infant longer than 45 minutes to allow more food to be small
778. A 40-year-old women is a gravida 2, para 2 and is current conceive. Her previous pregnancy resulted in the birth of a cleft lip and palate. The patient is anxious and concerned pregnancies and the nurse provides genetic counselling and Which foods would most effectively prevent recurrence a palate?
- A. **Green vegetables and citrus fruit**
 - B. Eggs, milk and dairy products
 - C. Wheat, corn, rice, oats and rye
 - D. Beef, chicken and yellow vegetables
779. A nurse is caring for a new-born with cleft lip. At which age would the nurse expect the doctor to perform?
- A. **3-6 months**
 - B. 6-10 months
 - C. 11-14 months
 - D. more than 14 months
780. A 2-day-old new-born is admitted to the nursery. While the nurse is administrating oral feeding, the milk returns through the child's nose and mouth and the infant become cyanotic. Which of the following condition the new-born should have?
- A. Anorectal malformation
 - B. **Tracheoesophageal fistula**
 - C. Cleft lip and palate
 - D. Cardiac condition
781. A full-term newborn admitted to the neonatal intensive care unit with diagnosis of tracheoesophageal fistula. Which of the following nursing measure to prevent pulmonary complication and improve good prognosis of the newborn condition?
- A. encourage exclusive breast feeding
 - B. insert nasogastric tube for gastric decompression
 - C. **intermittent suction by double-lumen catheter**
 - D. prepare for the insertion of tracheostomy

782. A nurse is caring for 14 month old immediately after a surgical repair of cleft palate. In which position of the following should the nurse put the child?
- A. Prone
 - B. LateralSupine
783. A 9-month-old child who has a repair cleft palate the nurse explaining mother on how she will give feeds to her child. Which of the following instruction can be expected to include feeling education?
- A. Open cup
 - B. Tea spoon
 - C. Bottle feed
 - D. Special bottle feed

Spina bifida

784. 26- Which of the following vitamin supplements can decrease the incidence of Neural tube defects such as anencephaly and spina bifida new-borns ?
- A. Vitamin A
 - B. Riboflavin
 - C. Folic Acid
 - D. Vitamin K
785. All of the following are types of spina bifida EXCEPT:
- A. Myelomeningocele
 - B. Hemophilia
 - C. Meningocele
 - D. Spina Bifida Occulta
786. While caring for a neonate with a meningococcal, the nurse should avoid positioning the child on the:
- A. Abdomen
 - B. Left side
 - C. Right side
 - D. Back
787. infant is born with spinal bifida Which of the following complication is always found in these Infants?
- A. Hydrocephalus
 - B. Craniosynostosis
 - C. Meningitis
 - D. Cerebral palsy
788. On the second day of hospitalization for ventriculoperitoneal shunt revision, a child with spina bifida developed hives, itching and wheezing. The nurse should determine if the patient has been exposed to:
- A. Peanuts
 - B. Strawberries
 - C. Eggs
 - D. Latex
789. Infant with spina bifida the nurse should monitor
- A. Head circumference
 - B. Abdomen circumference

790. Position for the baby with spina bifida is :

- A. prone position
- B. back
- C. supine

791. A full term infant is admitted to NICU with a diagnosis of Spina vital signs are stable. Which of the following positions is suitable for this infant?

- A. Supine position
- B. Semi-fowler position
- C. Prone position
- D. Sitting position

Hirschsprung's Disease :

792. **Definition** :is a congenital anomaly also known as congenital aganglionosis or aganglionic megacolon. The disease occurs as the result of an absence of ganglion cells in the rectum and other areas of the affected intestine

793. A 24 hours after delivery , the nurses noted that the newborn failed to pass meconium. This indicates which of the following condition?

- A. GERD
- B. Pyloric stenosis
- C. Failure to thrive
- D. Hirschsprung disease

794. A nurse is assessing a 6-month-old infant that has retar reduced responsiveness and interaction with the environment to smile or make eye contact. The nurse notices that the attempt to hold or comfort the crying infant. What diagnosis should the nurse anticipate?

- A. Celiac disease
- B. Failure to thrive
- C. Cystic fibrosis
- D. Growth hormone deficiency

795. A 3 days old newborn is diagnosed with Hirschsprung disease. The nurse is conducting a physical examination. Which of the following findings will alert the nurse to suspect this disease in the newborn?

- A. palpable sausage-shaped mass
- B. cyanosis of fingers and toes
- C. failure to pass meconium within 24-48 hours of life
- D. weight less than expected for height and age

796. A 9 month old child is diagnosed Hirschsprung disease scheduled for surgical operation which of the following should the nurse understand the purpose for surgical intervention?

- A. To remove the aganglion portion of the bowel to relieve obstruction
- B. To maintain optimum nutritional status growth the intestinal
- C. To stimulate intestinal adaptation with internal feeding
- D. To minimize complication related to the disease

797. A 9 month old child is diagnosed with Hirschsprung disease he is scheduled for surgical operation which of the following is the most common complication expected during this age ?

- A. Mechanical obstruction
- B. Entero colitis
- C. Pleural effusion
- D. Esophageal Artesia

798. An 8-month-old infant is admitted with Hirschsprung disease. Which of the following would be a significant finding in this infant?
- A. Depressed anterior fontanel
 - B. Polyuria, hematuria
 - C. Weight gain, edema
 - D. Failure to thrive, constipation**
799. The nurse is assessing a 2 -years-old child with Wilms surgery Which of the following should the nurse avoid?
- A. Putting the child in lateral position
 - B. Palpating the child's abdomen**
 - C. Putting the child in a private room
 - D. Provide mouth hygiene 30 minutes after meal

Intussusceptions

800. A nurse is taking a history from the infant parents who have a suspected diagnosis of intussusception. Which of the following assessment question would be most helpful?
- A. Do you breast feeding your child ?
 - B. What does your child's stool look like?**
 - C. How often your child urinating?
 - D. What is the colour of your child' urine?
801. A 5-month-old boy has been vomiting green coloured vomit He has intermittent abdominal pain during which he draws his chest, turns pale and cries forcefully. On observation, the in the stool which has a jelly-like consistency. Abdominal pal a long, tube-like mass. There is no fever, rash nor diarrhea are hyperactive in all quadrants which is the most likely form of initial treatment?
- A. Manual manipulation
 - B. Surgical resection
 - C. Normal saline enema**
 - D. Laparoscopy
802. A 7-month-old infant seen in the Emergency Department suffering from episodes of severe abdominal pain, and the infant's stool became like red jelly. Abdominal examination revealed palpable sausage-shaped mass in the right upper quadrant. Which of the following is the first line of therapeutic management for this infant?
- A. Non-surgical hydrostatic reduction**
 - B. Surgical simple reduction
 - C. Pyloromyotomy
 - D. Endorectal pull-Through
803. 7month-old infant seen in the Emergency Department suffering from episodes of severe abdominal pain, and the infant's stool become like red jelly. Abdominal examination revealed palpable sausage-shaped mass in the right upper quadrant which of the following is the best diagnosis?
- A. Hirschsprung disease
 - B. Hypertrophic pyloric stenosis
 - C. Infant colic
 - D. Intussuception**

804. Nurse is preparing to care for a child with a diagnosis of intussusception. The nurse reviews the child's record and expects to note which sign of this disorder documented?
- A. watery diarrhea
 - B. rib bone-like stools
 - C. profuse projectile vomiting
 - D. Red jelly stool**
805. A 5-year-old child was seen to the Emergency Department abdominal pain, palpable sausage-shaped mass, and Intussusception is suspected Which of the following is the best diagnostic evaluation to?
- A. X-ray
 - B. endoscopy
 - C. Rectal biopsy
 - D. Ultrasonograph**

Pyloric stenosis

806. A 5-week-old newborn was admitted to pediatric Ward with pyloric stenosis, the newborn has weight loss, and projectile vomiting during feeding . They scheduled surgical repair of pyloric stenosis Which of the following postoperative intervention for this
- A. IV fluid infant is retaining adequate amount by mouth**
 - B. Administration of proper analgesia until infant discomfort resolve
 - C. Start feeding immediately after postoperative
 - D. Vomiting is uncommon in the first 24-48 hrs
807. Surgery for pyloric stenosis movement?
- A. Pylorotomy
 - B. Pylorectomy
 - C. Pylorostomy
 - D. Pyloromyotomy**
808. PYLORIC stenosis peristalsis movement ?
- A. From right to left
 - B. From left to right**
809. 1 month-old infant is admitted to the surgical unit with hypertrophic pyloric stenosis and scheduled for the surgery. Which of the following is the findings of abdominal examination?
- A. palpable olive-like mass in the left side
 - B. palpable olive-like mass in the right side**
 - C. Palpable olive-like mass moved from left to right
 - D. Palpable olive-like mass moved from right to left
810. A 7-week-old infant boy is admitted with projectile vomiting decreased urine output, decreased bowel movements and weight loss. He has poor turgor and appears hungry. The nurse observes left-to right peristaltic waves after he vomits. The nurse would expect to find which of the following during the physical assessment?
- A. Hepato-splenomegaly
 - B. A palpable pyloric mass**
 - C. Lymphadenopathy
 - D. Bulging fontanelles

811. The nurse is assessing a child (an infant) with pyloric stenosis . which of the following is likely to note?
- Diarrhea
 - Projectile vomiting**
 - Swallowing difficulties
 - Currant jelly like stool
812. A 5 week old newborn admitted paediatric ward with pyloric stenosis the newborn has weight loss, and projectile vomiting after feeding. Which of the following abdominal organs are directly affected when stenosis diagnosed?
- Stomach and duodenum**
 - Stomach and oesophagus
 - Liver and spleen
 - Liver and bile duct
813. Tetralogy of Fallot where is it located :
- right heart side**
 - left heart side
814. The Foley Family is caring for their youngest child, Justin, who is suffering from tetralogy of Fallot. Which of the following positions is used for congenital heart condition?
- Semi fowler's position
 - knee chest position**
 - prone position
815. Which of the following instruction should be given to the parents about colostomy care?
- Use baby powder after stoma cleaning
 - Empty the pouch when it is completely full
 - Avoid tight diapers around the infant abdomen.**
 - Use baby wipes to clean the skin around the stoma
816. A Community Nursing nurse is working with a family in their home The parents complain that their eight year-old son is "wild" and that he never listens to them. They become upset at his antics to gain attention from the nurse and send him out of the room. Which of the following responses from the nurse is the most appropriate?
- " That was the right thing, it teaches him who is in charge"
 - " Don't worry, he is a boy and with time, he will grow out of this"
 - " Let's talk about how we can teach your child the right behavior and your expectations of him"**
 - " You are only reinforcing his behavior when you do that. Try putting him in time out instead"
817. 11- child is admitted to the pediatrics ward with acute lymphocytic LEUKEMIA , and has had joint pain for several weeks. Physical findings ude widespread ecchymosis, generalized lymphadenopathy, hepatosplenomegaly, and pallor. Lab work shows a low haemoglobin level, RBC count, low WBC count low haematocrit , and low platelets. Which high risk is expected?
- Infection**
 - Fractures
 - Dehydration
 - Hepatitis

818. A nurse is teaching a group of new mothers in the post-natal ward on how to manage breast engorgement after they are discharged to home. Which of the following statements by the mother will indicate to the nurse that they understood how to prevent engorgement of the breasts?
- breast feed every 4 hours
 - breast feeding during the day and bottle feeds at night
 - breast feed every 2-3 hours during the day and stop at night
 - D. breastfeed the baby every 2-3 hours during the day and night**
819. Which of the following is the most common site to obtain a capillary blood sample from Neonates?
- A. Heel**
 - Earlobe
 - Fingertip
820. Mother came to the clinic and afraid that her baby maybe will have meningitis as his brother already have
What should the nurse tell the mother?
- A. The vaccine is reducing the risk for this disease**
821. Temperature 41 c isHyperpyrexia
822. When checking rectal temperature how depth thermometer should be inserted
- 2cm
 - B. 3.5cm**
 - 4cm
 - 4.5
823. Patients with cardiac disease, the nurse should avoided which of the following place when taking temperature
- A. Rectal**
 - Oral
 - Axillary
 - Tympanic
824. A 15-year-old girl is admitted to the hospital with diarrhea. She has been repeatedly vomiting for now weak and lethargic. She is oriented to time to questions appropriately. The nurse prepares temperature using an electronic thermometer. Which measurement would be most appropriate?
- Oral
 - Rectal
 - Axillary
 - D. Tympanic**
825. Fastest and most sanitary thermometer is :
- Glass
 - Electronic
 - Chemical
 - D. Tympanic**

2. Rectal
a. Place the client in the Sims position.
b. The temperature is taken rectally when an accurate temperature cannot be obtained orally or when the client has nasal congestion, has had recent nasal surgery or had the jaws wired, has a nasogastric tube in place, is unable to keep the mouth closed, or is at risk for seizures.
c. The thermometer is lubricated and inserted into the rectum, toward the umbilicus, about 1.5 inches (3.8 cm) (no more than 0.5 inch [1.25 cm] in an infant).

Vaccination

جدول التطعيمات الوطني

التطعيم Vaccine	الزيارة Visit
• BCG • Hepatitis B	• درن • التهاب كبدي (ب) عند الولادة At Birth
• IPV • DTaP • Hepatitis B • Hib • Pneumococcal Conjugate (PCV)* • Rota**	• شلل أطفال معطل • الثلاثي البكتيري • التهاب الكبد (ب) • المستدمية النزلية • البكتيريا العقدية الرئوية* • فيروس الروتا** عمر شهرين 2 months
• IPV • DTaP • Hepatitis B • Hib • Pneumococcal Conjugate (PCV)* • Rota**	• شلل أطفال معطل • الثلاثي البكتيري • التهاب الكبد (ب) • المستدمية النزلية • البكتيريا العقدية الرئوية* • فيروس الروتا** عمر ٤ شهور 4 months
• OPV • IPV • DTaP • Hepatitis B • Hib • Pneumococcal Conjugate (PCV)* • Rota**	• شلل الأطفال الضموي • شلل أطفال معطل • الثلاثي البكتيري • التهاب الكبد (ب) • المستدمية النزلية • البكتيريا العقدية الرئوية* • فيروس الروتا** عمر ٦ شهور 6 months
• Measles • Meningococcal Conjugate quadrivalent (MCV4)	• الحصبة المفرد • الحمى الشوكية الرباعي المقترن عمر ٩ أشهر 9 months
• OPV • MMR • Pneumococcal Conjugate (PCV)* • Meningococcal Conjugate quadrivalent (MCV4)	• شلل الأطفال الضموي • الثلاثي الفيروسي • البكتيريا العقدية الرئوية* • الحمى الشوكية الرباعي المقترن عمر ١٢ شهر 12 months
• OPV • DTaP • Hib • MMR • Varicella • Hepatitis A • Hepatitis A	• شلل الأطفال الضموي • الثلاثي البكتيري • المستدمية النزلية • الثلاثي الفيروسي • الجدري المائي • التهاب الكبد (أ) • التهاب الكبد (أ) عمر ١٨ شهر 18 months
• OPV • DTaP (Td)*** • MMR • Varicella	• شلل الأطفال الضموي • الثلاثي البكتيري (الثنائي البكتيري)*** • الثلاثي الفيروسي • الجدري المائي عمر ٢٤ شهر 24 months
• OPV • DTaP (Td)*** • MMR • Varicella	• شلل الأطفال الضموي • الثلاثي البكتيري (الثنائي البكتيري)*** • الثلاثي الفيروسي • الجدري المائي عند دخول الصف الأول الابتدائي First class primary school age

826. Vaccine refrigerator temperature is ?

- A. 0-3
- B. 1-5
- C. 2-8**
- D. 3-9

827. How to storage vaccines in ICE

828. Vaccination refrigerator temperature **2-8 c**

829. 9 month vaccine :

- A. MCV4**
- B. Hepatitis A
- C. Hib
- D. MMR

830. 1 year vaccine :

- A. Opv**
- B. Hib
- C. DTap
- D. Hepatitis A and varicella

831. 18 month vaccine :


- A. Rota
- B. Hepatitis B
- C. Hepatitis A**

832. First dose of hepatitis B should be given when ?

- A. At birth**
- B. At 2 month
- C. At 6 month
- D. At 1 year

833. Measles vaccine administration route

- A. Intramuscular
- B. Subcutaneous**
- C. Intradermal
- D. Intravenous



834. DTap vaccine administration route

- A. Intramuscular**
- B. Subcutaneous
- C. Intradermal
- D. Intravenous

835. What of the following disease are prevented by MMR vaccine?

- A. Mumps-measles-Scarlet fever
- B. Mumps-measles-rotavirus
- C. Mumps-measles-rabies
- D. Mumps-measles-germen- measles**

836. A mother brought her 6-month-old healthy infant to the well-baby clinic Which immunization should the nurse anticipate to administer as per World Health Organization's recommendation?
- Varicella (Chicken pox)
 - Rotavirus and hepatitis
 - Measles, Mumps, Rubella
 - D. Diphtheria, Tetanus and pertussis**
837. 6month-old boy with hydrocephalus is admitted to the pediatric surgical Ward for ventriculoperitoneal Shunt (VPS) insertion. Which of the following findings should be of the most concern when assessing the child postoperative?
- Sunken fontanelle and irritability
 - decreased head circumference
 - C. poor feeding and pupillary change**
 - headache and excessive sleepiness
838. Which of the following routes is used to administered Diphtheria, Tetanus and pertussis DTP vaccine?
- Oral
 - B. Intramuscular**
 - Subcutaneous
 - Intradermal
839. The nurse are reviewing the immunized schedule if the 11-month of baby Which of the disease should the nurse expect the infant that here immunized against?
- Pertussis, Tetanus, polio, and varicella
 - B. Polio, Pertussis, Tetanus, and Diphtheria**
 - Varicella, polio, Tuberculosis, and pertussis
 - Measles, Mumps. Rubella, and Tuberculosis
840. Dexamethasone **accelerates maturation of fetal lung .**
841. Hirschsprung **disease failure of neonate to pass stool)Meconium) for 24 hours**
842. Apgar score done at **1 minute and 5 minutes** at birth
843. most common cause of diarrhoea in paediatric **virus rotavirus ,**
844. Meconium is..... **Dark green stool in newborn**
845. In determining the one minute APGAR score of a male infant the nurse assesses a heart rate of 120 beats per minute and respiratory rate of 44 per minute. He has flaccid muscle tone with slight flexion and resistance to straightening. He has a loud cry with colour is acrocyanotic What is the APGAR score for the infant
- 7
 - B. 8**
 - 9
 - 10

846. Five minutes post-birth, a neonate has a heart rate of 98, irregular breathing, actively moves all extremities, but has bluish hands and feet, as well as a weak and timid cry. Which is the correct APGAR assessment score?
- 9
 - 8
 - 7
 - 6**
847. A nurse documented assessment on a newborn as listed (see table) Points Indicator less than 98 /min Heart rate 28 /min irregular Respiratory rate Muscle tone Minimal flexion of the extremities Reflex irritability Grimace Body pink, extremities blue Colour
- What is the total Apgar Score?
- 0
 - 3
 - 5**
- ## Paediatrics
848. A 5 years old child brought to emergency room with fractured right forearm he had several bruises on his body but showed no signs of pain while palpating them. seemed scared and did not answer any questions asked Which of the following should the nurse do to gain the child's body trust?
- Arrange meal
 - Comfort measures
 - Show the playing area**
 - Therapeutic communication
849. During clinical assessment, the nurse suspected that a child has been psychological abuse. Which statement by the child is most likely support the suspicion?
- My parents tell me that iam stupid**
 - My parents hurts me to get attention from the doctors
 - My mother forces not give meals on times
 - My uncle show me picture of nude people
850. While a nurse is assessing vital signs of newborn infant first hour of delivery. HR 170 RR 70 TEM 36 . the nurse would interpret these finding as in the discharge instruction
- Anemia
 - Cold distress .**
 - Heart defects
 - Hyperglycemia
851. Toddler is admitted to the pediatric room with several episodes of diarrhea 3 days . the child is diagnosed with gastroenteritis. Which organism is responsible about the most diarrhea episodes in children?
- Rota**
 - Bacillus magisterium
 - Shigella
 - Staphylococcus

852. A child with a diagnosis of Tetralogy of Fallot is scheduled to be discharge from the hospital. the nurse who is planning discharge education should instruct the caregivers that during a hyper cyanotic spell, which position is most likely to benefit the child?
- A. Supine
 - B. Prone.
 - C. side-lying
 - D. Knee-chest**
853. A 12- year- old boy was brought to the Emergency respiratory arrest due to drowning. Cardiac resuscitation what is the major complication that might happen if treated after drowning quickly?
- A. Sepsis
 - B. Alkalosis
 - C. Acidosis**
 - D. Hypothermia
854. which of the following is the leading cause of injury for children who are more than five years old ?
- A. accidental suffocation
 - B. motor vehicle**
 - C. congenital anomalies
 - D. drowning
855. A 1-year-old girl admitted to podiatric medical unit significant weight loss, diminished mid-arm circumference diarrhea, and red hair. Which of the following type of malnutrition do the nurse suspect
- A. Marasmus
 - B. Spitting up
 - C. Kwashiorkor**
 - D. Rickets
856. 33 month-old infant is admitted to the Emergency Department (ED) with fractured arm. The mother indicated that while the child was crawling fell down the stairs and broke his arm. Which of the following observation would lead the nurse to suspect that this is a victim of abuse?
- A. Age inappropriate injury
 - B. Pattern and shape of the injury
 - C. Child is appearing malnourished
 - D. Improper explanation of the cause of injury**
857. 5-year-old child was brought to the Emergency Room with a fractured right forearm. He had several bruises on his body but showed no signs of an while palpating them. He seemed scared and did not answer any questions asked. Why should the nurse discuss this case with the nurse manager?
- A. Continuity of care
 - B. Rule out child abuse**
 - C. Psychological support
 - D. Fracture management
858. A four-year old child is seen in the Emergency Department with a spiral fracture of the left arm . The x-ray examination showed previously broken healed bones. What is your immediate action?
- A. Call social services to immediately arrange foster care for the child
 - B. Ask the child about the previous accidents and management
 - C. Report the child abuse to the local authorities**
 - D. Try to establish rapport and trust with the child's family

859. A 62-year-old woman admitted to the emergency department for the fourth time this year, each time the patient comes with severe injuries and bruises in the body. What is your responsibility as a nurse to prevent such incident to happen again?
- Reports the assault to the local police and write a report**
 - Provides information about safe shelter and support
 - Instructs the woman to move away from her home
 - Discharge the patient to a safe shelter
860. 5-year-old child admitted to the pediatric ward with fracture arm. While assessing the child, the nurse notices a bruise in the child's back. This is the third time in 1 month that the parent brings the child to hospital. Each time, the child family provides vague explanations for various injuries. Which of the following is the pediatric nurse's priority intervention?
- Prevent the parents from visiting the child
 - Question the parents about the child's injury
 - Encourage the child to tell the truth about his injury
 - Report suspicion of abuse to the proper hospital authorities**
861. A child with a ventriculoperitoneal shunt discharge is anticipated after 2 days, the nurse is teaching the parents about signs of shunt malfunction. Which of the following signs indicate shunt malfunction?
- Depressed fontanel
 - Vomiting, lethargy**
 - Increase heart rate
 - Hematuria
862. A 3-year-old child is seen to the emergency department experiencing a seizure at home. There is no previous history of seizure. The mother informs that she does not believe the child has epilepsy. Which of the following is the best response?
- No need to worry because epilepsy is easily treated
 - Very few children have actual epilepsy
 - The seizure may or may not be epilepsy**
 - Your child has had only one seizure
863. A 16-year-old girl was found unconscious in her home by her mother. She was brought to the hospital by ambulance. The patient has no previously known medical history. On arrival, the nurse performs an assessment for Level of consciousness and notes a Glasgow Coma Score of five. The nurse notes the patient's breath smells like acetone or fingernail paint remover. What is the priority therapeutic goal?
- Increase blood glucose
 - Increase serum osmolarity
 - Increase circulatory volume**
 - Decrease intracranial pressure
864. A patient has a defect with the beta cells of the Islet of Langerhans. Which of the following should the nurse most likely expect this patient to exhibit?
- Anemia
 - Appendicitis
 - Cholelithiasis
 - Hyperglycemia**