

Copyright
by
Chenyu Tian
2025

The Dissertation Committee for Chenyu Tian
certifies that this is the approved version of the following dissertation:

Simulation of partial melting material

Committee:

Todd Arbogast, Supervisor

Marc Hesse, Co-supervisor

Omar Ghattas

Patric Heimbach

Thorsten Becker

Simulation of partial melting material

by
Chenyu Tian

Dissertation

Presented to the Faculty of the Graduate School of
The University of Texas at Austin
in Partial Fulfillment
of the Requirements
for the Degree of

Doctor of Philosophy

The University of Texas at Austin
May 2025

Dedication

To all future Longhorn graduate students.

Epigraph

What starts here changes the world.

—University of Texas at Austin

Acknowledgments

Thanks to the previous authors of this template, and to the Graduate School for providing it.

Preface

Readers, this document represents a great deal of time and effort. Enjoy.
Cheers, Chenyu Tian.

Abstract

Simulation of partial melting material

Chenyu Tian, PhD
The University of Texas at Austin, 2025

SUPERVISORS: Todd Arbogast, Marc Hesse

Table of Contents

List of Tables	10
List of Figures	11
Chapter 1: Governing Equations	12
1.1 Mechanics of Mantle Dynamics	12
1.2 Conservation of species and energy	14
1.3 Eutectic Phase Behavior	15
1.4 Rescaling and Nondimensionalization	15
Works Cited	16

List of Tables

List of Figures

1.1	An representative volume element (RVE).	14
-----	---	----

Chapter 1: Governing Equations in Mantle Dynamics

1.1 Mechanics of Mantle Dynamics

Fluid dynamics of the mixture of two phase flow has been extensively discussed. We will discuss the three governing equations of mass and momentum referring to the similar discussion in Drew and Segel (1971), McKenzie (1984) and Katz (2022). Considering a conservation equation for given volumetric variable γ in a representative volume element V .

$$\frac{d}{dt} \int_V \gamma d\mathbf{v} = - \int_{\partial V} \mathbf{f} \cdot d\sigma + \int_V S d\mathbf{v} \quad (1.1)$$

where $\mathbf{f}$ represents flux travelling across the boundary of a representative volume element V . The minus sign indicates that the positive directions aligns with normal unit vector pointing outwards. S , source/sink term, represents the volumetric production rate of γ . We can apply Gauss's theorem to obtain differential form,

$$\frac{\partial \gamma}{\partial t} + \nabla \cdot \mathbf{f} = S \quad (1.2)$$

We also define the quantity of phase fraction rigorously with the help of an indicator function admitting that multiple phase cannot coexist at the same spatial point, which are regarded as immiscible phases. We define indicator function,

$$i(\mathbf{x}) = \begin{cases} 1 & \text{phase}_i \\ 0 & \text{not phase}_i \end{cases} \quad (1.3)$$

We then define macroscopic phase fraction function:

$$\phi_i = \frac{1}{\delta \mathbf{v}} \int_{\delta \mathbf{v}} i(\mathbf{x}) d\mathbf{v} \quad (1.4)$$

In the beginning of my discussion, We will discuss two immiscible phases liquid magma and solid mantle. Phase fraction of liquid magma is labelled as ϕ_l and phase fraction

of solid mantle is labelled as ϕ_s , where $\phi_l + \phi_s = 1$ is always attained. Phase fraction can also be regarded as volume fraction in macroscopic sense.

Substituting bulk density $\bar{\rho} = \phi_l \rho_l + \phi_s \rho_s$ for generalized volumetric variable γ in equation 1.1, we can obtain conservation law for bulk mass,

$$\frac{\partial \bar{\rho}}{\partial t} + \nabla \cdot \bar{\rho} \mathbf{v} = 0 \quad (1.5)$$

with the help of Boussinesq approximation ignoring compressibility of each individual phase, and extend Boussinesq approximation further applying to density difference between each phase except body force term. We can write continuous function for two phase mixture simply as,

$$\nabla \cdot \bar{\mathbf{v}} = 0 \quad (1.6)$$

Substituting phase averaged momentum $\bar{\rho} \mathbf{v}$, We then write conservation of momentum with phase averaged term,

$$\frac{\partial \bar{\rho} \mathbf{v}}{\partial t} + \nabla \cdot \overline{\rho \mathbf{v} \otimes \mathbf{v}} = \bar{\rho} \mathbf{g} + \nabla \cdot \bar{\boldsymbol{\tau}} - \nabla \bar{P} \quad (1.7)$$

where $\boldsymbol{\tau}$ represents deviatoric stress, Ignoring change of inertia term due to near zero Reynolds number in magma and mantle,

$$\bar{\rho} \mathbf{g} + \nabla \cdot \bar{\boldsymbol{\tau}} - \nabla \bar{P} = 0 \quad (1.8)$$

Insert more details later!

Combining equations, I finally arrive with the full flow mechanical system,

$$\phi_f (\mathbf{v}_l - \mathbf{v}_s) + \frac{k_{\phi_l}}{\mu_l} \nabla (p_l - p_f \mathbf{g}) = 0, \quad (1.9)$$

$$\nabla \cdot (\phi_l \mathbf{v}_l + \phi_s \mathbf{v}_s) = 0, \quad (1.10)$$

$$\nabla \bar{p} - \nabla \cdot \boldsymbol{\tau}(\mathbf{v}_s) = (\rho_s - \rho_l) \mathbf{g}, \quad (1.11)$$

$$\mu_s \nabla \cdot \mathbf{v}_s - \phi_f (p_l - p_s) = 0, \quad (1.12)$$

where the deviatoric stress of the mixture is

$$\tau(\mathbf{v}_s) = 2\mu_s\phi_s(\mathcal{D}\mathbf{v}_s - \frac{1}{3}\nabla \cdot \mathbf{v}_s\mathbf{I}) \quad (1.13)$$

and $\mathcal{D}\mathbf{v}_s = \frac{1}{2}(\nabla\mathbf{v}_s + \nabla\mathbf{v}_s^T)$ is the symmetric gradient. Moreover, the relative permeability is $k_{\phi_l} = k_0\phi_l^{2+2\Theta}$, where $\Theta \in [0, 1/2]$. μ_s and μ_l are dynamic viscosities for solid and liquid correspondingly.

1.2 Conservation of species and energy

Chemical compositions of mantle and magma is very complicate and very hard to trace at the same time. We simplify the model to a two-phase system with two species. We negelect detailed description of chemical reaction process and effect on the Gibbs free energies. Similar discussion can be found in extensive literatures like Spiegelman and Elliott (1993), Hewitt and Fowler (2008), Katz (2008) and Marc A. Hesse and Oza (2022). Both two species can be solid or liquid. Two species are immiscible as solid but miscible as liquid. I consider a homogenous density ρ_s for solid phase and ρ_l for liquid phase. Within the representative volume element V ,




Figure 1.1: An representative volume element (RVE).

$\gamma_{i,j}$ represents one quantity of the specie i presenting in phase j , where $i \in 1, 2$ and $j \in \{s, l\}$. The mass fraction $c_{i,j}$ of a given specie i in phase j is often regarded as concentration. Hence, we can write transport equations for a individual specie i

assuming a pseudo melting term Γ ,

$$\frac{\partial}{\partial t}(\phi_l \rho_l c_{i,l}) + \nabla \cdot (\phi_l \rho_l c_{i,l} \mathbf{v}_l - \phi_l \rho_l \mathcal{D}_i \nabla c_{i,l}) = \Gamma_i \quad (1.14)$$

$$\frac{\partial}{\partial t}(\phi_s \rho_s c_{i,s}) + \nabla \cdot (\phi_s \rho_s c_{i,s} \mathbf{v}_s - \phi_s \rho_s \mathcal{D}_i \nabla c_{i,s}) = -\Gamma_i \quad (1.15)$$

where we can safely get rid of dissusion term in solid phase. Summing up 1.14 and 1.15, we can write conservation equation in phase averaged volumetric bulk term,

$$\frac{\partial \overline{\rho c_i}}{\partial t} + \nabla \cdot (\overline{\rho c_i} \mathbf{v} - \phi_l \rho_l \mathcal{D}_i \nabla c_i) = 0 \quad (1.16)$$

where the phase averaged volumetric bulk term $\overline{\rho c_i} = \sum_{j \in \{s,l\}} \phi_j \rho_j c_{i,j}$ and $\overline{\rho c_i} \mathbf{v} = \sum_{j \in \{s,l\}} \phi_j \rho_j c_{i,j} \mathbf{v}_j$.

1.3 Eutectic Phase Behavior

Embracing the fact that Mid-ocean ridge forms at the place where temperature is generally lower, eutectic melting, where melting temperature is significantly lowered with the presence of the secondary specie, is a good model for modelling melting behavior at certain location. The binary eutectic phase behavior describes the chemical behavior of two immiscible crystals from a completely miscible solution. On a typical eutectic phase behavior diagram, we can locate any phase situation using enthalpy and composition information. I follow phase behavior details described in Marc A. Hesse and Oza (2022). Some adjustments are made for our partially molten mantle system.

- The composition of the second specie in the liquid formed by eutectic melting X_e is a function of pressure.
- Melting temperature T_m is a function of pressure.

Following the previous discussion,

1.4 Rescaling and Nondimensionalization

Works Cited

- Donald Drew and Lee Segel. Averaged equations for two-phase flows. *Studies in Applied Mathematics*, page 3, 1971.
- I.J Hewitt and A.C Fowler. Partial melting in an upwelling mantle column. *Proc. R. Soc. A*, 464:2467–2491, 2008.
- Richard F. Katz. Magma dynamics with the enthalpy method: Benchmark solutions and magmatic focusing at mid-ocean ridges. *Journal of Petrology*, 49: 2099–2121, 2008.
- Richard F. Katz. *The Dynamics of Partially Molten Rock*. Princeton University Press, 1st edition, 2022. ISBN 9780691232645.
- Steve D. Vance Marc A. Hesse, Jacob S. Jordan and Apurva V. Oza. Supporting information for "downward oxidant transport through europa's ice shell by density-driven brine percolation". *Geophysical Research Letters*, 2022.
- Dan Mckenzie. The generation and compaction of partially molten rock. *Journal of Petrology*, 25:713–765, 1984.
- Marc Spiegelman and Tim Elliott. Consequences of melt transport for uranium series disequilibrium in young lavas. *Earth and Planetary Science Letters*, 118: 1–20, 1993.