

Chapter 1: Numerical Methods

We introduce a locally mass conserving numerical framework. We solve the static mechanics system with conforming mixed finite element method, which is adjusted to be locally mass conservative. We solve the transport system with finite volume scheme, which is a mass conservative scheme.

1.1 Quadrilateral Mesh

We consider a bounded computational domain $\Omega \subset \mathbb{R}^2$. Here, $\partial\Omega$ denotes the Lipschitz-continuous boundary of the domain. Let $\mathcal{T}_h$ be a conforming mesh of quadrilaterals covering Ω . The quadrilateral $E \in \mathcal{T}_h$ is referred as element in finite element context, while referred as cell in finite volume context. Each quadrilateral E is assumed to be strictly convex, not degenerate into a triangle, line or point. For ease of implementation, we use a logically rectangular mesh, where each element(cell) is indexed using Cartesian coordinates $\{i, j\}$. We show a reference element(cell) $\hat{E}$ and a general quadrilateral element(cell) E in Figure 1.1. There exists a non-affine bijective and smooth map $\mathbf{F}_{\hat{E}} : \hat{E} \rightarrow E$ mapping the vertices of $\hat{E}$ to the vertices of E .




Figure 1.1: An exhibition of a reference square and a general non-degenerate quadrilateral element(cell).

The edges of element E are denoted by e_i , $i \in \{0, 1, 2, 3\}$ in the counterclockwise direction. The vertices are represented as $v_i = e_i \cap e_{i+1}$ in the sense of modulo. Here, $\boldsymbol{\nu}_i$ and $\boldsymbol{\tau}_i$ are used to represent the unit normal and the unit tangent vector respectively. The unit normal vector $\boldsymbol{\nu}_i$ points outwards the quadrilateral. The tangent $\boldsymbol{\tau}_i$ is generated by rotating $\boldsymbol{\nu}_i$ counterclockwise by right-hand rule.

In Figure 1.2, we illustrate a full logically rectangular quasi-uniform quadrilateral mesh $\mathcal{T}_h$, which is generated by randomly distort the interior vertex of a square mesh.




Figure 1.2: An exhibition of a full logically rectangular quasi-uniform quadrilateral mesh $\mathcal{Q}_h$.

1.1.1 Shape Regularity

We give an official definition of uniform shape regular of a quadrilateral mesh. For any quadrilateral $E \in \mathcal{T}_h$, we can get a set of four triangles T_i , $i \in \{0, 1, 2, 3\}$. We form the triangle T_i by excluding vertex v_i from the quadrilateral E , and use the remaining three vertex. We define the following parameters as

$$\begin{aligned} h_E &= \text{diameter of } E, \\ \rho_E &= 2 \min_{1 \leq i \leq 4} \{\text{diameter of largest circle inscribed in } T_i\}. \end{aligned} \tag{1.1}$$

We consider a mesh portion $\mathcal{T}_h$ is uniformly shape regular if there exists a shape regularity parameter σ_* independent of $\mathcal{T}_h$, such that the ratio

$$\rho_E/h_E \geq \sigma_* > 0, \tag{1.2}$$

for all quadrilaterals $E \in \mathcal{T}_h$.

1.2 Direct Serendipity Space

Serendipity element have the same accuracy of approximation with classical tensor product element while use smaller number of degrees of freedom. However, the classical serendipity space suffer from poor approximation on quadrilaterals. We define local shape functions directly on quadrilaterals according to Arbogast et al. (2022). H^1 and $H(\text{div})$ conforming direct mixed finite element space are constructed with the direct serendipity function space.

To begin with, we introduce some notations and distance auxilliary functions. Let $\mathbb{P}_r(\omega)$ denote the space of polynomials of degree up to r on $\omega \subset \mathbb{R}^d$. The dimension of $\mathbb{P}_r$ can be calculated as

$$\dim \mathbb{P}_r(\mathbb{R}^d) = \binom{r+2}{2} = \frac{(r+d)!}{r!d!}. \quad (1.3)$$

We define the linear polynomial $\lambda_i(\mathbf{x})$ giving the distance of $\mathbf{x} \in \mathbb{R}^2$ to edge e_i as

$$\lambda_i(\mathbf{x}) = -(\mathbf{x} - \mathbf{x}_i^*) \cdot \boldsymbol{\nu}_i, \quad i = 0, 1, 2, 3, \quad (1.4)$$

where $\mathbf{x}_i^* \in e_i$ is any point on the edge e_i . The value of distance function λ_i keeps positive as long as $\mathbf{x}$ is in the interior of the quadrilateral. We define two additional distance functions giving the distance of $\mathbf{x} \in \mathbb{R}^2$ to diagonal line d_i joining v_i with v_{i+2} , $i \in 0, 1$. Let $\boldsymbol{\nu}_{d,i}$ be the unit normal vector to the diagonal line d_i . The direction of the $\boldsymbol{\nu}_{d,i}$ is defined by counterclockwise rotating of the vector pointing from v_i to v_{i+2} . The resulting diagonal distance function is defined as

$$\lambda_{d,i}(\mathbf{x}) = -(\mathbf{x} - \mathbf{x}_{d,i}^*) \cdot \boldsymbol{\nu}_{d,i}, \quad i = 1, 2, \quad (1.5)$$

where $\mathbf{x}_{d,i}^*$ is any point on the diagonal line d_i . For simplicity, we use the coordinate of corner v_i in later computation. We denote the coordinates of corner v_i as $\mathbf{x}_{v,i}$.

Now we present set of nodal basis functions in first and second order direct serendipity space $\mathcal{DS}_1$ and $\mathcal{DS}_2$ defined by Arbogast et al. (2022).

1.2.1 Direct Serendipity Space $\mathcal{DS}_2$

We supplement polynomial space $\mathbb{P}_r(E)$, $r \geq 2$ with two linearly independent functions, $\phi_{s,1}(\mathbf{x})$ and $\phi_{s,2}(\mathbf{x})$. We denote the supplemental function space as $\mathbb{S}_r^{\mathcal{DS}}(E) = \text{span}\{\phi_{s,1}, \phi_{s,2}\}$, $r \geq 2$. We present the second order direct serendipity space as

$$\mathcal{DS}_2(E) = \mathbb{P}_2(E) \oplus \mathbb{S}_2^{\mathcal{DS}}(E). \quad (1.6)$$

The supplemental functions are defined as

$$\phi_{s,i}(\mathbf{x}) = \lambda_{i+1}(\mathbf{x})\lambda_{i+2}(\mathbf{x})R_{i,i+2}(\mathbf{x}), \quad i \in \{1, 2\}, \quad (1.7)$$

where i is in the sense of modulo. We use a simple definition for rational function $R_{i,i+2}$ as

$$R_{i,i+2}(\mathbf{x}) = \frac{\lambda_i(\mathbf{x}) - \lambda_{i+2}(\mathbf{x})}{\lambda_i(\mathbf{x}) + \lambda_{i+2}(\mathbf{x})}, \quad (1.8)$$

which satisfies the conditions as

$$R_{i,i+2}|_{e_i} = -1, \quad R_{i,i+2}|_{e_{i+2}} = 1. \quad (1.9)$$

We continue to define

$$R_i(\mathbf{x}) = \frac{1}{2} \left(1 - (-1)^{\lfloor i/2 \rfloor} R_{i,i+2}(\mathbf{x}) \right) = \frac{\lambda_{i+2}(\mathbf{x})}{\lambda_i(\mathbf{x}) + \lambda_{i+2}(\mathbf{x})}, \quad (1.10)$$

where i is understood (mod 2). Here, $R_i|_{e_i} = 1$, and $R_i|_{e_{i+2}} = 0$ by definition. In Figure 1.3, we draw R_0 on a general quadrilateral, which evaluates 1 on the edge e_0 , and evaluates 0 on the opposite edge e_2 .

The edge nodal basis functions for $\mathcal{DS}_2$ are constructed as

$$\varphi_{e,i}^2(\mathbf{x}) = \frac{\lambda_{i+1}(\mathbf{x})\lambda_{i+3}(\mathbf{x})R_i(\mathbf{x})}{\lambda_{i+1}(\mathbf{x}_{e,i})\lambda_{i+3}(\mathbf{x}_{e,i})R_i(\mathbf{x}_{e,i})}, \quad (1.11)$$

where $\mathbf{x}_{e,i}$ is the middle point of the edge e_i , the i index is understood in the sense of modulo of 4. In Figure 1.4, we present $\varphi_{e,0}^2$ on a general quadrilateral.




Figure 1.3: Rational function R_0 .




Figure 1.4: Edge nodal basis function $\varphi_{e,0}^2$ in $\mathcal{DS}_2$.

The vertex nodal basis functions for $\mathcal{DS}_2$ are constructed as

$$\varphi_{v,i}^2(\mathbf{x}) = \frac{\lambda_{i+2}(\mathbf{x})\lambda_{i+3}(\mathbf{x}) - \sum_{k=i}^{i+1} \lambda_{i+2}(\mathbf{x}_{e,i})\lambda_{i+3}(\mathbf{x}_{e,i})\varphi_{e,i}^2(\mathbf{x})}{\lambda_{i+2}(\mathbf{x}_{v,i})\lambda_{i+3}(\mathbf{x}_{v,i}) - \sum_{k=i}^{i+1} \lambda_{i+2}(\mathbf{x}_{e,i})\lambda_{i+3}(\mathbf{x}_{e,i})\varphi_{e,i}^2(\mathbf{x}_{v,i})}, \quad (1.12)$$

where $\mathbf{x}_{v,i}$ is the coordinates of the vertex node v_i . The i index is understood in the sence of module of 4 as well. In Figure 1.5, we show $\varphi_{v,0}^2$ on a general quadrilateral.




Figure 1.5: Vertex nodal basis function $\varphi_{v,0}^2$ in $\mathcal{DS}_2$.

We conclude that unisolvence follows directly from the constructed nodal basis

functions.

1.2.2 Direct Serendipity Space $\mathcal{DS}_1$

The supplemental space for $\mathcal{DS}_1$ differs from that for $\mathcal{DS}_2$. We present the first order serendipity as

$$\mathcal{DS}_1(E) = \mathbb{P}_1(E) \oplus \text{span}\{R\}, \quad (1.13)$$

where $R(\mathbf{x}_{v,0}) = R(\mathbf{x}_{v,2}) = -R(\mathbf{x}_{v,1}) = -R(\mathbf{x}_{v,3}) = 1$. The construction of rational function R is not unique. We present one way to construct $R(\mathbf{x})$ with edge nodal basis function (1.11) in $\mathcal{DS}_2$ as

$$R(\mathbf{x}) = r(\mathbf{x}) - \sum_{i=0}^3 r(\mathbf{x}_{e,i}) \varphi_{e,i}^2(\mathbf{x}), \quad (1.14)$$

where auxiliary function $r(\mathbf{x})$ is defined as

$$r(\mathbf{x}) = \frac{\lambda_2(\mathbf{x})\lambda_3(\mathbf{x})}{\lambda_2(\mathbf{x}_{v,0})\lambda_3(\mathbf{x}_{v,0})} - \frac{\lambda_0(\mathbf{x})\lambda_3(\mathbf{x})}{\lambda_0(\mathbf{x}_{v,1})\lambda_3(\mathbf{x}_{v,1})} + \frac{\lambda_0(\mathbf{x})\lambda_1(\mathbf{x})}{\lambda_0(\mathbf{x}_{v,2})\lambda_1(\mathbf{x}_{v,2})} - \frac{\lambda_1(\mathbf{x})\lambda_2(\mathbf{x})}{\lambda_1(\mathbf{x}_{v,3})\lambda_2(\mathbf{x}_{v,3})}. \quad (1.15)$$

In Figure 1.6, we plot the rational function on a general quadrilateral.




Figure 1.6: Rational function in $\mathcal{DS}_1$.

We define vertex nodal basis function accordingly as

$$\varphi_{v,i}^1(\mathbf{x}) = \frac{\lambda_{d,m}(\mathbf{x}) - \frac{1}{2}\lambda_{d,m}(\mathbf{x}_{v,i+2})(1 + (-1)^i R(\mathbf{x}))}{\lambda_{d,m}(\mathbf{x}_{v,i}) - \lambda_{d,m}(\mathbf{x}_{v,i+2})}, \quad (1.16)$$

where $m \equiv i + 1 \pmod{2}$, and index i is understood in the sense of modulo of 4. In Figure 1.7, we show the vertex nodal basis function $\varphi_{v,0}^1$ in $\mathcal{DS}_1$ on a general quadrilateral, which evaluates 1 at vertex v_0 and 0 at other vertex.




Figure 1.7: Vertex nodal basis function in $\mathcal{DS}_1$.

We first define an interpolation operator of function $v \in L^2(\Omega)$ onto $\mathcal{DS}_r(K_i)$, $r = 1, 2$. Here, K_i has different meanings with different types of basis functions. For each basis functions associated with edge node, we set $K_i = e_i$. For each basis functions associated with vertex node, we set K_i being a set of edges sharing the vertex v_i . Following the work by Arbogast et al. (2022), we define a interpolation operator $\mathcal{I}_h^r : W_p^l(E) \rightarrow \mathcal{DS}_r$ as

$$\mathcal{I}_h^r v(\mathbf{x}) = \sum_{i=1}^{N_r} \varphi_i(\mathbf{x}) \int_{K_i} \psi_i(\mathbf{y}) v(\mathbf{y}) d\mathbf{y} \in \mathcal{DS}_r, \quad (1.17)$$

where $1 \leq p \leq \infty$ and $l > 1/p$ (but $l \geq 1$ if $p = 1$), and $\psi(\mathbf{x})$ is the dual nodal basis such that

$$\int_{K_i} \psi_i(\mathbf{x}) \varphi_j(\mathbf{x}) d\mathbf{x} = \delta_{i,j}, \quad i, j \in \{0, 1, 2, \dots, N_r\}, \quad (1.18)$$

where N_r is the number of nodal basis defined on K_i . We present the following two lemmas regarding the constructed interpolation operator for future reference.

Lemma 1.1. (*Boundedness of the interpolation operator Arbogast et al. (2022)*)

Let $\mathcal{T}_h$ be uniformly shape regular (1.2) with shape regularity parameter σ_* . For each element $E \in \mathcal{T}_h$, There exists a bounded interpolation operator (1.17). For

every $E \in \mathcal{T}_h$, suppose that the basis functions of $\mathcal{DS}_r(E)$ are constructed using λ_{24} and λ_{13} such that the zero set $\mathcal{Z}_{i,i+2}$ intersects e_{i+3} and e_{i+1} . Moreover, assume that $R_{i,i+2}$ is uniformly differentiable functions of the vertices of E up to order m . Then for $r \geq 1$, $E \in \mathcal{T}_h$, $1 \leq q \leq \infty$, and any nonnegative integer m ,

$$\|\mathcal{J}_h^r v\|_{W_q^m} \leq C(\sigma_*, m, q) \sum_{k=0}^l h_E^{k-m+\frac{2}{q}-\frac{2}{p}} |v|_{W_p^k(E^*)}, \quad (1.19)$$

where $E^* = \cup_{F \in \mathcal{T}_h, F \cap E \neq \emptyset} F$ and $|\cdot|_{W_p^k}$ is the seminorm of k -th order derivatives.

Lemma 1.2. (Approximability of the interpolation operator Arbogast et al. (2022))

Under the assumptions of 1.1, there exists a constant $C = C(r, \sigma_*) > 0$ such that for all functions $v \in W_{l,p}(E^*)$, with $1 \leq p \leq \infty$ and $l > 1/p$ or ($l \geq 1$ if $p = 1$),

$$\|v - \mathcal{J}_h^r v\|_{W_p^m} \leq C h_E^{l-m} |v|_{W_p^l(E^*)}, \quad 0 \leq m \leq \min(l, r+1). \quad (1.20)$$

Moreover, there exists a constant $C = C(r, \sigma_*) > 0$, independent of $h = \max_{E \in \mathcal{T}_h} h_E$, such that for all functions $v \in W_p^l(\Omega)$,

$$\left(\sum_{E \in \mathcal{T}_h} \|v - \mathcal{J}_h^r v\|_{W_p^m}^p \right)^{1/p} \leq C h^{l-m} |v|_{W_p^l(\Omega)}, \quad 0 \leq m \leq \min(l, r+1). \quad (1.21)$$

1.3 Mixed Finite Element

We define H^1 and $H(\text{div})$ conforming mixed element space with the direct serendipity space, and test with simple stokes and darcy problems respectively.

1.3.1 $H(\text{div})$ Conforming Element

We have some choice for $H(\text{div})$ confirming space. For example, we can choose the famous Raviart-Thomas element by Raviart and Thomas (1977). The serendipity space is the precursor of the Brezzi-Douglas-Marini element by Brezzi et al. (1985). We recover Arbogast-Correa element by Arbogast and Correa (2016) if we use mapped direct serendipity space. We generated a reduced $H(\text{div})$ approximation, that $\nabla \cdot \mathbf{u}$ is approximated one less order than approximation of $\mathbf{u}$.

We write the rotated 2D exact sequence for energy space as

$$H^1 \xrightarrow{\text{curl}} H(\text{div}) \xrightarrow{\text{div}} L^2 \quad (1.22)$$

$$\cup \quad \quad \cup \quad \quad \cup \quad (1.23)$$

$$\mathcal{DS}_2 \xrightarrow{\text{curl}} \mathbf{V}_r^{r-1} \xrightarrow{\text{div}} \mathbb{P}_{r-1}. \quad (1.24)$$

We define the first order reduced $H(\text{div})$ conforming space with $\mathcal{DS}_2$ as

$$\mathbb{V}_1^0 = \mathbb{P}_1^2 \oplus \text{curl}\mathbb{S}_2^{\mathcal{DS}}. \quad (1.25)$$

The reduced space $\mathbb{V}_1^0$ can be given a global basis with the normal components of the shape functions merged continuously on the shared edge of two elements. The dimension of the derived space is

$$\dim(\mathbb{V}_1^0) = 2 \times 3 + 2 = 8. \quad (1.26)$$

We need two independent basis functions on each edge of a quadrilateral E , one of which has a vanishing average flux on the edge and the other one has a non-vanishing average flux on the edge. On each edge, we define two independent basis functions, $\varphi_{l,i}$ and $\varphi_{c,i}$, such that

$$\varphi_{c,i} \cdot \boldsymbol{\nu}_j|_{e_j} = \begin{cases} 0, & j \neq i, \\ 1, & j = i. \end{cases} \quad (1.27)$$

$$\varphi_{l,i} \cdot \boldsymbol{\nu}_j|_{e_j} = \begin{cases} 0, & j \neq i, \\ l_i(\mathbf{x}), & j = i, \end{cases} \quad (1.28)$$

where l_i is a non vanishing linear polynomial defined on edge e_i orthogonal to one, i.e., $\int_{e_i} l_i ds = 0$.

To begin with, we write curl of the auxiliary rational function R_i (1.8) as

$$\text{curl}R_i(\mathbf{x}) = \frac{\boldsymbol{\tau}_{i+2}(\mathbf{x})\lambda_i(\mathbf{x}) - \boldsymbol{\tau}_i(\mathbf{x})\lambda_{i+2}(\mathbf{x})}{(\lambda_i(\mathbf{x}) + \lambda_{i+2}(\mathbf{x}))^2}, \quad (1.29)$$

where the index i is understood with modulo of 4. We first present a construction of basis function $\varphi_{l,i}$ as

$$\tilde{\varphi}_{l,i} = \text{curl}(\lambda_{i+1}\lambda_{i+3}R_i) = (\boldsymbol{\tau}_{i+1}\lambda_{i+3} + \lambda_{i+1}\boldsymbol{\tau}_{i+3})R_i + \lambda_{i+1}\lambda_{i+3}\text{curl}R_i. \quad (1.30)$$

We then normalize it such that l_i varying from -1 to 1 as

$$\boldsymbol{\varphi}_{l,i} = \frac{\tilde{\boldsymbol{\varphi}}_{l,i}(\mathbf{x})|e_i|}{4\lambda_{i+1}(\mathbf{x}_{e,i})\lambda_{i+3}(\mathbf{x}_{e,i})R_i(\mathbf{x}_{e,i})}, \quad (1.31)$$

where the index i is understood in the sense of modulo 4, and edge length $|e_i| = |\mathbf{x}_{v,i} - \mathbf{x}_{v,i+3}|$. In Figure 1.8, we draw $\boldsymbol{\varphi}_{l,2}$ on the left element and $\boldsymbol{\varphi}_{l,0}$ on the right element sharing a edge, and show normal flux joining continuously as a linear function on the shared edge.




Figure 1.8: Linear basis function $\boldsymbol{\varphi}_l$.

The construction of a the constant basis function is a bit more complicated. We follow the process by Arbogast et al. (2022). We start by defining a linear nodal function $\phi_{v,i}^* \in \mathcal{DS}_2(E)$ such that $\phi_{v,i}^*(\mathbf{x}_{v,k}) = \delta_{ik}$. Its restriction to each edge of E is a linear function as

$$\phi_{v,i}^*(\mathbf{x}) = \varphi_{v,i}^2(\mathbf{x}) + \left[\frac{1}{2}\varphi_{e,i}^2 + \frac{1}{2}\varphi_{e,i+1}^2\right], \quad (1.32)$$

where the index i is understood in modulo of 4. Then define $\boldsymbol{\varphi}_{v,i}^*(\mathbf{x}) = \text{curl}\phi_{v,i}^*$ so that

$$\boldsymbol{\varphi}_{v,i}^* \cdot \boldsymbol{\nu}_j|_{e,j} = \begin{cases} 1/|e_i|, & j = i, \\ -1/|e_i|, & j = i + 1, \\ 0, & j = i + 2, i + 3. \end{cases} \quad (1.33)$$

We then define the "constant" basis function defined on the edge as

$$\boldsymbol{\varphi}_{e,i} = \frac{(\mathbf{x}_{v,i} - \mathbf{x}_{v,i+2}) \cdot \boldsymbol{\nu}_{i+1}|e_{i+1}|\boldsymbol{\varphi}_{v,i}^* + \mathbf{x} - \mathbf{x}_{v,i+2}}{(\mathbf{x}_{v,i} - \mathbf{x}_{v,i+2}) \cdot \boldsymbol{\nu}_{i+1}|e_{i+1}|/|e_i| + (\mathbf{x}_{v,i} - \mathbf{x}_{v,i+2}) \cdot \boldsymbol{\nu}_i}, \quad (1.34)$$

where the index i is again understood in the sense of modulo of 4. The defined basis function $\varphi_{e,i}$ has flux of constant value 1 on e_i and 0 on all the other edges. In Figure 1.9, we draw $\varphi_{c,2}$ on the left element and $\varphi_{c,0}$ on the right element sharing a edge, and show normal flux joining continuously as a constant 1 on the shared edge.




Figure 1.9: Constant basis function φ_c .

Lemma 1.3. *Stability and approximation properties Arbogast et al. (2022).*

Under the assumption of 1.1, there exists a constant $C > 0$, independent of $\mathcal{T}_h$ and $h > 0$, such that

$$\|\mathbf{v} - \pi\mathbf{v}\|_{0,\Omega} \leq C \|\mathbf{v}\|_{k,\Omega} \quad (1.35)$$

1.3.2 Test Darcy Problem

We test our $H(\text{div})$ conforming element on a simple darcy problem as

$$\begin{aligned} \mathbf{u} + \nabla p &= \mathbf{g} \\ \nabla \cdot \mathbf{u} &= 0 \end{aligned} \quad (1.36)$$

weak form

$$\begin{aligned} (\mathbf{u}, \mathbf{v})_\Omega - (p, \nabla \cdot \mathbf{v})_\Omega &= (\mathbf{g}, \mathbf{v})_\Omega - \langle p_0, v_n \rangle_{\Gamma_i} - \langle \mathbf{u}_0, \mathbf{v} \rangle_{\Gamma_u} \\ (\nabla \cdot \mathbf{u}, \omega) &= 0 \end{aligned} \quad (1.37)$$

Test space $\{\mathbf{v} \in H(\text{div}, \Omega) : v_n = 0 \text{ on } \Gamma_u\}$.

1.3.3 H^1 Conforming Element

Stable mixed finite element pair for Stokes problem has been widely researched. We are referring to the element discussed by Christine Bernardi (1985) and by Fortin

(1981). A more general type of element was discussed by Arbogast and Wheeler (2005).

Vertex basis mapping to sobolev space difficulty. Need Clément interpolation. Define Π operator two parts.

We consider the following space for each element $E \in \mathcal{T}_h$ as

$$V_1^{\text{BR}}(E) = (\mathcal{DS}_1(E))^2 \oplus \text{span}\{\mathbf{b}_0, \mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}, \quad (1.38)$$

where $\mathbf{b}_i$ is a quadratic bubble function defined on each edge e_i . We pick previously defined function $\varphi_{e,i} \in \mathcal{DS}_2$ as

$$\mathbf{b}_i = \varphi_{e,i}(\mathbf{x})\mathbf{n}_i \quad (1.39)$$

which is quadratic when restricted to edge e_i .

Lemma 1.4. *For an element $E \in \mathcal{T}_h$, $\mathbf{u}_h$ is uniquely defined by the following degrees of freedom:*

(i) *for corner point v_i and Cartesian direction unitnormal vectors $\{\mathbf{i}, \mathbf{j}\}$,*

$$DOF_{(v_i, \mathbf{i})}^{(i)}(\mathbf{u}_h) = \mathbf{u}_h(v_i) \cdot \mathbf{i}, \quad DOF_{(v_i, \mathbf{j})}^{(i)}(\mathbf{u}_h) = \mathbf{u}_h(v_i) \cdot \mathbf{j};$$

(ii) *for edge e_i ,*

$$DOF_{e_i}^{(ii)} = \langle \mathbf{u}_h \cdot \boldsymbol{\nu}_i, 1 \rangle,$$

Proof. We prove the lemma by construction. We pick nodal basis on vertex v_i in two Cartesian direction as

$$N_{v_i, x} = \begin{bmatrix} \psi_{v,i}(\mathbf{x}) \\ 0 \end{bmatrix}, \quad N_{v_i, y} = \begin{bmatrix} 0 \\ \psi_{v,i}(\mathbf{x}) \end{bmatrix} \quad (1.40)$$

We pick edge basis on edge e_i as

$$N_e = \varphi_{e,i} \boldsymbol{\nu}_i \quad (1.41)$$

Note that bubble functions are quadratic when restricted to edge e_i , while nodal basis functions are linear when restricted to edge e_i , therefore bubble functions cannot be in the same space as nodal basis function. We conclude that this element is well-defined, i.e., the degrees of freedom uniquely determine the function. $\square$

We need a π operator, which mappes $(H^1(E))^2$ onto $V_1^{\mathcal{BR}}$ for our spaces. For $\mathbf{v} \in (H^1(\Omega))^2$, let $\pi\mathbf{v} \in \mathbb{V}_1^{\mathcal{BR}}$ be defined as the interpolant of the degrees of freedom from 1.4. For each F , we require the following as

(i) For each corner point $v \in \partial F$

$$\pi\mathbf{v}(\mathbf{x}_v) \cdot \mathbf{i} = \mathcal{J}_{\mathcal{DS}_1} \mathbf{v}(\mathbf{x}_v) \cdot \mathbf{j}. \quad (1.42)$$

(ii) On each edge $e \subset \partial E$,

$$\langle \pi\mathbf{u} \cdot \boldsymbol{\nu}, \lambda_0 \rangle = \langle \mathbf{u} \cdot \boldsymbol{\nu}, \lambda_0 \rangle_e \quad \forall \lambda_0 \in P_0(e). \quad (1.43)$$

The interpolation operator $\mathcal{I}_{\mathcal{DS}_1}$ has been defined in 1.17. We claim that we can represent for $\mathbf{u} \in (H^1(\Omega))^2$,

$$\pi\mathbf{u} = \mathcal{I}_{\mathcal{DS}_1} \mathbf{u} + \sum_{i=0}^3 \alpha_i \mathbf{b}_i, \quad (1.44)$$

where the coefficient $\alpha_i = \left(\int_{e_i} (\mathbf{u} - \mathcal{J}_{\mathcal{DS}_1} \mathbf{u}) \cdot \boldsymbol{\nu}_i d\sigma \right) / \int_{e_i} \mathbf{b}_i \cdot \boldsymbol{\nu}_i d\sigma$. It is obvious that $\pi\mathbf{u} \in \mathbb{V}_1^{\mathcal{BR}}$. We now prove the approximability and stability of our projection operator and eventually derive the inf – sup condition.

Lemma 1.5. *Assuming that $u \in (H^1(\Omega))^2$.*

(i) *there exists a bounded linear operator π*

(ii) $\forall \mathbf{v} \in (H^1(\Omega))^2$

1.3.4 Test Stokes Problem

We test our modified Bernardi-Raugel elements on the incompressible isotropic elasticity problem with Dirichlet boundary condition:

Find the displacement $\mathbf{u}$ and pressure p such that

$$\begin{aligned} -\nabla \cdot \boldsymbol{\tau}(\mathbf{u}) + \nabla q &= \mathbf{f} \quad \text{in } \Omega, \\ \nabla \cdot \mathbf{u} &= 0 \quad \text{in } \Omega, \\ \mathbf{u} &= \mathbf{g} \quad \text{on } \partial\Omega, \end{aligned} \quad (1.45)$$

where $f \in L^2(\Omega)$, and $\boldsymbol{\tau}(\mathbf{u}) = 2(\mathcal{D}\mathbf{u} - 1/3\nabla \cdot \mathbf{u}\mathbf{I})$. The symmetrical gradient $\mathcal{D}\mathbf{u} = 1/2(\nabla\mathbf{u} + \nabla\mathbf{u}^T)$.

Define a simple choice of energy space

$$\begin{aligned}\mathbb{V} &= \{\mathbf{v} \in (H^1\Omega)^2 : \mathbf{v} = \mathbf{0} \text{ on } \partial\Omega\}, \\ \mathbb{U} &= \{\mathbf{u} \in (H^1\Omega)^2 : \mathbf{u} = \mathbf{g} \text{ on } \partial\Omega\} = \tilde{\mathbf{g}} + \mathbb{V},\end{aligned}\tag{1.46}$$

where $\tilde{\mathbf{g}}$ is a finite energy lift. We assume that $\mathbf{g} \in (H^{1/2}(\Omega))^d$ and understand as a continuous extension $\tilde{\mathbf{g}} \in (H^1(\Omega))^d$ from the boundary into the domain. We write the problem in weak form as:

Find $\mathbf{u} \in \mathbb{U}$ and $p \in L^2(\Omega)$ such that

$$\begin{aligned}(\boldsymbol{\tau}, \nabla\mathbf{v}) - (p, \nabla \cdot \mathbf{v}) &= (\mathbf{f}, \mathbf{v}), \quad \forall \mathbf{v} \in \mathbb{V}, \\ (\nabla \cdot \mathbf{u}, \omega) &= 0 \quad \forall \omega \in L^2(\Omega).\end{aligned}\tag{1.47}$$

We consider the test problem (1.47) defined on a square region $\Omega = [-1, 1]^2$. We manufacture a displacement solution $\mathbf{u} = [x^3y^2, -x^2y^3]$ and pressure $p = y - x$. The source term can be calculated as

$$\mathbf{f} = -2 \begin{bmatrix} 3xy^2 + x^3 \\ -3x^2y - y^3 \end{bmatrix} + \begin{bmatrix} -1 \\ 1 \end{bmatrix}.\tag{1.48}$$

Solution are computed on rectangular and quadrilateral meshes. The convergence rate versus the number of elements will be presented.

1.4 Coupled Degenerate System

Consider a degenerate scaled darcy-stokes coupled boundary-value problem:

Find velocity-pressure potential pair $(\check{\mathbf{v}}_r, \check{q}_l)$ and $(\check{\mathbf{v}}_s, \check{q})$ such that

$$\check{\mathbf{v}}_r + \phi^{1+\Theta}\nabla(\phi^{-1/2}\check{q}_l) = 0,\tag{1.49}$$

$$\phi^{-1/2}\nabla \cdot (\phi^{1+\Theta}\check{\mathbf{v}}_r) + \frac{1}{1-\phi}(\check{q}_l - \phi^{1/2}\check{q}) = 0,\tag{1.50}$$

$$\nabla\check{q} - \nabla \cdot \check{\boldsymbol{\tau}}(\check{\mathbf{v}}_s) = -(1-\phi)\mathbf{n},\tag{1.51}$$

$$\nabla \cdot \check{\mathbf{v}}_s - \frac{\phi^{1/2}}{1-\phi}(\check{q}_l - \phi^{1/2}\check{q}) = 0,\tag{1.52}$$

and

$$\begin{aligned}
\check{\mathbf{v}}_r &= \mathbf{g}_r & \text{on } \Gamma_{u,D}, \\
\check{\mathbf{v}}_s &= \mathbf{g}_s & \text{on } \Gamma_{u,S}, \\
p &= p_0 & \text{on } \Gamma_{i,D}, \\
\boldsymbol{\tau} \cdot \boldsymbol{\nu} - p &= \mathbf{t}_0 & \text{on } \Gamma_{i,S},
\end{aligned} \tag{1.53}$$

where $\Gamma_{u,D}$ and $\Gamma_{u,S}$ denote the part of boundary we assign essential boundary conditions for Darcy and Stokes problem respectively, and $\Gamma_{i,D}$ and $\Gamma_{i,S}$ represent the part of boundary we assign natural boundary conditions respectively. We have $\Gamma_{u,D} \cup \Gamma_{i,D} = \partial\Omega$ and $\Gamma_{u,S} \cup \Gamma_{i,S} = \partial\Omega$.

We follow the procedure by Arbogast et al. (2017) to discretize this boundary value problem. To begin with, a new Hilbert space is defined for this specific scaled formulation, where the scaled velocity $\check{\mathbf{v}}_r$ should lie in, so

$$\mathbb{V}_d^\phi = H_\phi(\text{div}; \Omega) = \left\{ \mathbf{v} \in (L^2(\Omega))^2 : \phi^{-1/2} \nabla \cdot (\phi^{1+\Phi} \mathbf{v}) \in L^2(\Omega) \right\}, \tag{1.54}$$

which equipped with the inner product as

$$(\mathbf{u}_\phi, \mathbf{v}_\phi)_{\mathbb{V}_r^\phi} = (\mathbf{u}_\phi, \mathbf{v}_\phi) + (\phi^{-1/2} \nabla \cdot (\phi^{1+\Theta} \mathbf{u}_\phi), \phi^{-1/2} \nabla \cdot (\phi^{1+\Theta} \mathbf{v}_\phi)). \tag{1.55}$$

The normal trace on $\partial\Omega$ is well-defined as well as

$$\left\| \phi^{1/2+\Theta} \mathbf{v}_\phi \cdot \boldsymbol{\nu} \right\|_{-1/2, \partial\Omega} < C_\Omega \left\{ \|\mathbf{v}_\phi\| + \left\| \phi^{-1/2} \nabla \cdot (\phi^{1+\Theta} \mathbf{v}_\phi) \right\| \right\}. \tag{1.56}$$

1.4.1 The weak formulation

We define the following function spaces

$$\begin{aligned}
\mathbb{V}_D^\phi &= \left\{ \mathbf{v} \in H_\phi(\text{div}; \Omega) : \phi^{1/2+\Theta} \mathbf{v} \cdot \boldsymbol{\nu} = 0 \text{ on } \Gamma_{u,D} \right\}, \\
\mathbb{W}_D &= L^2(\Omega), \\
\mathbb{V}_S &= \left\{ \mathbf{v} \in (H^1(\Omega))^2 : \mathbf{v} = \mathbf{0} \text{ on } \Gamma_{u,S} \right\}, \\
\mathbb{W}_S &= L^2(\Omega),
\end{aligned} \tag{1.57}$$

where each space is equipped with their natural norm. To impose essential boundary conditions, we assume that a function $\mathbf{g}_s = \check{\mathbf{v}}_{s,0}$ on $\Gamma_{u,S}$ and $\mathbf{g}_s = \mathbf{0}$ on $\partial\Omega/\Gamma_{u,S}$.

Then $\mathbf{g}_s \in (H^{1/2}(\partial\Omega))^2$ and extended it continuously from the boundary to the domain, so that the extension $\mathbf{g}_s \in (H^1(\Gamma_{u,S}))^2$ and $\|\mathbf{g}_{s,0}\|_1 \leq C \|\mathbf{v}_{s,0}\|_{1/2,\Gamma_{u,S}}$. Similarly, we can define a bounded extension of $\check{\mathbf{v}}_r \in \mathbb{V}_D^\phi$ such that

Scaled nondimensionalized weak form. Find $\check{\mathbf{v}}_r \in \mathbb{V}_D^\phi + \tilde{\mathbf{g}}_r$, $\check{q}_l \in \mathbb{W}_D$, $\check{\mathbf{v}}_s \in \mathbb{V}_S + \tilde{\mathbf{g}}_S$, and $\check{q} \in \mathbb{W}_S$ such that

$$(\check{\boldsymbol{\tau}}(\mathbf{v}_s), \nabla \boldsymbol{\psi}_s) - (\check{q}, \nabla \cdot \boldsymbol{\psi}_s) = -((1 - \phi)\mathbf{n}, \boldsymbol{\psi}_s) \quad \forall \boldsymbol{\psi}_s \in \mathbb{V}_S, \quad (1.58)$$

$$(\nabla \cdot \check{\mathbf{v}}_s, \omega) + \left(\frac{\phi^{1/2}}{1 - \phi} (\check{q}_l - \phi^{1/2} \check{q}), \omega \right) = 0 \quad \forall \omega \in \mathbb{W}_S, \quad (1.59)$$

$$(\check{\mathbf{v}}_r, \boldsymbol{\psi}_r) - (\check{q}_l, \phi^{-1/2} \nabla \cdot (\phi^{1+\Theta} \check{\mathbf{v}}_r)) = 0, \quad \forall \boldsymbol{\psi}_r \in \mathbb{V}_D^\phi, \quad (1.60)$$

$$(\phi^{-1/2} \nabla \cdot (\phi^{1+\Theta} \check{\mathbf{v}}_r), \omega_f) + \left(\frac{1}{1 - \phi} (\check{q}_l - \phi^{1/2} \check{q}), \omega_f \right) = 0 \quad \forall \omega_f \in \mathbb{W}_D. \quad (1.61)$$

The Existence and uniqueness of the solution has been proved by Arbogast et al. (2017). We will not repeat the proof here.

1.4.2 MFEM discretization

We discretize the formed degenerate system with previously defined $H(\text{div})$ and H^1 conforming space. We write the function spaces out as

$$\begin{aligned} \mathbb{V}_1^0 &= (\mathbb{P}_1)^2 \oplus \text{curl} \mathbb{S}_2^{\mathcal{DS}} \\ \mathbb{W} &= \mathbb{P}_0 \\ \mathbb{V}_1^{\mathcal{BR}} &= \mathcal{DS}_1 \oplus \text{span}\{\mathbf{b}_0, \mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\} \\ \mathbb{W} &= \mathbb{P}_0. \end{aligned} \quad (1.62)$$

We can impose essential boundary conditions

Scaled mixed finite element method. Find $\check{\mathbf{v}}_{r,h} \in \mathbb{V}_1^0 + \hat{\mathbf{g}}_r$, $\check{q}_{r,h} \in \mathbb{W}_D$, $\mathbf{v}_{s,h} \in$

$\mathbf{V}_S + \hat{\mathbf{g}}_s$ and $\check{q}_h \in \mathbb{W}_S$ such that

$$(\check{\boldsymbol{\tau}}(\mathbf{v}_{s,h}), \nabla \boldsymbol{\psi}_s) - (\check{q}_h, \nabla \cdot \boldsymbol{\psi}_s) = -((1 - \phi)\mathbf{n}, \boldsymbol{\psi}_s) \quad \forall \boldsymbol{\psi}_s \in \mathbb{V}_1^{\mathcal{BR}}, \quad (1.63)$$

$$(\nabla \cdot \check{\mathbf{v}}_{s,h}, \omega) - \left(\frac{\phi^{1/2}}{1 - \phi} (\check{q}_{l,h} - \phi^{1/2} \check{q}_h), \omega \right) = 0 \quad \forall \omega \in \mathbb{W}_0^{\mathcal{BR}}, \quad (1.64)$$

$$(\check{\mathbf{v}}_{r,h}, \boldsymbol{\psi}_r) - (\check{q}_{l,h}, \phi^{-1/2} \nabla \cdot (\phi^{1+\Theta} \boldsymbol{\psi}_{r,h})) = 0, \quad \forall \boldsymbol{\psi}_r \in \mathbb{V}_1^0, \quad (1.65)$$

$$(\phi^{-1/2} \nabla \cdot (\phi^{1+\Theta} \check{\mathbf{v}}_{r,h}), \omega_f) + \left(\frac{1}{1 - \phi} (\check{q}_{l,h} - \phi^{1/2} \check{q}_h), \omega_f \right) = 0 \quad \forall \omega_f \in \mathbb{W}_0. \quad (1.66)$$

1.4.3 Local Mass Conservation

Before we continue the discussion, we need to make sure the formulation is well-defined by not dividing by zero. The term $\phi^{-1/2}$ in the symmetric equations (1.65) and (1.66) causes trouble when there is no melting, i.e., $\phi = 0$. In order to deal with it, we first define a element-averaged $\bar{\phi}_E$ as

$$\bar{\phi}_E = \frac{1}{|E|} \int_E \phi(\mathbf{x}) d\mathbf{x}, \quad (1.67)$$

where $|E|$ is the area of element E . Let

$$\phi_E = \begin{cases} \bar{\phi}_E & \text{if } \bar{\phi}_E \neq 0, \\ 0 & \text{if } \bar{\phi}_E = 0. \end{cases} \quad (1.68)$$

We now replace the $\phi^{-1/2}$ with $\phi_E^{-1/2}$. Then we expand the integral with divergence theorem as

$$\begin{aligned} \int_E \phi_E^{-1/2} \nabla \cdot (\phi^{1+\Theta} \boldsymbol{\psi}_{r,h}) \omega_l &= \int_{\partial E} \phi_E^{-1/2} \phi^{1+\Theta} \boldsymbol{\psi}_{r,h} \cdot \boldsymbol{\nu} \omega_f ds \\ &\quad - \int_E \phi_E^{-1/2} \phi^{1+\Theta} \boldsymbol{\psi}_{r,h} \cdot \nabla \omega_l d\mathbf{x}. \end{aligned} \quad (1.69)$$

The second term vanishes when ω_l is constant on element E . We define an auxiliary auxiliary space ω_E begin 1 on E and 0 elsewhere. We sum equations (1.64) and (1.66) together assuming that $\omega = \omega_E \in \mathbb{W}_S$ and $\omega_l = \phi_E^{-1/2} \omega_E \in \mathbb{W}_D$. We can achieve local mass conservation of the phase averaged mixture if we replace the term $\phi^{1/2}$ in equation (1.64) with $\phi_E^{1/2}$ as

$$\int_E \nabla \cdot (\phi^{1+\Theta} \check{\mathbf{v}}_r + \check{\mathbf{v}}_s) d\mathbf{x} = 0. \quad (1.70)$$

We now propose our local mass conserved version as

$$(\check{\boldsymbol{\tau}}(\mathbf{v}_{s,h}), \nabla \boldsymbol{\psi}_s) - (\check{q}_h, \nabla \cdot \boldsymbol{\psi}_s) = -((1 - \phi)\mathbf{n}, \boldsymbol{\psi}_s) \quad \forall \boldsymbol{\psi}_s \in \mathbb{V}_1^{\mathcal{BR}}, \quad (1.71)$$

$$(\nabla \cdot \check{\mathbf{v}}_{s,h}, \omega) - \left(\frac{\phi_E^{1/2}}{1 - \phi} (\check{q}_{l,h} - \phi^{1/2} \check{q}_h), \omega \right) = 0 \quad \forall \omega \in \mathbb{W}_0^{\mathcal{BR}}, \quad (1.72)$$

$$(\check{\mathbf{v}}_{r,h}, \boldsymbol{\psi}_r) - (\check{q}_{l,h}, \phi_E^{-1/2} \nabla \cdot (\phi^{1+\Theta} \boldsymbol{\psi}_{r,h})) = 0, \quad \forall \boldsymbol{\psi}_r \in \mathbb{V}_1^0, \quad (1.73)$$

$$(\phi_E^{-1/2} \nabla \cdot (\phi^{1+\Theta} \check{\mathbf{v}}_{r,h}), \omega_f) + \left(\frac{1}{1 - \phi} (\check{q}_{l,h} - \phi^{1/2} \check{q}_h), \omega_f \right) = 0 \quad \forall \omega_f \in \mathbb{W}_0. \quad (1.74)$$

1.4.4 Test Run

We test our MFEM setup with two problems. First, we test with a manufactured solution. Second, we test a more complex yet a more realistic problem.

Problem setup 1:

We solve a manufactured system of equations of the rectangular domain $-1 < x < 1$,

$$\mathbf{v}_s = \quad (1.75)$$

Problem setup 2:

We solve the full system of equations on the rectangular domain $-160\text{km} < x < 160\text{km}$, $0, x, 160\text{km}$. The boundary conditions are define by imposing

$$\mathbf{v}_s = \quad (1.76)$$

1.5 Saddle Point System

Saddle point system is widely spread in technical and science scenarios, such as constrained optimization problem and has been widely researched. In our case, saddle point system is formed along with Mixed Finite Element discretization. In this section, we give a brief overview of characteristics of a saddle point system and

some common way to solve such system. A saddle point system is formed as

$$\begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{B}^T & \mathbf{C} \end{bmatrix} \begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix} = \begin{bmatrix} \mathbf{f} \\ \mathbf{g} \end{bmatrix}. \quad (1.77)$$

We made the following assumptions

- $\mathbf{A}$ is symmetric: $\mathbf{A} = \mathbf{A}^T$;
- $\mathbf{A}$ is nonsingular: $\mathbf{A}^{-1}$ exists;
- $\mathbf{C}$ is symmetric and positive semidefinite;
- $\mathbf{B}$ is rank deficient.

The null space of $\mathbf{B}$ matrix $\ker(\mathbf{B})$ in our darcy problem is not fixed due to evolution of melting (change of porosity ϕ). Considering the need of parallel, a proper iterative method is discussed.

1.5.1 SPS Induced by MFEM

Now we discuss the saddle point system (SPS) induced by MFEM method in details including the boundary conditions. To begin with, we regroup dofs according to the boundary condition.

$$\mathbf{A} = \begin{bmatrix} \mathbf{A}_K & \mathbf{A}_M \\ \mathbf{A}_M^T & \mathbf{M} \end{bmatrix}, \quad (1.78)$$

where $\mathbf{A}_K$ consists of interaction within interior dofs and natural boundary condition dofs, and $\mathbf{A}_M$ consists of interaction between interior dofs, natural boundary dofs and essential boundary dofs.

1.5.2 Degenerate Coupled Saddle Point Problem and Uzawa Iteration

The linear system formed from the equations (1.71), (1.72), (1.73) and (1.74) is

$$\begin{bmatrix} \mathbf{A}_s & -\mathbf{B}_s & \mathbf{0} & \mathbf{0} \\ \mathbf{B}_s^T & \mathbf{C}_s & \mathbf{0} & \mathbf{K} \\ \mathbf{0} & \mathbf{0} & \mathbf{A}_d & -\mathbf{B}_d \\ \mathbf{0} & \mathbf{K} & \mathbf{B}_d^T & \mathbf{C}_d \end{bmatrix} \begin{bmatrix} \mathbf{x}_s \\ \mathbf{y}_s \\ \mathbf{x}_d \\ \mathbf{y}_d \end{bmatrix} = \begin{bmatrix} \mathbf{f}_s \\ \mathbf{g}_s \\ \mathbf{f}_d \\ \mathbf{g}_d \end{bmatrix}, \quad (1.79)$$

wherein the solution represents the degrees of freedom of primary variables with respect to the bases of the finite element spaces. (Insert details of calculation of each blocked matrix). We form a double saddle point system by symmetrically permutate original linear system,

$$\begin{bmatrix} \mathbf{A}_s & \mathbf{0} & -\mathbf{B}_s & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_d & \mathbf{0} & -\mathbf{B}_d \\ \mathbf{B}_s^T & \mathbf{0} & \mathbf{C}_s & \mathbf{K} \\ \mathbf{0} & \mathbf{B}_d^T & \mathbf{K} & \mathbf{C}_d \end{bmatrix} \begin{bmatrix} \mathbf{x}_s \\ \mathbf{x}_d \\ \mathbf{y}_s \\ \mathbf{y}_d \end{bmatrix} = \begin{bmatrix} \mathbf{f}_s \\ \mathbf{f}_d \\ \mathbf{g}_s \\ \mathbf{g}_d \end{bmatrix}. \quad (1.80)$$

We group these block matrix to form a concise version,

$$\mathbf{A} = \begin{bmatrix} \mathbf{A}_s & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_d \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} -\mathbf{B}_s & \mathbf{0} \\ \mathbf{0} & -\mathbf{B}_d \end{bmatrix}, \quad \mathbf{C} = \begin{bmatrix} -\mathbf{C}_s & -\mathbf{K} \\ -\mathbf{K} & -\mathbf{C}_d \end{bmatrix}, \quad \mathbf{F} = \begin{bmatrix} \mathbf{f}_s \\ \mathbf{f}_d \end{bmatrix}, \quad \mathbf{G} = \begin{bmatrix} -\mathbf{g}_s \\ -\mathbf{g}_d \end{bmatrix}, \quad (1.81)$$

$$\begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{B}^T & \mathbf{C} \end{bmatrix} \begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix} = \begin{bmatrix} \mathbf{F} \\ \mathbf{G} \end{bmatrix}, \quad (1.82)$$

where $\mathbf{A}$ is a positive definite matrix. However, $\mathbf{B}$ is usually a singular matrix when no melting region presents. We will present a modified Uzawa iteration algorithm to deal with this coupled degenerate singular system. Uzawa algorithm applied to the case of discretizing Stokes problem with Mixed Finite Element has been discussed by Elman and Golub (1994) and Bramble et al. (1997),

Algorithm 1 Coupled Uzawa Iteration

- 1: Initialize $\mathbf{x}_0 = \mathbf{0}$ and $\mathbf{y}_0 = \mathbf{0}$
 - 2: Update $\mathbf{x}_{i+1} = \mathbf{x}_i + \mathbf{A}^{-1}(\mathbf{F} - (\mathbf{A}\mathbf{x}_i + \mathbf{B}\mathbf{y}_i))$
 - 3: Obtain $\mathbf{r} = \mathbf{B}^T\mathbf{x}_{i+1} + \mathbf{C}\mathbf{y}_i - \mathbf{G}$
 - 4: Obtain $\tilde{\mathbf{y}} = \operatorname{argmin}_{\mathbf{y} \in V} \|(\mathbf{B}^T\mathbf{A}^{-1}\mathbf{B} - \mathbf{C})^{-1}\tilde{\mathbf{y}} - \mathbf{r}\|$
 - 5: Update $\mathbf{y}_{i+1} = \mathbf{y}_i + \tilde{\mathbf{y}}$
-

Works Cited

T. Arbogast, M. A. Hesse, and A. L. Taicher. Mixed methods for two-phase Darcy-Stokes mixtures of partially melted materials with regions of zero porosity. *SIAM J. Sci. Comput.*, 39(2):B375–B402, 2017. DOI 10.1137/16M1091095.

Todd Arbogast and Maicon R. Correa. Two families of $H(\text{div})$ mixed finite elements on quadrilaterals of minimal dimension. *SIAM Journal on Numerical Analysis*, 54(6):3332–3356, 2016. doi: 10.1137/15M1013705. URL <https://doi.org/10.1137/15M1013705>.

Todd Arbogast and Mary F. Wheeler. A family of rectangular mixed elements with a continuous flux for second order elliptic problems. *SIAM Journal on Numerical Analysis*, 42(5):1914–1931, 2005. doi: 10.1137/S0036142903435247. URL <https://doi.org/10.1137/S0036142903435247>.

Todd Arbogast, Zhen Tao, and Chuning Wang. Direct serendipity and mixed finite elements on convex quadrilaterals. *Numerische Mathematik*, 150:929–974, 2022.

James H. Bramble, Joseph E. Pasciak, Apostol T, and Vassilev. Analysis of the inexact uzawa algorithm for saddle point problems. *SIAM. J. Numer. Anal.*, 34:1072–1092, 1997.

Franco Brezzi, Jim Douglas Jr., and L. D. Marini. Two families of mixed finite element for second order elliptic problems. *Numerische Mathematik*, 47:217–235, 1985.

Genevieve Raugel Christine Bernardi. Analysis of some finite elements for the stokes problem. *Math. of Comput.*, 44:71–79, 1985.

Howard C. Elman and Gene H. Golub. Inexact and preconditioned uzawa algorithms for saddle point problems. *SIAM. J. Numer. Anal.*, 31:1645–1661, 1994.

Michel Fortin. Old and new finite elements for incompressible flows. *International Journal for Numerical Methods in Fluids*, 1:347–364, 1981. URL <https://api.semanticscholar.org/CorpusID:122770101>.

P. A. Raviart and J. M. Thomas. A mixed finite element method for 2-nd order elliptic problems. In Ilio Galligani and Enrico Magenes, editors, *Mathematical Aspects of Finite Element Methods*, pages 292–315, Berlin, Heidelberg, 1977. Springer Berlin Heidelberg. ISBN 978-3-540-37158-8.