

Unsupervised Learning : PCA

Swati Mishra

Applications of Machine Learning (4AL3)

Fall 2024




ENGINEERING

Review

- Hierarchical Clustering
- Dendrograms
- Linkage Techniques
- Introduction to PCA, Projections


Projections



How do we project a sample in this space?




Dimensionality Reduction



Let Z_i be a linear combination of p

$$Z_i = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

Dimensionality Reduction




Let Z_1 be a linear combination of p

$$Z_i = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

This can be written as:

$$Z_m = \sum_{j=1}^p \phi_{jm}X_j$$

Dimensionality Reduction



Let Z_1 be a linear combination of p

$$Z_i = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$


This can be written as:

$$Z_m = \sum_{j=1}^p \phi_{jm}X_j$$

This can **also** be written as:

$$y_i = \theta_0 + \sum_{m=1}^M \theta_m Z_m + \epsilon_i$$

Dimensionality Reduction



Let Z_1 be a linear combination of p

$$Z_i = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

This can be written as:

$$Z_m = \sum_{j=1}^p \phi_{jm}X_j$$

This can **also** be written as:


$$y_i = \theta_0 + \sum_{m=1}^M \theta_m Z_m + \epsilon_i$$

The above equation can be solved using ordinary least squares

Principal Component Analysis

- Suppose that we wish to visualize n observations with measurements on a set of p features.
- Not all of p dimensions are equally interesting.
- PCA finds a low-dimensional representation of a data set that contains as much as possible of the variation.
- PCA seeks a small number of dimensions that are as interesting as possible.

Scatterplot Matrix




Principal Component Analysis

- First Principal Component:
 - Normalized linear combination of the features that has the largest variance.

$$Z = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

- ϕ_{11} = loadings of the PCA



Principal Component Analysis

- First Principal Component:
 - Normalized linear combination of the features that has the largest variance.


$$Z = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

- ϕ_{11} = loadings of the PCA

$$Z_m = \sum_{j=1}^p \phi_{jm}X_j$$

This can be written as: $y_i = \theta_0 + \sum_{m=1}^M \theta_m z_m + \epsilon_i$

- The above equation can be solved using ordinary least squares



Principal Component Analysis

- First Principal Component:
 - Normalized linear combination of the features that has the largest variance.


$$Z = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

- ϕ_{11} = loadings of the PCA

$$Z_m = \sum_{j=1}^p \phi_{jm}X_j$$

This can be written as: $y_i = \theta_0 + \sum_{m=1}^M \theta_m z_m + \epsilon_i$

- The above equation can be solved using ordinary least squares




Principal Component Analysis

- First Principal Component:
 - Normalized linear combination of the features that has the largest variance.

$$Z = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

- ϕ_{11} = loadings of the PCA

Let's assume: $\phi_{11} = 0.839$, $\phi_{21} = 0.544$




Principal Component Analysis

- First Principal Component:
 - Normalized linear combination of the features that has the largest variance.

$$Z = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

- ϕ_{11} = loadings of the PCA

Let's assume: $\phi_{11} = 0.839$, $\phi_{21} = 0.544$



$$Z_1 = 0.839 \times (pop - mean(pop)) + 0.544 \times (ad - mean(ad))$$


Principal Component Analysis

- First Principal Component:
 - Normalized linear combination of the features that has the largest variance.

$$Z = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

- ϕ_{11} = loadings of the PCA

Let's assume: $\phi_{11} = 0.839$, $\phi_{21} = 0.544$



$$Z_1 = 0.839 \times (pop - mean(pop)) + 0.544 \times (ad - mean(ad))$$

Principal Component Analysis

- First Principal Component:
 - Normalized linear combination of the features that has the largest variance.


$$Z = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

- ϕ_{11} = loadings of the PCA

Let's assume: $\phi_{11} = 0.839$, $\phi_{21} = 0.544$

$$\text{Then, } \phi_{11} \times \phi_{11} + \phi_{21} \times \phi_{21} = 1$$

$$Z_1 = 0.839 \times (\text{pop} - \text{mean}(\text{pop})) + 0.544 \times (\text{ad} - \text{mean}(\text{ad}))$$



Principal Component Analysis

- First Principal Component:
 - Normalized linear combination of the features that has the largest variance.


$$Z = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

- ϕ_{11} = loadings of the PCA

Let's assume: $\phi_{11} = 0.839$, $\phi_{21} = 0.544$

$$\text{Then, } \phi_{11} \times \phi_{11} + \phi_{21} \times \phi_{21} = 1$$

$$z_i = 0.839 \times (\text{pop}_i - \text{mean}(\text{pop})) + 0.544 \times (\text{adi} - \text{mean}(\text{ad}))$$



Principal Component Analysis

- First Principal Component:
 - Normalized linear combination of the features that has the largest variance.

$$Z = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$


- ϕ_{11} = loadings of the PCA

Let's assume: $\phi_{11} = 0.839$, $\phi_{21} = 0.544$

$$\text{Then, } \phi_{11} \times \phi_{11} + \phi_{21} \times \phi_{21} = 1$$

Principal component scores

$$z_i = 0.839 \times (\text{pop}_i - \text{mean}(\text{pop})) + 0.544 \times (\text{adi} - \text{mean}(\text{ad}))$$



Principal Component Analysis

- First Principal Component:
 - Normalized linear combination of the features that has the largest variance.

$$Z = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

- ϕ_{11} = loadings of the PCA

How do we compute principal components?



Principal Component Analysis

- First Principal Component:
 - Normalized linear combination of the features that has the largest variance.

$$Z = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

- ϕ_{11} = loadings of the PCA
- Normalized linear combination of the features $\sum_{j=1}^p \phi_{j1}^2 = 1$

Principal Component Analysis

- First Principal Component:
 - Normalized linear combination of the features that has the largest variance.

$$Z = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$

- ϕ_{11} = loadings of the PCA
- Normalized linear combination of the features $\sum_{j=1}^p \phi_{j1}^2 = 1$
- Solve the optimization problem:

$$\underset{\phi_{11}, \dots, \phi_{p1}}{\text{maximize}} \left\{ \frac{1}{n} \sum_{i=1}^n \left(\sum_{j=1}^p \phi_{j1} x_{ij} \right)^2 \right\} \text{ subject to } \sum_{j=1}^p \phi_{j1}^2 = 1.$$

Principal Component Analysis



- First Principal Component:
 - Normalized linear combination of the features that has the largest variance.

$$Z = \phi_{11}X_1 + \phi_{21}X_2 + \dots + \phi_{p1}X_p$$



- ϕ_{11} = loadings of the PCA
- Normalized linear combination of the features $\sum_{j=1}^p \phi_{j1}^2 = 1$
- Solve the optimization problem:

$$\underset{\phi_{11}, \dots, \phi_{p1}}{\text{maximize}} \left\{ \frac{1}{n} \sum_{i=1}^n \underbrace{\left(\sum_{j=1}^p \phi_{j1} x_{ij} \right)^2}_{Z_{i1}^2} \right\} \text{ subject to } \sum_{j=1}^p \phi_{j1}^2 = 1.$$

Principal Component Analysis



Principal Component Analysis




Principal Component Analysis

- First Principal Component (Z_1):
 - Normalized linear combination of the features that has the largest variance.
- **Second Principal Component:**
 - Normalized linear combination of the features that has the largest variance out of all linear combinations that are **uncorrelated** with Z_1 .

$$Z = \phi_{12}X_1 + \phi_{22}X_2 + \dots + \phi_{p2}X_p$$

- ϕ_{i2} = loadings of the PCA




Principal Component Analysis

- First Principal Component (Z_1):
 - Normalized linear combination of the features that has the largest variance.
- Second Principal Component (Z_2)
 - Normalized linear combination of the features that has the largest variance out of all linear combinations that are uncorrelated with Z_1 .

$$Z = \phi_{12}X_1 + \phi_{22}X_2 + \dots + \phi_{p2}X_p$$

- ϕ_{i2} = loadings of the PCA
- Constraining Z_2 to be uncorrelated with Z_1 is equivalent to constraining the direction ϕ_1 to be orthogonal to the direction of ϕ_2



Principal Component Analysis


- First Principal Component (Z_1):
 - Normalized linear combination of the features that has the largest variance.
- Second Principal Component:
 - Normalized linear combination of the features that has the largest variance out of all linear combinations that are uncorrelated with Z_1 .

$$Z = \phi_{12}X_1 + \phi_{22}X_2 + \dots + \phi_{p2}X_p$$

- ϕ_{i2} = loadings of the PCA

Second principal component scores

$$z_i = 0.544 \times (\text{pop}_i - \text{mean}(\text{pop})) - 0.839 \times (\text{adi} - \text{mean}(\text{ad}))$$



Principal Component Analysis

- First Principal Component (Z_1):
 - Normalized linear combination of the features that has the largest variance.
- Second Principal Component:
 - Normalized linear combination of the features that has the largest variance out of all linear combinations that are uncorrelated with Z_1 .



$$Z = \phi_{12}X_1 + \phi_{22}X_2 + \dots + \phi_{p2}X_p$$

- ϕ_{i2} = loadings of the PCA
- Constraining Z_2 to be uncorrelated with Z_1 is equivalent to constraining the direction ϕ_1 to be orthogonal to the direction of ϕ_2

$$\underset{\phi_{11}, \dots, \phi_{p1}}{\text{maximize}} \left\{ \frac{1}{n} \sum_{i=1}^n \left(\sum_{j=1}^p \phi_{j1} x_{ij} \right)^2 \right\} \text{ subject to } \sum_{j=1}^p \phi_{j1}^2 = 1.$$



Replace with ϕ_2

Principal Component Analysis





Principal Component Analysis

- For multi-dimensional data, there are multiple principal components.
- Principal components can be used to produce low-dimensional views.



Principal Component Analysis

- For multi-dimensional data, there are multiple principal components.
- Principal components can be used to produce low-dimensional views.
 - It is not feature selection. We consider linear combination of all features



Principal Component Analysis

- For multi-dimensional data, there are multiple principal components.
- Principal components can be used to produce low-dimensional views.
 - It is not feature selection. We consider linear combination of all features
- Eigen decomposition method is used to solve PCA.
 - Principal Component directions are computed using the order sequence of eigenvectors of the matrix $\mathbf{X}^T \mathbf{X}$

$$\mathbf{X}^T \mathbf{X} = \left(\mathbf{U} \mathbf{\Sigma} \mathbf{W}^T \right)^T \mathbf{U} \mathbf{\Sigma} \mathbf{W}^T = \mathbf{W} \mathbf{\Sigma}^2 \mathbf{W}^T$$

Principal Component Analysis

- For multi-dimensional data, there are multiple principal components.
- Principal components can be used to produce low-dimensional views.
 - It is not feature selection. We consider linear combination of all features
- Eigen decomposition method is used to solve PCA.
 - Principal Component directions are computed using the order sequence of eigenvectors of the matrix $\mathbf{X}^T \mathbf{X}$
- For a set of p features and n observations, there are at most $\min(n - 1, p)$ principal components. We use the smallest subset, determined by **proportion of variance**.

Principal Component Analysis

- For multi-dimensional data, there are multiple principal components.
- Principal components can be used to produce low-dimensional views.
 - It is not feature selection. We consider linear combination of all features
- Eigen decomposition method is used to solve PCA.
 - Principal Component directions are computed using the order sequence of eigenvectors of the matrix $\mathbf{X}^T \mathbf{X}$
- For a set of p features and n observations, there are at most $\min(n - 1, p)$ principal components. We use the smallest subset, determined by **proportion of variance**.
- Before PCA is performed, the variables should be centered to have mean zero.
 - Scaling heavily impacts PCA result

Principal Component Analysis

- For multi-dimensional data, there are multiple principal components.
- Principal components can be used to produce low-dimensional views.
 - It is not feature selection. We consider linear combination of all features
- Eigen decomposition method is used to solve PCA.
 - Principal Component directions are computed using the order sequence of eigenvectors of the matrix $\mathbf{X}^T \mathbf{X}$
- For a set of p features and n observations, there are at most $\min(n - 1, p)$ principal components. We use the smallest subset, determined by **proportion of variance**.
- Before PCA is performed, the variables should be centered to have mean zero.
 - Scaling heavily impacts PCA result

Popularly used in Recommender systems!



Readings

Required Readings:

Introduction to Statistical Learning

- Chapter 6 – Section 6.3 page 253 – 259
- Chapter 12 – Section 12.2 page 504 – 515

Supplemental Readings (Not required but recommended):

Deep Learning

- Chapter 5 – Section 5.8 page 147 – 150

Thank You
