Question 1 (1 point) Saved
Leo Kanner was the first music therapist to publish research about the benefits of music therapy for supporting communication and social interactions with people diagnosed with ASD.
☐ True
False
Question 2 (1 point) Saved
The development and stages of PD are similar for all individuals.
True
False
Question 3 (1 point) ✓ Saved
The 'refrigerator mother' is a theory still being explored as playing a role in the development of ASD.
True
False

Question 4 (1 point) Saved	
What was/were the outcome(s) of the study "Music Therapy in Parkinson's Dise An Integrative Method for Motor and Emotional Rehabilitation"?	ease:
Music therapy demonstrated no improvement in HM	
Music therapy demonstrated no improvement to the UPDRS scores	
Music therapy demonstrated significant improvement in UPDRS scores	
Music therapy demonstrated significant improvement in speech scores	
Question 5 (1 point) ✓ Saved	
Which of the following best describes Parkinson's Disease?	
A muscle disorder that impacts movement	
A genetic disease	
A neurological disorder that effects cognition	
A neurological disorder effecting the dopaminergic system	
Question 6 (1 point) Saved	
Autism presents itself the same in all people diagnosed with Autism.	
True	
False	

Question 7 (1 point)
The average age for the onset of PD is 60yrs old, however, someone can be diagnosed as early as their 30's.
True
○ False
Question 8 (1 point)
In a music therapy session, a client diagnosed with ASD would typically prefer the use of pre-recorded music over live music.
☐ True
False
Question 9 (1 point) Saved
Structure in a music therapy session is not important when working with an individual with Autism. In fact, predicability in a session can often evoke anxiety for an individual of the autism spectrum.
○ True
False

Question 10 (1 point	t) ✓ Saved
Which of the follow Disease?	ving best describes the reading assigned about Parkinson's
A pilot study	
A systematic re	view
A randomized of	control trial
A case study	
Question 11 (1 point	t) ✓ Saved
Which goal(s) is/are	e typical to working with clients diagnosed with ASD?
improving com	munication
assisting with s	ocial interactions
helping manage	e behavioural issues
all of the above	
Question 12 (1 point	t)
implementation of	sic Therapy in Parkinson's Disease" the results do not suggest the music therapy as part of symptom management for PD, as there is ce to support its efficacy.
○ True	
False	A &

Question 13 (1 point) ✓ Saved
Which of the following is a motor symptom of Parkinson's disease?
resting tremors
ognitive decline
depression
Oloss of smell
Question 14 (1 point) ✓ Saved
Understanding the best approach to supporting individuals with ASD has been complicated by the fact that it was not differentiated from schizophrenia in the DSM until approximately when?
the 1980s
the 2000s
the 1960s
O the 1990s

Que	stion 15 (1 point) Saved
Au	itism is best described as
	an acquired brain injury
	developmental delay specific to children
	a result of poor parenting
	a complex developmental disability
_	
Que	stion 16 (1 point) Saved
thi	child diagnosed with ASD is referred to music therapy to support self-expression rough music. Which approach/model would a music therapist most likely draw on?
C	Cognitive Behavioural Approach
•	Nordoff-Robbins
C	A.O.M
0) GIM

"Music Therapy in Parkinson's Disease: An Integrative Method for Motor and Emotional Rehabilitation" had measurable results supporting music therapy.
True
○ False
Question 18 (1 point) Saved
Autism is an example of an acquired brain injury.
☐ True
False
Question 19 (1 point) ✓ Saved
In the systematic review of music therapy and parkinson's disease, research was NO predominately focused on
Motor
Speech
Cognition
 Social communication
Question 20 (1 point) Saved
If an individual is diagnosed with Parkinson's disease, their symptoms will remain relatively the same over time.
○ True
False

Question 17 (1 point)

Saved