



© International Baccalaureate Organization 2024

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organisation du Baccalauréat International 2024

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organización del Bachillerato Internacional, 2024

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

Mathematics: analysis and approaches

Higher level

Paper 3

6 May 2024

Zone A afternoon | **Zone B** afternoon | **Zone C** afternoon

1 hour

Instructions to candidates

- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Answer all the questions in the answer booklet provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **mathematics: analysis and approaches HL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[55 marks]**.

Answer **all** questions in the answer booklet provided. Please start each question on a new page. Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. Solutions found from a graphic display calculator should be supported by suitable working. For example, if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

1. [Maximum mark: 24]

If two functions $f(x)$ and $g(x)$ are differentiable, then their product is differentiable and the two functions satisfy the product rule: $(f(x)g(x))' = f'(x)g(x) + g(x)f'(x)$.

In this question, you will meet examples of pairs of differentiable functions, $f(x)$ and $g(x)$, that also satisfy $(f(x)g(x))' = f'(x)g'(x)$.

In part (a), consider $f(x) = \frac{1}{(2-x)^2}$, where $x \in \mathbb{R}$, $x \neq 2$, and $g(x) = x^2$, where $x \in \mathbb{R}$.

(a) (i) Find an expression for $f'(x)$. [2]

(ii) Show that $f'(x)g'(x) = \frac{4x}{(2-x)^3}$. [2]

(iii) Show that $f(x)g'(x) + g(x)f'(x) = \frac{4x}{(2-x)^3}$. [4]

In parts (b) and (c), consider two non-constant functions, $f(x)$ and $g(x)$, where $f(x) > 0$ and $g(x) \neq g'(x)$.

(b) By rearranging the equation $f(x)g'(x) + g(x)f'(x) = f'(x)g'(x)$, show that $\frac{f'(x)}{f(x)} = \frac{g'(x)}{g'(x) - g(x)}$. [2]

(This question continues on the following page)

(Question 1 continued)

- (c) Hence, by integrating both sides of $\frac{f'(x)}{f(x)} = \frac{g'(x)}{g'(x) - g(x)}$, show that $f(x) = Ae^{\left(\int \frac{g'(x)}{g'(x) - g(x)} dx\right)}$, where A is an arbitrary positive constant. [2]

The result from part (c) can be used to find pairs of functions, $f(x)$ and $g(x)$, which satisfy **both** of the following:

$$(f(x)g(x))' = f(x)g'(x) + g(x)f'(x) \text{ and } (f(x)g(x))' = f'(x)g'(x).$$

In parts (d) and (e), use the result in part (c) with $A = 1$.

- (d) Consider $g(x) = xe^x$.

Find $f(x)$ such that $f(x)$ and $g(x)$ satisfy the above two equations. [5]

- (e) Consider $g(x) = \sin x + \cos x$.

Find $f(x)$ such that $f(x)$ and $g(x)$ satisfy the above two equations over the domain $0 < x < \pi$.

Give your answer in the form $f(x) = \sqrt{e^x h(x)}$, where $h(x)$ is a function to be determined. [7]

2. [Maximum mark: 31]

This question asks you to find the probability of graphs of randomly generated quadratic functions having a specified number of x -intercepts.

In parts (a) – (f), consider quadratic functions, $f(x) = ax^2 + bx + c$, whose coefficients, a , b and c , are randomly generated in turn by rolling an unbiased six-sided die three times and reading off the value shown on the uppermost face of the die.

For example, rolling a 2, 3 and 5 in turn generates the quadratic function $f(x) = 2x^2 + 3x + 5$.

- (a) Explain why there are 216 possible quadratic functions that can be generated using this method. [1]

- (b) The set of coefficients, $a = 1$, $b = 4$ and $c = 4$, is randomly generated to form the quadratic function $f(x) = x^2 + 4x + 4$.

Verify that this graph of f has only one x -intercept. [2]

- (c) By considering the discriminant, or otherwise, show that the probability of the graph of such a randomly generated quadratic function having only one x -intercept is $\frac{5}{216}$. [6]

Now consider randomly generated quadratic functions whose corresponding graphs have two **distinct** x -intercepts.

- (d) By considering the discriminant, determine the set of possible values of ac . [3]

- (e) (i) For the case where $ac = 1$, show that there are four quadratic functions whose corresponding graphs have two distinct x -intercepts. [1]

- (ii) For the case where $ac = 2$, show that there are eight quadratic functions whose corresponding graphs have two distinct x -intercepts. [2]

Let p be the probability of the graph of such a randomly generated quadratic function having two distinct x -intercepts.

- (f) Using the approach started in part (e), or otherwise, find the value of p . [6]

(This question continues on the following page)

(Question 2 continued)

In parts (g) and (h), consider a randomly generated quadratic function, $f(x) = x^2 + 2Zx + 1$, where the continuous random variable $Z \sim N(0, 1)$.

- (g) Find the probability that the graph of f has two x -intercepts. [3]

The continuous random variables, X_1 and X_2 , represent the x -intercepts of the graph of f where $X_1 = -Z - \sqrt{Z^2 - 1}$ and $X_2 = -Z + \sqrt{Z^2 - 1}$.

- (h) Given that the graph of f has two x -intercepts, X_1 and X_2 , find the probability that both X_1 and X_2 are greater than 0.5. [7]
-