

# HW II

EE 546: Mathematics of High-dimensional Data

University of Southern California

Assigned on: October 5, 2020

Due date: by 7PM October 18, 2020

---

The purpose of this homework is two fold:

- (1) Give you some practice with concentration inequalities.
- (2) Help you refresh some results on operator norms.

1- **Concentration.** Let  $X_1, X_2, \dots, X_m$  be i.i.d. Gaussian random variables with  $\mathcal{N}(0, 1)$  distribution. Also let  $\mathbf{a}_r$  be i.i.d. random Gaussian vectors distributed as  $\mathcal{N}(\mathbf{0}, \mathbf{I})$  and independent of  $X_1, X_2, \dots, X_m$ . Let  $\mathbf{y}$  be a fixed vector. We are interested in understanding the random variable

$$Z = \frac{1}{m} \sum_{r=1}^m X_r |X_r| \mathbf{sign}(X_r + \mathbf{a}_r^T \mathbf{y})$$

- (a) What is the expected value of  $Z$ ?

**Hint:** You are allowed to use the following identity without proof. For  $X$  and  $Y$  independent  $\mathcal{N}(0, 1)$  random variables we have

$$\mathbb{E}[X^2 \text{sgn}(X) \text{sgn}(\alpha X + \beta Y)] = \frac{2}{\pi} \tan^{-1} \left( \frac{\alpha}{\beta} \right) + \frac{2}{\pi} \frac{\alpha \beta}{\alpha^2 + \beta^2}.$$

- (b) How well is the random variable  $Z$  concentrated? That is, can you bound  $\mathbb{P}\{|Z - \mathbb{E}[Z]| \geq t\}$ ?

**Hint:** What kind of a random variable is  $X_r |X_r| \mathbf{sign}(X_r + \mathbf{a}_r^T \mathbf{y})$ ?

2- **Operator norms.** Remember that the  $\ell_2$  operator norm of  $\mathbf{A} \in \mathbb{R}^{m \times n}$  is defined as

$$\|\mathbf{A}\| := \max \{ \|\mathbf{A}\mathbf{x}\|_{\ell_2} : \|\mathbf{x}\|_{\ell_2} = 1 \}.$$

- (i) Prove that the operator norm can alternatively be written in the following two forms

$$\begin{aligned} \|\mathbf{A}\| &= \max \{ \langle \mathbf{x}, \mathbf{A}\mathbf{y} \rangle : \mathbf{x} \in \mathbb{R}^m, \mathbf{y} \in \mathbb{R}^n, \text{ and } \|\mathbf{x}\|_{\ell_2} = \|\mathbf{y}\|_{\ell_2} = 1 \}, \\ &= \sigma_1(\mathbf{A}), \end{aligned}$$

where  $\sigma_1(\mathbf{A}) \geq \sigma_2(\mathbf{A}) \geq \dots \geq \sigma_n(\mathbf{A})$  are the singular values of  $\mathbf{A}$ .

- (ii) Prove that

$$\sum_{s=1}^r \sigma_s(\mathbf{A}) = \max \{ \text{trace}(\mathbf{U}^T \mathbf{A} \mathbf{V}) : \mathbf{U} \in \mathbb{R}^{m \times r}, \mathbf{V} \in \mathbb{R}^{n \times r}, \text{ and } \mathbf{U}^T \mathbf{U} = \mathbf{V}^T \mathbf{V} = \mathbf{I}_r \}.$$

where  $\sigma_1(\mathbf{A}) \geq \sigma_2(\mathbf{A}) \geq \dots \geq \sigma_n(\mathbf{A})$  are the singular values of  $\mathbf{A}$  and  $\mathbf{I}_r$  is the  $r \times r$  identity matrix.

3- Prove the following upper and lower bounds

$$\|\mathbf{A}\| \leq \sqrt{m} \max_{i \in \{1, 2, \dots, m\}} \left( \sum_{j=1}^n A_{ij}^2 \right)^{1/2},$$

$$\|\mathbf{A}\| \geq \frac{1}{\sqrt{mn}} \sum_{i=1}^m \left| \sum_{j=1}^n A_{ij} \right|.$$

Give examples of matrices such that these bounds are satisfied with equality.

4- Prove that

$$\|\mathbf{A}\|_F = \left( \sum_{i=1}^{\min(m,n)} \sigma_i^2(\mathbf{A}) \right)^{1/2},$$

where  $\|\mathbf{A}\|_F = \sqrt{\sum_{i=1}^m \sum_{j=1}^n A_{ij}^2}$  is the Frobenius norm of the matrix  $\mathbf{A}$ . Deduce that

$$\|\mathbf{A}\| \leq \|\mathbf{A}\|_F \leq \sqrt{\text{rank}(\mathbf{A})} \|\mathbf{A}\|.$$