Braided-Symmetric Derivations of Algebras

Abderrahim Adrabi[a,b]

[a]Faculty of Sciences, Mohammed V University in Rabat, Morocco;
[b]Faculty of Educational Sciences, Mohammed V University in Rabat, Morocco

ARTICLE HISTORY
Preprint version 54, compiled 2026/10/28

ABSTRACT
We study braided-symmetric derivations of an algebra over a field. These are maps that satisfy the Leibniz rule for the product symmetrized by an involution of $A \otimes A$. The choice of involution decides which derivations appear. The identity involution gives the ordinary derivations when 2 is invertible. The swap gives the Jordan derivations. We then twist the swap by an involutive automorphism. Let the algebra be commutative and reduced, with 2 invertible. Then the twisted maps that commute with the automorphism are exactly the derivations that commute with it. An example with nilpotent elements shows that the reduced hypothesis is needed, and an example in characteristic 2 shows that the invertibility of 2 is needed.

KEYWORDS
Braided-symmetric derivations; twisted Jordan derivations; Jordan derivations; Leibniz defect

AMS CLASSIFICATION
MSC 2020: 16W25; 16E40; 13N15

1. Introduction

Throughout, K is a field and an algebra means a unital associative K-algebra. We take all maps between algebras to be K-linear. We say so when a result cited below holds more generally for additive maps on rings.

A *derivation* of an algebra A is a linear map $D\colon A \to A$ with $D(xy) = D(x)y + xD(y)$ for all $x, y \in A$. The notion goes back to Jacobson [9, Part I], who studied the derivations of an arbitrary nonassociative algebra. Hochschild [7, §§2 and 4] defined the cocycles of an associative algebra, and its 1-cocycles are the derivations. A *Jordan derivation* of A is a linear map with $D(x \circ y) = D(x) \circ y + x \circ D(y)$ for all $x, y \in A$, where $x \circ y = xy + yx$. Herstein [5, p. 1104] introduced this notion. His result needs three notions for a ring R. We call R *prime* if $aRb = 0$ implies $a = 0$ or $b = 0$ [2, p. 1003]. We call R *semiprime* if $aRa = 0$ implies $a = 0$ [2, p. 1003]. We call R 2-*torsion free* if $2a = 0$ implies $a = 0$ [2, p. 1003]. So every prime ring is semiprime. Herstein proved that every Jordan derivation of a prime ring of characteristic different from 2 is a derivation [5, Theorem 3.1]. Every derivation is a Jordan derivation, and the converse is false in general [2, p. 1003]. Cusack [4, Corollary 5] extended this result of Herstein to 2-torsion free semiprime rings, and Brešar [2, Theorem 1] gave an independent proof.

CONTACT A. Adrabi. Email: abderrahim.adrabi@um5r.ac.ma

Benkovič [1, §1] treats derivations, Jordan derivations and Lie derivations as f-derivations, for $f = x_1x_2$, $x_1 \circ x_2$ and $[x_1, x_2]$, so that one identity for a multilinear polynomial f covers all three. We obtain them from an identity of a different kind. Our identity keeps the polynomial x_1x_2, and an involution of $A \otimes A$ acts before we multiply. Here the involution may use extra structure of A, such as an automorphism. The identity is then no longer an f-derivation identity; Section 2 explains why.

Let A be an algebra over K with product μ. Let $\sigma\colon A \otimes A \to A \otimes A$ be a K-linear map with $\sigma^2 = \mathrm{id}$, for instance the swap $\tau(x \otimes y) = y \otimes x$. It gives a symmetrized product

$$x \cdot_\sigma y = \mu(x \otimes y) + \mu(\sigma(x \otimes y)).$$

For $\sigma = \tau$ this is the Jordan product $x \circ y$. Put $\mu_\sigma = \mu + \mu \circ \sigma$. Since $\sigma^2 = \mathrm{id}$, this gives

$$\mu_\sigma \circ \sigma = \mu_\sigma.$$

Example 3.3 shows that a twisted version of this product can fail to be associative and to have a unit.

We call a linear map $D\colon A \to A$ a *braided-symmetric derivation* for σ if

$$D(x \cdot_\sigma y) = D(x) \cdot_\sigma y + x \cdot_\sigma D(y). \tag{1}$$

So (1) is the Leibniz rule for the symmetrized product. Other authors have studied *braided derivations*, which use a braiding operator directly [11, §2] [10, §4.1, Definition 14] [12, §3, equation (3.9)]. Remark 3.2 compares the two identities. Our identity uses only μ, σ and D.

We now describe the main results. Our first step is a dictionary. Remark 3.4 shows that the identity gives exactly the derivations when 2 is invertible, and the swap gives exactly the Jordan derivations.

Our second step twists the swap by an involutive automorphism or anti-automorphism α of A. Proposition 4.2 describes the twisted class through the defect $E_D(x, y) = D(xy) - D(x)y - xD(y)$. Corollary 4.5 shows that, on 2-torsion-free semiprime rings, the Jordan maps in the twisted class are the derivations commuting with α.

Theorem 5.1 treats commutative reduced algebras and needs no Jordan hypothesis. Example 5.2 shows that the reduced hypothesis is needed, and Example 5.3 shows the same for the invertibility of 2. Proposition 5.4 shows that the twisted class and the α-derivations on one algebra do not contain each other.

We follow this order. Section 2 recalls the classical notions and contains no new results. Section 3 defines braided-symmetric derivations and treats the identity and the swap. Section 4 treats the twisted swap through the Leibniz defect. Section 5 treats reduced commutative algebras and compares the twisted class with α-derivations. We place each example right after the result it illustrates.

2. Preliminaries

This section recalls the classical notions we need, and none of it is new. We treat five kinds of derivation and then triangular algebras.

From now on, A is a K-algebra with unit 1. We write $[x, y] = xy - yx$ for the commutator, $x \circ y = xy + yx$ for the Jordan product, and $Z(A)$ for the centre. The centre is the set of elements z with $zx = xz$ for all $x \in A$. We call an element x *nilpotent* if $x^n = 0$ for some $n \geq 1$. A commutative algebra is *reduced* if its only nilpotent element is zero [3, 1.4.12].

A derivation of an algebra is as defined by Jacobson [9, Part I]: a linear map D with $D(xy) = D(x)y + xD(y)$. Every derivation satisfies $D(1) = 0$, since $D(1) = D(1 \cdot 1) = 2D(1)$. For $g \in A$, the map $\mathrm{ad}_g(x) = [g, x]$ is an *inner derivation*, called a principal derivation by Weibel [13, §9.2]. We write $\mathrm{Der}(A)$ and $\mathrm{Inn}(A)$ for these two spaces.

We add a word on the characteristic for Jordan derivations. A *Jordan derivation* in the sense of Herstein [5, p. 1104] is an additive map with $D(x^2) = D(x)x + xD(x)$. We write $\mathrm{JDer}(A)$ for the space of linear maps with $D(x \circ y) = D(x) \circ y + x \circ D(y)$ for all x, y. Replacing x by $x + y$ gives the condition $D(x \circ y) = D(x) \circ y + x \circ D(y)$ in every characteristic. Setting $y = x$ in this condition gives $2D(x^2) = 2(D(x)x + xD(x))$. So the two conditions agree when 2 is invertible in K, and we use the $\circ$ form throughout. When $\mathrm{char}\, K = 2$ and A is commutative, $x \circ y = 0$, and every linear map satisfies the $\circ$ condition.

For an automorphism α of A, an *α-derivation* satisfies $D(xy) = D(x)\alpha(y) + \alpha(x)D(y)$ [8, §1]. We write $\mathrm{Der}_\alpha(A)$ for this space.

We now compare the Jordan derivations with the derivations. Every derivation is a Jordan derivation [2, p. 1003], so $\mathrm{Der}(A) \subseteq \mathrm{JDer}(A)$.

We recall a polynomial way to unify these maps. A *multilinear polynomial* in variables $x_1, \ldots, x_n$ that do not commute has the following form [1, §1, (1.1)].

$$f = \sum_{\varpi} \kappa_\varpi \, x_{\varpi(1)} \cdots x_{\varpi(n)}.$$

Here ϖ runs over the permutations of $\{1, \ldots, n\}$, and the coefficients κ_ϖ lie in K [1, §1, (1.1)].

Fix a nonzero multilinear polynomial f in n variables. An *f-derivation* satisfies the identity [1, §1, (1.2)]

$$D(f(x_1, \ldots, x_n)) = \sum_{i=1}^{n} f(x_1, \ldots, D(x_i), \ldots, x_n)$$

for all $x_1, \ldots, x_n$ [1, §1, (1.2)]. Derivations, Jordan derivations and Lie derivations are the f-derivations for $f = x_1x_2$, $x_1 \circ x_2$ and $[x_1, x_2]$ [1, §1]. In each case f uses only sums, products and scalars, so it makes sense in every algebra.

The twisted products of Section 4 are different. They have the form

$$xy + \alpha(y)\alpha(x)$$

for a fixed automorphism or anti-automorphism α of A. Here the expression applies α to the variables, and α is not a polynomial operation. So it is not of the form $f(x, y)$ for any polynomial f. Hence the results on f-derivations do not apply directly.

We use three involutions of $A \otimes A$: the identity, the swap $\tau(x \otimes y) = y \otimes x$, and the α-twisted map $\sigma_\alpha(x \otimes y) = \alpha(y) \otimes \alpha(x)$ of Section 4.

Let $\sigma \colon A \otimes A \to A \otimes A$ be a K-linear map with $\sigma^2 = \mathrm{id}$. We write $x \cdot_\sigma y = xy + \mu(\sigma(x \otimes y))$. We ask for no further axiom. The derivations below are called

braided-symmetric because σ plays the role of the braiding operator in the work on braided derivations cited above.

3. Braided-symmetric derivations

This section defines braided-symmetric derivations and identifies them for the identity and the swap.

Definition 3.1. Let A be a K-algebra and let σ be a K-linear involution of $A \otimes A$. A *braided-symmetric derivation* for σ is a K-linear map $D\colon A \to A$ that satisfies identity (1), that is, $D(x \cdot_\sigma y) = D(x) \cdot_\sigma y + x \cdot_\sigma D(y)$ for all $x, y \in A$. We write $\mathrm{BSDer}_\sigma(A)$ for the set of braided-symmetric derivations for σ. This set is a K-subspace of $\mathrm{End}_K(A)$, since the condition is linear in D. For an automorphism α of A, we write $\mathrm{BSDer}_\sigma(A)^\alpha$ for the subset of maps D with $D\alpha = \alpha D$.

The name is close to an older one, and the next remark separates the two.


Remark 3.2. Braided derivations [11, §2] [10, §4.1, Definition 14] form a different notion from braided-symmetric derivations, although both use a braiding. Weber [12, §3, equation (3.9)] defines a braided derivation X of a braided commutative module algebra A over a triangular Hopf algebra (H, R) by a Leibniz rule for the product ab itself:

$$X(ab) = X(a)b + (R_1^{-1} \triangleright a)\big((R_2^{-1} \triangleright X)(b)\big).$$

The braiding enters through the R-matrix. Definition 3.1 asks instead for the ordinary Leibniz rule for the new product $\mu + \mu \circ \sigma$.

Example 3.3 shows that the product $\cdot_\sigma$ can fail to be associative and unital, even when A is commutative.

Example 3.3. Let $A = K \times K$ with componentwise operations, and let $\alpha(a, b) = (b, a)$. Then α is an automorphism with $\alpha^2 = \mathrm{id}$, and $\sigma_\alpha(x \otimes y) = \alpha(y) \otimes \alpha(x)$ is an involution of $A \otimes A$. Since A is commutative, $x \cdot_{\sigma_\alpha} y = xy + \alpha(xy)$. For $x = (x_1, x_2)$ and $y = (y_1, y_2)$ this gives $x \cdot_{\sigma_\alpha} y = (x_1y_1 + x_2y_2)(1, 1)$. Put $p = (1, 0)$ and $q = (0, 1)$. Then $(p \cdot_{\sigma_\alpha} p) \cdot_{\sigma_\alpha} q = (1, 1) \cdot_{\sigma_\alpha} q = (1, 1)$, while $p \cdot_{\sigma_\alpha} (p \cdot_{\sigma_\alpha} q) = p \cdot_{\sigma_\alpha} 0 = 0$. The two bracketings are the two trees



where each dot is the product $\cdot_{\sigma_\alpha}$ and each label is the value on that edge. So $\cdot_{\sigma_\alpha}$ is not associative. Every product $u \cdot_{\sigma_\alpha} y$ lies in $K(1, 1)$, so $u \cdot_{\sigma_\alpha} p \neq p$ for every u, and $\cdot_{\sigma_\alpha}$ has no unit.

The next remark records what Definition 3.1 says for the two involutions that use no structure of A.

Remark 3.4. Let A be a K-algebra and let σ be an involution of $A \otimes A$. Identity (1) for σ is the Leibniz rule for the product $\cdot_\sigma$. So $\mathrm{BSDer}_\sigma(A)$ is the set of derivations of the nonassociative algebra $(A, \cdot_\sigma)$, in the sense of Jacobson [9, Part I].

For the identity, $x \cdot_{\mathrm{id}} y = 2xy$, so the identity reads

$$2D(xy) = 2\big(D(x)y + xD(y)\big).$$

If $\operatorname{char} K \neq 2$, we divide by 2 and get $\mathrm{BSDer}_{\mathrm{id}}(A) = \mathrm{Der}(A)$. If $\operatorname{char} K = 2$, then $x \cdot_{\mathrm{id}} y = 0$, so $\mathrm{BSDer}_{\mathrm{id}}(A)$ contains every linear map. So it is larger than $\mathrm{Der}(A)$. For example, take the linear map with $D(1) = 1$ and $D = 0$ on a complement of $K1$. Then D is not a derivation, since a derivation satisfies $D(1) = D(1 \cdot 1) = 2D(1) = 0$ in characteristic 2.

For the swap τ we get $x \cdot_\tau y = xy + yx = x \circ y$. So the defining identity is the $\circ$ form of the Jordan identity from Section 2. Hence $\mathrm{BSDer}_\tau(A) = \mathrm{JDer}(A)$ in every characteristic.

Every derivation is a Jordan derivation [2, p. 1003]. So in every characteristic

$$\mathrm{Der}(A) \subseteq \mathrm{BSDer}_\tau(A) = \mathrm{JDer}(A).$$

When $\operatorname{char} K \neq 2$, this gives $\mathrm{BSDer}_{\mathrm{id}}(A) \subseteq \mathrm{BSDer}_\tau(A)$. This is an inclusion between the spaces of two involutions on one algebra.

4. Twisted involutions and the Leibniz defect

Remark 3.4 treated the identity and the swap, which use no structure of A. This section twists the swap by an involution α of A and treats it with one tool, the Leibniz defect.

We recall the notions used here. An *automorphism* of A is a bijective K-linear map $\alpha\colon A \to A$ with $\alpha(xy) = \alpha(x)\alpha(y)$ and $\alpha(1) = 1$. An *anti-automorphism* of A is a bijective K-linear map $\alpha\colon A \to A$ with $\alpha(xy) = \alpha(y)\alpha(x)$ and $\alpha(1) = 1$. By $(xy)^T = y^T x^T$, the transpose is an anti-automorphism of $M_n(K)$. In both cases α is *involutive* if $\alpha^2 = \mathrm{id}$. An involutive anti-automorphism is an involution of the ring A in the sense of Herstein [6, §1.2], that is, a map $*$ with $a^{**} = a$, $(a+b)^* = a^* + b^*$ and $(ab)^* = b^* a^*$.

We need $\alpha^2 = \mathrm{id}$ for the following reason. Applying σ_α below twice gives

$$x \otimes y \mapsto \alpha^2(x) \otimes \alpha^2(y),$$

so σ_α is an involution of $A \otimes A$ exactly when α is involutive. When 2 is invertible, an involutive α splits $A = A_+ \oplus A_-$, where $A_\pm = \{x \in A : \alpha(x) = \pm x\}$. Indeed, for $x \in A$,

$$x = \tfrac{1}{2}\big(x + \alpha(x)\big) + \tfrac{1}{2}\big(x - \alpha(x)\big).$$

We call the elements of A_+ *symmetric* and the elements of A_- *skew*.

Let α be involutive, of either kind. Then $\sigma_\alpha(x \otimes y) = \alpha(y) \otimes \alpha(x)$ is a K-linear involution of $A \otimes A$. So σ_α defines a symmetrized product as in Section 2. The product

it defines is

$$x \cdot_{\sigma_\alpha} y = xy + \alpha(y)\alpha(x). \tag{2}$$

For $\alpha = \mathrm{id}$ the product μ_{σ_α} is the Jordan product. If α is an automorphism, then $\alpha(y)\alpha(x) = \alpha(yx)$. If α is an anti-automorphism, then $\alpha(y)\alpha(x) = \alpha(xy)$.

Definition 4.1. The *Leibniz defect* of a linear map $D\colon A \to A$ is the bilinear map $E_D\colon A \times A \to A$ given, for $x, y \in A$, by

$$E_D(x, y) = D(xy) - D(x)y - xD(y).$$

The name is ours, and the map is classical: Cusack [4, §2] writes it $d(a, b) = D(ab) - aDb - (Da)b$ and studies Jordan derivations through it. We recall the Hochschild complex of A with coefficients in A [13, §9.1]. An *n-cochain* is a K-multilinear map from A^n to A. So a 0-cochain is an element of A, and a 1-cochain is a linear map $D\colon A \to A$. The *coboundary* δ sends each n-cochain to an $(n+1)$-cochain, and $\delta \circ \delta = 0$ [13, §9.1]. An *n-cocycle* is an n-cochain f with $\delta f = 0$. The n-th *Hochschild cohomology* $HH^n(A, A)$ is the space of n-cocycles modulo the images δg of the $(n-1)$-cochains g [13, §9.1].

On a 1-cochain D the coboundary is $(\delta D)(x, y) = xD(y) - D(xy) + D(x)y$ [7, §2, Definition 2.1]. So $E_D = -\delta D$. Thus D is a derivation exactly when $E_D = 0$, so derivations are the 1-cocycles. Since $\delta \circ \delta = 0$, the defect is a 2-cocycle: for all $a, b, c \in A$,

$$aE_D(b, c) - E_D(ab, c) + E_D(a, bc) - E_D(a, b)c = 0. \tag{3}$$

This says that E_D is a cocycle of the Hochschild complex [13, §9.1].

The defect also describes the other classes of Section 2. Expanding $D(x \circ y)$ shows that D is a Jordan derivation exactly when $E_D(x, y) + E_D(y, x) = 0$ for all x, y.

Proposition 4.2. *Let A be a unital associative K-algebra, let α be an involutive automorphism or anti-automorphism of A, and let $D\colon A \to A$ be a linear map with $D\alpha = \alpha D$.*

(i) If α is an automorphism, then $D \in \mathrm{BSDer}_{\sigma_\alpha}(A)$ if and only if $E_D(x, y) + \alpha(E_D(y, x)) = 0$ for all $x, y \in A$.

(ii) If α is an anti-automorphism, then $D \in \mathrm{BSDer}_{\sigma_\alpha}(A)$ if and only if $E_D(x, y) + \alpha(E_D(x, y)) = 0$ for all $x, y \in A$.

In both cases every derivation that commutes with α lies in $\mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha$.

Proof. Write $E = E_D$. By Remark 3.4, $D \in \mathrm{BSDer}_{\sigma_\alpha}(A)$ means $D(x \cdot_{\sigma_\alpha} y) = D(x) \cdot_{\sigma_\alpha} y + x \cdot_{\sigma_\alpha} D(y)$ for all x, y. We compute the difference of the two sides in each case.

For (i), the product (2) is $x \cdot_{\sigma_\alpha} y = xy + \alpha(yx)$. Since $D\alpha = \alpha D$, the left side is $D(xy) + \alpha(D(yx))$. The right side is $D(x)y + \alpha(yD(x)) + xD(y) + \alpha(D(y)x)$. We group the terms without α and the terms with α. The first group gives $D(xy) - D(x)y - xD(y) = E(x, y)$. Since α is linear, the second group gives $\alpha\big(D(yx) - D(y)x - yD(x)\big) = \alpha(E(y, x))$. So the left side minus the right side is $E(x, y) + \alpha(E(y, x))$, which proves (i).

The proof of (ii) follows the same steps as the proof of (i). Here the product (2) is

$x \cdot_{\sigma_\alpha} y = xy + \alpha(xy)$. Since $D\alpha = \alpha D$, the left side is

$$D(xy) + \alpha(D(xy)),$$

and the right side is

$$D(x)y + \alpha(D(x)y) + xD(y) + \alpha(xD(y)).$$

The same grouping gives the difference

$$E(x, y) + \alpha(E(x, y)),$$

which proves (ii).

If D is a derivation, then $E = 0$, and both conditions hold. □

We now read the two conditions, assuming that 2 is invertible.

In case (ii), the equation $E_D + \alpha(E_D) = 0$ says $\alpha(E_D) = -E_D$. So E_D takes values in A_-, the skew elements of α. In case (i), we write $E_D = E_\mathrm{s} + E_\mathrm{a}$. Here $E_\mathrm{s}(x, y) = \frac{1}{2}(E_D(x, y)+E_D(y, x))$ is symmetric. And $E_\mathrm{a}(x, y) = \frac{1}{2}(E_D(x, y)-E_D(y, x))$ is antisymmetric.

We swap x and y in the condition of (i). Then we take the sum and the difference of the two equations. So the condition holds exactly when E_s takes values in A_- and E_a takes values in A_+. Now let $\alpha = \mathrm{id}$. Then $A_- = 0$ and $A_+ = A$, so the condition says $E_\mathrm{s} = 0$. This is the Jordan condition, so Proposition 4.2(i) recovers the swap case of Remark 3.4.

Remark 4.3. Let α be an involutive automorphism of A, and let $D \in \mathrm{Der}(A)$. Then $E_D = 0$. By (2), the two sides of the σ_α-identity differ by

$$D(\alpha(yx)) - \alpha(D(yx)).$$

Since A is unital, the products yx span A. So $D \in \mathrm{BSDer}_{\sigma_\alpha}(A)$ if and only if $D\alpha = \alpha D$, and then $D \in \mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha$.

Next we look at the inner derivations ad_g in this class. The condition $\mathrm{ad}_g \alpha = \alpha \mathrm{ad}_g$ means

$$[g - \alpha(g), \alpha(x)] = 0 \qquad \text{for all } x.$$

Since α is bijective, this holds if and only if $g-\alpha(g) \in Z(A)$. Let $\mathrm{Inn}_\alpha(A)$ be the space of the maps ad_g with $g-\alpha(g) \in Z(A)$. These are the inner derivations in $\mathrm{BSDer}_{\sigma_\alpha}(A)$, so $\mathrm{Inn}_\alpha(A) = \mathrm{Inn}(A) \cap \mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha$. The quotient $\mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha / \mathrm{Inn}_\alpha(A)$ is the space of *twisted outer derivations.*

Cusack [4, Corollary 5] and Brešar [2, Theorem 1] proved that a Jordan derivation of a 2-torsion free semiprime ring is a derivation.

Theorem 4.4 ([2, Theorem 1])**.** *Let R be a 2-torsion free semiprime ring. Then every Jordan derivation $D\colon R \to R$ is a derivation.*

The next corollary applies Theorem 4.4 to the twisted class.

Corollary 4.5. *Let A be a 2-torsion-free semiprime K-algebra with an involutive automorphism α. Then* $\mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha \cap \mathrm{JDer}(A) = \mathrm{Der}(A)^\alpha$*, where* $\mathrm{Der}(A)^\alpha$ *is the set of derivations commuting with α.*

Proof. Every derivation is a Jordan derivation by Remark 3.4. With Proposition 4.2, this gives $\mathrm{Der}(A)^\alpha \subseteq \mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha \cap \mathrm{JDer}(A)$. Conversely, every Jordan derivation of A is a derivation by Theorem 4.4, which applies since a K-linear map is additive. So a map in the intersection is a derivation that commutes with α. □

5. Reduced commutative algebras and twisted derivations

For commutative algebras the Jordan hypothesis in Corollary 4.5 can be dropped. Recall from Section 2 that a reduced commutative algebra has no nonzero nilpotent element.

Theorem 5.1. *Let A be a commutative reduced K-algebra with 2 invertible in K, and let α be an involutive automorphism of A. Then* $\mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha = \mathrm{Der}(A)^\alpha$.

Proof. Every derivation commuting with α lies in $\mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha$ by Proposition 4.2. Conversely, let $D \in \mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha$. Write $E = E_D$. Since A is commutative, E is symmetric. So E takes values in A_- by Proposition 4.2(i).

Since D commutes with α, it maps each of A_+ and A_- into itself. So if x and y lie in the same part, then $E(x, y)$ lies in

$$A_+ \cap A_- = 0.$$

Now let $x \in A_+$ and $y \in A_-$, and put $z = E(x, y)$. Then z lies in A_-, and so do xy, zx and xz, while zy lies in A_+. Apply the cocycle identity (3) to $(a, b, c) = (z, x, y)$:

$$zE(x, y) - E(zx, y) + E(z, xy) - E(z, x)y = 0.$$

The pairs (zx, y) and (z, xy) lie in A_-, so E vanishes on them, and we get $z^2 = E(z, x)y$. Apply (3) to $(a, b, c) = (x, z, y)$:

$$xE(z, y) - E(xz, y) + E(x, zy) - E(x, z)y = 0.$$

The pairs (z, y) and (xz, y) lie in A_-, and the pair (x, zy) lies in A_+, so E vanishes on all three and $E(x, z)y = 0$. Since E is symmetric, $z^2 = E(z, x)y = E(x, z)y = 0$. So $z = 0$ because A is reduced. Since E is bilinear and symmetric and $A = A_+ \oplus A_-$, we get $E = 0$. Hence D is a derivation. □

The mixed-parity step uses the cocycle identity twice. Example 5.2 shows that the reduced hypothesis cannot be dropped, and Example 5.3 shows the same for the invertibility of 2.

Example 5.2. Let $A = \mathbb{F}_5[u, v]/(u^2, v^2)$. It has basis $\{1, u, v, uv\}$. Let α fix u and send v to $-v$. Let D send uv to uv and the other three basis elements to 0. Then D commutes with α. It satisfies the Leibniz rule for $x \cdot_{\sigma_\alpha} y$ on all basis pairs. So $D \in \mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha$. But D is not a derivation, since

$$D(uv) = uv \qquad \text{while} \qquad D(u)v + uD(v) = 0.$$

Direct computation on the four basis elements verifies all three statements. Here $z = E(u,v) = uv$ is a nonzero element with $z^2 = 0$.

The next example shows that the invertibility of 2 is needed in Theorem 5.1.

Example 5.3. Let $A = \mathbb{F}_2 \times \mathbb{F}_2$, which is commutative and reduced, and let α swap the two factors. Put $p = (1,0)$, $q = (0,1)$ and $e = (1,1)$, and let D be the linear map with $D(p) = D(q) = e$, so that $D(e) = 0$. Then D commutes with α. On the basis pairs (p,p), (q,q) and (p,q), both sides of the Leibniz rule for $x \cdot_{\sigma_\alpha} y = xy + \alpha(xy)$ are zero. So $D \in \mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha$. But D is not a derivation: $D(p) = D(p^2) = 2pD(p) = 0$ would hold for a derivation, while $D(p) = e$.

Our second natural candidate for the twisted class is the space $\mathrm{Der}_\alpha(A)$ of α-derivations. It was defined for automorphisms in Section 2. It differs from $\mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha$ in both directions, and one algebra shows both failures.

Proposition 5.4. *Let $A = \mathbb{F}_3[\varepsilon]/(\varepsilon^3)$, and let α be the automorphism with $\alpha(\varepsilon) = -\varepsilon$. Then*

$$\mathrm{Der}_\alpha(A) \not\subseteq \mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha \qquad \textit{and} \qquad \mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha \not\subseteq \mathrm{Der}_\alpha(A).$$

Proof. The algebra A has basis $\{1, \varepsilon, \varepsilon^2\}$. The map α fixes 1 and ε^2 and sends ε to $-\varepsilon$. So it is an involutive automorphism with $A_+ = \mathbb{F}_3 1 + \mathbb{F}_3 \varepsilon^2$ and $A_- = \mathbb{F}_3 \varepsilon$. Since A is commutative, E_D is symmetric for every linear map D. Let D be a linear map with $D\alpha = \alpha D$. By Proposition 4.2(i), D lies in $\mathrm{BSDer}_{\sigma_\alpha}(A)$ exactly when E_D takes values in A_-. We give one map for each non-inclusion. Both maps are diagonal on the basis, so both commute with α.

For the first non-inclusion, let $D(a + b\varepsilon + c\varepsilon^2) = b\varepsilon + c\varepsilon^2$. We check that D is an α-derivation on pairs of basis elements. Pairs that contain 1 give 0 on both sides, since $D(1) = 0$ and $\alpha(1) = 1$. For the pair $(\varepsilon, \varepsilon)$, the left side is $D(\varepsilon^2) = \varepsilon^2$. The right side is $D(\varepsilon)\alpha(\varepsilon) + \alpha(\varepsilon)D(\varepsilon) = -2\varepsilon^2 = \varepsilon^2$. The remaining pairs have product 0, and each term on the right side is a multiple of $\varepsilon^3 = 0$. So $D \in \mathrm{Der}_\alpha(A)$. On the other hand, $E_D(\varepsilon, \varepsilon) = D(\varepsilon^2) - 2\varepsilon D(\varepsilon) = \varepsilon^2 - 2\varepsilon^2 = 2\varepsilon^2$. This element is nonzero and lies in A_+, so it does not lie in A_-. Hence $D \notin \mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha$.

The second non-inclusion uses a derivation.

Let d be the linear map

$$d(a + b\varepsilon + c\varepsilon^2) = b\varepsilon + 2c\varepsilon^2.$$

It is a derivation, since $d(\varepsilon^i) = i\varepsilon^i$ and exponents add under multiplication. So $E_d = 0$, and $d \in \mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha$. On the other hand,

$$d(\varepsilon^2) - d(\varepsilon)\alpha(\varepsilon) - \alpha(\varepsilon)d(\varepsilon) = 2\varepsilon^2 + 2\varepsilon^2 = \varepsilon^2 \neq 0.$$

Hence $d \notin \mathrm{Der}_\alpha(A)$. □

The two classes can also meet, and the next example shows both cases.

Example 5.5. Let $A = M_2(\mathbb{F}_3)$ and let α be the transpose, an anti-automorphism. Then $\mathrm{Der}_\alpha(A)$ is not defined, since it needs an automorphism. Take an inner derivation ad_g with g skew for the transpose. It commutes with α and has $E = 0$, so it lies in $\mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha$ by Proposition 4.2.

Now let $A = K[\varepsilon]/(\varepsilon^2)$ with $\alpha(\varepsilon) = -\varepsilon$. The map $D(a+b\varepsilon) = b\varepsilon$ is a derivation and an α-derivation, and it commutes with α. So it lies in both $\mathrm{Der}_\alpha(A)$ and $\mathrm{BSDer}_{\sigma_\alpha}(A)^\alpha$ by Proposition 4.2.

References

[1] Benkovič, D.: A note on f-derivations of triangular algebras. Aequationes Math. **89**, 1207–1211 (2015).

[2] Brešar, M.: Jordan derivations on semiprime rings. Proc. Amer. Math. Soc. **104**(4), 1003–1006 (1988).

[3] Chambert-Loir, A.: (Mostly) Commutative Algebra. Universitext. Springer, Cham (2021).

[4] Cusack, J.M.: Jordan derivations on rings. Proc. Amer. Math. Soc. **53**(2), 321–324 (1975).

[5] Herstein, I.N.: Jordan derivations of prime rings. Proc. Amer. Math. Soc. **8**(6), 1104–1110 (1957).

[6] Herstein, I.N.: Rings with Involution. Chicago Lectures in Mathematics. University of Chicago Press, Chicago (1976).

[7] Hochschild, G.: On the cohomology groups of an associative algebra. Ann. of Math. **46**(1), 58–67 (1945).

[8] Hosseini, A., Karizaki, M.M.: Functional equations characterizing σ-derivations. Acta Sci. Math. (Szeged) **85**, 431–440 (2019).

[9] Jacobson, N.: Abstract derivation and Lie algebras. Trans. Amer. Math. Soc. **42**(2), 206–224 (1937).

[10] Huru, H.L.: Quantizations of braided derivations. I: Monoidal categories. Lobachevskii J. Math. **24**, 13–42 (2006).

[11] Różański, J.: Bicharacters, braids and Jacobi identity. arXiv:q-alg/9611029 (1996).

[12] Weber, T.: Braided Cartan calculi and submanifold algebras. J. Geom. Phys. **156**, 103612 (2020). `https://doi.org/10.1016/j.geomphys.2020.103612`

[13] Weibel, C.A.: An Introduction to Homological Algebra. Cambridge Stud. Adv. Math. 38, Cambridge University Press, Cambridge (1994).