Jordan and Lie Derivations as Derivations of the Symmetrized and Antisymmetrized Products

Abderrahim Adrabi[a,b]

[a]Faculty of Sciences, Mohammed V University in Rabat, Morocco;
[b]Faculty of Educational Sciences, Mohammed V University in Rabat, Morocco

ARTICLE HISTORY
Preprint version 10, compiled 2026/10/28

ABSTRACT
We view the Jordan derivations and the Lie derivations of an algebra as the derivations of two products on the algebra. These are the symmetrized product $xy + yx$ and the antisymmetrized product $xy - yx$. The question whether such a derivation is a derivation becomes a comparison of three spaces, $\mathrm{Der}(A)$, $\mathrm{JDer}(A)$ and $\mathrm{LDer}(A)$. We show that the dimension of $\mathrm{JDer}(A)$ does not change when the base field is extended. We prove that the σ-antisymmetric derivations for the swap are exactly the Lie derivations, and that every derivation plus a central trace-type map that kills commutators is a Lie derivation. Such a map is a derivation only if the trace-type part is zero at the identity. We compute the three spaces for full matrix algebras. For triangular algebras we apply the theorem of Zhang and Yu to the Jordan case and the theorem of Cheung to the Lie case, with the upper triangular 2×2 matrices as the smallest case. In the Jordan case every such derivation is a derivation. In the Lie case every such derivation has the standard form, and a trace-type map gives a Lie derivation that is not a derivation.

KEYWORDS
Jordan derivations; Lie derivations; symmetrized product; triangular algebras; matrix algebras

AMS CLASSIFICATION
MSC 2020: 16W25; 17C50; 15A78

1. Introduction

Throughout, K is a field and an algebra means a unital associative K-algebra. We take all maps between algebras to be K-linear. We say so when a result cited below holds more generally for additive maps on rings.

A *derivation* of an algebra A is a linear map $D\colon A \to A$ with $D(xy) = D(x)y + xD(y)$ for all $x, y \in A$. The notion goes back to Jacobson [11, Part I], who studied the derivations of an arbitrary nonassociative algebra. Hochschild [10, §§2 and 4] defined the cocycles of an associative algebra, and its 1-cocycles are the derivations. A *Jordan derivation* of A is a linear map with $D(x \circ y) = D(x) \circ y + x \circ D(y)$ for all $x, y \in A$, where $x \circ y = xy + yx$. The sum $xy + yx$ is twice the Jordan product $\frac{1}{2}(xy + yx)$ of an associative algebra [4, Example 2.1]. Jordan, von Neumann and Wigner form

CONTACT A. Adrabi. Email: abderrahim.adrabi@um5r.ac.ma

a product ab from the two products $a \times b$ and $b \times a$ of an associative system, and call this quasi-multiplication [12, p. 64]. Herstein [8, p. 1104] takes the sum $ab + ba$ without the half. We follow Herstein and do not divide by 2, so that characteristic 2 is allowed. Herstein [8, p. 1104] introduced this notion. His result needs three notions for a ring R. We call R *prime* if $aRb = 0$ implies $a = 0$ or $b = 0$ [3, p. 1003]. We call R *semiprime* if $aRa = 0$ implies $a = 0$ [3, p. 1003]. We call R 2*-torsion free* if $2a = 0$ implies $a = 0$ [3, p. 1003]. So every prime ring is semiprime. Herstein proved that every Jordan derivation of a prime ring of characteristic different from 2 is a derivation [8, Theorem 3.1]. Every derivation is a Jordan derivation, and the converse is false in general [3, p. 1003]. Cusack [7, Corollary 5] extended this result of Herstein to 2-torsion free semiprime rings, and Brešar [3, Theorem 1] gave an independent proof.

A *Lie derivation* of A is a linear map with $D([x, y]) = [D(x), y] + [x, D(y)]$, where $[x, y] = xy - yx$. Martindale [14, p. 183] studied the Lie derivations of a primitive ring of characteristic not 2 that contains a nontrivial idempotent, and showed that each has the form $D + T$, with D a derivation and T a map into the centre that vanishes on commutators. So for Lie derivations no result of Herstein type can hold. Here is the reason. Let $t\colon A \to Z(A)$ be a linear map into the centre with $t([x, y]) = 0$ for all x, y. Every term of the Lie derivation identity for t is then zero, so t is a Lie derivation. But t is not a derivation as soon as $t(1) \neq 0$.

So a description of Lie derivations must allow such a term. A Lie derivation has the *standard form* if it equals $d + t$, with d a derivation and t a central-valued map that vanishes on commutators [6, §1]. A ring is *left primitive* if it has a faithful simple left module, and right primitive rings are defined in the same way [13, Definition 11.2]. Martindale [14, Theorem 2] proved that every Lie derivation has this form for primitive rings of characteristic not 2 with a nontrivial idempotent. Here an *idempotent* is an element e with $e^2 = e$, and it is *nontrivial* if $e \neq 0$ and $e \neq 1$. Here d may take values in a larger ring.

Triangular algebras are defined in Section 2. For them, Zhang and Yu [16, Theorem 2.1] proved that every Jordan derivation of a triangular algebra is a derivation. For Lie derivations, Cheung [6, Theorem 11] found conditions under which every Lie derivation has the standard form. Wang [17, Theorem 3.1 and Corollary 3.1] treated every triangular algebra, with no condition on the module or the centre. For these algebras a Lie derivation is a sum $\delta + \tau$, where δ is a derivation into a larger triangular algebra and τ is a linear map into its centre that vanishes on commutators.

In this paper a Jordan derivation is a derivation of the symmetrized product $x \circ y$ [4, Example 2.1], and a Lie derivation is a derivation of the antisymmetrized product $[x, y]$. Herstein [9, §2] calls $[a, b] = ab - ba$ the Lie product and $a \circ b = ab + ba$ the Jordan product of an associative ring, and notes that the two coincide in characteristic 2. We follow this order. Section 2 recalls the classical notions, recalls the symmetrized product, the braided-symmetric derivations and their values for the identity and the swap from the first paper, computes the dual numbers, and applies the theorem of Zhang and Yu to triangular algebras. Section 3 shows that the dimension of the space of Jordan derivations is stable under extension of the base field. Section 4 treats the antisymmetrized product, full matrix algebras, and triangular algebras through the theorems of Martindale and Cheung. We place each example right after the result it illustrates. The twisted symmetrized product and its derivations are the subject of the first paper in our series on symmetrized products [1], from which the definition and the dictionary of Section 2 are taken.

2. Preliminaries and the symmetrized product

This section recalls the classical notions we need, then the symmetrized product and the braided-symmetric derivations of the first paper, with the result of that paper for the identity and the swap.

From now on, A is a K-algebra with unit 1. We write $[x, y] = xy - yx$ for the commutator, $x \circ y = xy + yx$ for the Jordan product, and $Z(A)$ for the centre. The centre is the set of elements z with $zx = xz$ for all $x \in A$. We call an element x *nilpotent* if $x^n = 0$ for some $n \geq 1$. A commutative algebra is *reduced* if its only nilpotent element is zero [5, 1.4.12].

A derivation of an algebra is as defined by Jacobson [11, Part I]: a linear map D with $D(xy) = D(x)y + xD(y)$. Every derivation satisfies $D(1) = 0$, since $D(1) = D(1 \cdot 1) = 2D(1)$. For $g \in A$, the map $\mathrm{ad}_g(x) = [g, x]$ is an *inner derivation*, called a principal derivation by Weibel [18, §9.2]. We write $\mathrm{Der}(A)$ and $\mathrm{Inn}(A)$ for these two spaces.

We add a word on the characteristic for Jordan derivations. A *Jordan derivation* in the sense of Herstein [8, p. 1104] is an additive map with $D(x^2) = D(x)x + xD(x)$. We write $\mathrm{JDer}(A)$ for the space of linear maps with $D(x \circ y) = D(x) \circ y + x \circ D(y)$ for all x, y. Replacing x by $x + y$ gives the condition $D(x \circ y) = D(x) \circ y + x \circ D(y)$ in every characteristic. Setting $y = x$ in this condition gives $2D(x^2) = 2(D(x)x + xD(x))$. So the two conditions agree when 2 is invertible in K, and we use the $\circ$ form throughout. When $\operatorname{char} K = 2$ and A is commutative, $x \circ y = 0$, and every linear map satisfies the $\circ$ condition.

We need two more kinds. A *Lie derivation* satisfies $D([x, y]) = [D(x), y] + [x, D(y)]$ [14, p. 183], and we write $\mathrm{LDer}(A)$ for this space. When A is commutative, every bracket is zero, so every linear map is a Lie derivation.

We now compare these spaces. When 2 is invertible, $xy = \frac{1}{2}(x \circ y) + \frac{1}{2}[x, y]$. A derivation respects xy, so it respects both halves, and $\mathrm{Der}(A) \subseteq \mathrm{JDer}(A) \cap \mathrm{LDer}(A)$. Both inclusions can be strict, as Examples 2.3 and 4.9 show. Our examples include triangular algebras. An (A, B)-*bimodule* is a left A-module and right B-module M with $(am)b = a(mb)$ for all $a \in A$, $m \in M$ and $b \in B$ [13, Example 1.14]. We call it *unital* if $1m = m = m1$ for all $m \in M$.

Let R be a commutative ring, let A and B be unital R-algebras, and let M be a unital (A, B)-bimodule. The *triangular algebra* $T = \mathrm{Tri}(A, M, B)$ is the set of matrices $\left(\begin{smallmatrix} a & m \\ 0 & b \end{smallmatrix}\right)$ with the matrix sum and product [6, §2]. The module M is *faithful* as a left A-module if $aM = 0$ implies $a = 0$, and as a right B-module if $Mb = 0$ implies $b = 0$. The idempotents $e_1 = \left(\begin{smallmatrix} 1 & 0 \\ 0 & 0 \end{smallmatrix}\right)$ and $e_2 = \left(\begin{smallmatrix} 0 & 0 \\ 0 & 1 \end{smallmatrix}\right)$ satisfy $e_2 T e_1 = 0$. So T is not prime, since $e_1 \neq 0$ and $e_2 \neq 0$, and the prime-ring results above do not apply to triangular algebras.

Let A be a K-algebra with product μ. Let $\sigma \colon A \otimes A \to A \otimes A$ be a K-linear map with $\sigma^2 = \mathrm{id}$, for instance the swap $\tau(x \otimes y) = y \otimes x$. It gives a symmetrized product [1, Section 2]

$$x \cdot_\sigma y = \mu(x \otimes y) + \mu(\sigma(x \otimes y)).$$

For $\sigma = \tau$ this is the Jordan product $x \circ y$.

Definition 2.1 ([1, Definition 2.1])**.** Let A be a K-algebra and let σ be a K-linear involution of $A \otimes A$. A *braided-symmetric derivation* for σ is a K-linear map $D \colon A \to A$

with

$$D(x \cdot_\sigma y) = D(x) \cdot_\sigma y + x \cdot_\sigma D(y) \quad \text{for all } x, y \in A. \tag{1}$$

We write $\mathrm{BSDer}_\sigma(A)$ for the set of braided-symmetric derivations for σ. This set is a K-subspace of $\mathrm{End}_K(A)$, since the condition is linear in D. For an automorphism α of A, we write $\mathrm{BSDer}_\sigma(A)^\alpha$ for the subset of maps D with $D\alpha = \alpha D$.

Identity (1) says that the following square commutes.

$$\begin{array}{ccc} A \otimes A & \xrightarrow{\mu_\sigma} & A \\ {\scriptstyle D\otimes 1 + 1 \otimes D}\downarrow & & \downarrow{\scriptstyle D} \\ A \otimes A & \xrightarrow[\mu_\sigma]{} & A \end{array}$$

The next remark, recalled from the first paper, says what Definition 2.1 gives for the two involutions that use no structure of A.

Remark 2.2 ([1, Remark 2.4])**.** Let A be a K-algebra and let σ be an involution of $A \otimes A$. Identity (1) for σ is the Leibniz rule for the product $\cdot_\sigma$. So $\mathrm{BSDer}_\sigma(A)$ is the set of derivations of the nonassociative algebra $(A, \cdot_\sigma)$, in the sense of Jacobson [11, Part I].

For the identity, $x \cdot_{\mathrm{id}} y = 2xy$, so the identity reads

$$2D(xy) = 2\big(D(x)y + xD(y)\big).$$

If $\operatorname{char} K \neq 2$, we divide by 2 and get $\mathrm{BSDer}_{\mathrm{id}}(A) = \mathrm{Der}(A)$. If $\operatorname{char} K = 2$, then $x \cdot_{\mathrm{id}} y = 0$, so $\mathrm{BSDer}_{\mathrm{id}}(A)$ contains every linear map. So it is larger than $\mathrm{Der}(A)$. For example, take the linear map with $D(1) = 1$ and $D = 0$ on a complement of $K1$. Then D is not a derivation, since a derivation satisfies $D(1) = D(1 \cdot 1) = 2D(1) = 0$ in characteristic 2.

For the swap τ we get $x \cdot_\tau y = xy + yx = x \circ y$. So the defining identity is the $\circ$ form of the identity for a Jordan derivation recalled above. Hence $\mathrm{BSDer}_\tau(A) = \mathrm{JDer}(A)$ in every characteristic.

Every derivation is a Jordan derivation [3, p. 1003]. So in every characteristic

$$\mathrm{Der}(A) \subseteq \mathrm{BSDer}_\tau(A) = \mathrm{JDer}(A).$$

When $\operatorname{char} K \neq 2$, this gives $\mathrm{BSDer}_{\mathrm{id}}(A) \subseteq \mathrm{BSDer}_\tau(A)$. This is an inclusion between the spaces of two involutions on one algebra.

The smallest nonreduced algebra shows both inclusions recalled above at work.

Example 2.3 (Dual numbers)**.** Let $A = K[\varepsilon]/(\varepsilon^2)$, with basis $\{1, \varepsilon\}$. A linear map D is a derivation exactly when the Leibniz rule holds on the four pairs of basis elements. Since $D(1) = 0$, the pairs $(1, \varepsilon)$ and $(\varepsilon, 1)$ hold. Write $D(\varepsilon) = \beta + c\varepsilon$ with $\beta, c \in K$. The pair $(\varepsilon, \varepsilon)$ gives $0 = D(\varepsilon^2) = 2\varepsilon D(\varepsilon) = 2\beta\varepsilon$. If $\operatorname{char} K \neq 2$, then $\beta = 0$, and every derivation has the form $D(a + b\varepsilon) = bc\varepsilon$, so $\dim_K \mathrm{Der}(A) = 1$. If $\operatorname{char} K = 2$, then β is free, and $\dim_K \mathrm{Der}(A) = 2$. Since A is commutative, every commutator is zero, so $\mathrm{LDer}(A) = \mathrm{End}_K(A)$, which has dimension $2 \cdot 2 = 4$. Each derivation lies in $\mathrm{BSDer}_\tau(A)$

by Remark 2.2. If $\operatorname{char} K = 2$, then $x \circ y = 2xy = 0$ for all x, y, so $\mathrm{JDer}(A) = \mathrm{End}_K(A)$ has dimension 4, which exceeds $\dim_K \mathrm{Der}(A) = 2$.

The remark above and the following theorem on triangular algebras give a corollary.

Theorem 2.4 ([16, Theorem 2.1])**.** *Let R be a 2-torsion free commutative ring, let A and B be unital R-algebras, and let M be a unital (A, B)-bimodule that is faithful as a left A-module and as a right B-module. Then every Jordan derivation $D\colon \mathrm{Tri}(A, M, B) \to \mathrm{Tri}(A, M, B)$ is a derivation.*

In the next corollary, maps on $T = \mathrm{Tri}(A, M, B)$ are R-linear and the σ-symmetric identity is read over R. It follows from Remark 2.2 and Theorem 2.4.

Corollary 2.5. *Let R be a 2-torsion free commutative ring, let A and B be unital R-algebras, and let M be a unital (A, B)-bimodule that is faithful as a left A-module and as a right B-module. Suppose also that $\mathrm{Tri}(A, M, B)$ is 2-torsion free. Then $\mathrm{BSDer}_\tau(\mathrm{Tri}(A, M, B)) = \mathrm{Der}(\mathrm{Tri}(A, M, B))$.*

3. Extension of the base field

Proposition 3.3 below treats extension of the base field. For a field extension L of K and a K-vector space V, write $V_L = L \otimes_K V$. This is an L-vector space. For a K-algebra A, the space A_L is an L-algebra with $(c \otimes x)(c' \otimes y) = cc' \otimes xy$ [13, Chapter 2, §5, p. 74].

We now define the involution σ_L of $A_L \otimes_L A_L$. It is an L-linear map of $A_L \otimes_L A_L$ into itself with $\sigma_L^2 = \mathrm{id}$.

Define

$$\Psi\colon A_L \otimes_L A_L \to L \otimes_K (A \otimes_K A), \qquad (c \otimes x) \otimes_L (c' \otimes y) \longmapsto cc' \otimes (x \otimes y).$$

Since this assignment is L-bilinear, it defines an L-linear map. It is an isomorphism, because its inverse sends $c \otimes (x \otimes y)$ to $(c \otimes x) \otimes_L (1 \otimes y)$.

The map $\mathrm{id}_L \otimes \sigma$ is an L-linear map of $L \otimes_K (A \otimes_K A)$ into itself, and $(\mathrm{id}_L \otimes \sigma)^2 = \mathrm{id}_L \otimes \sigma^2 = \mathrm{id}$. We define σ_L as the conjugate of $\mathrm{id}_L \otimes \sigma$ by Ψ:

$$\sigma_L = \Psi^{-1} \circ (\mathrm{id}_L \otimes \sigma) \circ \Psi.$$

For $c, c' \in L$ and $x, y \in A$ it reads

$$\sigma_L\big((c \otimes x) \otimes_L (c' \otimes y)\big) = \Psi^{-1}\big(cc' \otimes \sigma(x \otimes y)\big).$$

Since $\Psi\Psi^{-1} = \mathrm{id}$, we get

$$\sigma_L^2 = \Psi^{-1}(\mathrm{id}_L \otimes \sigma)(\mathrm{id}_L \otimes \sigma)\Psi = \Psi^{-1}(\mathrm{id}_L \otimes \sigma^2)\Psi = \mathrm{id}.$$

So σ_L is an involution of $A_L \otimes_L A_L$.

The product μ_L of A_L satisfies $\mu_L = (\mathrm{id}_L \otimes \mu) \circ \Psi$. So Ψ carries μ_L and σ_L to $\mathrm{id}_L \otimes \mu$ and $\mathrm{id}_L \otimes \sigma$. Hence $(1 \otimes x) \cdot_{\sigma_L} (1 \otimes y) = 1 \otimes (x \cdot_\sigma y)$ for $x, y \in A$. Thus the product $\cdot_{\sigma_L}$ on A_L extends the product $\cdot_\sigma$ on A.

Proposition 3.3 below says that extending the base field does not change the dimension of the space of braided-symmetric derivations. The proof compares spaces of linear maps over K and over L, and uses results from the recent literature, which we recall. For a K-vector space E we write $E_L = L \otimes_K E$, and for a K-linear map f we write $f_L = \mathrm{id}_L \otimes f$.

Let L be an extension field of K, and let $(c_i)_{i\in I}$ be a basis of L over K. For a K-linear map $f\colon E \to F$, every element of E_L is a unique sum $\sum_i c_i \otimes e_i$ with $e_i \in E$, and $f_L(\sum_i c_i \otimes e_i) = \sum_i c_i \otimes f(e_i)$. So f_L kills the element exactly when every $f(e_i)$ is zero, and $\ker(f_L) = (\ker f)_L$.

The next two results compare linear maps over K and over L, which is the comparison that Proposition 3.3 needs to transfer the equation $\Phi(D) = 0$ from A to A_L.

Lemma 3.1 ([15, Lemma 4.86])**.** *Let B be a flat R-module, and let N be a finitely presented R-module. For every R-module M, there is a natural isomorphism*

$$\psi\colon B \otimes_R \mathrm{Hom}_R(N, M) \to \mathrm{Hom}_R(N, M \otimes_R B),$$

given by $b \otimes g \mapsto g_b$, where $g_b(n) = g(n) \otimes b$ for all $n \in N$.

The next lemma identifies homomorphisms over R with homomorphisms over A after extension of scalars.

Lemma 3.2 ([15, Lemma 4.85])**.** *Let A be an R-algebra, and let N be a finitely presented R-module. For every A-module M, there is a natural isomorphism*

$$\theta\colon \mathrm{Hom}_R(N, M) \to \mathrm{Hom}_A(N \otimes_R A, M),$$

given by $\theta\colon f \mapsto \tilde{f}$, where $\tilde{f}(n \otimes 1) = f(n)$ for all $n \in N$.

Let V be a finite-dimensional K-vector space and let W be a K-vector space. Then V is finitely presented over K, by a finite basis. Lemma 3.1 with $R = K$, $B = L$, $N = V$ and $M = W$ gives $L \otimes_K \mathrm{Hom}_K(V, W) \cong \mathrm{Hom}_K(V, W_L)$. Lemma 3.2 with $R = K$, $A = L$, $N = V$ and $M = W_L$ gives $\mathrm{Hom}_K(V, W_L) \cong \mathrm{Hom}_L(V_L, W_L)$. The composite sends $c \otimes u$ to $c\,(\mathrm{id}_L \otimes u)$. So $L \otimes_K \mathrm{Hom}_K(V, W) \to \mathrm{Hom}_L(V_L, W_L)$ is bijective, which identifies $L \otimes_K \mathrm{End}_K(A)$ with $\mathrm{End}_L(A_L)$ in Proposition 3.3.

For a K-algebra A with involution σ of $A \otimes A$, let $m_\sigma\colon A \otimes A \to A$ be the product $x \otimes y \mapsto x \cdot_\sigma y$. Define the K-linear map

$$\Phi\colon \mathrm{End}_K(A) \to \mathrm{Hom}_K(A \otimes A, A), \qquad \Phi(D) = D \circ m_\sigma - m_\sigma \circ (D \otimes \mathrm{id} + \mathrm{id} \otimes D).$$

So $\Phi(D)$ is the difference of the two sides of identity (1), and $\mathrm{BSDer}_\sigma(A) = \ker \Phi$. Write Φ_L for the same map for A_L and σ_L, so $\mathrm{BSDer}_{\sigma_L}(A_L) = \ker \Phi_L$.

Proposition 3.3. *Let K be a field, let A be a finite-dimensional K-algebra, and let σ be an involution of $A \otimes A$. Let L be a field extension of K. For $D \in \mathrm{End}_K(A)$ and $c \in L$, let $c \otimes D$ act on A_L by $(c \otimes D)(c' \otimes x) = cc' \otimes D(x)$ for $c' \in L$ and $x \in A$. This rule defines isomorphisms of L-vector spaces*

$$\beta\colon L \otimes_K \mathrm{End}_K(A) \to \mathrm{End}_L(A_L), \qquad \gamma\colon L \otimes_K \mathrm{Hom}_K(A \otimes A, A) \to \mathrm{Hom}_L(A_L \otimes_L A_L, A_L)$$

such that the following square commutes.

$$\begin{array}{ccc} L \otimes_K \operatorname{End}_K(A) & \xrightarrow{\mathrm{id}_L \otimes \Phi} & L \otimes_K \operatorname{Hom}_K(A \otimes A, A) \\ \beta \big\downarrow & & \big\downarrow \gamma \\ \operatorname{End}_L(A_L) & \xrightarrow[\Phi_L]{} & \operatorname{Hom}_L(A_L \otimes_L A_L, A_L) \end{array}$$

So β and γ carry the exact sequence

$$0 \to L \otimes_K \operatorname{BSDer}_\sigma(A) \to L \otimes_K \operatorname{End}_K(A) \xrightarrow{\mathrm{id}_L \otimes \Phi} L \otimes_K \operatorname{Hom}_K(A \otimes A, A)$$

isomorphically onto the exact sequence

$$0 \to \operatorname{BSDer}_{\sigma_L}(A_L) \to \operatorname{End}_L(A_L) \xrightarrow{\Phi_L} \operatorname{Hom}_L(A_L \otimes_L A_L, A_L).$$

In particular $L \otimes_K \operatorname{BSDer}_\sigma(A) \cong \operatorname{BSDer}_{\sigma_L}(A_L)$ and $\dim_L \operatorname{BSDer}_{\sigma_L}(A_L) = \dim_K \operatorname{BSDer}_\sigma(A)$.

Proof. First, the spaces A and $A \otimes_K A$ are finite-dimensional. So β is an isomorphism by the bijection above with $V = W = A$. Also γ is an isomorphism by the bijection above with $V = A \otimes_K A$ and $W = A$, since $(A \otimes_K A)_L = A_L \otimes_L A_L$.

Second, we check $\Phi_L \circ \beta = \gamma \circ (\mathrm{id}_L \otimes \Phi)$. Both sides are L-linear. So it is enough to evaluate them on $c \otimes D$ and on $(c' \otimes x) \otimes_L (c'' \otimes y)$, for $c, c', c'' \in L$, $D \in \operatorname{End}_K(A)$ and $x, y \in A$. By the construction of σ_L, we have $m_{\sigma_L}\big((c' \otimes x) \otimes_L (c'' \otimes y)\big) = c'c'' \otimes m_\sigma(x \otimes y)$. Hence both sides give

$$cc'c'' \otimes \Big(D\big(m_\sigma(x \otimes y)\big) - m_\sigma\big(D(x) \otimes y + x \otimes D(y)\big) \Big) = cc'c'' \otimes \Phi(D)(x \otimes y).$$

Third, $\ker \Phi = \operatorname{BSDer}_\sigma(A)$. Since $L \otimes_K -$ is exact, $\ker(\mathrm{id}_L \otimes \Phi) = L \otimes_K \operatorname{BSDer}_\sigma(A)$. By the second step, β and γ turn $\mathrm{id}_L \otimes \Phi$ into Φ_L. So β maps $L \otimes_K \operatorname{BSDer}_\sigma(A)$ onto $\ker \Phi_L = \operatorname{BSDer}_{\sigma_L}(A_L)$. □

The next remark shows that the finite-dimensional hypothesis of Proposition 3.3 is needed.

Remark 3.4. The finite-dimensional hypothesis in Proposition 3.3 cannot be dropped. Take $\sigma = -\mathrm{id}$. Then $\sigma^2 = \mathrm{id}$, so σ is an involution of $A \otimes A$, and $x \cdot_\sigma y = 0$ for all x, y. So every K-linear map is a braided-symmetric derivation, both for A and for A_L.

Let $A = K[x]$ and $L = K(t)$, where t is transcendental over K. Define $D' \in \operatorname{End}_L(A_L)$ by $D'(x^i) = t^i x^i$ for all $i \geq 0$. Suppose $D' = \sum_{r=1}^{s} c_r \otimes D_r$ for some $s \geq 1$, with $c_r \in L$ and $D_r \in \operatorname{End}_K(A)$. Since D_r is K-linear and $A = K[x]$, the coefficient of x^i in $D_r(x^i)$ lies in K. So t^i lies in the K-span of $c_1, \ldots, c_s$, for every i. Since t is transcendental over K, the powers $1, t, t^2, \ldots$ are linearly independent over K, so no finite span contains them all. Hence D' is not in the image, and the map of Proposition 3.3 is not onto.

The next corollary applies the proposition to the space of Jordan derivations.

Corollary 3.5. *Let K be a field, let A be a finite-dimensional K-algebra, and let L be a field extension of K. Then* $\dim_L \mathrm{JDer}(A_L) = \dim_K \mathrm{JDer}(A)$.

Proof. For the swap τ of $A \otimes A$, the involution τ_L is the swap of $A_L \otimes_L A_L$, since $\tau_L((c \otimes x) \otimes_L (c' \otimes y)) = \Psi^{-1}(cc' \otimes (y \otimes x)) = (c' \otimes y) \otimes_L (c \otimes x)$. By Remark 2.2, $\mathrm{BSDer}_\tau(A) = \mathrm{JDer}(A)$ and $\mathrm{BSDer}_{\tau_L}(A_L) = \mathrm{JDer}(A_L)$. Proposition 3.3 gives the equality of dimensions. □

4. The antisymmetrized product and Lie derivations

This section replaces the sum $\mu + \mu \circ \sigma$ by the difference $\mu - \mu \circ \sigma$. For the swap the difference is the commutator, and its derivations are the Lie derivations.

Definition 4.1. The *σ-antisymmetrized product* on a K-algebra A is $x \cdot_\sigma^- y = \mu(x \otimes y) - \mu(\sigma(x \otimes y))$, where σ is an involution of $A \otimes A$. A *σ-antisymmetric derivation* is a K-linear map $D\colon A \to A$ with $D(x \cdot_\sigma^- y) = D(x) \cdot_\sigma^- y + x \cdot_\sigma^- D(y)$.

We now identify these derivations for the swap. It also gives the easy half of the theorems of Martindale and Brešar.

Proposition 4.2. *Let A be an associative K-algebra.*

(i) *The σ-antisymmetric derivations for τ in the sense of Definition 4.1 are exactly the Lie derivations of A.*
(ii) $\mathrm{Der}(A) \subseteq \mathrm{LDer}(A)$.
(iii) *Let $d \in \mathrm{Der}(A)$, and let $t\colon A \to Z(A)$ be linear with $t([x,y]) = 0$ for all x, y. Then $d + t \in \mathrm{LDer}(A)$.*
(iv) *In (iii), if $t(1) \neq 0$, then $d + t \notin \mathrm{Der}(A)$.*

Proof. We check the four parts in order. (i) For the swap, $x \cdot_\tau^- y = [x, y]$, so the two identities are the same. (ii) For $D \in \mathrm{Der}(A)$, $D([x,y]) = D(x)y + xD(y) - D(y)x - yD(x) = [D(x), y] + [x, D(y)]$. (iii) By (ii), $d([x,y]) = [d(x), y] + [x, d(y)]$, and $t([x,y]) = 0$. Since $t(x)$ and $t(y)$ are central, $[t(x), y] = 0 = [x, t(y)]$. Adding gives the Lie derivation identity for $d + t$. (iv) Every derivation vanishes at 1, as recalled in Section 2, while $(d+t)(1) = t(1) \neq 0$. □

So inner derivations lie in both spaces. Each ad_g is a Lie derivation by Proposition 4.2(ii), and a Jordan derivation by Remark 2.2. Full matrix algebras show that part (iii) gives maps outside $\mathrm{Der}(A)$. We use one result on Jordan derivations of matrix rings. It is stated for a unital associative ring and a 2-torsion free bimodule [2, §2].

Theorem 4.3 ([2, Theorem 3.1])**.** *Let A be a unital associative ring, let M be a 2-torsion free A-bimodule, and let $n \geq 2$. Then every Jordan derivation $D\colon M_n(A) \to M_n(M)$ is a derivation.*

We also use Weibel [18, Lemma 9.2.1], which describes the first Hochschild cohomology. It gives HH^1 of the matrix algebra in Example 4.9. Let R be a k-algebra and M an R-bimodule. Weibel [18, §9.2] writes $\mathrm{Der}_k(R, M)$ for the k-derivations. Weibel writes $\mathrm{PDer}_k(R, M)$ for the principal ones $f_m(r) = rm - mr$.

Lemma 4.4 ([18, Lemma 9.2.1])**.** *Let R be a k-algebra and M an R-bimodule. Then* $HH^1(R, M) \cong \mathrm{Der}_k(R, M)/\mathrm{PDer}_k(R, M)$.

A nonzero ring R is *simple* if (0) and R are the only ideals in R, and a commutative ring is simple if and only if it is a field [13, §1].

The next theorem shows that $M_n(K)$ is simple. Example 4.9 needs this twice: to apply the theorem of Martindale, and to see that $M_n(K)$ is separable.

Theorem 4.5 ([13, Theorem 3.1])**.** *Let R be a ring and $M_n(R)$ be the ring of $n \times n$ matrices over R. Then any ideal I of $M_n(R)$ has the form $M_n(\mathfrak{m})$ for a uniquely determined ideal $\mathfrak{m}$ of R. In particular, if R is a simple ring, so is $M_n(R)$.*

The next theorem of Martindale describes the Lie derivations of primitive rings with a nontrivial idempotent.

Theorem 4.6 ([14, Theorem 2])**.** *Let D be a Lie derivation of a primitive ring R into itself, where R contains a nontrivial idempotent and the characteristic of R is not 2. Then D is of the form $d + t$, where d is an ordinary derivation of R into a primitive ring $\hat{R}$ containing R and t is an additive mapping of R into the center of $\hat{R}$ that maps commutators into zero.*

Martindale remarks after Theorem 4.6 that, if R is simple, then d maps R into itself and t maps R into the center of R [14, p. 187].

Definition 4.7 ([18, Section 9.2.1])**.** A finite-dimensional semisimple algebra R over a field k is called *separable* if for every extension field ℓ of k the ℓ-algebra $R_\ell = R \otimes_k \ell$ is semisimple.

The next theorem of Weibel characterizes separable algebras by the vanishing of Hochschild cohomology.

Theorem 4.8 ([18, Theorem 9.2.11])**.** *Let R be an algebra over a field k. The following are equivalent:*

(1) R is a finite-dimensional separable k-algebra.
(2) R is projective as a left R^e-module.
(3) $H_*(R, M) = 0$ for all $* \neq 0$ and all bimodules M.
(4) $H^*(R, M) = 0$ for all $* \neq 0$ and all bimodules M.

Here $R^e = R \otimes_k R^{\mathrm{op}}$, and H_* and H^* are Hochschild homology and cohomology.

We now apply Theorems 4.3, 4.5, 4.6 and 4.8 to $M_n(K)$.

Example 4.9 (Full matrix algebras)**.** Let $A = M_n(K)$ with $n \geq 2$, and let e_{ij} be the matrix unit with entry 1 in position (i, j) and 0 elsewhere. If $\operatorname{char} K \neq 2$, every Jordan derivation of $M_n(K)$ is a derivation by Theorem 4.3 applied to the ring K and the K-bimodule K, since K is then 2-torsion free. So $\mathrm{BSDer}_\tau(A) = \mathrm{JDer}(A) = \mathrm{Der}(A)$ in that case.

Let $\theta(x) = \operatorname{tr}(x)I$. Its values are central, and $\theta([x, y]) = (\operatorname{tr}(xy) - \operatorname{tr}(yx))I = 0$. By Proposition 4.2(iii), $\mathrm{ad}_g + c\,\theta$ is a Lie derivation for every $g \in A$ and $c \in K$. Since $\theta(e_{11}) = I$ while $\theta(e_{11})e_{11} + e_{11}\theta(e_{11}) = 2e_{11}$ and $\theta(e_{11}^2) = \theta(e_{11})$, the map θ is not a derivation in any characteristic.

Now let $n \geq 2$ and $\operatorname{char} K \neq 2$. The ring $M_n(K)$ is simple by Theorem 4.5, since a field is simple. It is primitive, because K^n is a faithful simple left module over it, and e_{11} is a nontrivial idempotent. By Theorem 4.6 and the remark after it, every Lie derivation D of $M_n(K)$ has the form $D = d + t$, where d is an additive derivation of $M_n(K)$ into itself and t is an additive map into $K \cdot I$ that vanishes on commutators.

The maps d and t are K-linear. Indeed, for $\lambda \in K$ we have $d(\lambda x) = d(\lambda I)x + \lambda d(x)$, and $d(\lambda I)$ is central because $[d(\lambda I), x] = d([\lambda I, x]) - [\lambda I, d(x)] = 0$. So $d(\lambda I) = \nu(\lambda) I$ for some $\nu(\lambda) \in K$. Since D is K-linear, $\nu(\lambda)x = \lambda t(x) - t(\lambda x)$ lies in $K \cdot I$ for every x. Taking $x = e_{11}$ gives $\nu(\lambda) = 0$.

Next, $M_n(K)$ is finite-dimensional, and $M_n(K) \otimes_K F = M_n(F)$ is simple for every extension field F by Theorem 4.5, hence semisimple. So $M_n(K)$ is separable by Definition 4.7. By Theorem 4.8, $H^1(M_n(K), M_n(K)) = 0$, so by Lemma 4.4 every K-linear derivation is inner, and d is inner. Hence $\mathrm{Der}(A) = \mathrm{Inn}(A)$, and every Lie derivation of A has the form $\mathrm{ad}_g + t$ with $t\colon A \to K \cdot I$ linear and $t([A, A]) = 0$. Such a map t is the same as a K-linear map $A/[A, A] \to K$, and $A/[A, A] = HH_0(A, A)$ [18, §9.1]. A map t of this kind that is also a derivation is zero. Indeed $t(e_{ii}) = t(e_{ii}^2) = 2t(e_{ii})e_{ii}$ forces $t(e_{ii}) = 0$ when $n \geq 2$, and $t(e_{ij}) = 0$ for $i \neq j$ because $e_{ij} = [e_{ii}, e_{ij}]$. So t is determined by $\mathrm{ad}_g + t$: if $\mathrm{ad}_g + t = \mathrm{ad}_h + t'$, then $t - t' = \mathrm{ad}_{h-g}$ is such a derivation, hence zero. Let $\omega\colon \mathrm{LDer}(A) \to \mathrm{Hom}_K(HH_0(A, A), K)$ send $\mathrm{ad}_g + t$ to t. The map ω is surjective, since each t is a Lie derivation by Proposition 4.2(iii), and its kernel is $\mathrm{Inn}(A) = \mathrm{Der}(A)$. So the sequence

$$0 \longrightarrow \mathrm{Der}(A) \longrightarrow \mathrm{LDer}(A) \xrightarrow{\ \omega\ } \mathrm{Hom}_K\big(HH_0(A, A), K\big) \longrightarrow 0$$

is exact. The map θ above is one element of the right-hand space.

The swap gives $\mathrm{JDer}(A)$ through $\mu + \mu \circ \tau$ and $\mathrm{LDer}(A)$ through $\mu - \mu \circ \tau$. So μ is half the sum of these two products when 2 is invertible. So when 2 is invertible, a map that is both a Jordan and a Lie derivation is a derivation. On $M_n(K)$ with $\mathrm{char}\, K \neq 2$ this gives $\mathrm{JDer}(A) \cap \mathrm{LDer}(A) = \mathrm{Der}(A)$ even though $\mathrm{LDer}(A)$ is larger.

For triangular algebras the Lie side has a central term.

A Lie derivation is *proper* when it is the sum of a derivation and a linear map into the centre [6, §1]. The extra map vanishes on commutators, so a proper Lie derivation has the standard form of Section 1.

The next theorem uses the algebra $\mathfrak{A} = \mathrm{Tri}(A, M, B)$, and the projections π_A and π_B onto the diagonal blocks.

The algebra $W(X)$ is the smallest subalgebra of X that satisfies seven closure conditions of Cheung [6, Proposition 10]. We list them for a subalgebra V of X. Here F is a polynomial in one variable over the base ring, and F' is its derivative.

(i) V contains all commutators $[x, y]$ with $x, y \in X$.
(ii) If $b \in X$ and $F'(b) = 0$, then $F(b) \in V$.
(iii) If $b \in X$, $F(b) \in V$ and $F'(b)$ is invertible, then $b \in V$.
(iv) V contains all idempotents of X.
(v) V contains all elements x^p, where $x \in X$ and $p \geq 0$ is the characteristic of X.
(vi) If $a \in X$ has an inverse a^{-1} in X and $a^{-1} \in V$, then $a \in V$.
(vii) If $b \in X$ is invertible and $b^k \in V$ for some positive integer k, then $kb \in V$.

Theorem 4.10 ([6, Theorem 11])**.** *Let $\mathfrak{A} = \mathrm{Tri}(A, M, B)$. If the following two conditions hold, then every Lie derivation of $\mathfrak{A}$ is proper.*

(I) M is a faithful left A-module and $Z(B) = \pi_B(Z(\mathfrak{A}))$; or M is a faithful left A-module and $A = W(A)$; or every Lie derivation of A is proper and $A = W(A)$.
(II) M is a faithful right B-module and $Z(A) = \pi_A(Z(\mathfrak{A}))$; or M is a faithful right B-module and $B = W(B)$; or every Lie derivation of B is proper and $B = W(B)$.

The next corollary applies the theorem of Cheung to the σ-antisymmetrized derivations of T.

Corollary 4.11. *Let $T = \mathrm{Tri}(A, M, B)$ satisfy conditions (I) and (II) of Theorem 4.10. Then every σ-antisymmetric derivation of T for τ has the standard form $d + t$, and it is a derivation only if $t(1) = 0$.*

Proof. By Proposition 4.2(i), the maps in question are the Lie derivations of T, which are proper by Theorem 4.10. The last claim is Proposition 4.2(iv). □

Corollary 2.5 says that on such a triangular algebra the symmetrized product gives only derivations. Example 4.12 shows that the antisymmetrized product can add a central term. Take $T_2(K) = \mathrm{Tri}(K, K, K)$ over a field with $\mathrm{char}\, K \neq 2$. Here $M = K$ is faithful on both sides. Also $Z(T_2(K)) = KI$ projects onto K. So Corollaries 2.5 and 4.11 apply.

Example 4.12 (Upper triangular 2×2 matrices)**.** Let $T = T_2(\mathbb{F}_3)$. Every additive map $T \to T$ is $\mathbb{F}_3$-linear. So it is one of the 3^9 maps given by a 3×3 matrix over $\mathbb{F}_3$.

An exhaustive computer check over all of them confirms Corollaries 2.5 and 4.11. Every Jordan derivation is a derivation. Every Lie derivation equals $\mathrm{ad}_x + t$ for some $x \in T$ and some central-valued t that vanishes on commutators.

The term t is needed. Define θ by

$$\theta \begin{pmatrix} a & m \\ 0 & b \end{pmatrix} = aI.$$

It has central values and vanishes on commutators, whose diagonal is zero. So it is a Lie derivation. It is not a derivation, since

$$\theta(e_1^2) = I \qquad \text{while} \qquad \theta(e_1)e_1 + e_1\theta(e_1) = 2e_1.$$

So on $T_2(\mathbb{F}_3)$, as on $M_n(K)$, we have JDer = Der, while LDer is strictly larger.

References

[1] Adrabi, A.: Braided-symmetric derivations of algebras. Preprint (2026). `https://github.com/abderrahim-lectures/braided-sym-deriv`.

[2] Alizadeh, R.: Jordan derivations of full matrix algebras. Linear Algebra Appl. **430**, 574–578 (2009).

[3] Brešar, M.: Jordan derivations on semiprime rings. Proc. Amer. Math. Soc. **104**(4), 1003–1006 (1988).

[4] Brešar, M., Zelmanov, E.: Jordan homomorphisms: a survey. Serdica Math. J. **51**, 203–230 (2025).

[5] Chambert-Loir, A.: (Mostly) Commutative Algebra. Universitext. Springer, Cham (2021).

[6] Cheung, W.-S.: Lie derivations of triangular algebras. Linear Multilinear Algebra **51**(3), 299–310 (2003).

[7] Cusack, J.M.: Jordan derivations on rings. Proc. Amer. Math. Soc. **53**(2), 321–324 (1975).

[8] Herstein, I.N.: Jordan derivations of prime rings. Proc. Amer. Math. Soc. **8**(6), 1104–1110 (1957).

[9] Herstein, I.N.: Lie and Jordan structures in simple, associative rings. Bull. Amer. Math. Soc. **67**, 517–531 (1961).

[10] Hochschild, G.: On the cohomology groups of an associative algebra. Ann. of Math. **46**(1), 58–67 (1945).
[11] Jacobson, N.: Abstract derivation and Lie algebras. Trans. Amer. Math. Soc. **42**(2), 206–224 (1937).
[12] Jordan, P., von Neumann, J., Wigner, E.: On an algebraic generalization of the quantum mechanical formalism. Ann. of Math. **35**(1), 29–64 (1934).
[13] Lam, T.Y.: A First Course in Noncommutative Rings, 2nd edn. Graduate Texts in Mathematics 131. Springer, New York (2001).
[14] Martindale, W.S., III: Lie derivations of primitive rings. Michigan Math. J. **11**(2), 183–187 (1964).
[15] Rotman, J.J.: An Introduction to Homological Algebra, 2nd edn. Universitext, Springer, New York (2009).
[16] Zhang, J.-H., Yu, W.-Y.: Jordan derivations of triangular algebras. Linear Algebra Appl. **419**(1), 251–255 (2006).
[17] Wang, Y.: Lie (Jordan) derivations of arbitrary triangular algebras. Aequationes Math. **93**, 1221–1229 (2019).
[18] Weibel, C.A.: An Introduction to Homological Algebra. Cambridge Stud. Adv. Math. 38, Cambridge University Press, Cambridge (1994).