

A field-measurable regularized-Coulomb sliding law: the $s_N(N)$ master curve, an N_c inversion, and a tidal-admittance ungrounding early-warning

Abdelrahman Saifelden¹

¹Alexandria Higher Institute of Engineering and Technology (AIET), Alexandria, Egypt

Abstract. Ice-stream sliding weakens as the bed approaches flotation, a behaviour large-scale models impose by hand (Joughin, Smith & Schoof 2019 add an ad-hoc near-flotation ramp to a regularized-Coulomb law). We show this weakening is a **closed-form pole** of the regularized-Coulomb law itself and, crucially, that it is **measurable from surface velocity alone**. Solving $\tau_b = CN(u/(u+u_0))^{1/m}$ at the driving stress gives, exactly, the sliding-sensitivity master curve $|s_N|(N) = m/(1-(N_c/N)^m)$ with $N_c = \tau_a/C$ — a simple pole at the flotation threshold N_c , verified to 1.4×10^{-4} against the numeric s_N . The curve makes three independent surface observables read the **same** law: a lake-drainage step ($du/u = |s_N| \cdot dN/N$), a continuous ocean-thermal-forcing gating slope, and the **tidal admittance** of velocity (whose $2f/1f$ harmonic ratio diverges toward N_c). A drainage-step population inverts for N_c to **0.2/0.4/1.0 %** at 5/10/20 % amplitude noise; the tidal probe recovers m and the dimensionless flotation proximity $R = (N_c/N)^m$ from velocity alone (answering the “no knowledge of N ” objection), with the tidal forcing amplitude *measured*, not assumed (CATS2008 across the BedMachine grounding zone: median tidal pressure swing 0.30 % of overburden, rising to 0.45 % within 2–4 km of the line). Because the basal-drag stiffness $\partial\tau_b/\partial u \propto (1-R)^2/R \rightarrow 0$ at N_c , an ice stream nearing ungrounding should show rising variance and lag-1 autocorrelation of surface speed; we **operationalize** the established critical-slowness early-warning (Scheffer et al. 2009; Dakos et al. 2008; Rosier et al. 2021) as the continuous reading $R(t) = (N_c/N)^m \rightarrow 1$ from tidal admittance. Read through the same lens, three real MacAyeal-lake drainages sit far from the N_c pole and show no critical-slowness signature — a correct **true negative** on stable ice (the detectors fire on planted surges/CSD). The contribution is the field-measurable law and its inversion, not a new early-warning concept; absolute s_N remains uncalibrated and the strong test awaits a stream actually nearing flotation.

1 Introduction

The basal sliding law sets how fast an ice stream flows for a given driving stress and effective pressure $N = p_i - p_w$ (Schoof 2005; Gagliardini et al. 2007). Near the grounding line $N \rightarrow 0$ and sliding becomes Coulomb-limited; regularized-Coulomb laws (Joughin, Smith & Schoof 2019; Tulaczyk et al. 2000; Gudmundsson 2011) capture this but introduce the near-flotation weakening through an *ad hoc* ramp and a fixed velocity scale u_0 . The difficulty is that N is rarely known: it depends on the unobserved subglacial water pressure.

The tools we use, not claim, are the regularized-Coulomb friction law (Schoof 2005; Joughin et al. 2019), the nonlinear tidal
 25 (MSf) velocity response of ice streams (Gudmundsson 2011), and critical-slowness-down early-warning theory, *including its*
ice-sheet application (Scheffer et al. 2009; Dakos et al. 2008; Rosier et al. 2021; Boers & Rypdal 2021).

What is new, stated against the closest prior work. Joughin, Smith & Schoof (2019) *impose* near-flotation weakening with
 a hand-tuned ramp; we *derive* it as a closed-form pole $|s_N| = m/(1 - (N_c/N)^m)$ of the regularized-Coulomb law and turn
 N_c into a **measured** quantity. Rosier et al. (2021) established critical-slowness-down indicators (lag-1 autocorrelation, variance,
 30 Kendall- τ) for grounding-line flux ahead of marine-ice-sheet tipping; we do not re-claim the early-warning but give a specific
operational route — reading $R(t) = (N_c/N)^m \rightarrow 1$ from the continuous tidal admittance of *surface velocity*, which needs no
 knowledge of N .

2 The $s_N(N)$ master curve

2.1 $s_N(N)$ as the common observable. A lake drainage is an in-situ effective- pressure *step* whose surge amplitude $du/u =$
 35 $|s_N| \cdot (dN/N)$ measures the sliding sensitivity $s_N = d\ln u_b / d\ln N$; ocean thermal forcing lowers N *continuously*, so the gating
 slope $d\ln u_*/dT$ measures the **same** s_N ($+0.46 \rightarrow +0.62/^\circ C$, direct BedMachine N); ocean tides oscillate N at known
 amplitude, a third probe (§4). All three read one curve.

2.2 The closed-form master curve. Solving the regularized-Coulomb law $\tau_b = CN(u/(u + u_0))^{1/m}$ at $\tau_b = \tau_d$ gives, ex-
 40 to 1.4×10^{-4} against the numeric s_N . This *derives* the near-flotation weakening that Joughin, Smith & Schoof (2019) impose
 by hand (ad hoc $h_{af} < h_T$ ramp + fixed u_0). (validation/synthetic/sn_master_curve.py)

3 Inversion — measuring N_c and m

A drainage-step population recovers the flotation threshold N_c to **0.2 / 0.4 / 1.0 %** at 5 / 10 / 20 % amplitude noise; m is
 degenerate with the per-event drop far from flotation. The recovery is unbiased and sign-faithful under noise and null under
 45 permutation (synthetic plant-and-recover).

4 The tidal third probe — N_c , m , and R from velocity alone

Ocean **tides** give a continuous probe: the fundamental velocity admittance equals $|s_N|$ and the 2f/1f harmonic ratio diverges
 toward N_c , so admittance + harmonics + the known tidal amplitude recover m and the dimensionless flotation proximity
 $R = (N_c/N)^m$ **from surface velocity alone** — a sliding-law reading of the Gudmundsson nonlinear MSf response, answering
 50 the “no knowledge of N ” objection. The tidal forcing amplitude is **measured**, not assumed: sampling the CATS2008 tide
 model (USAP-DC 601235) across the BedMachine grounding zone (within 10 km of the line; $n = 1108$ valid cells) gives a tidal
 ocean-pressure swing with median **0.30 % of ice overburden** (median tide $\eta \approx 0.84m$), rising monotonically from 0.21 % at
 8–10 km to 0.45 % at 2–4 km from the grounding line (validation/synthetic/tidal_admittance_probe.py,

validation/reports/tidal_forcing_gz.json). **Contamination robustness (synthetic).** The two known non-sliding contributors at tidal frequencies — flexure leakage and tidally induced grounding-line migration, the mechanism Rosier & Gudmundsson (2020) identify as the only one strong enough to carry the observed MSf response, with Minchew et al. (2017) placing the strongest signal at the grounding zone — are quantified and screened rather than assumed away: a memoryless sliding response to a cosine tide is a Chebyshev expansion, so *every* velocity harmonic is phase-locked to the tide ($\phi_k \in \{0, \pi\}$; clean synthetic closure residual 10^{-15} rad). Lagged GL-migration rectification injects a 2f component that can fake flotation proximity (+46% bias on R at 2% modulation) but breaks 2f phase closure by 0.2–0.9 rad — flagged and screened before inversion; lagged (viscoelastic) flexure is flagged the same way at the fundamental, while exactly in-phase flexure is phase-invisible (20% R bias at 2% leakage) and is capped by station placement upstream of the flexure zone. Red (AR-1) noise from concurrent forcing at 1% of $\ln u$ over a 40-cycle record leaves ± 0.03 (68%) scatter on $R \approx 0.22$. (validation/synthetic/tidal_admittance_contamination.py.)

5 An ungrounding early-warning (operationalizing established CSD theory)

The N_c fold makes the basal drag stiffness $\partial\tau_b/\partial u \propto (1-R)^2/R \rightarrow 0$ (critical slowing down), so an ice stream approaching ungrounding should show **rising variance and lag-1 autocorrelation of its surface speed** (Kendall- $\tau \approx 0.54$ in an OU demonstration with N declining toward N_c). Critical-slowness early-warning is established (Scheffer et al. 2009; Dakos et al. 2008) and has already been applied to marine-ice-sheet tipping — Rosier et al. (2021) detected rising lag-1 autocorrelation and variance in grounding-line flux ahead of Pine Island tipping, and Boers & Rypdal (2021) in Greenland surface melt. The contribution here is **not** the early-warning concept but a specific **operational route**: reading $R(t) = (N_c/N)^m \rightarrow 1$ from the continuous tidal admittance of surface velocity. A single-snapshot **spatial** analogue (variance + along-flow correlation length rising toward the grounding line) is also derived (exact Lyapunov solve). **The discriminating prediction** is stated for the next near-flotation event: a stream approaching ungrounding must show the 2f/1f admittance ratio rising along the pole law $(\varepsilon/4)mR/(1-R) + \dots \rightarrow \infty$ *together* with rising variance and lag-1 autocorrelation, with the phase-closure screen (§above) clean; observing the CSD indicators *without* the admittance divergence (or vice versa) falsifies the sliding-law reading. The natural watch list is the Amundsen sector (Thwaites and Pine Island grounding zones, where Rosier et al. 2021 place the tipping indicators and BedMachine puts the bed closest to flotation). (validation/synthetic/spatial_ews.py)

The fold exponents and a calibration-free spectral gauge (derived). The warning here is not generic: the s_N pole is the Schoof (2007) grounding-line saddle-node, so the critical-slowness exponents are fixed, not fitted. Near $N = N_c$ the restoring eigenvalue $\lambda \propto (1-R)^2/R \sim (N - N_c)^2$, giving $|s_N| \propto (N - N_c)^{-1}$, variance $\propto (N - N_c)^{-2}$ and $AC1 \rightarrow 1$ (recovered exponents $-1.00, -2.00, +2.00$). The same rate has a frequency-domain face: the velocity spectrum is a Lorentzian $S(\omega) = 2D/(\lambda^2 + \omega^2)$ whose corner $f_c = \lambda/2\pi \propto (N - N_c)^2$ *reddens* toward ungrounding (the spectral early-warning of Kuehn 2011; Bury et al. 2020). Crucially the fluctuation–dissipation identity $\text{Var} \cdot 2\pi f_c = D$ holds independent of N , so the noise corner f_c is a **calibration-free** flotation-proximity gauge (it falls as $(N - N_c)^2$ regardless of the unknown noise strength D); combined with the tidal-admittance *response* (§4) it even returns D itself — a response/fluctuation pair from one velocity

record. (Derived consistency relations, verified on OU realizations: corner recovered to 3%, FDT product constant to 5×10^{-17} ; nr3_misi_fold_ews, nr8_spectral_fdt_ews.)

6 The framework on the same real lakes

90 Reading open CryoSat-2 drainage + ITS_LIVE velocity (3 MacAyeal lakes) through this lens (validation/external/
lake_lag_sn_ews.py, 5 tests): across the 3 resolved drainage events $|\Delta v/v| \leq 2\%$, mixed sign, near the $\sim 1\%$ year-to-
year noise floor — **0 positive $>2\sigma$ surge detections** — so via the master curve these trunk lakes sit **far from the N_c pole**; and
no lake shows the joint critical-slowness signature (rising variance *and* high AC1), so the early-warning correctly returns
a **true negative** on stable ice (unit tests confirm the detectors *do* fire on planted surges/CSD, so the null is real). Both probes
95 agree: far from ungrounding.

7 Estimator calibration, scope, and falsification

7.1 Calibration. Every estimator has a synthetic plant-and-recover harness in `pytest`: the master-curve closed form (1.4×10^{-4}), the N_c inversion (0.2/0.4/1.0 %), the tidal admittance/harmonic recovery, and the CSD detectors (fire on planted surges/CSD, null on stable OU).

100 **7.2 Scope and falsification.** Absolute s_N is uncalibrated (per-event dN/N is a lumped-storage lower bound); the robust claims are the sign/shape and the *threshold* N_c . With $n \approx 8$ annual points per lake the CSD test is weak; a strong test needs the dense quarterly/GPS series and a stream actually nearing flotation. Falsifiers: a co-located dN +GPS population whose $|s_N(N)|$ does not follow $m/(1 - (N_c/N)^m)$; a stream nearing flotation with no variance/AC1 rise; tidal admittance/harmonics not steepening toward the grounding line.

105 8 Reproduce

```
pip install -r requirements.txt
python glaciers/validation/synthetic/sn_master_curve.py           # $2 s_N(N) master curve + in
python glaciers/validation/synthetic/tidal_admittance_probe.py    # $4 tidal third probe
python glaciers/validation/synthetic/spatial_ews.py               # $5 spatial early-warning
110 python glaciers/validation/external/lake_lag_sn_ews.py         # $6 framework on real lakes
pytest glaciers/tests/test_validation_synthetic.py -v             # $7 estimator calibration
```

Code and data availability. All analysis code and derived products are in the public repository at <https://github.com/abdh0saifelden2-debug/K>, and the results regenerate from the scripts above. The study uses the following open datasets:

- BedMachine Antarctica v4 (Morlighem, 2025, <https://doi.org/10.5067/POJQI54A45HX>; see Morlighem et al., 2020).

- 115 – CATS2008 circum-Antarctic tide model: USAP-DC 601235 (Howard et al., 2019, <https://doi.org/10.15784/601235>).
- ITS_LIVE surface velocities (Gardner et al., 2025, <https://doi.org/10.5067/JQ6337239C96>; see Gardner et al., 2018) and CryoSat-2 lake-drainage dates.

 BedMachine and ITS_LIVE were obtained from the NASA NSIDC DAAC; CATS2008 from the U.S. Antarctic Program Data Center (<https://www.usap-dc.org>).

120 *Author contributions.* A. Saifelden conceived the study, performed all computations and analysis, and wrote the manuscript.

Competing interests. The author declares that there are no competing interests.

References

- Boers, N., & Rypdal, M. (2021). Critical slowing down suggests that the western Greenland Ice Sheet is close to a tipping point. *Proceedings of the National Academy of Sciences* 118(21), e2024192118. doi:10.1073/pnas.2024192118
- 125 Dakos, V., Scheffer, M., van Nes, E. H., Brovkin, V., Petoukhov, V., & Held, H. (2008). Slowing down as an early warning signal for abrupt climate change. *Proceedings of the National Academy of Sciences* 105(38), 14308–14312. doi:10.1073/pnas.0802430105
- Gagliardini, O., Cohen, D., Råback, P., & Zwinger, T. (2007). Finite-element modeling of subglacial cavities and related friction law. *Journal of Geophysical Research: Earth Surface* 112, F02027. doi:10.1029/2006JF000576
- Gardner, A. S., Moholdt, G., Scambos, T., Fahnestock, M., Ligtenberg, S., van den Broeke, M., & Nilsson, J. (2018). Increased West Antarctic and unchanged East Antarctic ice discharge over the last 7 years. *The Cryosphere* 12, 521–547. doi:10.5194/tc-12-521-2018
- 130 Gardner, A., Fahnestock, M., Greene, C. A., Kennedy, J. H., Liukis, M., Lopez, L., & Scambos, T. (2025). MEaSURES ITS_LIVE Regional Glacier and Ice Sheet Surface Velocities, Version 2 (NSIDC-0776) [Data set]. *NASA NSIDC DAAC*. doi:10.5067/JQ6337239C96
- Gudmundsson, G. H. (2011). Ice-stream response to ocean tides and the form of the basal sliding law. *The Cryosphere* 5, 259–270. doi:10.5194/tc-5-259-2011
- 135 Howard, S. L., Erofeeva, S., & Padman, L. (2019). CATS2008: Circum-Antarctic Tidal Simulation version 2008. *U.S. Antarctic Program Data Center*. doi:10.15784/601235
- Joughin, I., Smith, B. E., & Schoof, C. G. (2019). Regularized Coulomb friction laws for ice sheet sliding: Application to Pine Island Glacier, Antarctica. *Geophysical Research Letters* 46, 4764–4771. doi:10.1029/2019GL082526
- Minchew, B. M., Simons, M., Riel, B., & Milillo, P. (2017). Tidally induced variations in vertical and horizontal motion on Rutford Ice Stream, West Antarctica, inferred from remotely sensed observations. *Journal of Geophysical Research: Earth Surface* 122(1), 167–190. doi:10.1002/2016JF003971
- 140 Morlighem, M., Rignot, E., Binder, T., Blankenship, D. D., Drews, R., Eagles, G., et al. (2020). Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet. *Nature Geoscience* 13, 132–137. doi:10.1038/s41561-019-0510-8
- Morlighem, M. (2025). MEaSURES BedMachine Antarctica, Version 4 (NSIDC-0756) [Data set]. *NASA National Snow and Ice Data Center DAAC*. doi:10.5067/POJQI54A45HX
- 145 Padman, L., Fricker, H. A., Coleman, R., Howard, S., & Erofeeva, L. (2002). A new tide model for the Antarctic ice shelves and seas. *Annals of Glaciology* 34, 247–254. doi:10.3189/172756402781817752
- Rosier, S. H. R., & Gudmundsson, G. H. (2020). Exploring mechanisms responsible for tidal modulation in flow of the Filchner–Ronne Ice Shelf. *The Cryosphere* 14(1), 17–37. doi:10.5194/tc-14-17-2020
- 150 Rosier, S. H. R., Reese, R., Donges, J. F., De Rydt, J., Gudmundsson, G. H., & Winkelmann, R. (2021). The tipping points and early warning indicators for Pine Island Glacier, West Antarctica. *The Cryosphere* 15, 1501–1516. doi:10.5194/tc-15-1501-2021
- Scheffer, M., Bascompte, J., Brock, W. A., Brovkin, V., Carpenter, S. R., Dakos, V., Held, H., van Nes, E. H., Rietkerk, M., & Sugihara, G. (2009). Early-warning signals for critical transitions. *Nature* 461, 53–59. doi:10.1038/nature08227
- Schoof, C. (2005). The effect of cavitation on glacier sliding. *Proceedings of the Royal Society A* 461, 609–627. doi:10.1098/rspa.2004.1350
- 155 Siegfried, M. R., & Fricker, H. A. (2021). Illuminating active subglacial lake processes with ICESat-2 laser altimetry. *Geophysical Research Letters* 48, e2020GL091089. doi:10.1029/2020GL091089
- Tulaczyk, S., Kamb, W. B., & Engelhardt, H. F. (2000). Basal mechanics of Ice Stream B, West Antarctica: 1. Till mechanics. *Journal of Geophysical Research: Solid Earth* 105(B1), 463–481. doi:10.1029/1999JB900329