

Outline today

- Why probabilistic modeling?
- Basics of Probability
 - Random variable
 - Probability distribution
 - Dependent and independent variables
 - Correlation vs. Causation
- Bayes rule
- Probabilistic Graphical Models
- Probabilistic Robotics

Random Variable

- X is a random variable if it can take different values with different probabilities and can be observed with an experiment

Random Variable

- X is a random variable if it can take different values with different probabilities and can be observed with an experiment
- X can be continuous or discrete
 - Outcome of a dice → Discrete random variable
 - Height of adult population of a country → Continuous random variable

Random Variable


- X is a random variable if it can take different values with different probabilities and can be observed with an experiment
- X can be continuous or discrete
 - Outcome of a dice → Discrete random variable
 - Height of adult population of a country → Continuous random variable
- Sample space: all possible values that a random variable can take
 - $X = \text{outcome of a dice} \rightarrow S = \{1, 2, 3, 4, 5, 6\} \rightarrow \text{Discrete}$
 - $X = \text{Height of adult population} \rightarrow S = [1 \text{ foot}, 9 \text{ feet}] \rightarrow \text{Continuous}$

Random Variable

- X is a random variable if it can take different values with different probabilities and can be observed with an experiment
- X can be continuous or discrete
 - Outcome of a dice → Discrete random variable
 - Height of adult population of a country → Continuous random variable
- Sample space: all possible values that a random variable can take
 - X=outcome of a dice → $S=\{1,2,3,4,5,6\}$ → Discrete
 - X=Height of adult population → $S=[1 \text{ feet}, 9 \text{ feet}]$ → Continuous
- $P(X=x)$: Probability of an event (or probability of X taking the value x)
 - $P(X=x) \leq 1$
 - $\sum_x P(X = x) = 1$



Probability distribution

- Function $P(X=x)$ (or directly $P(x)$)
 - When we plot it:
 - x axis: the sample space
 - y axis: the probability of each sample, $P(x)$






Probability distribution

- Function $P(X=x)$ (or directly $P(x)$)
 - When we plot it:
 - x axis: the sample space
 - y axis: the probability of each sample, $P(x)$



Probability distribution

- Function $P(X=x)$ (or directly $P(x)$)
 - When we plot it:
 - x axis: the sample space
 - y axis: the probability of each sample, $P(x)$




Mean of a Random Variable

- “Mean” is also called “Expected Value”
- The mean value depends on:
 - The numeric value of each sample
 - The probability of each sample
- Gives you the value that is most likely to happen if we repeat the experiment
- Formally: $E(X) = \sum_x x \cdot P(x)$

Mean of a Random Variable


- $E(X) = \sum_x x \cdot P(x)$




$$E(x) = \sum_x x \cdot P(x) = 1 \cdot \frac{1}{6} + 2 \cdot \frac{1}{6} + 3 \cdot \frac{1}{6} + 4 \cdot \frac{1}{6} + 5 \cdot \frac{1}{6} + 6 \cdot \frac{1}{6} = 3.5$$

Mean of a Random Variable

- $E(X) = \sum_x x \cdot P(x)$




$$E(x) = \sum_x x \cdot P(x) = 1 \cdot \frac{1}{6} + 2 \cdot \frac{1}{6} + 3 \cdot \frac{1}{6} + 4 \cdot \frac{1}{6} + 5 \cdot \frac{1}{6} + 6 \cdot \frac{1}{6} = 3.5$$



$$E(x) = \sum_x x \cdot P(x) = 1 \cdot 0.5 + 2 \cdot 0.1 + 3 \cdot 0.1 + 4 \cdot 0.1 + 5 \cdot 0.1 + 6 \cdot 0.1 = 2.5$$


Informed Decisions based on Probabilistic Modeling



Instapoll



Informed decisions based on probabilistic modeling




Mean: \$3.5

Law of Large Numbers

- The average of the outcome of an experiment repeated many times tends to the expected value (mean)
- It gets closer, the more we repeat the experiment

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{X_i}{n} = E(X)$$



Dependent and Independent Variables

- Let's assume X and Y are two random variables
 - $X \rightarrow$ outcome of a coin toss, $Y \rightarrow$ outcome of a dice roll
 - $X \rightarrow$ develop lung cancer, $Y \rightarrow$ be a smoker
- $P(X,Y)$ is called the joint probability
 - $P(X=\text{"tails"}, Y=\text{"6"})$
 - $P(X=\text{"develop lung cancer"}, Y=\text{"smoker"})$

Dependent and Independent Variables

- Let's assume X and Y are two random variables
 - $X \rightarrow$ outcome of a coin toss, $Y \rightarrow$ outcome of a dice toss
 - $X \rightarrow$ develop lung cancer, $Y \rightarrow$ be a smoker
- $P(X,Y)$ is called the joint probability
 - $P(X=\text{"tails"}, Y=\text{"6"})$
 - $P(X=\text{"develop lung cancer"}, Y=\text{"smoker"})$
- $P(X|Y)$ is called a conditional probability, probability of X conditioned on knowing Y
 - $P(X|Y=\text{"6"}) \rightarrow$ Probability of outcome of a coin toss given that the dice toss was "6"
 - $P(X|Y=\text{"smoker"}) \rightarrow$ Probability of developing lung cancer given that is a smoker

Dependent and Independent Variables

- X and Y are said to be independent variables if the distribution of X is not influenced by the value taken by Y and vice versa. In that case:
 - the probability of a joint event is the product of the probability of one event and the probability of the other
 - $P(X=\text{"tails"}, Y=\text{"6"}) = P(X=\text{"tails"}) \cdot P(Y=\text{"6"})$
 - the conditional probability of one knowing the other is the same as the original probability
 - $P(X \mid Y=\text{"6"}) = P(X)$

Dependent and Independent Variables

- X and Y are said to be independent variables if the distribution of X is not influenced by the value taken by Y and vice versa. In that case:
 - the probability of a joint event is the product of the probability of one event and the probability of the other
 - $P(X=\text{"tails"}, Y=\text{"6"}) = P(X=\text{"tails"}) \cdot P(Y=\text{"6"})$
 - the conditional probability of one knowing the other is the same as the original probability
 - $P(X \mid Y=\text{"6"}) = P(X)$
- X and Y are said to be dependent variables if knowing the value of one of the events affects the probability distribution over the other event. In that case:
 - $P(X \mid Y=\text{"smoker"}) \neq P(X)$

Correlation vs. Causality

- Correlation:
 - Measurement of “how much two random variables are dependent on each other”
 - Correlation = 0 \rightarrow Variables are independent
 - Correlation \neq 0 \rightarrow Variables are dependent
- Correlation is not causation
 - We can mathematically proof that two variables are correlated but that doesn't mean one is causing the other!
 - Careful! This is sometimes the door to “biases” and “prejudices”

Correlation vs. Causality

ICE CREAM SALES



SUNBURN

Correlation vs. Causality

ICE CREAM SALES



SUNBURN

Correlation vs. Causality

ICE CREAM SALES



SUNBURN

Correlation vs. Causality

ICE CREAM SALES



SUNBURN

Correlation vs. Causality

