

# Symplectic geometry and Toric Varieties

## Part I: A walk through Symplectic geometry

Abdullah Ahmed<sup>1</sup>

<sup>1</sup>Department of Mathematics  
Lahore University of Management Sciences, Pakistan  
Supervisor: Dr. Shaheen Nazir

LUMS, February 16, 2024

# Table of Contents

- 1 Preliminaries from Differential topology
- 2 Symplectic structure
- 3 Contact structure
- 4 Complex structure

# Table of Contents

1 Preliminaries from Differential topology

2 Symplectic structure

3 Contact structure

4 Complex structure

# Quick recap of definitions

- A  $C^\infty$  manifold  $M$  is a second countable, Hausdorff, locally Euclidean space with an atlas  $\mathfrak{A} = \{U_\alpha \subset M, V_\alpha \subset \mathbb{R}^n, \phi_\alpha : U_\alpha \rightarrow V_\alpha\}$  with  $\cup_\alpha U_\alpha = M$ ,  $\phi_\alpha$  are homeomorphisms, and  $\phi_\alpha^{-1} \circ \phi_\beta$  are smooth.

# Quick recap of definitions

- A  $C^\infty$  manifold  $M$  is a second countable, Hausdorff, locally Euclidean space with an atlas  $\mathfrak{A} = \{U_\alpha \subset M, V_\alpha \subset \mathbb{R}^n, \phi_\alpha : U_\alpha \rightarrow V_\alpha\}$  with  $\cup_\alpha U_\alpha = M$ ,  $\phi_\alpha$  are homeomorphisms, and  $\phi_\alpha^{-1} \circ \phi_\beta$  are smooth.
- The *tangent space*  $T_p M$  at a point  $p \in M$  is defined to be the vector space of all derivations at  $p$ . It can be identified with the basis  $\{\frac{\partial}{\partial x_i}\}_{i=1}^n$  where  $\{x_i\}_{i=1}^n$  are the coordinates at  $p$ . The dual vector space  $T_p^* M$  is called the *cotangent space* at  $p$ .

# Quick recap of definitions

- A  $C^\infty$  manifold  $M$  is a second countable, Hausdorff, locally Euclidean space with an atlas  $\mathfrak{A} = \{U_\alpha \subset M, V_\alpha \subset \mathbb{R}^n, \phi_\alpha : U_\alpha \rightarrow V_\alpha\}$  with  $\cup_\alpha U_\alpha = M$ ,  $\phi_\alpha$  are homeomorphisms, and  $\phi_\alpha^{-1} \circ \phi_\beta$  are smooth.
- The *tangent space*  $T_p M$  at a point  $p \in M$  is defined to be the vector space of all derivations at  $p$ . It can be identified with the basis  $\{\frac{\partial}{\partial x_i}\}_{i=1}^n$  where  $\{x_i\}_{i=1}^n$  are the coordinates at  $p$ . The dual vector space  $T_p^* M$  is called the *cotangent space* at  $p$ .
- The *tangent bundle*  $TM = \sqcup_p T_p M$  over an  $n$ -dimensional manifold  $M$ , is a  $2n$ -dimensional manifold with coordinates  $\{x_i, c_i\}_{i=1}^n$  where  $c_i$  are the coefficients of  $v \in T_{p=(x_1, \dots, x_n)} M$ . The *cotangent bundle*  $T^* M = \sqcup_p T_p^* M$  is constructed similarly.

# Quick recap of definitions

- A  $C^\infty$  manifold  $M$  is a second countable, Hausdorff, locally Euclidean space with an atlas  $\mathfrak{A} = \{U_\alpha \subset M, V_\alpha \subset \mathbb{R}^n, \phi_\alpha : U_\alpha \rightarrow V_\alpha\}$  with  $\cup_\alpha U_\alpha = M$ ,  $\phi_\alpha$  are homeomorphisms, and  $\phi_\alpha^{-1} \circ \phi_\beta$  are smooth.
- The *tangent space*  $T_p M$  at a point  $p \in M$  is defined to be the vector space of all derivations at  $p$ . It can be identified with the basis  $\{\frac{\partial}{\partial x_i}\}_{i=1}^n$  where  $\{x_i\}_{i=1}^n$  are the coordinates at  $p$ . The dual vector space  $T_p^* M$  is called the *cotangent space* at  $p$ .
- The *tangent bundle*  $TM = \sqcup_p T_p M$  over an  $n$ -dimensional manifold  $M$ , is a  $2n$ -dimensional manifold with coordinates  $\{x_i, c_i\}_{i=1}^n$  where  $c_i$  are the coefficients of  $v \in T_{p=(x_1, \dots, x_n)} M$ . The *cotangent bundle*  $T^* M = \sqcup_p T_p^* M$  is constructed similarly.
- A field of  $n$ -covectors on  $M$  is called a *differential  $n$ -form*. The collection is denoted by  $\Omega^n(M)$ . The exterior derivative  $d : \Omega^n(M) \rightarrow \Omega^{n+1}(M)$  is given by  $d(\sum_I f_I dx_I) = \sum_{I,j} \frac{\partial f_I}{\partial x_j} dx_j \wedge dx_I$ .

# Quick recap of definitions

- A  $C^\infty$  manifold  $M$  is a second countable, Hausdorff, locally Euclidean space with an atlas  $\mathfrak{A} = \{U_\alpha \subset M, V_\alpha \subset \mathbb{R}^n, \phi_\alpha : U_\alpha \rightarrow V_\alpha\}$  with  $\cup_\alpha U_\alpha = M$ ,  $\phi_\alpha$  are homeomorphisms, and  $\phi_\alpha^{-1} \circ \phi_\beta$  are smooth.
- The *tangent space*  $T_p M$  at a point  $p \in M$  is defined to be the vector space of all derivations at  $p$ . It can be identified with the basis  $\{\frac{\partial}{\partial x_i}\}_{i=1}^n$  where  $\{x_i\}_{i=1}^n$  are the coordinates at  $p$ . The dual vector space  $T_p^* M$  is called the *cotangent space* at  $p$ .
- The *tangent bundle*  $TM = \sqcup_p T_p M$  over an  $n$ -dimensional manifold  $M$ , is a  $2n$ -dimensional manifold with coordinates  $\{x_i, c_i\}_{i=1}^n$  where  $c_i$  are the coefficients of  $v \in T_{p=(x_1, \dots, x_n)} M$ . The *cotangent bundle*  $T^* M = \sqcup_p T_p^* M$  is constructed similarly.
- A field of  $n$ -covectors on  $M$  is called a *differential  $n$ -form*. The collection is denoted by  $\Omega^n(M)$ . The exterior derivative  $d : \Omega^n(M) \rightarrow \Omega^{n+1}(M)$  is given by  $d(\sum_I f_I dx_I) = \sum_{I,j} \frac{\partial f_I}{\partial x_j} dx_j \wedge dx_I$ .
- The  $k$ th de Rham cohomology of  $M$  is  $H^k(M) := \frac{\{\text{closed } k\text{-forms on } M\}}{\{\text{exact } k\text{-forms on } M\}}$ .



# Table of Contents

- 1 Preliminaries from Differential topology
- 2 Symplectic structure**
- 3 Contact structure
- 4 Complex structure


# Motivation from Classical Mechanics

The origin of the ideas of symplectic geometry can be traced back to the works of Lagrange and Hamilton in Classical Mechanics in the 19th century.

# Motivation from Classical Mechanics

The origin of the ideas of symplectic geometry can be traced back to the works of Lagrange and Hamilton in Classical Mechanics in the 19th century.


- Consider a point particle moving inside the surface of a sphere.



# Motivation from Classical Mechanics

The origin of the ideas of symplectic geometry can be traced back to the works of Lagrange and Hamilton in Classical Mechanics in the 19th century.


- Consider a point particle moving inside the surface of a sphere.



# Motivation from Classical Mechanics

The origin of the ideas of symplectic geometry can be traced back to the works of Lagrange and Hamilton in Classical Mechanics in the 19th century.

- Consider a point particle moving inside the surface of a sphere.




- Hamilton's equations and the laws of physics impose restrictions on the structure of the phase space.

# Motivation from Classical Mechanics

The origin of the ideas of symplectic geometry can be traced back to the works of Lagrange and Hamilton in Classical Mechanics in the 19th century.

- Consider a point particle moving inside the surface of a sphere.



- Hamilton's equations and the laws of physics impose restrictions on the structure of the phase space.
- This resulting structure is exactly equivalent to the symplectic structure.

# Symplectic Vector Space

## Theorem

*Let  $\Omega$  be a skew-symmetric bilinear form on a real vector space  $V$ . Then there exists a basis  $\{e_1, \dots, e_n, f_1, \dots, f_n, u_1, \dots, u_k\}$  of  $V$  such that:*

- (i)  $\Omega(e_i, e_j) = \Omega(f_i, f_j) = \Omega(u_i, \cdot) = 0$*
- (ii)  $\Omega(e_i, f_j) = \delta_{ij}$ .*

# Symplectic Vector Space

## Theorem

Let  $\Omega$  be a skew-symmetric bilinear form on a real vector space  $V$ . Then there exists a basis  $\{e_1, \dots, e_n, f_1, \dots, f_n, u_1, \dots, u_k\}$  of  $V$  such that:

- (i)  $\Omega(e_i, e_j) = \Omega(f_i, f_j) = \Omega(u_i, \cdot) = 0$
- (ii)  $\Omega(e_i, f_j) = \delta_{ij}$ .

## Definition

Let  $\Omega$  be a skew-symmetric bilinear form on a real vector space  $V$ . Then  $(V, \Omega)$  is called a symplectic vector space if and only if  $\text{span}\{u_i\}_{i=1}^k = \{0\}$  (in other words,  $\Omega$  is non-degenerate).



# Symplectic Vector Space

## Theorem

Let  $\Omega$  be a skew-symmetric bilinear form on a real vector space  $V$ . Then there exists a basis  $\{e_1, \dots, e_n, f_1, \dots, f_n, u_1, \dots, u_k\}$  of  $V$  such that:

- (i)  $\Omega(e_i, e_j) = \Omega(f_i, f_j) = \Omega(u_i, \cdot) = 0$
- (ii)  $\Omega(e_i, f_j) = \delta_{ij}$ .

## Definition

Let  $\Omega$  be a skew-symmetric bilinear form on a real vector space  $V$ . Then  $(V, \Omega)$  is called a symplectic vector space if and only if  $\text{span}\{u_i\}_{i=1}^k = \{0\}$  (in other words,  $\Omega$  is non-degenerate).

- In particular, all symplectic vector spaces are even-dimensional.

# Symplectic Vector Space

## Theorem

Let  $\Omega$  be a skew-symmetric bilinear form on a real vector space  $V$ . Then there exists a basis  $\{e_1, \dots, e_n, f_1, \dots, f_n, u_1, \dots, u_k\}$  of  $V$  such that:

- (i)  $\Omega(e_i, e_j) = \Omega(f_i, f_j) = \Omega(u_i, \cdot) = 0$
- (ii)  $\Omega(e_i, f_j) = \delta_{ij}$ .

## Definition

Let  $\Omega$  be a skew-symmetric bilinear form on a real vector space  $V$ . Then  $(V, \Omega)$  is called a symplectic vector space if and only if  $\text{span}\{u_i\}_{i=1}^k = \{0\}$  (in other words,  $\Omega$  is non-degenerate).

- In particular, all symplectic vector spaces are even-dimensional.
- Example: Any even-dimensional vector space  $V$  can be given a symplectic structure.

# Symplectic Manifolds

## Definition

*A manifold  $M$  is symplectic iff it is equipped with a differential 2-form  $\omega$  (called the symplectic form) such that:*

- (i)  $\omega$  is closed, i.e,  $d\omega = 0$ , and*
- (ii)  $\omega_p$  is non-degenerate for all  $p \in M$ .*

# Symplectic Manifolds

## Definition

*A manifold  $M$  is symplectic iff it is equipped with a differential 2-form  $\omega$  (called the symplectic form) such that:*

- (i)  $\omega$  is closed, i.e,  $d\omega = 0$ , and*
- (ii)  $\omega_p$  is non-degenerate for all  $p \in M$ .*

- In particular, (ii) is equivalent to  $M$  being even-dimensional with  $\omega^n$  as a volume form.

# Symplectic Manifolds

## Definition

*A manifold  $M$  is symplectic iff it is equipped with a differential 2-form  $\omega$  (called the symplectic form) such that:*

- (i)  $\omega$  is closed, i.e,  $d\omega = 0$ , and*
- (ii)  $\omega_p$  is non-degenerate for all  $p \in M$ .*

- In particular, (ii) is equivalent to  $M$  being even-dimensional with  $\omega^n$  as a volume form.
- Example:  $\mathbb{R}^{2n}$  with coordinates  $(x_1, \dots, x_n, y_1, \dots, y_n)$  has a symplectic form  $\omega = \sum_i dx_i \wedge dy_i$ .

# Symplectic Manifolds

## Definition

*A manifold  $M$  is symplectic iff it is equipped with a differential 2-form  $\omega$  (called the symplectic form) such that:*

- (i)  $\omega$  is closed, i.e,  $d\omega = 0$ , and*
- (ii)  $\omega_p$  is non-degenerate for all  $p \in M$ .*

- In particular, (ii) is equivalent to  $M$  being even-dimensional with  $\omega^n$  as a volume form.
- Example:  $\mathbb{R}^{2n}$  with coordinates  $(x_1, \dots, x_n, y_1, \dots, y_n)$  has a symplectic form  $\omega = \sum_i dx_i \wedge dy_i$ .
- Example: Every orientable surface has its area form as a symplectic form.

# Symplectic Manifolds

## Definition

*A manifold  $M$  is symplectic iff it is equipped with a differential 2-form  $\omega$  (called the symplectic form) such that:*

- (i)  $\omega$  is closed, i.e,  $d\omega = 0$ , and*
- (ii)  $\omega_p$  is non-degenerate for all  $p \in M$ .*

- In particular, (ii) is equivalent to  $M$  being even-dimensional with  $\omega^n$  as a volume form.
- Example:  $\mathbb{R}^{2n}$  with coordinates  $(x_1, \dots, x_n, y_1, \dots, y_n)$  has a symplectic form  $\omega = \sum_i dx_i \wedge dy_i$ .
- Example: Every orientable surface has its area form as a symplectic form.
- Does a symplectic form  $\omega$  always exist on an even-dimensional manifold?

# Symplectic Manifolds

## Definition

A manifold  $M$  is symplectic iff it is equipped with a differential 2-form  $\omega$  (called the symplectic form) such that:

- (i)  $\omega$  is closed, i.e,  $d\omega = 0$ , and
- (ii)  $\omega_p$  is non-degenerate for all  $p \in M$ .

- In particular, (ii) is equivalent to  $M$  being even-dimensional with  $\omega^n$  as a volume form.
- Example:  $\mathbb{R}^{2n}$  with coordinates  $(x_1, \dots, x_n, y_1, \dots, y_n)$  has a symplectic form  $\omega = \sum_i dx_i \wedge dy_i$ .
- Example: Every orientable surface has its area form as a symplectic form.
- Does a symplectic form  $\omega$  always exist on an even-dimensional manifold? **No, consider the Mobius strip.**



# Symplectic Manifolds

## Definition

*A manifold  $M$  is symplectic iff it is equipped with a differential 2-form  $\omega$  (called the symplectic form) such that:*

- (i)  $\omega$  is closed, i.e,  $d\omega = 0$ , and*
- (ii)  $\omega_p$  is non-degenerate for all  $p \in M$ .*

- In particular, (ii) is equivalent to  $M$  being even-dimensional with  $\omega^n$  as a volume form.
- Example:  $\mathbb{R}^{2n}$  with coordinates  $(x_1, \dots, x_n, y_1, \dots, y_n)$  has a symplectic form  $\omega = \sum_i dx_i \wedge dy_i$ .
- Example: Every orientable surface has its area form as a symplectic form.
- Does a symplectic form  $\omega$  always exist on an even-dimensional manifold? **No, consider the Mobius strip.**
- Does a symplectic form  $\omega$  always exists on even-dimensional orientable manifolds?

# Symplectic Manifolds

## Definition

A manifold  $M$  is symplectic iff it is equipped with a differential 2-form  $\omega$  (called the symplectic form) such that:

- (i)  $\omega$  is closed, i.e,  $d\omega = 0$ , and
- (ii)  $\omega_p$  is non-degenerate for all  $p \in M$ .

- In particular, (ii) is equivalent to  $M$  being even-dimensional with  $\omega^n$  as a volume form.
- Example:  $\mathbb{R}^{2n}$  with coordinates  $(x_1, \dots, x_n, y_1, \dots, y_n)$  has a symplectic form  $\omega = \sum_i dx_i \wedge dy_i$ .
- Example: Every orientable surface has its area form as a symplectic form.
- Does a symplectic form  $\omega$  always exist on an even-dimensional manifold? **No, consider the Mobius strip.**
- Does a symplectic form  $\omega$  always exists on even-dimensional orientable manifolds? **No, consider  $S^{2n}$  for  $n > 1$ .**

# Cotangent Bundle

Recall the *pullback map* on differential forms:

# Cotangent Bundle

Recall the *pullback map* on differential forms:

Let  $\phi : M \rightarrow N$  be smooth. Then the pullback  $\phi^* : \Omega^n(N) \rightarrow \Omega^n(M)$  is defined as  $(\phi^*\omega)_p(v_1, \dots, v_n) = \omega_{\phi(p)}(\phi_*v_1, \dots, \phi_*v_n)$ .

# Cotangent Bundle

Recall the *pullback map* on differential forms:

Let  $\phi : M \rightarrow N$  be smooth. Then the pullback  $\phi^* : \Omega^n(N) \rightarrow \Omega^n(M)$  is defined as  $(\phi^*\omega)_p(v_1, \dots, v_n) = \omega_{\phi(p)}(\phi_*v_1, \dots, \phi_*v_n)$ .

Considering the structure of the cotangent bundle:

# Cotangent Bundle

Recall the *pullback map* on differential forms:

Let  $\phi : M \rightarrow N$  be smooth. Then the pullback  $\phi^* : \Omega^n(N) \rightarrow \Omega^n(M)$  is defined as  $(\phi^*\omega)_p(v_1, \dots, v_n) = \omega_{\phi(p)}(\phi_*v_1, \dots, \phi_*v_n)$ .

Considering the structure of the cotangent bundle:

## Definition

*The Liouville 1-form  $\alpha$  is an intrinsic 1-form defined on  $T^*M$  given by  $\alpha_{(x,\eta)} = \pi^*\eta$ , where  $\pi : T^*M \rightarrow M$  is the canonical projection  $(x, \eta) \mapsto x$ .*

# Cotangent Bundle

Recall the *pullback map* on differential forms:

Let  $\phi : M \rightarrow N$  be smooth. Then the pullback  $\phi^* : \Omega^n(N) \rightarrow \Omega^n(M)$  is defined as  $(\phi^*\omega)_p(v_1, \dots, v_n) = \omega_{\phi(p)}(\phi_*v_1, \dots, \phi_*v_n)$ .

Considering the structure of the cotangent bundle:

## Definition

*The Liouville 1-form  $\alpha$  is an intrinsic 1-form defined on  $T^*M$  given by  $\alpha_{(x,\eta)} = \pi^*\eta$ , where  $\pi : T^*M \rightarrow M$  is the canonical projection  $(x, \eta) \mapsto x$ .*

- In local coordinates,  $\alpha = \sum_i \eta_i dx_i$ , with  $d\alpha = \sum_i dx_i \wedge d\eta_i$ .

# Cotangent Bundle

Recall the *pullback map* on differential forms:

Let  $\phi : M \rightarrow N$  be smooth. Then the pullback  $\phi^* : \Omega^n(N) \rightarrow \Omega^n(M)$  is defined as  $(\phi^*\omega)_p(v_1, \dots, v_n) = \omega_{\phi(p)}(\phi_*v_1, \dots, \phi_*v_n)$ .

Considering the structure of the cotangent bundle:

## Definition

*The Liouville 1-form  $\alpha$  is an intrinsic 1-form defined on  $T^*M$  given by  $\alpha_{(x,\eta)} = \pi^*\eta$ , where  $\pi : T^*M \rightarrow M$  is the canonical projection  $(x, \eta) \mapsto x$ .*

- In local coordinates,  $\alpha = \sum_i \eta_i dx_i$ , with  $d\alpha = \sum_i dx_i \wedge d\eta_i$ .
- $d^2\alpha = 0$  and  $(d\alpha)^n = (n!)dx_1 \wedge \dots \wedge dx_n \wedge d\eta_1 \wedge \dots \wedge d\eta_n$  is nowhere-vanishing.



# Cotangent Bundle

Recall the *pullback map* on differential forms:

Let  $\phi : M \rightarrow N$  be smooth. Then the pullback  $\phi^* : \Omega^n(N) \rightarrow \Omega^n(M)$  is defined as  $(\phi^*\omega)_p(v_1, \dots, v_n) = \omega_{\phi(p)}(\phi_*v_1, \dots, \phi_*v_n)$ .

Considering the structure of the cotangent bundle:

## Definition

*The Liouville 1-form  $\alpha$  is an intrinsic 1-form defined on  $T^*M$  given by  $\alpha_{(x,\eta)} = \pi^*\eta$ , where  $\pi : T^*M \rightarrow M$  is the canonical projection  $(x, \eta) \mapsto x$ .*

- In local coordinates,  $\alpha = \sum_i \eta_i dx_i$ , with  $d\alpha = \sum_i dx_i \wedge d\eta_i$ .
- $d^2\alpha = 0$  and  $(d\alpha)^n = (n!)dx_1 \wedge \dots \wedge dx_n \wedge d\eta_1 \wedge \dots \wedge d\eta_n$  is nowhere-vanishing.
- $\omega = d\alpha$  defines a natural symplectic structure on any cotangent bundle.

# Cotangent Bundle

Recall the *pullback map* on differential forms:

Let  $\phi : M \rightarrow N$  be smooth. Then the pullback  $\phi^* : \Omega^n(N) \rightarrow \Omega^n(M)$  is defined as  $(\phi^*\omega)_p(v_1, \dots, v_n) = \omega_{\phi(p)}(\phi_*v_1, \dots, \phi_*v_n)$ .

Considering the structure of the cotangent bundle:

## Definition

*The Liouville 1-form  $\alpha$  is an intrinsic 1-form defined on  $T^*M$  given by  $\alpha_{(x,\eta)} = \pi^*\eta$ , where  $\pi : T^*M \rightarrow M$  is the canonical projection  $(x, \eta) \mapsto x$ .*

- In local coordinates,  $\alpha = \sum_i \eta_i dx_i$ , with  $d\alpha = \sum_i dx_i \wedge d\eta_i$ .
- $d^2\alpha = 0$  and  $(d\alpha)^n = (n!)dx_1 \wedge \dots \wedge dx_n \wedge d\eta_1 \wedge \dots \wedge d\eta_n$  is nowhere-vanishing.
- $\omega = d\alpha$  defines a natural symplectic structure on any cotangent bundle.
- Single most important symplectic manifold.

# Symplectomorphisms

## Definition

*Two symplectic manifolds  $(M_1, \omega_1)$  and  $(M_2, \omega_2)$  are symplectomorphic if there exists a diffeomorphism  $\phi : M_1 \rightarrow M_2$  with  $\phi^* \omega_2 = \omega_1$ . The map  $\phi$  is called a symplectomorphism.*

# Symplectomorphisms

## Definition

*Two symplectic manifolds  $(M_1, \omega_1)$  and  $(M_2, \omega_2)$  are symplectomorphic if there exists a diffeomorphism  $\phi : M_1 \rightarrow M_2$  with  $\phi^*\omega_2 = \omega_1$ . The map  $\phi$  is called a symplectomorphism.*

- Example: Every diffeomorphism  $\phi : M \rightarrow N$  naturally lifts to a symplectomorphism  $\tilde{\phi} : (T^*M, d\alpha_1) \rightarrow (T^*N, d\alpha_2)$ , with  $\tilde{f}(x, \eta) = (f(x), (f^{-1})^*\eta)$ .

# Symplectomorphisms

## Definition

*Two symplectic manifolds  $(M_1, \omega_1)$  and  $(M_2, \omega_2)$  are symplectomorphic if there exists a diffeomorphism  $\phi : M_1 \rightarrow M_2$  with  $\phi^*\omega_2 = \omega_1$ . The map  $\phi$  is called a symplectomorphism.*

- Example: Every diffeomorphism  $\phi : M \rightarrow N$  naturally lifts to a symplectomorphism  $\tilde{\phi} : (T^*M, d\alpha_1) \rightarrow (T^*N, d\alpha_2)$ , with  $\tilde{\phi}(x, \eta) = (\phi(x), (\phi^{-1})^*\eta)$ .
- Natural question: How does one check for the existence of a symplectomorphism between arbitrary symplectic manifolds?

# Lagrangian submanifolds

## Definition

*Let  $(M, \omega)$  be a  $2n$ -dimensional symplectic manifold. An  $n$ -dimensional submanifold  $i : Y \hookrightarrow M$  with  $i^*\omega = \omega|_Y = 0$  is called a Lagrangian submanifold of  $M$ .*

# Lagrangian submanifolds

## Definition

*Let  $(M, \omega)$  be a  $2n$ -dimensional symplectic manifold. An  $n$ -dimensional submanifold  $i : Y \hookrightarrow M$  with  $i^*\omega = \omega|_Y = 0$  is called a Lagrangian submanifold of  $M$ .*

- Example:  $M$  embedded into  $T^*M$  as  $i(x) = (x, \mu_x)$  for a fixed  $\mu \in \Omega^1(M)$  is Lagrangian iff  $\mu$  is closed.

# Lagrangian submanifolds

## Definition

*Let  $(M, \omega)$  be a  $2n$ -dimensional symplectic manifold. An  $n$ -dimensional submanifold  $i : Y \hookrightarrow M$  with  $i^*\omega = \omega|_Y = 0$  is called a Lagrangian submanifold of  $M$ .*

- Example:  $M$  embedded into  $T^*M$  as  $i(x) = (x, \mu_x)$  for a fixed  $\mu \in \Omega^1(M)$  is Lagrangian iff  $\mu$  is closed.
- If  $\phi : (M_1, \omega_1) \rightarrow (M_2, \omega_2)$  is a symplectomorphism, then the graph  $\{(x, \phi(x)) | x \in M_1\} \subset (M_1 \times M_2, \tilde{\omega} = \pi_1^*\omega_1 - \pi_2^*\omega_2)$  is Lagrangian.



# Lagrangian submanifolds

## Definition

Let  $(M, \omega)$  be a  $2n$ -dimensional symplectic manifold. An  $n$ -dimensional submanifold  $i : Y \hookrightarrow M$  with  $i^*\omega = \omega|_Y = 0$  is called a Lagrangian submanifold of  $M$ .

- Example:  $M$  embedded into  $T^*M$  as  $i(x) = (x, \mu_x)$  for a fixed  $\mu \in \Omega^1(M)$  is Lagrangian iff  $\mu$  is closed.
- If  $\phi : (M_1, \omega_1) \rightarrow (M_2, \omega_2)$  is a symplectomorphism, then the graph  $\{(x, \phi(x)) | x \in M_1\} \subset (M_1 \times M_2, \tilde{\omega} = \pi_1^*\omega_1 - \pi_2^*\omega_2)$  is Lagrangian.  
*Proof:*  $i^*(\tilde{\omega}) = (\pi_1 \circ i)^*\omega_1 - (\pi_2 \circ i)^*\omega_2 = (id)^*\omega_1 - \phi^*\omega_2 = 0$ .

# Lagrangian submanifolds

## Definition

Let  $(M, \omega)$  be a  $2n$ -dimensional symplectic manifold. An  $n$ -dimensional submanifold  $i : Y \hookrightarrow M$  with  $i^*\omega = \omega|_Y = 0$  is called a *Lagrangian submanifold* of  $M$ .

- Example:  $M$  embedded into  $T^*M$  as  $i(x) = (x, \mu_x)$  for a fixed  $\mu \in \Omega^1(M)$  is Lagrangian iff  $\mu$  is closed.
- If  $\phi : (M_1, \omega_1) \rightarrow (M_2, \omega_2)$  is a symplectomorphism, then the graph  $\{(x, \phi(x)) | x \in M_1\} \subset (M_1 \times M_2, \tilde{\omega} = \pi_1^*\omega_1 - \pi_2^*\omega_2)$  is Lagrangian.  
*Proof:*  $i^*(\tilde{\omega}) = (\pi_1 \circ i)^*\omega_1 - (\pi_2 \circ i)^*\omega_2 = (id)^*\omega_1 - \phi^*\omega_2 = 0$ .
- Converse also holds, thus giving a criterion for a diffeomorphism to be a symplectomorphism.

# Lagrangian submanifolds

## Definition

Let  $(M, \omega)$  be a  $2n$ -dimensional symplectic manifold. An  $n$ -dimensional submanifold  $i : Y \hookrightarrow M$  with  $i^*\omega = \omega|_Y = 0$  is called a *Lagrangian submanifold* of  $M$ .

- Example:  $M$  embedded into  $T^*M$  as  $i(x) = (x, \mu_x)$  for a fixed  $\mu \in \Omega^1(M)$  is Lagrangian iff  $\mu$  is closed.
- If  $\phi : (M_1, \omega_1) \rightarrow (M_2, \omega_2)$  is a symplectomorphism, then the graph  $\{(x, \phi(x)) | x \in M_1\} \subset (M_1 \times M_2, \tilde{\omega} = \pi_1^*\omega_1 - \pi_2^*\omega_2)$  is Lagrangian.  
*Proof:*  $i^*(\tilde{\omega}) = (\pi_1 \circ i)^*\omega_1 - (\pi_2 \circ i)^*\omega_2 = (id)^*\omega_1 - \phi^*\omega_2 = 0$ .
- Converse also holds, thus giving a criterion for a diffeomorphism to be a symplectomorphism.
- Lagrangian submanifolds provide a geometric method of studying symplectomorphisms, and is especially effective in fixed point theory.

# Important results

## Theorem (Moser)

*For a compact manifold  $M$ , let  $S_c = \{\text{symplectic forms } \omega \text{ on } M \mid [\omega] = c\}$  for some  $c \in H_{de Rham}^2(M)$ . Then all symplectic forms on a path-connected component of  $S_c$  are symplectomorphic.*

# Important results

## Theorem (Moser)

*For a compact manifold  $M$ , let  $S_c = \{\text{symplectic forms } \omega \text{ on } M \mid [\omega] = c\}$  for some  $c \in H_{de Rham}^2(M)$ . Then all symplectic forms on a path-connected component of  $S_c$  are symplectomorphic.*

## Theorem (Darboux Theorem, Weinstein)

*Let  $(M, \omega)$  be a symplectic manifold and let  $p \in M$ . Then there exists a coordinate chart  $(U, x_1, \dots, x_n, y_1, \dots, y_n)$  centered at  $p$  such that  $\omega|_U = \sum_i dx_i \wedge dy_i$ .*

# Important results

## Theorem (Moser)

*For a compact manifold  $M$ , let  $S_c = \{\text{symplectic forms } \omega \text{ on } M \mid [\omega] = c\}$  for some  $c \in H_{de\ Rham}^2(M)$ . Then all symplectic forms on a path-connected component of  $S_c$  are symplectomorphic.*

## Theorem (Darboux Theorem, Weinstein)

*Let  $(M, \omega)$  be a symplectic manifold and let  $p \in M$ . Then there exists a coordinate chart  $(U, x_1, \dots, x_n, y_1, \dots, y_n)$  centered at  $p$  such that  $\omega|_U = \sum_i dx_i \wedge dy_i$ .*

- No local invariants exist for symplectic manifolds.

# Important results

## Theorem (Moser)

*For a compact manifold  $M$ , let  $S_c = \{\text{symplectic forms } \omega \text{ on } M \mid [\omega] = c\}$  for some  $c \in H_{de\ Rham}^2(M)$ . Then all symplectic forms on a path-connected component of  $S_c$  are symplectomorphic.*

## Theorem (Darboux Theorem, Weinstein)

*Let  $(M, \omega)$  be a symplectic manifold and let  $p \in M$ . Then there exists a coordinate chart  $(U, x_1, \dots, x_n, y_1, \dots, y_n)$  centered at  $p$  such that  $\omega|_U = \sum_i dx_i \wedge dy_i$ .*

- No local invariants exist for symplectic manifolds.
- In 1971, Weinstein also proved that all Lagrangian embeddings are locally zero sections.

# Important results

## Theorem (Moser)

*For a compact manifold  $M$ , let  $S_c = \{\text{symplectic forms } \omega \text{ on } M \mid [\omega] = c\}$  for some  $c \in H_{de Rham}^2(M)$ . Then all symplectic forms on a path-connected component of  $S_c$  are symplectomorphic.*

## Theorem (Darboux Theorem, Weinstein)

*Let  $(M, \omega)$  be a symplectic manifold and let  $p \in M$ . Then there exists a coordinate chart  $(U, x_1, \dots, x_n, y_1, \dots, y_n)$  centered at  $p$  such that  $\omega|_U = \sum_i dx_i \wedge dy_i$ .*

- No local invariants exist for symplectic manifolds.
- In 1971, Weinstein also proved that all Lagrangian embeddings are locally zero sections.



# Table of Contents

- 1 Preliminaries from Differential topology
- 2 Symplectic structure
- 3 Contact structure**
- 4 Complex structure

# Contact Manifolds

- Consider a tangent hyperplane (codimension 1 subspace)  
 $\ker \alpha_p = H_p \subset T_p M$  for some covector  $\alpha_p$ .

# Contact Manifolds

- Consider a tangent hyperplane (codimension 1 subspace)  
 $\ker \alpha_p = H_p \subset T_p M$  for some covector  $\alpha_p$ .
- A smooth field of tangent hyperplanes  $H : p \rightarrow H_p \subset T_p M$  is, at least locally, defined as  $\ker \alpha$  for some 1-form  $\alpha$ .

# Contact Manifolds

- Consider a tangent hyperplane (codimension 1 subspace)  
 $\ker \alpha_p = H_p \subset T_p M$  for some covector  $\alpha_p$ .
- A smooth field of tangent hyperplanes  $H : p \rightarrow H_p \subset T_p M$  is, at least locally, defined as  $\ker \alpha$  for some 1-form  $\alpha$ .

## Definition

*A field of tangent hyperplanes  $H$  on  $M$  defines a contact structure iff  $\alpha \wedge (d\alpha)^n$  is nowhere-vanishing for any locally defining 1-form  $\alpha$ . The pair  $(M, H)$  is then called a contact manifold, and  $\alpha$  a local contact form.*

# Contact Manifolds

- Consider a tangent hyperplane (codimension 1 subspace)  $\ker \alpha_p = H_p \subset T_p M$  for some covector  $\alpha_p$ .
- A smooth field of tangent hyperplanes  $H : p \rightarrow H_p \subset T_p M$  is, at least locally, defined as  $\ker \alpha$  for some 1-form  $\alpha$ .

## Definition

*A field of tangent hyperplanes  $H$  on  $M$  defines a contact structure iff  $\alpha \wedge (d\alpha)^n$  is nowhere-vanishing for any locally defining 1-form  $\alpha$ . The pair  $(M, H)$  is then called a contact manifold, and  $\alpha$  a local contact form.*

- Equivalently,  $d\alpha|_H$  is symplectic, and thus  $M$  is odd-dimensional.

# Contact Manifolds

- Consider a tangent hyperplane (codimension 1 subspace)  $\ker \alpha_p = H_p \subset T_p M$  for some covector  $\alpha_p$ .
- A smooth field of tangent hyperplanes  $H : p \rightarrow H_p \subset T_p M$  is, at least locally, defined as  $\ker \alpha$  for some 1-form  $\alpha$ .

## Definition

*A field of tangent hyperplanes  $H$  on  $M$  defines a contact structure iff  $\alpha \wedge (d\alpha)^n$  is nowhere-vanishing for any locally defining 1-form  $\alpha$ . The pair  $(M, H)$  is then called a contact manifold, and  $\alpha$  a local contact form.*

- Equivalently,  $d\alpha|_H$  is symplectic, and thus  $M$  is odd-dimensional.
- As such, contact manifolds are the odd-dimensional analogue of symplectic manifolds.

# Contact Manifolds

- Consider a tangent hyperplane (codimension 1 subspace)  $\ker \alpha_p = H_p \subset T_p M$  for some covector  $\alpha_p$ .
- A smooth field of tangent hyperplanes  $H : p \rightarrow H_p \subset T_p M$  is, at least locally, defined as  $\ker \alpha$  for some 1-form  $\alpha$ .

## Definition

*A field of tangent hyperplanes  $H$  on  $M$  defines a contact structure iff  $\alpha \wedge (d\alpha)^n$  is nowhere-vanishing for any locally defining 1-form  $\alpha$ . The pair  $(M, H)$  is then called a contact manifold, and  $\alpha$  a local contact form.*

- Equivalently,  $d\alpha|_H$  is symplectic, and thus  $M$  is odd-dimensional.
- As such, contact manifolds are the odd-dimensional analogue of symplectic manifolds.
- $\mathbb{R}^{2n+1}$  with coordinates  $(x_1, \dots, x_n, y_1, \dots, y_n, z)$  is a contact manifold with global contact form  $\alpha = \sum_i x_i dy_i + dz$ .

# Contact Manifolds

- Consider a tangent hyperplane (codimension 1 subspace)  $\ker \alpha_p = H_p \subset T_p M$  for some covector  $\alpha_p$ .
- A smooth field of tangent hyperplanes  $H : p \rightarrow H_p \subset T_p M$  is, at least locally, defined as  $\ker \alpha$  for some 1-form  $\alpha$ .

## Definition

*A field of tangent hyperplanes  $H$  on  $M$  defines a contact structure iff  $\alpha \wedge (d\alpha)^n$  is nowhere-vanishing for any locally defining 1-form  $\alpha$ . The pair  $(M, H)$  is then called a contact manifold, and  $\alpha$  a local contact form.*

- Equivalently,  $d\alpha|_H$  is symplectic, and thus  $M$  is odd-dimensional.
- As such, contact manifolds are the odd-dimensional analogue of symplectic manifolds.
- $\mathbb{R}^{2n+1}$  with coordinates  $(x_1, \dots, x_n, y_1, \dots, y_n, z)$  is a contact manifold with global contact form  $\alpha = \sum_i x_i dy_i + dz$ .
- Any compact orientable 3-manifold admits a contact structure (Martinet, 1971).



# Projectivized Cotangent Bundle

Just as how every manifold  $M$  has  $T^*M$  as an associated symplectic structure, there exists a canonical associated contact structure as well.

# Projectivized Cotangent Bundle

Just as how every manifold  $M$  has  $T^*M$  as an associated symplectic structure, there exists a canonical associated contact structure as well.

## Definiton

Let  $\sim$  be an equivalence relation on  $T^*M \setminus \{\text{zero section}\}$  given by  $(p, \zeta) \sim (p, \zeta')$  iff  $\zeta' = \lambda\zeta$  for some  $\lambda \in \mathbb{R}^\times$ . Then the projectivized cotangent bundle is given as  $\mathbb{P}^*M = (T^*M \setminus \{\text{zero section}\}) / \sim$ .

# Projectivized Cotangent Bundle

Just as how every manifold  $M$  has  $T^*M$  as an associated symplectic structure, there exists a canonical associated contact structure as well.

## Definiton

Let  $\sim$  be an equivalence relation on  $T^*M \setminus \{\text{zero section}\}$  given by  $(p, \zeta) \sim (p, \zeta')$  iff  $\zeta' = \lambda\zeta$  for some  $\lambda \in \mathbb{R}^\times$ . Then the projectivized cotangent bundle is given as  $\mathbb{P}^*M = (T^*M \setminus \{\text{zero section}\}) / \sim$ .

- $\mathbb{P}^*M$  has the differential structure of  $M \times \mathbb{RP}^{n-1}$ .

# Projectivized Cotangent Bundle

Just as how every manifold  $M$  has  $T^*M$  as an associated symplectic structure, there exists a canonical associated contact structure as well.

## Definiton

Let  $\sim$  be an equivalence relation on  $T^*M \setminus \{\text{zero section}\}$  given by  $(p, \zeta) \sim (p, \zeta')$  iff  $\zeta' = \lambda\zeta$  for some  $\lambda \in \mathbb{R}^\times$ . Then the projectivized cotangent bundle is given as  $\mathbb{P}^*M = (T^*M \setminus \{\text{zero section}\}) / \sim$ .

- $\mathbb{P}^*M$  has the differential structure of  $M \times \mathbb{RP}^{n-1}$ .
- $\mathbb{P}^*M$  has a contact structure defined by the family of local contact forms  $\alpha_i = dx_i + \sum_{j \neq i} \zeta_j dx_j$  on  $U_i = \{(p, [\zeta]) \in \mathbb{P}^*M \mid \zeta_i \neq 0\}$ .

# Projectivized Cotangent Bundle

Just as how every manifold  $M$  has  $T^*M$  as an associated symplectic structure, there exists a canonical associated contact structure as well.

## Definition

Let  $\sim$  be an equivalence relation on  $T^*M \setminus \{\text{zero section}\}$  given by  $(p, \zeta) \sim (p, \zeta')$  iff  $\zeta' = \lambda\zeta$  for some  $\lambda \in \mathbb{R}^\times$ . Then the projectivized cotangent bundle is given as  $\mathbb{P}^*M = (T^*M \setminus \{\text{zero section}\}) / \sim$ .

- $\mathbb{P}^*M$  has the differential structure of  $M \times \mathbb{RP}^{n-1}$ .
- $\mathbb{P}^*M$  has a contact structure defined by the family of local contact forms  $\alpha_i = dx_i + \sum_{j \neq i} \zeta_j dx_j$  on  $U_i = \{(p, [\zeta]) \in \mathbb{P}^*M \mid \zeta_i \neq 0\}$ .
- Example: For  $M = \mathbb{R}^3$ ,  $\mathbb{P}^*M \cong \mathbb{R}^3 \times \mathbb{RP}^2$ .

# Projectivized Cotangent Bundle

Just as how every manifold  $M$  has  $T^*M$  as an associated symplectic structure, there exists a canonical associated contact structure as well.

## Definition

Let  $\sim$  be an equivalence relation on  $T^*M \setminus \{\text{zero section}\}$  given by  $(p, \zeta) \sim (p, \zeta')$  iff  $\zeta' = \lambda\zeta$  for some  $\lambda \in \mathbb{R}^\times$ . Then the projectivized cotangent bundle is given as  $\mathbb{P}^*M = (T^*M \setminus \{\text{zero section}\}) / \sim$ .

- $\mathbb{P}^*M$  has the differential structure of  $M \times \mathbb{RP}^{n-1}$ .
- $\mathbb{P}^*M$  has a contact structure defined by the family of local contact forms  $\alpha_i = dx_i + \sum_{j \neq i} \zeta_j dx_j$  on  $U_i = \{(p, [\zeta]) \in \mathbb{P}^*M \mid \zeta_i \neq 0\}$ .
- Example: For  $M = \mathbb{R}^3$ ,  $\mathbb{P}^*M \cong \mathbb{R}^3 \times \mathbb{RP}^2$ .
- Example: For  $M = \mathbb{T}^2$ ,  $\mathbb{P}^*M \cong \mathbb{T}^2 \times \mathbb{RP}^1$ .

# Local contact structure

We can also classify the local behavior of contact manifolds in a very similar way to symplectic manifolds by using the following technique:

# Local contact structure

We can also classify the local behavior of contact manifolds in a very similar way to symplectic manifolds by using the following technique:

## Theorem (Symplectization)

*Let  $(M, H)$  be contact manifold with a contact form  $\alpha$ . Define  $\tilde{M} = M \times \mathbb{R}$ , with  $\tau$  being the additional coordinate on  $\mathbb{R}$ . Then with  $\pi : \tilde{M} \rightarrow M$  given by  $\pi(p, \tau) = p$ ,  $\omega = d(e^\tau \pi^* \alpha)$  is symplectic on  $\tilde{M}$ .*



# Local contact structure

We can also classify the local behavior of contact manifolds in a very similar way to symplectic manifolds by using the following technique:

## Theorem (Symplectization)

*Let  $(M, H)$  be contact manifold with a contact form  $\alpha$ . Define  $\tilde{M} = M \times \mathbb{R}$ , with  $\tau$  being the additional coordinate on  $\mathbb{R}$ . Then with  $\pi : \tilde{M} \rightarrow M$  given by  $\pi(p, \tau) = p$ ,  $\omega = d(e^\tau \pi^* \alpha)$  is symplectic on  $\tilde{M}$ .*

- For a given contact manifold  $(M, H)$ , we apply the original Darboux Theorem to its symplectization to locally classify contact forms on  $M$ :

# Local contact structure

We can also classify the local behavior of contact manifolds in a very similar way to symplectic manifolds by using the following technique:

## Theorem (Symplectization)

*Let  $(M, H)$  be contact manifold with a contact form  $\alpha$ . Define  $\tilde{M} = M \times \mathbb{R}$ , with  $\tau$  being the additional coordinate on  $\mathbb{R}$ . Then with  $\pi : \tilde{M} \rightarrow M$  given by  $\pi(p, \tau) = p$ ,  $\omega = d(e^\tau \pi^* \alpha)$  is symplectic on  $\tilde{M}$ .*

- For a given contact manifold  $(M, H)$ , we apply the original Darboux Theorem to its symplectization to locally classify contact forms on  $M$ :

## Theorem (Darboux, Contact version)

*Let  $(M, H)$  be a contact manifold and  $p \in M$ . Then there exists a coordinate chart  $(U, x_1, \dots, x_n, y_1, \dots, y_n, z)$  at  $p$  such that  $\alpha = \sum_i x_i dy_i + dz$  is a local contact form for  $H$ .*

# Table of Contents

- 1 Preliminaries from Differential topology
- 2 Symplectic structure
- 3 Contact structure
- 4 Complex structure**

# Complex linear algebra

## Definition

A complex structure on a real vector space  $V$  is a map  $J : V \rightarrow V$  such that  $J^2 = -Id$ .

# Complex linear algebra

## Definition

A complex structure on a real vector space  $V$  is a map  $J : V \rightarrow V$  such that  $J^2 = -Id$ .

- Example: Any even-dimensional real vector space admits a complex structure.

# Complex linear algebra

## Definition

A complex structure on a real vector space  $V$  is a map  $J : V \rightarrow V$  such that  $J^2 = -Id$ .

- Example: Any even-dimensional real vector space admits a complex structure.
- A complex structure  $J$  on a symplectic vector space  $(V, \Omega)$  is  $\Omega$ -compatible if  $G_J(u, v) := \Omega(u, Jv)$  is a positive inner product on  $V$ .

# Complex linear algebra

## Definition

A complex structure on a real vector space  $V$  is a map  $J : V \rightarrow V$  such that  $J^2 = -Id$ .

- Example: Any even-dimensional real vector space admits a complex structure.
- A complex structure  $J$  on a symplectic vector space  $(V, \Omega)$  is  $\Omega$ -compatible if  $G_J(u, v) := \Omega(u, Jv)$  is a positive inner product on  $V$ .

## Theorem

Let  $(V, \Omega)$  be a symplectic vector space. Then there exists a compatible complex structure  $J$  on  $V$ .

# Complex linear algebra

## Definition

A complex structure on a real vector space  $V$  is a map  $J : V \rightarrow V$  such that  $J^2 = -Id$ .

- Example: Any even-dimensional real vector space admits a complex structure.
- A complex structure  $J$  on a symplectic vector space  $(V, \Omega)$  is  $\Omega$ -compatible if  $G_J(u, v) := \Omega(u, Jv)$  is a positive inner product on  $V$ .

## Theorem

Let  $(V, \Omega)$  be a symplectic vector space. Then there exists a compatible complex structure  $J$  on  $V$ .

- For a symplectic basis  $\{e_i, f_i\}_{i=1}^n$ , set  $J(e_i) = f_i$  and  $J(f_i) = -e_i$ .



# Complex linear algebra

## Definition

A complex structure on a real vector space  $V$  is a map  $J : V \rightarrow V$  such that  $J^2 = -Id$ .

- Example: Any even-dimensional real vector space admits a complex structure.
- A complex structure  $J$  on a symplectic vector space  $(V, \Omega)$  is  $\Omega$ -compatible if  $G_J(u, v) := \Omega(u, Jv)$  is a positive inner product on  $V$ .

## Theorem

Let  $(V, \Omega)$  be a symplectic vector space. Then there exists a compatible complex structure  $J$  on  $V$ .

- For a symplectic basis  $\{e_i, f_i\}_{i=1}^n$ , set  $J(e_i) = f_i$  and  $J(f_i) = -e_i$ .
- Alternatively, there is a basis-independent method of generating a  $J$  using polar decomposition.

# Complex linear algebra

## Definition

A complex structure on a real vector space  $V$  is a map  $J : V \rightarrow V$  such that  $J^2 = -Id$ .

- Example: Any even-dimensional real vector space admits a complex structure.
- A complex structure  $J$  on a symplectic vector space  $(V, \Omega)$  is  $\Omega$ -compatible if  $G_J(u, v) := \Omega(u, Jv)$  is a positive inner product on  $V$ .

## Theorem

Let  $(V, \Omega)$  be a symplectic vector space. Then there exists a compatible complex structure  $J$  on  $V$ .

- For a symplectic basis  $\{e_i, f_i\}_{i=1}^n$ , set  $J(e_i) = f_i$  and  $J(f_i) = -e_i$ .
- Alternatively, there is a basis-independent method of generating a  $J$  using polar decomposition.
- The converse compatibility also always exists.

# Almost complex manifolds

These notions can be somewhat generalized to manifold structures.

# Almost complex manifolds

These notions can be somewhat generalized to manifold structures.

## Definition

*An almost complex structure on a manifold  $M$  is a smooth field  $J$  of complex structures  $J_p : T_p M \rightarrow T_p M$ .*

# Almost complex manifolds

These notions can be somewhat generalized to manifold structures.

## Definition

*An almost complex structure on a manifold  $M$  is a smooth field  $J$  of complex structures  $J_p : T_p M \rightarrow T_p M$ .*

- An almost complex structure  $J$  on a symplectic manifold  $(M, \omega)$  is  $\omega$ -compatible if  $p \mapsto g_p : T_p M \times T_p M \rightarrow \mathbb{R}$  with  $g_p(v, u) = (v, J_p u)$  is a Riemannian metric on  $M$ . Then  $(\omega, J, g)$  is a *compatible triple*.

# Almost complex manifolds

These notions can be somewhat generalized to manifold structures.

## Definition

*An almost complex structure on a manifold  $M$  is a smooth field  $J$  of complex structures  $J_p : T_p M \rightarrow T_p M$ .*

- An almost complex structure  $J$  on a symplectic manifold  $(M, \omega)$  is  $\omega$ -compatible if  $p \mapsto g_p : T_p M \times T_p M \rightarrow \mathbb{R}$  with  $g_p(v, u) = (v, J_p u)$  is a Riemannian metric on  $M$ . Then  $(\omega, J, g)$  is a *compatible triple*.
- For a compatible triple,  $\omega(u, v) = g(Ju, v)$  and  $J(u) = \tilde{g}^{-1}(\tilde{\omega}^{-1}(u))$ .

# Almost complex manifolds

These notions can be somewhat generalized to manifold structures.

## Definition

*An almost complex structure on a manifold  $M$  is a smooth field  $J$  of complex structures  $J_p : T_p M \rightarrow T_p M$ .*

- An almost complex structure  $J$  on a symplectic manifold  $(M, \omega)$  is  $\omega$ -compatible if  $p \mapsto g_p : T_p M \times T_p M \rightarrow \mathbb{R}$  with  $g_p(v, u) = \langle v, J_p u \rangle$  is a Riemannian metric on  $M$ . Then  $(\omega, J, g)$  is a compatible triple.
- For a compatible triple,  $\omega(u, v) = g(Ju, v)$  and  $J(u) = \tilde{g}^{-1}(\tilde{\omega}^{-1}(u))$ .
- Any symplectic manifold has compatible almost complex structures.

# Almost complex manifolds

These notions can be somewhat generalized to manifold structures.

## Definition

*An almost complex structure on a manifold  $M$  is a smooth field  $J$  of complex structures  $J_p : T_p M \rightarrow T_p M$ .*

- An almost complex structure  $J$  on a symplectic manifold  $(M, \omega)$  is  $\omega$ -compatible if  $p \mapsto g_p : T_p M \times T_p M \rightarrow \mathbb{R}$  with  $g_p(v, u) = \omega(v, J_p u)$  is a Riemannian metric on  $M$ . Then  $(\omega, J, g)$  is a *compatible triple*.
- For a compatible triple,  $\omega(u, v) = g(Ju, v)$  and  $J(u) = \tilde{g}^{-1}(\tilde{\omega}^{-1}(u))$ .
- Any symplectic manifold has compatible almost complex structures.
- The converse is not necessarily true because of closedness (e.g.,  $S^6$ ).



# Almost complex manifolds

These notions can be somewhat generalized to manifold structures.

## Definition

*An almost complex structure on a manifold  $M$  is a smooth field  $J$  of complex structures  $J_p : T_p M \rightarrow T_p M$ .*

- An almost complex structure  $J$  on a symplectic manifold  $(M, \omega)$  is  $\omega$ -compatible if  $p \mapsto g_p : T_p M \times T_p M \rightarrow \mathbb{R}$  with  $g_p(v, u) = \omega(v, J_p u)$  is a Riemannian metric on  $M$ . Then  $(\omega, J, g)$  is a *compatible triple*.
- For a compatible triple,  $\omega(u, v) = g(Ju, v)$  and  $J(u) = \tilde{g}^{-1}(\tilde{\omega}^{-1}(u))$ .
- Any symplectic manifold has compatible almost complex structures.
- The converse is not necessarily true because of closedness (e.g.,  $S^6$ ).
- That said,  $\mathfrak{J}(M, \omega)$  and  $\Omega(M, J)$  are path-connected.

# Almost complex manifolds

These notions can be somewhat generalized to manifold structures.

## Definition

*An almost complex structure on a manifold  $M$  is a smooth field  $J$  of complex structures  $J_p : T_p M \rightarrow T_p M$ .*

- An almost complex structure  $J$  on a symplectic manifold  $(M, \omega)$  is  $\omega$ -compatible if  $p \mapsto g_p : T_p M \times T_p M \rightarrow \mathbb{R}$  with  $g_p(v, u) = \omega(v, J_p u)$  is a Riemannian metric on  $M$ . Then  $(\omega, J, g)$  is a *compatible triple*.
- For a compatible triple,  $\omega(u, v) = g(Ju, v)$  and  $J(u) = \tilde{g}^{-1}(\tilde{\omega}^{-1}(u))$ .
- Any symplectic manifold has compatible almost complex structures.
- The converse is not necessarily true because of closedness (e.g.,  $S^6$ ).
- That said,  $\mathfrak{J}(M, \omega)$  and  $\Omega(M, J)$  are path-connected.

# Almost complex differential structure

Recall the complexification of a real vector space  $V$ .

# Almost complex differential structure

Recall the complexification of a real vector space  $V$ .

## Definition

*The complexification  $V^{\mathbb{C}} = V \otimes_{\mathbb{R}} \mathbb{C}$  of a real vector space  $V$  is a complex vector space with complex multiplication defined by setting  $i(v \otimes c) = v \otimes (ic)$  for  $c \in \mathbb{C}$ .*

# Almost complex differential structure

Recall the complexification of a real vector space  $V$ .

## Definition

*The complexification  $V^{\mathbb{C}} = V \otimes_{\mathbb{R}} \mathbb{C}$  of a real vector space  $V$  is a complex vector space with complex multiplication defined by setting  $i(v \otimes c) = v \otimes (ic)$  for  $c \in \mathbb{C}$ .*

- The complexification of the tangent bundle  $TM$  (fiber-wise), and defining  $J(v \otimes c) = Jv \otimes c$ , allows us to decompose  $TM \otimes \mathbb{C}$ .

# Almost complex differential structure

Recall the complexification of a real vector space  $V$ .

## Definition

*The complexification  $V^{\mathbb{C}} = V \otimes_{\mathbb{R}} \mathbb{C}$  of a real vector space  $V$  is a complex vector space with complex multiplication defined by setting  $i(v \otimes c) = v \otimes (ic)$  for  $c \in \mathbb{C}$ .*

- The complexification of the tangent bundle  $TM$  (fiber-wise), and defining  $J(v \otimes c) = Jv \otimes c$ , allows us to decompose  $TM \otimes \mathbb{C}$ .
- $TM \otimes \mathbb{C} = T_{1,0} \oplus T_{0,1}$ , where  $T_{1,0} = (+i)$ -eigenspace of  $J = \{v \otimes 1 - Jv \otimes i \mid v \in TM\}$  and  $T_{0,1} = (-i)$ -eigenspace of  $J = \{v \otimes 1 + Jv \otimes i \mid v \in TM\}$ .

# Almost complex differential structure

Recall the complexification of a real vector space  $V$ .

## Definition

*The complexification  $V^{\mathbb{C}} = V \otimes_{\mathbb{R}} \mathbb{C}$  of a real vector space  $V$  is a complex vector space with complex multiplication defined by setting  $i(v \otimes c) = v \otimes (ic)$  for  $c \in \mathbb{C}$ .*

- The complexification of the tangent bundle  $TM$  (fiber-wise), and defining  $J(v \otimes c) = Jv \otimes c$ , allows us to decompose  $TM \otimes \mathbb{C}$ .
- $TM \otimes \mathbb{C} = T_{1,0} \oplus T_{0,1}$ , where  $T_{1,0} = (+i)$ -eigenspace of  $J = \{v \otimes 1 - Jv \otimes i \mid v \in TM\}$  and  $T_{0,1} = (-i)$ -eigenspace of  $J = \{v \otimes 1 + Jv \otimes i \mid v \in TM\}$ .
- Similarly,  $T^*M \otimes \mathbb{C} = T^{1,0} \oplus T^{0,1}$  where  $T^{1,0} = (T_{1,0})^* = \{\zeta \otimes 1 - (\zeta \circ J) \otimes i \mid \zeta \in T^*M\}$  and  $T^{0,1} = (T_{0,1})^* = \{\zeta \otimes 1 + (\zeta \circ J) \otimes i \mid \zeta \in T^*M\}$ .

# Almost complex differential structure (continued)

- The above decomposition allows us to decompose  $\Omega^k(M; \mathbb{C})$ .



# Almost complex differential structure (continued)

- The above decomposition allows us to decompose  $\Omega^k(M; \mathbb{C})$ .
- Set  $\Lambda^{l,m} = (\Lambda^l T^{1,0}) \wedge (\Lambda^m T^{0,1})$ , then  $\Lambda^k(T^*M \otimes \mathbb{C}) = \bigoplus_{l+m=k} \Lambda^{l,m}$ .

# Almost complex differential structure (continued)

- The above decomposition allows us to decompose  $\Omega^k(M; \mathbb{C})$ .
- Set  $\Lambda^{l,m} = (\Lambda^l T^{1,0}) \wedge (\Lambda^m T^{0,1})$ , then  $\Lambda^k(T^*M \otimes \mathbb{C}) = \bigoplus_{l+m=k} \Lambda^{l,m}$ .
- Now setting  $\Omega^{l,m}$  = sections of  $\Lambda^{l,m}$ , we have the final decomposition of *complex-valued forms*  $\Omega^k(M; \mathbb{C}) = \bigoplus_{l+m=k} \Omega^{l,m}$ .

# Almost complex differential structure (continued)

- The above decomposition allows us to decompose  $\Omega^k(M; \mathbb{C})$ .
- Set  $\Lambda^{l,m} = (\Lambda^l T^{1,0}) \wedge (\Lambda^m T^{0,1})$ , then  $\Lambda^k(T^*M \otimes \mathbb{C}) = \bigoplus_{l+m=k} \Lambda^{l,m}$ .
- Now setting  $\Omega^{l,m}$  = sections of  $\Lambda^{l,m}$ , we have the final decomposition of *complex-valued forms*  $\Omega^k(M; \mathbb{C}) = \bigoplus_{l+m=k} \Omega^{l,m}$ .
- The exterior derivative  $d : \Omega^k(M; \mathbb{C}) \rightarrow \Omega^{k+1}(M; \mathbb{C})$  is defined as usual, but has different projections.

# Almost complex differential structure (continued)

- The above decomposition allows us to decompose  $\Omega^k(M; \mathbb{C})$ .
- Set  $\Lambda^{l,m} = (\Lambda^l T^{1,0}) \wedge (\Lambda^m T^{0,1})$ , then  $\Lambda^k(T^*M \otimes \mathbb{C}) = \bigoplus_{l+m=k} \Lambda^{l,m}$ .
- Now setting  $\Omega^{l,m}$  = sections of  $\Lambda^{l,m}$ , we have the final decomposition of *complex-valued forms*  $\Omega^k(M; \mathbb{C}) = \bigoplus_{l+m=k} \Omega^{l,m}$ .
- The exterior derivative  $d : \Omega^k(M; \mathbb{C}) \rightarrow \Omega^{k+1}(M; \mathbb{C})$  is defined as usual, but has different projections.
- $\partial := \pi^{l+1,m} \circ d : \Omega^{l,m}(M) \rightarrow \Omega^{l+1,m}$  and  $\bar{\partial} := \pi^{l,m+1} \circ d : \Omega^{l,m}(M) \rightarrow \Omega^{l,m+1}$ .

# Almost complex differential structure (continued)

- The above decomposition allows us to decompose  $\Omega^k(M; \mathbb{C})$ .
- Set  $\Lambda^{l,m} = (\Lambda^l T^{1,0}) \wedge (\Lambda^m T^{0,1})$ , then  $\Lambda^k(T^*M \otimes \mathbb{C}) = \bigoplus_{l+m=k} \Lambda^{l,m}$ .
- Now setting  $\Omega^{l,m}$  = sections of  $\Lambda^{l,m}$ , we have the final decomposition of *complex-valued forms*  $\Omega^k(M; \mathbb{C}) = \bigoplus_{l+m=k} \Omega^{l,m}$ .
- The exterior derivative  $d : \Omega^k(M; \mathbb{C}) \rightarrow \Omega^{k+1}(M; \mathbb{C})$  is defined as usual, but has different projections.
- $\partial := \pi^{l+1,m} \circ d : \Omega^{l,m}(M) \rightarrow \Omega^{l+1,m}$  and  $\bar{\partial} := \pi^{l,m+1} \circ d : \Omega^{l,m}(M) \rightarrow \Omega^{l,m+1}$ .
- $\bar{\partial}^2 = \partial^2 = 0$ , and hence allows for the formation of the *Dolbeault Cohomology*.

# Complex differential structure

Recall that a *complex manifold* of dimension  $n$  is defined analogously to a real manifold, except that it is locally biholomorphic to  $\mathbb{C}^n$ , and the transition maps are required to be biholomorphic instead of just smooth.

# Complex differential structure

Recall that a *complex manifold* of dimension  $n$  is defined analogously to a real manifold, except that it is locally biholomorphic to  $\mathbb{C}^n$ , and the transition maps are required to be biholomorphic instead of just smooth.

## Theorem

*Every complex manifold has a canonical almost complex structure  $J$ .*

# Complex differential structure

Recall that a *complex manifold* of dimension  $n$  is defined analogously to a real manifold, except that it is locally biholomorphic to  $\mathbb{C}^n$ , and the transition maps are required to be biholomorphic instead of just smooth.

## Theorem

*Every complex manifold has a canonical almost complex structure  $J$ .*

- In particular, if  $M$  has complex coordinates  $\{z_j\}_{j=1}^n$  in some nbhd.  $U$  of  $p \in M$ , and thus real coordinates  $\{x_j, y_j\}_{j=1}^n$  with  $z_j = x_j + iy_j$ , then  $J_p$  is defined by  $J_p(\frac{\partial}{\partial x_j}|_p) = \frac{\partial}{\partial y_j}|_p$  and  $J(\frac{\partial}{\partial y_j}|_p) = -\frac{\partial}{\partial x_j}|_p$ .



# Complex differential structure

Recall that a *complex manifold* of dimension  $n$  is defined analogously to a real manifold, except that it is locally biholomorphic to  $\mathbb{C}^n$ , and the transition maps are required to be biholomorphic instead of just smooth.

## Theorem

*Every complex manifold has a canonical almost complex structure  $J$ .*

- In particular, if  $M$  has complex coordinates  $\{z_j\}_{j=1}^n$  in some nbhd.  $U$  of  $p \in M$ , and thus real coordinates  $\{x_j, y_j\}_{j=1}^n$  with  $z_j = x_j + iy_j$ , then  $J_p$  is defined by  $J_p(\frac{\partial}{\partial x_j}|_p) = \frac{\partial}{\partial y_j}|_p$  and  $J(\frac{\partial}{\partial y_j}|_p) = -\frac{\partial}{\partial x_j}|_p$ .
- The complexification in this case  $T_p M \otimes \mathbb{C} = \mathbb{C}$ -span of  $\{\frac{\partial}{\partial x_j}|_p, \frac{\partial}{\partial y_j}|_p\}$ .

# Complex differential structure

Recall that a *complex manifold* of dimension  $n$  is defined analogously to a real manifold, except that it is locally biholomorphic to  $\mathbb{C}^n$ , and the transition maps are required to be biholomorphic instead of just smooth.

## Theorem

*Every complex manifold has a canonical almost complex structure  $J$ .*

- In particular, if  $M$  has complex coordinates  $\{z_j\}_{j=1}^n$  in some nbhd.  $U$  of  $p \in M$ , and thus real coordinates  $\{x_j, y_j\}_{j=1}^n$  with  $z_j = x_j + iy_j$ , then  $J_p$  is defined by  $J_p(\frac{\partial}{\partial x_j}|_p) = \frac{\partial}{\partial y_j}|_p$  and  $J(\frac{\partial}{\partial y_j}|_p) = -\frac{\partial}{\partial x_j}|_p$ .
- The complexification in this case  $T_p M \otimes \mathbb{C} = \mathbb{C}$ -span of  $\{\frac{\partial}{\partial x_j}|_p, \frac{\partial}{\partial y_j}|_p\}$ .
- The decomposition from before becomes:  
 $(T_{1,0})_p = \mathbb{C}$ -span $\{\frac{\partial}{\partial z_j} = \frac{1}{2}(\frac{\partial}{\partial x_j}|_p - i\frac{\partial}{\partial y_j}|_p)\}$  and  
 $(T_{0,1})_p = \mathbb{C}$ -span $\{\frac{\partial}{\partial \bar{z}_j} = \frac{1}{2}(\frac{\partial}{\partial x_j}|_p + i\frac{\partial}{\partial y_j}|_p)\}$

# Complex differential structure

Recall that a *complex manifold* of dimension  $n$  is defined analogously to a real manifold, except that it is locally biholomorphic to  $\mathbb{C}^n$ , and the transition maps are required to be biholomorphic instead of just smooth.

## Theorem

*Every complex manifold has a canonical almost complex structure  $J$ .*

- In particular, if  $M$  has complex coordinates  $\{z_j\}_{j=1}^n$  in some nbhd.  $U$  of  $p \in M$ , and thus real coordinates  $\{x_j, y_j\}_{j=1}^n$  with  $z_j = x_j + iy_j$ , then  $J_p$  is defined by  $J_p(\frac{\partial}{\partial x_j}|_p) = \frac{\partial}{\partial y_j}|_p$  and  $J(\frac{\partial}{\partial y_j}|_p) = -\frac{\partial}{\partial x_j}|_p$ .
- The complexification in this case  $T_p M \otimes \mathbb{C} = \mathbb{C}$ -span of  $\{\frac{\partial}{\partial x_j}|_p, \frac{\partial}{\partial y_j}|_p\}$ .
- The decomposition from before becomes:  
 $(T_{1,0})_p = \mathbb{C}$ -span $\{\frac{\partial}{\partial z_j} = \frac{1}{2}(\frac{\partial}{\partial x_j}|_p - i\frac{\partial}{\partial y_j}|_p)\}$  and  
 $(T_{0,1})_p = \mathbb{C}$ -span $\{\frac{\partial}{\partial \bar{z}_j} = \frac{1}{2}(\frac{\partial}{\partial x_j}|_p + i\frac{\partial}{\partial y_j}|_p)\}$
- This definition allows for a natural definition for a basis of 1-forms:  
 $T^{1,0} = \mathbb{C}$ -span $\{dz_j\}$  and  $T^{0,1} = \mathbb{C}$ -span $\{d\bar{z}_j\}$ , and thus  $d = \partial + \bar{\partial}$ .

# Integrability

Both symplectic and complex manifold structures admit almost complex structures. To connect the two, we explore the following phenomenon.

# Integrability

Both symplectic and complex manifold structures admit almost complex structures. To connect the two, we explore the following phenomenon.

## Definition

*An almost complex structure  $J$  on a manifold  $M$  is called integrable if it is induced by a complex manifold structure on  $M$ .*

# Integrability

Both symplectic and complex manifold structures admit almost complex structures. To connect the two, we explore the following phenomenon.

## Definition

*An almost complex structure  $J$  on a manifold  $M$  is called integrable if it is induced by a complex manifold structure on  $M$ .*

## Theorem (Newlander-Nirenberg, 1957)

*Let  $(M, J)$  be an almost complex manifold, and the Nijenhuis tensor  $\Re(v, w) = [Jv, Jw] - J[v, Jw] - J[Jv, w] - [v, w]$ . Then  $J$  is integrable if and only if  $\Re \equiv 0 \Leftrightarrow d = \partial + \bar{\partial}$ .*

# Integrability

Both symplectic and complex manifold structures admit almost complex structures. To connect the two, we explore the following phenomenon.

## Definition

*An almost complex structure  $J$  on a manifold  $M$  is called integrable if it is induced by a complex manifold structure on  $M$ .*

## Theorem (Newlander-Nirenberg, 1957)

*Let  $(M, J)$  be an almost complex manifold, and the Nijenhuis tensor  $\mathfrak{N}(v, w) = [Jv, Jw] - J[v, Jw] - J[Jv, w] - [v, w]$ . Then  $J$  is integrable if and only if  $\mathfrak{N} \equiv 0 \Leftrightarrow d = \partial + \bar{\partial}$ .*

- In particular, one can see that any orientable surface is symplectic, hence almost complex, and by the theorem above, also a complex manifold.

# Kähler Manifolds

## Definition

*A Kähler manifold is a symplectic manifold  $(M, \omega)$  with an integrable compatible almost complex structure.  $\omega$  is then called a Kähler form.*



# Kähler Manifolds

## Definition

*A Kähler manifold is a symplectic manifold  $(M, \omega)$  with an integrable compatible almost complex structure.  $\omega$  is then called a Kähler form.*

- In particular a Kähler form is a  $(1, 1)$ -form that is  $\partial$ - and  $\bar{\partial}$ -closed, and on a coordinate chart  $(U, z_1, \dots, z_n)$  looks like 
$$\omega = \frac{i}{2} \sum_{j,k} b_{j,k} dz_j \wedge d\bar{z}_k$$
 with  $b_{jk}(p)$  being a positive definite hermitian matrix for all  $p \in M$ .

# Kähler Manifolds

## Definition

*A Kähler manifold is a symplectic manifold  $(M, \omega)$  with an integrable compatible almost complex structure.  $\omega$  is then called a Kähler form.*

- In particular a Kähler form is a  $(1, 1)$ -form that is  $\partial$ - and  $\bar{\partial}$ -closed, and on a coordinate chart  $(U, z_1, \dots, z_n)$  looks like 
$$\omega = \frac{i}{2} \sum_{j,k} b_{j,k} dz_j \wedge d\bar{z}_k$$
 with  $b_{jk}(p)$  being a positive definite hermitian matrix for all  $p \in M$ .
- Example:  $\mathbb{C}^n$  with Kähler form  $\omega = \frac{i}{2} \sum_j dz_j \wedge \bar{z}_j$ .

# Kähler Manifolds

## Definition

*A Kähler manifold is a symplectic manifold  $(M, \omega)$  with an integrable compatible almost complex structure.  $\omega$  is then called a Kähler form.*

- In particular a Kähler form is a  $(1, 1)$ -form that is  $\partial$ - and  $\bar{\partial}$ -closed, and on a coordinate chart  $(U, z_1, \dots, z_n)$  looks like 
$$\omega = \frac{i}{2} \sum_{j,k} b_{j,k} dz_j \wedge d\bar{z}_k$$
 with  $b_{jk}(p)$  being a positive definite hermitian matrix for all  $p \in M$ .
- Example:  $\mathbb{C}^n$  with Kähler form  $\omega = \frac{i}{2} \sum_j dz_j \wedge \bar{z}_j$ .

## Theorem (Banyaga)

*Let  $(M, \omega)$  be a compact Kähler manifold with  $[\omega]_{de Rham} = c$ . Then the set of all symplectic forms on  $M$  with de Rham class  $c$  is path-connected.*

# Kähler Manifolds

## Definition

*A Kähler manifold is a symplectic manifold  $(M, \omega)$  with an integrable compatible almost complex structure.  $\omega$  is then called a Kähler form.*

- In particular a Kähler form is a  $(1, 1)$ -form that is  $\partial$ - and  $\bar{\partial}$ -closed, and on a coordinate chart  $(U, z_1, \dots, z_n)$  looks like 
$$\omega = \frac{i}{2} \sum_{j,k} b_{j,k} dz_j \wedge d\bar{z}_k$$
 with  $b_{jk}(p)$  being a positive definite hermitian matrix for all  $p \in M$ .
- Example:  $\mathbb{C}^n$  with Kähler form  $\omega = \frac{i}{2} \sum_j dz_j \wedge \bar{z}_j$ .

## Theorem (Banyaga)

*Let  $(M, \omega)$  be a compact Kähler manifold with  $[\omega]_{de\ Rham} = c$ . Then the set of all symplectic forms on  $M$  with de Rham class  $c$  is path-connected.*

- In particular,  $[\omega_1] = [\omega_2] \Rightarrow (M, \omega_1) \cong (M, \omega_2)$ .

# Local Kähler structure

## Definition

Let  $M$  be a complex manifold. A function  $f \in C^\infty(M; \mathbb{R})$  is strictly plurisubharmonic if on each local chart  $(U, z_1, \dots, z_n)$ , the matrix  $\left[ \frac{\partial^2 f}{\partial z_j \partial \bar{z}_k}(p) \right]$  is positive-definite for all  $f \in U$ .

# Local Kähler structure

## Definition

*Let  $M$  be a complex manifold. A function  $f \in C^\infty(M; \mathbb{R})$  is strictly plurisubharmonic if on each local chart  $(U, z_1, \dots, z_n)$ , the matrix  $\left[ \frac{\partial^2 f}{\partial z_j \partial \bar{z}_k}(p) \right]$  is positive-definite for all  $f \in U$ .*

- In particular,  $\omega = \frac{i}{2} \partial \bar{\partial} f$  is Kähler, and  $f$  is called a Kähler potential.

# Local Kähler structure

## Definition

Let  $M$  be a complex manifold. A function  $f \in C^\infty(M; \mathbb{R})$  is strictly plurisubharmonic if on each local chart  $(U, z_1, \dots, z_n)$ , the matrix  $\left[ \frac{\partial^2 f}{\partial z_j \partial \bar{z}_k}(p) \right]$  is positive-definite for all  $f \in U$ .

- In particular,  $\omega = \frac{i}{2} \partial \bar{\partial} f$  is Kähler, and  $f$  is called a Kähler potential.
- Every Kähler form locally looks like  $\frac{i}{2} \partial \bar{\partial} f$  where  $f$  is strictly plurisubharmonic.

# Local Kähler structure

## Definition

Let  $M$  be a complex manifold. A function  $f \in C^\infty(M; \mathbb{R})$  is strictly plurisubharmonic if on each local chart  $(U, z_1, \dots, z_n)$ , the matrix  $\left[ \frac{\partial^2 f}{\partial z_j \partial \bar{z}_k}(p) \right]$  is positive-definite for all  $f \in U$ .

- In particular,  $\omega = \frac{i}{2} \partial \bar{\partial} f$  is Kähler, and  $f$  is called a Kähler potential.
- Every Kähler form locally looks like  $\frac{i}{2} \partial \bar{\partial} f$  where  $f$  is strictly plurisubharmonic.
- Example: The above principle is used to construct the *Fubini-Study* Kähler form on  $\mathbb{CP}^n$ , with Kähler potential  $f(z) = \log(|z|^2 + 1)$ .



Thank you![2][1]



V.I Arnold.

*Mathematical Methods of Classical Mechanics.*

Springer-Verlag, 1974.



Annas Cannas de Silva.

*Lectures on Symplectic Geometry*, volume 1764 of *Lecture Notes in Mathematics*.

Springer-Verlag, 2001.