

BROILER FARMING MANAGEMENT GUIDELINE

Standard Operating Procedures & Environmental Guidelines

1. Housing & Brooding Management

- **Pre-Arrival Prep:** Clean, wash, and sanitize the house at least 10–14 days prior to chick arrival. Ensure fresh, dry bedding (wood shavings or straw) is laid to a depth of 5–8 cm.
- **Brooding Setup:** Pre-heat the brooding area 24 hours before delivery so the floor surface temperature reaches 30°C to 32°C.
- **Chick Placement:** Immediately provide clean water with electrolytes or glucose to reduce transport stress before opening feed pans.

2. Environmental & Temperature Guidelines

Age Period	Target Temperature (°C)	Target Temperature (°F)	Relative Humidity (%)	Ventilation Strategy
Day 1 – 3	32 – 34 °C	90 – 93 °F	60 – 70 %	Minimum (Air exchange only)
Day 4 – 7	30 – 32 °C	86 – 90 °F	60 – 70 %	Minimum ventilation
Week 2 (Days 8–14)	28 – 30 °C	82 – 86 °F	50 – 60 %	Transition ventilation
Week 3 (Days 15–21)	25 – 27 °C	77 – 81 °F	50 – 60 %	Transition / Tunnel
Week 4+ (Day 22–Market)	20 – 22 °C	68 – 72 °F	50 – 65 %	Tunnel cooling as needed

3. Nutrition & Feeding Schedule

- **Starter Phase (Days 0–10):** High-protein feed (22–23% Crude Protein) in crumble form to encourage early gut development and rapid initial growth.
- **Grower Phase (Days 11–24):** Balanced nutrition (20–21% Crude Protein) in mini-pellet form to support bone frame development and muscle growth.
- **Finisher Phase (Days 25 to Market):** High-energy feed (18–19% Crude Protein) in pellet form to maximize weight gain and meat yield.
- **Water Access:** Keep clean, fresh water available 24/7. Flow rates should scale from 20–30 mL/min for chicks up to 80–100 mL/min for mature birds.

4. Biosecurity & Health Management

- **Access Control:** Restrict unauthorized personnel and vehicles from entering the farm perimeter.
- **Sanitation:** Maintain functional footbaths containing fresh disinfectant at every house entrance.
- **All-In / All-Out:** Run strict single-age batch cycles to allow complete cleaning and disinfection between flocks.

Standard Vaccination Protocol

Age	Target Disease	Vaccine Type	Administration Method
Day 1 (Hatchery)	Marek's Disease & IB / ND	Live / HVT Vector	Subcutaneous / In-Ovo / Spray
Days 7 – 10	Newcastle Disease (ND) + IB	Live (B1 / LaSota)	Drinking Water or Eye Drop

Age	Target Disease	Vaccine Type	Administration Method
Day 14	Infectious Bursal Disease (Gumboro)	Intermediate Plus Live	Drinking Water
Day 28	Newcastle Disease Booster	Live (LaSota) / Killed	Drinking Water or Injection

5. Key Performance Indicators (KPIs)

Performance Metrics

- **Feed Conversion Ratio (FCR):** Calculate as total feed consumed (kg) divided by total live weight gained (kg). Target: < 1.55.
- **Mortality Rate:** Target < 3–4% total mortality over the entire production cycle.