


* MALE REPRODUCTIVE SYSTEM.

- Male reproductive system have mainly two external genitalia i.e. Scrotum, ^{testes} and penis.
- The internal genitalia is mainly Testes.
- Male R.S is ^{also} called as urinogenital system.
- male reproductive system has 3 Accessory glands viz; Seminal vesicle, prostate gland and Cowper's gland which is also called as Bulbourethral gland.
- It has Accessory ducts. Those are as follows;
 1. Seminiferous tubule.
 2. vasa deferentia.
 3. Epididymis.
 4. Ejaculatory duct.
 5. urethra.



- Accessory glands from where "semen" production occurs.

① Seminal vesicle: → 60-65% of semen produces from here only.

→ Its pH is 7.4 i.e Alkaline.

→ It is whitish in colour.

→ It contains Fructose for energy and Fibrogin for clotting of sperm in vagina. and prostaglandin for reverse peristalsis.

② prostate Gland: → 20-25% of semen produces from prostate gland.

→ It is acidic in Nature. because its pH is 6.8.

→ Contains calcium ions (Ca^{++}), zinc, Acid phosphates, citric acid. They are present to neutralize the acidic medium of vagina.

→ It is yellowish in colour.

③ Cowper's gland or Bulbourethral gland:

→ 10-15% of semen produces from here.

→ Its pH is 7.8 to neutralize the acidic medium of male's urethra.

→ It is watery in colour.


→ lubrication for smooth copulation.

* Semen = Sperm + Seminal plasma.

- All the Secretion of male reproductive system from accessory glands.


- For 1 Time Ejaculation.
- Male produces 2-3 ml of Semen.
- In which 60% should be normal in size. And 40% should be vigorously Active.
- In 2-3 ml of Semen there is 200-300 million Sperm cells.

* Testes:



- Each seminiferous Tubule contain 1-4 Testicular lobule.
- Testes is covered by Tunica vaginalis, Tunica Albuginea and Tunica vascularis.
- Tunica albuginea is a thick white fibrous connective tissue called capsule of testis.
- Tunica vascularis is the inner layer.
- It consist of blood vessels and Connective Tissues.
- Facilitates blood Supply to the testes.
- Tunica vaginalis overlying tunica albuginea.
- There are 2 layers of Tunica vaginalis:
 1. visceral $\rightarrow$ overlies the Tunica albuginea.
 2. Parietal $\rightarrow$ lines the scrotal cavity.
- Thin Fluid layer separates the 2 sections and reduces friction betⁿ the testes and Scrotum.

- Scrotum:



Q: Name the layers of tissue present in ureter.

→ The ureter has 3 layers;

- Mucosa (Innermost layer)
- Muscularis (Middle layer)
- Adventitia (Outer layer)