

Wood Surface Defects

Dataset Root

Total Images	Classes	Annotations
4,000	8	9,211

Qtz	
Quartzity	
Mineral deposits in wood grain	
Before	After
116	445
Gain: +329	
MINORITY	

LK	
Live_Knot	
Living branches bedded in wood	
Before	After
3242	3969
Gain: +727	
MAJORITY	

Mar	
Marrow	
Soft central core of wood	
Before	After
166	455
Gain: +289	
MINORITY	

Res	
Resin	
Resin pockets or gum spot	
Before	After
541	682
Gain: +141	
MAJORITY	

DK	
Dead_Knot	
Dead branches in wood	
Before	After
2372	2854
Gain: +482	
MAJORITY	

K+C	
Knot_with_crack	
Knot with radial crack	
Before	After
412	448
Gain: +36	
MAJORITY	

K-M	
Knot_missing	
Knot that has fallen out	
Before	After
97	420
Gain: +323	
MINORITY	

Crk	
Crack	
Split or crack in wood	
Before	After
410	574
Gain: +164	
MAJORITY	

Note: Minority classes (Quartzity, Marrow, Knot_missing) augmented via class-aware oversampling to match median (~412/class). Training set: 3,200 before to 3,981 after augmentation (+781 images).