

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/364947327>

# Teenage pregnancy and its predictors in Africa: A systematic review and meta-analysis

Article in International Journal of Health Sciences · October 2022

---

CITATIONS

6

READS

1,007

7 authors, including:



Addis Eyeberu

Haramaya University

83 PUBLICATIONS 242 CITATIONS

[SEE PROFILE](#)



Tamirat Getachew

Haramaya University

69 PUBLICATIONS 314 CITATIONS

[SEE PROFILE](#)



Addisu Sertsu

Haramaya University

34 PUBLICATIONS 96 CITATIONS

[SEE PROFILE](#)



Mekonnen Sisay

University of Gondar

39 PUBLICATIONS 627 CITATIONS

[SEE PROFILE](#)

# Teenage pregnancy and its predictors in Africa: A systematic review and meta-analysis

Addis Eyeberu<sup>1</sup>,  
 Tamirat Getachew<sup>1</sup>,  
 Addisu Sertsu<sup>1</sup>,  
 Mekonnen Sisay<sup>2</sup>,  
 Yohannes Baye<sup>1</sup>,  
 Adera Debella<sup>1</sup>,  
 Addisu Alemu<sup>3</sup>

<sup>1</sup>School of Nursing and Midwifery, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia, <sup>2</sup>School of Pharmacy, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia,  
<sup>3</sup>School of Public Health, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia

**Address for correspondence:**

Addis Eyeberu, School of Nursing and Midwifery, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia.  
 E-mail: addis7373@gmail.com

## ABSTRACT

**Objective:** Although teenage pregnancy has declined in the last decade, it remains a major public health issue in Africa. Maternal mortality is common among teenagers due to their increased risk of obstetric and medical complications. In Africa, there is a lack of robust and comprehensive data on the prevalence and predictors of teenage pregnancy. As a result, this systematic review and meta-analysis were carried out to summarize evidence that will assist concerned entities in identifying existing gaps and proposing strategies to reduce teenage pregnancy in Africa.

**Methods:** The review is registered by the international prospective register of systematic reviews (CRD42021275013). This search included all published and unpublished observational studies written in English between August 23, 2016, and August 23, 2021. The articles were searched using databases (PubMed, CINHAL [EBSCO], EMBASE, POPLINE, Google Scholar, DOAJ, Web of Sciences, MEDLINE, Cochrane Library, and SCOPUS). Data synthesis and statistical analysis were conducted using STATA version 14 software. Forest plots were used to present the pooled prevalence and odds ratio (OR) with a 95% confidence interval (CI) of meta-analysis using the random effect model.

**Results:** A total of 43,758 teenagers (aged 13–19) were included in 23 studies. In Africa, the overall pooled prevalence of teenage pregnancy was 30% (95% CI: 17–43). Western Africa had the highest prevalence of teenage pregnancy 33% (95% CI: 10–55). Age (18–19) (OR = 2.99 [95% CI = 1.124–7.927]), wealth index (OR = 1.84 [95% CI = 1.384–2.433]), and marital status (OR = 6.02 [95% CI = 2.348–15.43]) were predictors of teenage pregnancy in Africa.

**Conclusion:** In Africa, nearly one-third of teenagers become pregnant. Teenage pregnancy was predicted by age (18–19), wealth index, and marital status. Strengthening interventions aimed at increasing teenagers' economic independence, reducing child marriage, and increasing contraceptive use among married teenagers can help to prevent teenage pregnancy.

**WEBSITE:** ijhs.org.sa

**ISSN:** 1658-3639

**PUBLISHER:** Qassim University

**Keywords:** Africa, meta-analysis, predictors, systematic review, Teenage pregnancy

## Introduction

Teenage pregnancy is the occurrence of pregnancy within the ages of 13–19 years.<sup>[1]</sup> It is a global health issue that bears many sequelae of serious health, and social and economic problems for individuals, families, and communities.<sup>[2]</sup> Every year, approximately 21 million girls aged 15–19 years living in the developing countries become pregnant; nearly half of all teenage pregnancies are unintended, and more than half end in abortion.<sup>[3]</sup> The Centers for Disease Control and Prevention also reported that among mothers aged 15–19 in the US, 72.7% reported that they had an unintended pregnancy that resulted in a live birth.<sup>[4]</sup>

Teenage pregnancy is a major health issue in Africa. Even though teenage pregnancy has declined in the past few decades, it remains a health concern and priority for many African countries. There are few studies assessing the overall burden of teen pregnancy in African countries. Those that have been conducted showed that the prevalence of teen pregnancy was between 18% in Kenya to 29% in Malawi and Zambia.<sup>[5]</sup> Further, a study done in South Africa found that nearly half of female teens in Africa experienced an unintended pregnancy, with 26.3% of those unintended pregnancies ending in abortions.<sup>[6]</sup>

Teenage pregnancy is related to maternal mortality and morbidity. A study done in Nigeria showed that 33% of

maternal mortality attributes to teenage pregnancy.<sup>[7]</sup> Teenagers faced a higher risk for eclampsia, anemia, hemorrhage, cephalopelvic disproportion, prolonged labor, and cesarean section.<sup>[8]</sup> Teenagers are also more likely to develop obstetrical and medical complications such as preterm deliveries, iron deficiency anemia, and perineal tears.<sup>[9]</sup> It is suggested that increasing contraceptive use can prevent unwanted teenage pregnancies.<sup>[10]</sup> However, less than half (44.7%) of teenage mothers aged 15–19 who had an unintended pregnancy reported using contraception.<sup>[3]</sup> Similarly, contraceptive use remained low in Sub-Saharan Africa (18.87%).<sup>[11]</sup> In Nigeria, less than half (45.3%) of sexually active teenage girls used a modern contraceptive method.<sup>[12]</sup> Contraception is used by 31.8% of Ethiopian teenagers.<sup>[13]</sup> Meeting the unmet need for modern contraception among teenagers would prevent 6 million unintended pregnancies per year, averting 2.1 million unplanned births, 3.2 million abortions, and 5600 maternal deaths.<sup>[3]</sup>

Teenage pregnancy is also associated with perinatal health issues. Teenage mothers developed more adverse perinatal complications, such as preterm births, stillbirths, neonatal deaths, and delivered low birth weight babies.<sup>[14,15]</sup> Newborns born to teen mothers were admitted to the neonatal intensive care unit due to low birth weight, sepsis, respiratory distress syndrome, hyperbilirubinemia, and asphyxia.<sup>[11-13,16]</sup> Teenage pregnancy is needed to be tackled to prevent perinatal health issues.

Preventing teenage pregnancies is Africa's top priority. Investing in teenagers' health allows them to grow into healthy adults who can contribute positively to society. It is critical to reduce the burden of teenage pregnancies in the developing countries to reduce its sequels. As a result, reducing and preventing teenage pregnancies are a key component of sustainable development goal 3.<sup>[17]</sup> Furthermore, African countries such as Ethiopia have several policies aimed at improving adolescent health and preventing teen pregnancy.

Identifying the predictors of teenage pregnancy is important for planning appropriate intervention. Even though there are related studies conducted in Africa, they are inconsistent and not inclusive that illustrating the full degree of the larger impact. Furthermore, they are not recent pieces of evidence<sup>[18-20]</sup> and they did not use meta-analysis methods to pool the prevalence and predictors of teenage pregnancy.<sup>[18,19,21]</sup> Therefore, this systematic review and meta-analysis aimed to estimate the pooled prevalence of teenage pregnancy and summarize the predictors of teenage pregnancy in Africa. The summarized evidence obtained from this study could help interested entities to identify existing gaps and propose strategies to reduce teenage pregnancy in Africa.

## Methods

### Protocol and registration

This review was based on the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guideline [Additional File 1].<sup>[22]</sup> The review is registered by the international prospective register of systematic reviews (CRD42021275013).

### Eligibility criteria

Articles that met the following inclusion criteria were included in the systematic review and meta-analysis. (i) Study design: Observational studies (cross-sectional, case-control, and cohort studies); (ii) populations: Teen girls (13–19 years) residing in one of the countries in Africa; (iii) study setting: Institutional or community-based studies in rural or urban settings in Africa; (iv) language: Studies published in English; and (v) time: Studies published in the selected databases between August 23, 2016, and August 23, 2021. The duration of the study was 5 years.

Studies published before August 23, 2016, interventional studies, reviews, commentaries, editorials, case series/reports, conference abstracts, personal opinions, non-teenage participants, and patient stories were excluded from the study.

### Information sources and search strategy

Published articles were searched on the major databases (namely, PubMed, CINHAL (EBSCO), EMBASE, POPLINE, Google Scholar, DOAJ, Web of Sciences, MEDLINE, Cochrane Library, SCOPUS, Joanna Briggs Institute EBP (JBI EBP), Global Index Medicus, African journals online, Google search, and MedNar) using a combination of Boolean logic operators (AND, odds ratio (OR), and NOT), Medical Subject Headings (MeSH), and keywords.

The articles were searched using a combination of search terms (AND, OR, and NOT) Boolean (Search) operators. The search strategy includes ("teenage pregnancy "[Title/Abstract] AND "Africa "[MeSH Terms]), ("teenage pregnancy "[exactly on title] AND "Africa "[any field]), and ("teenage pregnancy "[all field] AND "Africa "[all filed]). Non-electronic sources are used combined with a direct Google search, Google Scholar, and MedNar. The search was also conducted by combining the aforementioned search terms with the names of all African countries. Furthermore, the investigators manually searched for gray literature and other relevant data sources such as email and unpublished thesis/papers with planned coverage dates. The CINHAL search strategy is outlined in Additional File 2.

Then, across all databases, all identified keywords and index terms were checked. Finally, all identified articles' reference lists were searched for further articles.

## Study selection

All search articles were exported to the EndNote X8 citation manager and duplicate studies were excluded. Then, these were screened carefully by reading the title and abstract. The authors (AE, TG, AS, and MS) screened the titles and abstracts of the identified articles using the eligibility criteria. Then, the full-text articles available in English were further evaluated based on objectives, methods, population, and key findings (prevalence/magnitude and predictors/determinants/factors associated with teenage pregnancy). The overall study selection process is presented using the PRISMA statement flow diagram<sup>[23]</sup> [Figure 1].

## Data extraction

After the selection of appropriate articles, the authors (AE, TG, AS, MS, YB, AA, and AD) independently extracted the data. A pre-defined Microsoft Excel 2016 format was used to extract the data from selected studies under the following heading: Author and year, region, country of study, setting, study design, sample size, study subject, data collection methods, the primary outcome of interest, and predictors of teenage pregnancy. The accuracy of the data extraction was verified by comparing the results of the two independent extracted data.

The quantitative data (the total sample size (N) and the frequency of the event (n) (number of teenagers who were pregnant), and effect size of predictors of teenage pregnancy) were extracted from the included articles and summarized using Microsoft Excel 2016 for meta-analysis and synthesis.


## Measurement and data items

Teenage pregnancy is measured as the “percentage of women aged 13–19 who have given birth or are pregnant with their first child.”<sup>[24-31]</sup> The prevalence of teenage pregnancy can also be measured as the percentage of pregnant teen women from all women in institutions,<sup>[32]</sup> who attended health institutions for delivery services<sup>[15,33]</sup> or antenatal care services<sup>[34]</sup> during a specific period. Therefore, this review included all studies which used either of the above definitions.

The wealth index is a composite measure of a household’s cumulative living standard. It is calculated using easy-to-collect data on a household’s ownership of selected assets, such as televisions and bicycles; materials used for housing construction; and types of water access and sanitation facilities.<sup>[35-37]</sup> The wealth index is categorized as low versus high.

Residency is referred to as a place in which a teenager resides. It was classified as rural versus urban. Teenagers who had a history of marital status are categorized as ever married versus not married. Teenagers currently married or divorced or previously married or separated were categorized as ever married. Family planning utilization is also classified as contraceptive user versus contraceptive nonuser.

Educational status is defined as the status of teenagers in education or training defined in law or legislation. It was classified as having no education, completed primary education, completed secondary education, and completed college and above education.



**Figure 1:** PRISMA flow diagram shows the selection process of included articles for systematic review and meta-analysis

## Risk of bias

The authors critically evaluated the risk of bias from individual studies. To minimize the risk of bias, a comprehensive search (electronic/database search and manual search) for published, unpublished, institutional, or community-based studies was carried out. The authors work cooperatively to reduce bias. All worked together in setting a schedule for the selection of articles based on the clear objectives and eligibility criteria, deciding the quality of the articles, regularly evaluating the review process, and extracting and compiling the data. Bias was explored using different statistical tests.<sup>[38]</sup>

The methodological reputability and quality of the findings of the included studies were critically evaluated using the quality assessment tool for observational studies (cross-sectional, case-control, and cohort studies) developed by the Joanna Briggs Institute.<sup>[39]</sup> The two groups of authors, Group 1 (TG, AS, AD, and MS) and Group 2 (AE, AA, and YB), independently evaluated the quality of the studies. The mean score of the two groups was taken for a final decision. The differences in the inclusion of the studies were resolved by consensus. The included studies were evaluated against each indicator of the tool and categorized as high, moderate, and low quality. A high-quality score is above 80%, moderate-quality is 60–80%, and low-quality is below 60%. Studies with a score greater than or equal to 60% were included. This critical appraisal was conducted to assess the internal validity (systematic error) and external validity (generalizability) of the studies and to reduce the risk of biases [Additional File 3].

## Statistical analysis

Data synthesis and statistical analysis were conducted using STATA 14 software. Forest plots were used to show the prevalence of teenage pregnancy in Africa. The random effect model of analysis was adopted as a method of meta-analysis because it reduces the heterogeneity of included studies. Subgroup analyses were also conducted by different study characteristics such as sub-regions of Africa (Northern, West, East, South, and Central Africa), study type (community based or institution based), and publication year. Moreover, the meta-analysis regression was conducted to identify the sources of heterogeneity among studies. The predictors of teenage pregnancy were presented using ORs with a 95% confidence interval (CI).

A meta-analysis of observational studies was carried out, based on the recommendations of the  $I^2$  statistic described by Higgins and Thompson (an  $I^2$  of 75/100% and above suggesting considerable heterogeneity).<sup>[40]</sup>  $P$ -value for  $I^2$  statistics  $<0.05$  was used to determine the presence of heterogeneity.

The investigators checked for potential publication bias through visual inspection of a funnel plot and Egger's regression test. Publication bias was assumed for  $P < 0.05$ . The results of the review were reported according to the PRISMA guidelines.

The findings of the included studies were first presented using a narrative synthesis and followed by a meta-analysis chart.

## Results

### Description of review studies

A total of articles 5418 were identified through the major medical and health electronic databases and other relevant sources. From all identified studies, 251 articles were removed due to duplication while 5167 studies were reserved for further screening. Of these, 5095 were excluded after being screened according to titles and abstracts. Of the 72 remaining articles, 49 studies were excluded due to studies did not present the outcome of interest, being qualitative studies, and non-teenage populations. Finally, 23 studies that fulfilled the eligibility criteria were included in the systematic review and meta-analysis [Figure 1].

### Characteristics of included studies

Of all studies, more than half 14 (60.9%) of studies were from East Africa.<sup>[10,24,26-30,41-48]</sup> Seven studies were from West Africa,<sup>[32-34,48-51]</sup> one study from North Africa,<sup>[15]</sup> and one study from South Africa.<sup>[25]</sup> The majority 18 (78.3%) of the included studies were cross-sectional studies,<sup>[10,15,25-29,32-34,42-47,50,51]</sup> while four studies were case-control studies<sup>[24,30,41,49]</sup> and one study was a mixed study.<sup>[48]</sup> The sample size of the included studies ranged from a minimum of 161 in a study conducted in Nigeria<sup>[50]</sup> to a maximum of 17,234 in a study conducted in East African countries.<sup>[47]</sup> A total of 43,758 teenagers (13–19 years) were included in the 23 studies. Thirteen of the included studies reported both prevalence and predictors of teenage pregnancy.<sup>[10,25,26,28,29,32,34,42-44,46,47,51]</sup> Five studies reported only the prevalence of teenage pregnancy<sup>[15,33,45,48,50]</sup> and the other five studies only report the predictors of teenage pregnancy.<sup>[24,27,30,41,49]</sup> The general characteristics of the studies selected for the meta-analysis are outlined in Table 1.

### Prevalence of teenage pregnancy

The pooled prevalence of teenage pregnancy ranged from 1% to 80% in studies done in Uganda and Ethiopia, respectively.<sup>[42,43]</sup> The overall pooled prevalence of teenage pregnancy in Africa was 30% (95% CI: 17–43) [Figure 2].

According to subgroup analysis by study region, West African countries had the highest prevalence of teenage pregnancy (33%, 95% CI: 10–55), while North African countries had the lowest prevalence of teenage pregnancy (26%, 95% CI: 24–29) [Figure 3].

Based on subgroup analysis by setting in which studies are conducted, the highest prevalence (30%, 95% CI: 13–47) of teenage pregnancy was seen in studies conducted in community-based settings, while the prevalence of teenage pregnancy among institutional-based studies was 28% (95% CI: 15–41) [Figure 4].

**Table 1:** General characteristics of the studies selected for the systematic review and meta-analysis in Africa, 2021



| S. No. | Author and years                         | Country  | Setting           | Design            | Data collection method     | Sample size | The primary outcome of interest                        | Prevalence (%) | Factors  |
|--------|--|----------|-------------------|-------------------|----------------------------|-------------|--|----------------|--|
| 1.     | Ayele et al., 2018 <sup>[41]</sup>       | Ethiopia | Community based   | Case – control    | Interview                  | 414         | Determinants of teenage pregnancy                      |                | Lower monthly income below \$25, lower monthly income below \$25–50, aged 18–19 years, being married, not communicating with parents on reproductive health issues, and having a history of maternal teenage pregnancy |
| 2.     | Birhanu et al., 2019. <sup>[42]</sup>    | Ethiopia | Community based   | Cross - sectional | DHS chart review           | 2679        | Teenage pregnancy                                      | 79.6           | Being sexually active before age 15, being married before age 15, rural residency, and lower contraceptive users   |
| 3.     | Asare et al., 2019 <sup>[49]</sup>       | Ghana    | Community based   | Case – control    | Interview                  | 245         | Factors associated with adolescent pregnancy           |                | Place of residence (urban), occupation (unemployed), and economic status (low)   |
| 4.     | Byonanebye et al., 2020 <sup>[43]</sup>  | Uganda   | Community based   | Cross - sectional | DHS chart review           | 2026        | Risk factors of teenage pregnancy                      | 8.1            | Marital status (married), living with a partner or separated, wealth status (poor), rural residency, wealth status (middle), marital status- living with the parent, and marital status-separated                      |
| 5.     | Donatus et al., 2018 <sup>[34]</sup>     | Cameron  | Institution based | Cross - sectional | Interview                  | 293         | Factors associated with teenage pregnancy              | 60.75          | Low contraceptive use, socioeconomic status, and physical violence   |
| 6.     | Bedzo and Manortey, 2019 <sup>[51]</sup> | Ghana    | Community based   | Cross - sectional | Interview                  | 223         | Prevalence and predictors of teenage pregnancy         | 43.95          | Older age (age group 18–19), teenagers being influenced by social media activities, and teenagers being influenced by their peers  |
| 7.     | Gabriel et al., 2020 <sup>[32]</sup>     | Cameron  | Community based   | Cross - sectional | Self-administered          | 250         | Prevalence and associated factors of teenage pregnancy | 5.2            | Multiple partners  |
| 8.     | Kassa et al., 2021 <sup>[44]</sup>       | Ethiopia | Community based   | Cross - section   | Interview                  | 343         | Prevalence and associated factors of teenage pregnancy | 25.4           | Residence urban, early age at 1 <sup>st</sup> sexual intercourse, not knowing family planning methods, not utilizing contraception, and did not know a place where contraceptives are provided                         |
| 9.     | Kefale et al., 2020 <sup>[27]</sup>      | Ethiopia | Community based   | Cross - sectional | DHS chart review           | 3381        | Factors associated with teenage pregnancy              |                | Age (being 17, 18, and 19 years old), uneducated, primary educated, being married, and communities with a higher proportion of poor  |
| 10.    | Kawo et al., 2019 <sup>[26]</sup>        | Ethiopia | Community based   | Cross - sectional | Chart review/ EDHS         | 2252        | Determinants of teenage pregnancy                      | 16.3           | Age at first sexual intercourse, marital status (married), education level (secondary), wealth index (middle), and contraceptive non-use   |
| 11.    | Menon et al., 2018 <sup>[48]</sup>       | Zambia   | Community based   | Mixed design      | Interview, FGD, IDIS, KIIS | 1,434       | Prevalence of teenage pregnancy and child marriage     | 48             |  |


(Contd...)

**Table 1:** (Continued)

| S. No. | Author and years                             | Country                | Setting           | Design            | Data collection method | Sample size | The primary outcome of interest                           | Prevalence (%) | Factors   |
|--------|--|------------------------|-------------------|-------------------|------------------------|-------------|---|----------------|---|
| 12.    | Mezmur et al., 2021 <sup>[29]</sup>          | Ethiopia               | Community based   | Cross - sectional | Interview              | 2258        | Teenage pregnancy   | 30.2           | Age (16–17 and 17–18 years old), not being in school, lack of formal education, being married, parental divorce, having an elder sister who had a history of teenage pregnancy, and not knowing the fertile period in the menstrual cycle |
| 13.    | Indongo, 2020 <sup>[25]</sup>                | Namibia                | Institution based | Cross - sectional | Chart review           | 1857        | Factors influencing teenage pregnancy                     | 20.4           | Age, age at first sex, education level, and household wealth status   |
| 14.    | Ochen et al., 2019 <sup>[30]</sup>           | Uganda                 | Community based   | Case – control    | Interview              | 495         | Predictors of teenage pregnancy                           |                | Having not had multiple sexual partners, frequent sex of less than 2, contraceptive use, and not being married  |
| 15.    | Onwubuariri and Kasso, 2018. <sup>[33]</sup> | Nigeria                | Institution based | Cross - sectional | Chart review           | 198         | Prevalence and predisposing factors                       | 1.5            |   |
| 16.    | Eldaboly et al., 2020. <sup>[15]</sup>       | Egypt                  | Institution based | Cross - sectional | Interview              | 848         | Prevalence of teenage pregnancy and its outcomes          | 26.4           |   |
| 17.    | Mathewos et al., 2018 <sup>[28]</sup>        | Ethiopia               | Institution based | Cross - sectional | Interview              | 578         | Prevalence of teenage pregnancy                           | 7.7            | Being Grade 11 student, Grade 12 students, not knowing the exact time to take emergency contraceptives, substance use, living with either biological parent, and poor parent-daughter interaction   |
| 18.    | Udoh et al., 2019. <sup>[50]</sup>           | Nigeria                | Institution based | Cross - sectional | Interview              | 161         | Prevalence of teenage pregnancy                           | 45.34          |   |
| 19.    | Omoro et al., 2017. <sup>[45]</sup>          | Kenya                  | Community based   | Cross - sectional | Interview              | 1592        | Teenage pregnancy   | 23.3           |   |
| 20.    | Uwizeye et al., 2020 <sup>[46]</sup>         | Rwanda                 | Community -Based  | Cross - Sectional | Chart review           | 2578        | Household-related factors associated with teen pregnancy  | 7.23           | Marital status, age of household head, household size, number of bedrooms given the size of the household, and educational level of the household head  |
| 21.    | Worku et al., 2021. <sup>[47]</sup>          | East African countries | Community based   | Cross - sectional | Chart review/ DHS      | 17, 234     | Prevalence and associated factors of adolescent pregnancy | 54.6           | Being age 18–19 years, using contraceptives, being employed girls, being spouse/head within the family, and being from higher community level contraceptive utilization   |
| 22.    | Habitu et al., 2018 <sup>[10]</sup>          | Ethiopia               | Community based   | Cross - sectional | Interview              | 514         | Prevalence of teenage pregnancy                           | 28.6           | Age, rural residence, contraceptive nonuse, and parental marital status (divorce)   |
| 23.    | Geda, 2019 <sup>[24]</sup>                   | Ethiopia               | Community based   | Case - control    | Chart review/ EDHS     | 1905        | Determinants of teenage pregnancy                         |                | Educational level, wealth index, knowledge of ovulatory cycle after period ended, knowledge of family planning method, contraceptive use, and reasons for not using contraceptives  |

DHS: Demographic health surveillance, EDHS: Ethiopia demographic health surveillance, FGD: Focused group discussion, IDIs: In-depth interviews, KIIs: Key informant interviews

**Figure 2:** The pooled prevalence of teenage pregnancy in Africa, 2021**Figure 3:** Subgroup analysis of the pooled prevalence of teenage pregnancy in Africa based on region, 2021

**Figure 4:** Subgroup analysis of the pooled prevalence of teenage pregnancy in Africa based on region, 2021

## Meta-regression to check the heterogeneity

A meta-regression analysis was conducted since there was statistically significant heterogeneity, with  $I^2$ -square test statistics  $<0.05$ . The goal of the analysis was to identify the source of heterogeneity so that the findings could be correctly interpreted. The meta-regression analysis, however, found no significant variables that could explain the heterogeneity. There were no statistically significant study level covariates: Sample size, publication year, or study setting. As a result, the heterogeneity can be explained by factors other than those discussed in this review [Table 2]. It may be a result of compiling different studies that have been done with different methodology.


## Publication bias

To observe publication bias, a visual inspection of the funnel plot was carried out [Figure 5]. It is asymmetric on the funnel plot. The prevalence rate among the various studies is different because the social realities are different and the study methodology also has been different, Egger's test showed that there is no small study effect publication bias ( $P = 0.134$ ).

## Predictors of teenage pregnancy in Africa

Age (below 15 versus 18–19)

A total of five studies were included to predict the association between age (18–19 years) and teenage pregnancy. Four studies

**Figure 5:** Funnel plot of prevalence of teenage pregnancy in Africa, 2021**Table 2:** Meta-regression analysis to check heterogeneity, 2021

| Variables        | Coefficients | SE        | P     |
|------------------|--------------|-----------|-------|
| Publication year | -0.0145666   | 0.0468326 | 0.760 |
| Sample size      | 0.0000159    | 0.0000141 | 0.276 |
| Setting          | -0.0228503   | 0.1177025 | 0.849 |

included in this meta-analysis.<sup>[27,28,41,47]</sup> found a significant association between age and teenage pregnancy while the one article<sup>[32]</sup> found an insignificant association with teenage

pregnancy. The pooled meta-analysis showed that there was a significant association between age (18–19 years) and teenage pregnancy. Teenagers whose age is 18–19 years were almost 3 times more likely to be pregnant than teenagers below the age of 15 ( $OR = 2.99$ , 95% CI = 1.124–7.927,  $P = 0.00001$ ). The heterogeneity test indicated  $I^2 = 91.4\%$ , and the random model was employed for analysis [Figure 6].

### Wealth index (low vs. higher)

In this meta-analysis, five studies were included to see the association between wealth index and teenage pregnancy. Three articles included in this meta-analysis<sup>[24,43,49]</sup> were found significantly associated with teenage pregnancy, while two articles<sup>[26,27]</sup> found an insignificant association with teenage pregnancy. The pooled meta-analysis showed that there was a significant association between wealth index and teenage pregnancy. Teenagers who had a low wealth index were 1.84 times more likely to become pregnant than teenagers who had a higher wealth index ( $OR = 1.84$ , 95% CI = 1.384–2.433) [Figure 7].

### Place of residency (urban vs. rural)

In this meta-analysis, six studies were included to see the association between residency and teenage pregnancy. Three of the included articles<sup>[10,42,43]</sup> were found significantly associated with teenage pregnancy, while three of the articles<sup>[24,27,47]</sup> found an insignificant association with teenage pregnancy. The pooled

meta-analysis showed that there was no significant association between residency and teenage pregnancy ( $OR = 1.35$ , 95% CI = 0.85–2.15). A random model was employed for analysis [Figure 8].

### Marital status (married vs. not married)

In this meta-analysis, nine studies were included to see the association between marital status and teenage pregnancy. Six of the included articles<sup>[26,27,41,43,45,46]</sup> were found significantly associated with teenage pregnancy, while three articles<sup>[10,28,49]</sup> found an insignificant association with teenage pregnancy. The pooled meta-analysis showed that there was a significant association between marital status and teenage pregnancy. Teenagers who were married were 6 times more likely to become pregnant than teenagers who were not married ( $OR = 6.02$ , 95% CI = 2.348–15.43) [Figure 9].

### Educational status (primary educational status vs. no educational status)

A total of five studies were included to predict the association between educational status and teenage pregnancy. Four of the included studies<sup>[24,26,27,47]</sup> found a significant association between educational status and teenage pregnancy while the rest of one article<sup>[10]</sup> found an insignificant association with teenage pregnancy. The pooled meta-analysis showed that there was no significant association between educational status (primary educational status vs. no educational status)




Figure 6: Association between age and teenage pregnancy in Africa, 2021




Figure 7: Association between wealth index and teenage pregnancy in Africa, 2021

and teenage pregnancy ( $OR = 1.2$  (95% CI = 0.57–2.51,  $P = 0.00001$ ) [Figure 10].


### Contraceptive non-use

In this meta-analysis, six studies were included to see the association between contraceptive non-use and teenage pregnancy. Three of the included articles<sup>[10,26,44]</sup> were found significantly associated with teenage pregnancy, while three of the articles<sup>[27,42,49]</sup> found an insignificant association with teenage pregnancy. The pooled meta-analysis showed that there was no significant association between contraceptive non-use and teenage pregnancy ( $OR = 1.66$  (95% CI = 0.747–3.69). A random model was employed for analysis [Figure 11].


## Discussion

This comprehensive study provides valuable information on the prevalence and predictors of teenage pregnancy in Africa. In this review, a total of 23 studies done in different parts of Africa were included. Age, marital status, and wealth index were significantly associated with teenage pregnancy. Teenage pregnancy is a health issue because it poses maternal and perinatal risks. Maternal mortality is highest among teenagers as well as infant and child mortality rates were high among children born to teenage mothers.<sup>[52]</sup>


Evidence showed that low-income countries had the highest prevalence of teenage pregnancies.<sup>[2]</sup> Similarly, this study




**Figure 8:** Association between residency and teenage pregnancy in Africa, 2021



**Figure 9:** Association between marital status and teenage pregnancy in Africa, 2021



**Figure 10:** Association between educational status and teenage pregnancy in Africa, 2021



**Figure 11:** Association between contraceptive use and teenage pregnancy in Africa, 2021

pointed out that the prevalence of teenage pregnancy in Africa was high (30% [95% CI: 17–43]). The pooled prevalence of teenage pregnancy in Africa is higher than the pooled prevalence of teenage pregnancy in Latin America (6.5%), South (4.5%), and East Asia (0.7%).<sup>[53]</sup> The possible justification may associate with the unavailability of contraceptive services, poor attitude toward contraceptive, poor knowledge of teenagers on their sexual health, and high prevalence of sexual assault and sexual violence in the low income country than in developed countries.<sup>[20,54,55]</sup> A similar study conducted in sub-Saharan African countries showed that the prevalence of teenage pregnancy was high.<sup>[56]</sup> Teenage pregnancy may be more prevalent due to a lack of access to contraceptive services, teenagers' limited decision-making capacity, the community's negative attitude toward contraceptive use, and peer pressure.<sup>[57-61]</sup> To alleviate the burden of teenage pregnancies, it is strongly recommended to involve prominent community leaders who play a key role in preventing early pregnancy in culturally acceptable ways. In addition, well-integrated sexuality-related education is linked to a contraceptive provision in the curriculum, as well as the formulation of laws and policies to improve access to contraceptive services.<sup>[52,62,63]</sup>

In this study, teenagers aged 18–19 were nearly 3 times more likely to become pregnant than teenagers under the age of 15. This could be because the majority of teenagers become sexually active when they reach the age of 18. This may also be related to the fact that child marriage at the age of 18 is common (28%) in developing regions, which predisposes teenagers to pregnancy.<sup>[3]</sup> This association between age and pregnancy has an implication for policymakers to work on setting strategies for preventing child marriage to reduce the burden of teenage pregnancy.

According to this study, a teenager who was married was 6 times more likely to become pregnant than teenagers who were not married. This finding was supported by evidence that showed that marriage was one of the predictors of teenage pregnancy.<sup>[52]</sup> This could be due to cultural pressure and peer

pressure to have a child soon after marriage. Another reason could be that teenagers lacked knowledge and access to sexual and reproductive health services, including family planning, making them more likely to become pregnant.<sup>[64,65]</sup> The majority of 90% of all teenage births in Africa occur within marriage. In Africa, the proportion of teenage marriage was 54% and it is leveled as high compared to other worlds.<sup>[66]</sup> This results in the high prevalence of teenage pregnancy.<sup>[67,68]</sup> Furthermore, evidence showed that the median age for marriage in Africa was 15–19 which affects the children's holistic live.<sup>[69]</sup> Evidence recommended that strengthening existing efforts on contraceptive usage among teenagers can avert teenage pregnancy.<sup>[64]</sup> It is advisable to encourage political leaders, planners, and community leaders to formulate and enforce laws and policies to prohibit the marriage of girls before 18 years of age.<sup>[52]</sup>

This study pointed out that teenagers who had a lower wealth index were 1.84 times more likely to become pregnant than teenagers who had a higher wealth index. This is in line with a systematic review and multilevel study done in sub-Saharan African countries which revealed that the wealth index is associated with teenage pregnancy and childbearing<sup>[70,71]</sup> The possible justification could be the improvement in the economic and cultural level makes the girl have more possibilities and life projects, decreasing the importance of traditional projects such as marriage and motherhood. Moreover, teenagers with higher wealth indexes are more likely to postpone marriage and early sexual activity. Evidence showed that teenagers who had a high wealth index had a lower desire to engage in sexual relationships with boys in exchange for cash and gifts, and a higher motivation to attend school.<sup>[72]</sup>

## Strengths and limitations

The investigators used extensive and comprehensive search strategies from multiple databases. Published, unpublished studies, and gray literature were included in the study. Studies were evaluated for methodological quality using a standardized tool. The literature search was systematic and assessed all related studies within the desired scope. It

appears that this is the first publication of a meta-analysis from across the African continent on the prevalence and risk factors involved with teenage pregnancy. It is possible that relevant publications, for example, publications reported in non-English language and local languages must have been missed. Studies with abstracts were the only ones included. This may affect the finding's inclusiveness. The observed heterogeneity of the study is also the drawback of this meta-analysis.

## Conclusion

Teenage pregnancy is still a major public health problem in Africa with a higher prevalence. Nearly a third of teenagers become pregnant in Africa. Age (18–19), wealth index, and marital status were the predictors of teenage pregnancy in Africa. Strengthening interventions that aimed to increase the economic independence of teenagers, decreasing child marriage, and contraceptive utilization among married teenagers can avert the problem of teenage pregnancy.

## Authors' Declaration Statements

### Ethics approval and consent to participate

Not applicable due to the nature of the study.

### Consent for publication

Not applicable.

### Availability of data and materials

The data that support the review findings of this study are available on submitting a reasonable request to the corresponding author.

### Competing Interests

The authors declared that have no competing interests.

### Funding

None.

### Authors' contributions

AE, TG, AS, and MS conceived and designed the review. AE\*, TG, and AS carried out the draft of the manuscript, and AE\* is the PI of the review. AE\*, TG, and AS developed the search strings. AE, TG, MS, YB, AD, and AA screened and selected studies. AS, MS, AA, AD, YB, and AE extracted the data and evaluated the quality of the studies. AE, YB, AD, AA, and MS carried out analysis and interpretation. AE, TG, AS, MS, YB, AD, and AA rigorously reviewed the manuscript. All authors read and approved the final version of the manuscript.

## Acknowledgments

We would like to thank the College of Health and Medical Sciences, Haramaya University, for the non-financial support.

## References

- UNICEF. Empowerment of Adolescent Girls and Preventing Teenage Pregnancies and Population Management and Planning in Developing Nations. New York: UNICEF; 2016.
- World Health Organization. Adolescent Pregnancy Evidence Brief-World Health Organization 2019. Geneva: World Health Organization; 2019.
- Darroch JE, Woog V, Bankole A, Ashford LS. Adding it up: Costs and Benefits of Meeting the Contraceptive Needs of Adolescents; 2016. Available from: <https://www.guttmacher.org/adding-it-up> [Last accessed on 2022 Oct 19].
- Centers for Disease Control and Prevention. Teen Pregnancy Prevention. United States: Centers for Disease Control and Prevention; 2009. Available from: <https://www.cdc.gov/winnablebattles/teenpregnancy/index.htm> [Last accessed on 2022 Oct 19].
- Wado YD, Sully EA, Mumah JN. Pregnancy and early motherhood among adolescents in five East African countries: A multi-level analysis of risk and protective factors. BMC Pregnancy Childbirth 2019;19:59.
- Ajayi AI, Ezege HC. Association between sexual violence and unintended pregnancy among adolescent girls and young women in South Africa. BMC Public Health 2020;20:1370.
- Adeyinka DA, Oladimeji O, Adekanbi TI, Adeyinka FE, Falope Y, Aimakhu C. Outcome of adolescent pregnancies in southwestern Nigeria: A case-control study. J Matern Fetal Neonatal Med 2010;23:785-9.
- Moraes AN, Likwa RN, Nzala SH. A retrospective analysis of adverse obstetric and perinatal outcomes in adolescent pregnancy: The case of luapula province, Zambia. Matern Health Neonatol Perinatol 2018;4:20.
- Lisonkova S, Haslam MD, Dahlgren L, Chen I, Synnes AR, Lim KI. Maternal morbidity and perinatal outcomes among women in rural versus urban areas. CMAJ. 2016;188(17-18):E456-E465.
- Habitu YA, Yalew A, Bisetegn TA. Prevalence and factors associated with teenage pregnancy, Northeast Ethiopia, 2017: A cross-sectional study. J Pregnancy 2018;2018:1714527.
- Abebe AM, Fitie GW, Jember DA, Reda MM, Wake GE. Teenage pregnancy and its adverse obstetric and perinatal outcomes at Lemlem Karl hospital, Tigray, Ethiopia, 2018. Biomed Res Int 2020;2020:3124847.
- Tamambang RF, Njim T, Njie AE, Mbuagbaw L, Mafuta A, Tchana M, *et al.* Adolescent deliveries in urban Cameroon: A retrospective analysis of the prevalence, 6-year trend and adverse outcomes. BMC Res Notes 2018;11:469.
- Eldaboly S, Allam N, Ibrahim M, Abo-Elhassan H. Prevalence and outcome of teenage pregnancy among attendants of labour room in Bassion general hospital-Egypt (cross section study). J Recent Adv Med 2021;2:166-72.
- Mukhopadhyay P, Chaudhuri RN, Paul B. Hospital-based perinatal outcomes and complications in teenage pregnancy in India. J Health Popul Nutr 2010;28:494-500.
- Papri FS, Khanam Z, Ara S, Panna MB. Adolescent pregnancy: Risk factors, outcome and prevention. Chattogram Maa Shishu Hosp Med Coll J 2016;15:53-6.
- Mezmur H, Assefa N, Alemayehu T. An increased adverse fetal outcome has been observed among teen pregnant women in rural

- Eastern Ethiopia: A comparative cross-sectional study. *Glob Pediatr Health* 2021;8:2333794x21999154.
17. World Health Organization. Progress Towards the SDGs: A Selection of Data from World Health Statistics 2018. Geneva: World Health Organization; 2018.
  18. Yakubu I, Salisu WJ. Determinants of adolescent pregnancy in sub-Saharan Africa: A systematic review. *Reprod Health* 2018;15:15.
  19. Sama CB, Ngasa SN, Dzekem BS, Choukem SP. Prevalence, predictors and adverse outcomes of adolescent pregnancy in sub-Saharan Africa: A protocol of a systematic review. *Syst Rev* 2017;6:247.
  20. Kassa GM, Arowojolu AO, Odukogbe AA, Yalew AW. Prevalence and determinants of adolescent pregnancy in Africa: A systematic review and Meta-analysis. *Reprod Health* 2018;15:195.
  21. Gunawardena N, Fantaye AW, Yaya S. Predictors of pregnancy among young people in sub-Saharan Africa: A systematic review and narrative synthesis. *BMJ Glob Health* 2019;4:e001499.
  22. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and metaanalyses: The PRISMA statement. *PLoS Med* 2009;6:e1000097.
  23. Stewart LA, Clarke M, Rovers M, Riley RD, Simmonds M, Stewart G, et al. Preferred reporting items for systematic review and meta-analyses of individual participant data: The PRISMA-IPD statement. *JAMA* 2015;313:1657-65.
  24. Geda Y. Determinants of teenage pregnancy in Ethiopia: A case-control study. 2019. *Curr Med Issues* 2019;17:112.
  25. Indongo N. Analysis of factors influencing teenage pregnancy in Namibia. *Med Res Arch* 2020;8. Doi:10.18103/mra.v8i6.2102.
  26. Kawo KN. Determinants of teenage pregnancy in rural Ethiopia. *J Health Med Nurs* 2019;68. Doi:10.7176/JHMN/68-02.
  27. Kefale B, Yalew M, Damtie Y, Adane B. A multilevel analysis of factors associated with teenage pregnancy in Ethiopia. *Int J Womens Health* 2020;12:785-93.
  28. Mathewos S, Mekuria A. Teenage pregnancy and its associated factors among school adolescents of Arba Minch Town, Southern Ethiopia. *Ethiop J Health Sci* 2018;28:287-98.
  29. Mezmur H, Assefa N, Alemayehu T. Teenage pregnancy and its associated factors in Eastern Ethiopia: A community-based study. *Int J Womens Health* 2021;13:267-78.
  30. Ochen AM, Chi PC, Lawoko S. Predictors of teenage pregnancy among girls aged 13-19 years in Uganda: A community based case-control study. *BMC Pregnancy Childbirth* 2019;19:211.
  31. World Health Organization. Adolescent Pregnancy. Geneva: World Health Organization; 2004.
  32. Gabriel ES, Halle-Ekane GE, Isah M, Wirsiy FS, Lebaga MX, Siysila ND, et al. Prevalence and associated factors of teenage pregnancy among secondary and high school students in the Tiko health District, South West Region, Cameroon. *J Biosci Med* 2020;8:99-113.
  33. Kasso T, Onwubuariri T. Teenage pregnancy: Prevalence, pattern and predisposing factors in a tertiary hospital, Southern Nigeria. *Asian J Med Health* 2020;2020:1-5.
  34. Donatus L, Sama DJ, Tsoka-Gwegweni JM, Cumber SN. Factors associated with adolescent school girl's pregnancy in Kumbo East health District North West region Cameroon. *Pan Afr Med J* 2018;31:138.
  35. Pirani E. Wealth Index. Encyclopedia of Quality of Life and Well-Being Research. Dordrecht: Springer; 2014.
  36. Rutstein SO, Staveteig S. Making the Demographic and Health Surveys Wealth Index Comparable. Vol. 9. Rockville, MD: ICF International Rockville; 2014.
  37. Gondwe KW, Walker RJ, Mkandawire-Valhmu L, Dressel A, Ngui EM, Kako PM, et al. Predictors of wealth index in Malawi-analysis of Malawi demographic Health Survey 2004–2015/16. *Public Health Pract (Oxf)* 2021;2:100059.
  38. Shil X, Nie C, Shi S, Wang T, Yang H, Zhou Y, Song X. Effect comparison between egger's test and begg's test in publication bias diagnosis in meta-analyses: Evidence from a pilot survey. *Int J Res Stud Biosci* 2017;5:14-20.
  39. Porritt K, Gomersall J, Lockwood C. JBI's systematic reviews: Study selection and critical appraisal. *Am J Nurs* 2014;114:47-52.
  40. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539-58.
  41. Ayele BG, Gebregzabher TG, Hailu TK, Assefa BA. Determinants of teenage pregnancy in Degua Tembien District, Tigray, Northern Ethiopia: A community-based case-control study. *PLoS One* 2018;13:e0200898.
  42. Birhanu BE, Kebede DL, Kahsay AB, Belachew AB. Predictors of teenage pregnancy in Ethiopia: A multilevel analysis. *BMC Public Health* 2019;19:601.
  43. Byonanebye J, Brazauskas R, Tumwesigye N, Young S, May T, Cassidy L. Geographic variation and risk factors for teenage pregnancy in Uganda. *Afr Health Sci* 2020;20:1898-907.
  44. Kassa BG, Belay HG, Ayele AD. Teenage pregnancy and its associated factors among teenage females in Farta Woreda, Northwest, Ethiopia, 2020: A community-based cross-sectional study. *Popul Med* 2021;3:1-8.
  45. Omoro T, Otieno G, Gray SC, Mbeda C, Phillips-Howard PA, Hayes TD, et al. Teen pregnancy in rural western Kenya: A public health issue. *Int J Adolesc Youth* 2017;23:1-10.
  46. Uwizeye D, Muhyiteto R, Kantarama E, Wiehler S, Murangwa Y. Prevalence of teenage pregnancy and the associated contextual correlates in Rwanda. *Heliyon* 2020;6:e05037.
  47. Worku MG, Tessema ZT, Teshale AB, Tesema GA, Yeshaw Y. Prevalence and associated factors of adolescent pregnancy (15-19 years) in East Africa: A multilevel analysis. *BMC Pregnancy Childbirth* 2021;21:253.
  48. Menon JA, Kusanthan T, Mwaba SO, Juanola L, Kok MC. "Ring" your future, without changing diaper-can preventing teenage pregnancy address child marriage in Zambia? *PLoS One* 2018;13:e0205523.
  49. Asare BY, Baafi D, Dwumfour-Asaree B, Adamad AR. Factors associated with adolescent pregnancy in the Sunyani municipality of Ghana. *Int J Afr Nurs Sci* 2019;10:87-91.
  50. Udoji SB, Iyanam VE, Uche NN, Udonwa NE. Teenage pregnancy: Family and social characteristics and risk factors in Etinan, SubUrban Area of South-South Nigeria. *SSRG Int J Med Sci* 2019;6:7-14.
  51. Bedzo JY, Manortey S. Factors influencing teenage pregnancy in the lower Manya Krobo municipality in the Eastern region of Ghana: A cross-sectional study. *Open Access Library J* 2019;6:1-17.
  52. World Health Organization. World Health Organization Guidelines on Preventing Early Pregnancy and Poor Reproductive Outcomes Among Adolescents in Developing Countries. Geneva: World Health Organization; 2011.
  53. United Nations Department of Economic and Social Affairs. SDG Indicators: Global Database. New York: United Nations Department of Economic and Social Affairs; 2017.
  54. McCurdy RJ, Schnatz PF, Weinbaum PJ, Zhu J. Contraceptive use in adolescents in Sub-Saharan Africa: Evidence from demographic and health surveys. *Conn Med* 2014;78:261-72.
  55. Darroch JE, Woog V, Bankole A, Ashford LS. Adding it up: Costs and Benefits of Meeting the Contraceptive Needs of Adolescents. New York: Guttmacher Institute; 2016.
  56. Ahinkorah BO, Kang M, Perry L, Brooks F, Hayen A. Prevalence of first adolescent pregnancy and its associated factors in sub-Saharan Africa: A multi-country analysis. *PLoS One* 2021;16:e0246308.

57. Krug JK, Mevissen FE, Flore KA, Ruiter RA. Girls cannot be trusted: Young men's perspectives on contraceptive decision making and sexual relationships in Bolgatanga, Ghana. *Eur J Contracept Reprod Health Care* 2018;23:139-46.
58. Machira K, Palamuleni ME. Health care factors influencing teen mothers' use of contraceptives in Malawi. *Ghana Med J* 2017;51:88-93.
59. Madlala ST, Sibya MN, Ngxongo TS. Perceptions of young men at the free state school of nursing with regards to teenage pregnancy. *Afr J Prim Health Care Fam Med* 2018;10:e1-7.
60. Marshall SA, Driver N, Allison MK. Attitudes towards contraception: focus groups with Arkansas teenagers and parents. *Sex Educ* 2021;21:161-75.
61. Mashia EO, Van Wyk NC, Leech R. Support of adolescents to resist peer pressure and coercion to sexual activity. *Int Nurs Rev* 2019;66:416-24.
62. Harden A, Brunton G, Fletcher A, Oakley A. Teenage pregnancy and social disadvantage: Systematic review integrating controlled trials and qualitative studies. *BMJ* 2009;339:b4254.
63. Chandra-Mouli V, Camacho AV, Michaud PA. WHO guidelines on preventing early pregnancy and poor reproductive outcomes among adolescents in developing countries. *J Adolesc Health* 2013;52:517-22.
64. Ahinkorah BO, Hagan JE Jr, Seidu AA, Sambah F, Adoboi F, Schack T, *et al.* Female adolescents' reproductive health decision-making capacity and contraceptive use in sub-Saharan Africa: What does the future hold? *PLoS One* 2020;15:e0235601.
65. Davids EL, Kredo T, Gerritsen AA, Mathews C, Slingers N, Nyirenda M, *et al.* Adolescent girls and young women: Policy-to-implementation gaps for addressing sexual and reproductive health needs in South Africa. *S Afr Med J* 2020;110:855-7.
66. Koski A, Clark S, Nandi A. Has child marriage declined in sub-Saharan Africa? An analysis of trends in 31 countries. *Popul Dev Rev* 2017;43:7-29.
67. Melesse DY, Cane RM, Mangombe A, Ijadunola MY, Manu A, Bamgbose E, *et al.* Inequalities in early marriage, childbearing and sexual debut among adolescents in sub-Saharan Africa. *Reprod Health* 2021;18:117.
68. Raj A, Saggurti N, Balaiah D, Silverman JG. Prevalence of child marriage and its effect on fertility and fertility-control outcomes of young women in India: A cross-sectional, observational study. *Lancet* 2009;373:1883-9.
69. Garenne M. Age at marriage and modernisation in sub-Saharan Africa. *Southern African J Demogr* 2004;9:59-79.
70. Magadi MA. Multilevel determinants of teenage childbearing in sub-Saharan Africa in the context of HIV/AIDS. *Health Place* 2017;46:37-48.
71. Odimegwu C, Mkwananzi S. Factors associated with teen pregnancy in sub-Saharan Africa: A multi-country cross-sectional study. *Afr J Reprod Health* 2016;20:94-107.
72. Milimo J, Zulu JM, Svanemyr J, Munsaka E, Mweemba O, Sandøy IF. Economic support, education and sexual decision making among female adolescents in Zambia: A qualitative study. *BMC Public Health*