1. The probability of a leap year selected at random contain 53					
Sunday is:					
(a) 53/ 366	(b) 1/7	(c) 2/7	(d) 53/365		
2. A bag contains	3 red and 2 b	olue marbles. A r	marble is drawn at		
random. The prob	ability of drav	wing a black ball	is:		
(a) 3/5	(b) 2/5	(9) 0/5	(d) 1/5		
3. The probability	that it will ra	in tomorrow is 0	.85. What is the		
probability that it					
			(d) none of these		
-	-		ed from the numbers		
(1, 2, 3,,15)	_				
(a) 1/5	(b) 4/5	(c) 2/15	(d) 1/3		
5. What are the to					
. ,	` '	(Z) 8	` '		
	-	number selecte	d at random from the		
numbers (1,2,3,					
			(d) none of these		
7. The sum of the	probability o	f an event and n	on event is :		
(a) 2	b) 1 (c)	0 (d) none	e of these.		
•		are given; choos	e the correct answer		
for that which is n			4.00		
(a) 0.15	(b) 2/7	(x) 7/5	(d) none of these.		
		nultaneously, tha	n the probability of		
getting at least tw	o heads, is:		(1) 4 (0)		
(a) 1/4	(b) 3/8	(C) ½	(d) 1/8		
10. A letter is cho					
♦ ASSASSINATIO					
(a) 6/13	(b) //13	(c) 1	(d) none of these.		
44 A Ji	Final Alexan				
	-		ting an even number.		
(A) 2/3	(B) I	(C) 5/6	D) 1/2		
12. Two coins are thrown at the same time. Find the probability of					
getting both heads		(D) 5			
(A) 3/4 (P) 1/4	(C) 1/2	(D) 0			
13. Two dice are thrown simultaneously. The probability of getting a					

1

sum of 9 is:

(A) 1/10	(B) 3/10	(2) 1/9	(D) 4/9				
14. 100 cards are numbered from 1 to 100. Find the probability of							
(A) 3/4	ime number. (B) 27/50	(Q) 1/4	(D) 2°	9/100			
	ontains 5 red ba a blue ball is do a bag is: (B) 10	uble that of a r		-			
taken out at non-defecti	f 600 bulbs contrained from to the result is:	his box. Then	the probabilit				
17. Cards marked with numbers 2 to 101 are placed in a box and mixed thoroughly. One card is drawn from this box randomly, then the probability that the number on card is a perfect square. (A) 9/100 (B) 1/10 (C) 3/10 (D) 19/100							
18. What is (A) 1/7	the probability (B) 53/366	of getting 53 (C) 2/7	Mondays in a (D) 7/3				
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13							
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312, then the probability that it will point to an odd number is: (A) 1/6 (B) 1/12 (C) 7/12 (D) 5/12							
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) 3/4 (B) 1/2 (C) 1 (D) 1/4							

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:							
	(B) 31/365	-	(D) 1/133225				
23. A number x is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that $x^2 < 2$ is? (A) $1/5$ (B) $2/5$ (C) $3/5$ (D) $4/5$							
a marble is dra red is 2/3, then		om the jar, the pr white marbles in t	d others are white. If obability that it is the jar is:				
Then the proba	is selected at ran ability that it is a 4/25 (C) 1/25	multiple of 3 and	0 natural numbers. I 4 is:				
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?							
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	$\frac{4}{21}$				
	_	_	nes are 50, 70, 82,				
	e standard deviat b) 25.49		d) 25.69				
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.							
a) 13, 15	b) 13, 18	c) 18, 15	d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is							
a) $\frac{1}{2}$ 30. X is a varia	b) $\frac{1}{3}$ ate between 0 and c) 7 c)	d 3. The value of	$\begin{array}{c} \text{d) } ^{1}/_{6} \\ \textbf{E(X^{2}) is } \underline{\hspace{1cm}} . \\ \begin{array}{c} 9 \end{array}$				
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?							

32.Out of probability	_	alues, which	one is not poss	ible in
		P(x) = 3		
c) $P(x) = 0$.	b) ∑ x 5 <mark>d) P(</mark>	(x) = -0.5		
	= 2 and E(z) = 4	•	•	CC at a state of
a) 2	b) 6	c) 0	a) ins	ufficient data
34.The cov	variance of two	independer	nt random variab	le is
a) 1	<u>(a) 0</u>	c) - 1	d) Un	defined
35.If Σ P(x a) 0	b) 1	, the value o		sufficient data
, ,	0.5 and x = 4, b) 0.5	then E(x) = 6 c) 4	? <mark>d) 2</mark>	
37.In a dis is always?	crete probabili	ty distributio	on, the sum of all	probabilities
a) 0	b) Infinite	c) 1	d) Und	lefined
38.If the p variance.	robability of hi	tting the tar	get is 0.4, find m	ean and
a) 0.4, 0.24	b) 0.6,	0.24	c) 0.4, 0.16	d) 0.6, 0.16
target is 60 a) 0.6, 0.24 40. Find th	o% and if 10 both by 6, 2.4 e mean of toss	ombs are dro 4 c) 0 sing 8 coins.		
a) 2 41. What i	s the mean and	c) 8 d variance fo	d) 1 or standard norm	al distribution?

c) 5

d) 7

a) 3 b) 4

a) Mean is 0 c) Mean is 0	and varia and varia	nce is 1 nce is ∞	b) Mea d) Me	n is 1 and an is ∞ aı	variance nd variand	is 0 ce is 0
42.Variance a) E(X)	e of a rand b) E(X			given by 2) - (E(X))		. d) (E(X))2
43.Mean of a random variable X is given by a) E(X)						
44.Mean of a a) 0	a constan by a	t 'a' is	c) a/2	_ ·	d) 1	
45.Variance of a constant 'a' is a) 0 b) a c) a/2 d) 1						
46.Find the I	mean and	variance	e of X?			
X	0	1	2	3	4	
f(x)	1/9	2/9	3/9	2/9	1/9	
a) 2, 4/3	b) 3	3, 4/3	(c) 2, 2/3		d) 3, 2/3
47. Find the expectation of a random variable X?						

	Х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) ().5	,	þ) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by



c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- (a) P(X = x) = nCx px qx
- \dot{b}) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2 $\sqrt[4]{\sqrt{npq}}$