

List 3: Discrete Fourier Transformation (DFT) and Fast Fourier Transformation (FFT)


Exercise 1 Discrete Fourier Transformation (DFT).


The spectra Xa[k] till Xe[k] were calculated with DFT. Match each spectrum with the corresponding discrete time functions from x1(t) till x6(t) and fill out the table below.

Spek- trum	time function	Why do these functions match each other?	N	T _s [s] Sampling time or tstep	F _s [Hz] Sampling frequency	f _{step} [Hz] frequency step	(N.T _s) time window ¹ [s]

Zurich University of Applied Sciences Page 1/4 SiSy_exer3_dft.docx


 $^{^{1}}$ The time window (N.Ts) is also called the observation window, which means how long values of the time function x(t) are sampled and stored, before being used to calculate the corresponding X[k] coefficients with the DFT.





Exercise 2 FFT and Sampling

The plot of a time continuous function x(t) is given below. The function x(t) should be sampled and its spectrum calculated with the FFT (algorithm for the implementation of the DFT).



- (a) How long must be the time window in order to get a spectrum with a frequency resolution (or frequency step) of f_{step} = 0.25Hz ?
- (b) What is the minimum value for the sampling frequency F_s according to the sampling theorem, which avoids aliasing effects?

Exercise 3 Discrete Fourier Transformation

The spectrum of a periodic square pulse was calculated with FFT and the amplitude values are plotted below. Determine the following parameters and justify your answer with a short statement.

- (a) The sampling frequency (F_s):
- (b) The lenght of the time window (N.T_s):
- (c) The period of the periodic square pulse (T_0) :
- (d) The width of the pulses in the time domain (tau):

