


L_p

xz -vlak (Eye)



$$I' = pA' + (1-p)B'$$

vgl L_p : $z = \frac{b}{a}x - b$ want $\begin{cases} z = -b \text{ in } x = 0 \\ x = a \text{ in } z = 0 \end{cases}$

geldig voor A, B en I :

$$\begin{cases} z_A = \frac{b}{a}x_A - b \Rightarrow \frac{x_A}{z_A} = \frac{a}{b} + \frac{a}{z_A} \\ z_B = \frac{b}{a}x_B - b \Rightarrow \frac{x_B}{z_B} = \frac{a}{b} + \frac{a}{z_B} \\ z_I = \frac{b}{a}x_I - b \Rightarrow \frac{x_I}{z_I} = \frac{a}{b} + \frac{a}{z_I} \end{cases}$$

Verder $x_I' = p x_A' + (1-p) x_B'$ (gegeven)

en $x_I' = \frac{dx_I}{-z_I}$, $x_A' = \frac{dx_A}{-z_A}$, $x_B' = \frac{dx_B}{-z_B}$

$\Rightarrow x_I' = \frac{da}{-b} + \frac{ad}{-z_I}$, $x_A' = \frac{ad}{-b} + \frac{ad}{-z_A}$, $x_B' = \frac{ad}{-b} + \frac{ad}{-z_B}$

$\Rightarrow \frac{1}{z_I} = p/z_A + (1-p)/z_B$