# Discrete Mathematics Week 5

Abeyah Calpatura

### 4.1

#### Exercises

Abeyah Calpatura #7, 13, 24, 28

#7 Solution: There exist real numbers a and b such that  $\sqrt{a+b} = \sqrt{a} + \sqrt{b}$ .

$$a = 16 \text{ and } b = 0$$
  
 $\sqrt{16 + 0} = \sqrt{16} + \sqrt{0}$   
 $4 = 4$ 

#13 Solution: For every integer n, if n is odd, then  $\frac{n-1}{2}$  is odd.

**Negation:** There exists an integer n such that n is odd and  $\frac{n-1}{2}$  is even.

Counterexample: n = 1

$$\frac{1-1}{2} = 0$$
 which is even.

**Conclusion:** The statement is false.

#24 Solution: For every real number x, if x > 1, then  $x^2 > x$ .

Quantification Implicit If x is a real number and x > 1, then  $x^2 > x$ . First sentence of proof: "Suppose x is a real number greater than 1." Last sentence of proof: " $x^2 > x$ "

#28 Solution:

- **a.**  $\forall$  integers m and n, if m and n are odd, then m+n is even, as  $\forall$  odd integers m and n, m+n is even, and as If m and n are any odd integers, then m+n is even.
- **b.** By **definition of odd**, m = 2r + 1 and n = 2s + 1 for some integers r and s.

Then

$$m + n = (2r + 1) + (2s + 1)$$
 by **substitution**

$$=2r+2s+2$$

$$= 2(r+s+1)$$
 by algebra

Let u = r + s + 1. Then u is an integer because r, s, and 1 are integers and because **definition of** 

## 4.2

## Exercises

Abeyah Calpatura #2, 5, 17, 20

#2 Solution: For every integer m, if m is even then 3m+5 is odd.

answer

#5 Solution: question

answer

#17 Solution: question

answer

#20 Solution: question

answer