

WebGPU Geant4-DNA: browser-native, kernel-fused Monte Carlo track-structure simulation for radiobiology, validated against Geant4 11.4.1

Ahmet Barış Günaydın
Independent researcher | webgpudna.com

June 2, 2026

Abstract

We present a WebGPU/WGSL re-implementation of Geant4-DNA track-structure physics and Karamitros Independent-Reaction-Time (IRT) radiolysis chemistry that runs entirely in a web browser, with no CUDA toolchain and no installation. The simulation uses a *kernel-fusion* architecture: one GPU thread per primary electron executes the full particle history inside a single fused compute dispatch, eliminating per-step dispatch overhead. We do not transpile Geant4; we re-implement its physics models from the Geant4-DNA source and the G4EMLOW 8.8 tabulated cross sections, and use a normally-compiled Geant4 11.4.1 as a validation oracle on a falsifiable, artifact-backed ladder. Cross sections match the reference to peak ratios of 0.975 to 1.000; the continuous-slowing-down range at 10 keV is $0.988\times$ Geant4; and the fused tracking is $455\times$ faster than Geant4 single-thread ($280\times$ vs. an 8-thread build) on an Apple M2 Pro. We also report the project’s honest limits: a 27 % cascade-ionisation deficit traceable to an intentional excitation cross-section choice, and the finding that the *inter-track* chemistry contribution—which closes most of the residual 1 μ s gap against Geant4’s `chem6`—is *not* reachable browser-native, because cross-primary IRT is $O(N^2)$ for a point source and GPU step-by-step chemistry is bounded by the encounter radius to $\sim 50\,000$ time steps. These negative results are quantified and motivate a native runtime as the path forward. All experiments are reproducible via committed JSON artifacts.

1 Introduction

Geant4-DNA [9, 2, 10] is the CNRS/IN2P3-coordinated extension of Geant4 [1] for discrete, event-by-event Monte Carlo track-structure simulation in liquid water at radiobiologically relevant (eV–keV) energies, together with the subsequent radiolysis chemistry. It is the de-facto reference for mechanistic modelling of radiation-induced DNA damage.

GPU acceleration of this class of simulation is an active area: gMicroMC [14] and MPEXS-DNA [13] target track-structure on the GPU, while GGEMS [6] and GPUMCD [7] target macroscopic dose. All of these are built on CUDA or OpenCL and require a workstation-class install. To our knowledge there is no open-source GPU Monte Carlo for radiation transport that runs *browser-native*—i.e. on the cross-vendor WebGPU API, with zero install, on any machine with a modern browser. That accessibility gap is the motivation here: a browser-native engine is trivially shareable, reproducible without a toolchain, and usable for teaching and rapid validation.

Our contributions are: (i) a complete WebGPU/WGSL re-implementation of the Geant4-DNA DNA_Opt2 physics chain plus Karamitros IRT chemistry; (ii) a kernel-fusion architecture that

[Figure placeholder: Phase A/B/C pipeline + buffer map]

Generate from ARCHITECTURE.md: Phase A one fused dispatch (1 thread/primary) writing the `rad_buf` and secondary buffer; Phase B wavefront over secondaries; Phase C IRT worker; atomic voxel grid for dose/radical deposition.

Figure 1: Three-phase pipeline. Phase A is fully fused; B and C cannot be.

yields large speedups over single- and multi-threaded Geant4; (iii) a research-grade, falsifiable validation ladder against compiled Geant4 11.4.1, with every claim backed by a committed artifact; and (iv) an honest characterisation of where the browser-native approach reaches its limits, including a quantified negative result on the inter-track chemistry contribution.

2 Methods

2.1 Re-implementation, not transpilation

None of Geant4’s runtime executes on the GPU. Each physics model is re-implemented as flat WGSL arithmetic, using the Geant4-DNA C++ source as the algorithmic specification (sampling routines and formulae) and the G4EMLOW 8.8 data files as the cross-section tables, converted once into WGSL constant arrays. A normally-compiled Geant4 11.4.1 / G4EMLOW 8.8 install serves as the *validation oracle*: we run the `dnaphysics` example single-threaded, dump its ntuples, and compare the WGSL output against them—the same validate-against-compiled-Geant4 methodology used by gMicroMC, MPEXS-DNA and GGEMS.

2.2 Kernel-fusion architecture

The pipeline has three phases (Fig. 1): **Phase A (primaries)** is a single fused `@compute` dispatch with one thread per primary electron; that thread runs the entire history—Born ionisation, Emfietzoglou excitation, elastic and vibrational scattering—in a `for` loop, with no per-step dispatch. **Phase B (secondaries)** is a wavefront stepper (one dispatch per physics step advancing all live secondaries), which cannot be fused because the secondary count is unknown until Phase A completes. **Phase C (chemistry)** runs the Karamitros 9-reaction IRT in a Web Worker off the main thread (a GPU spatial-hash alternative exists but is less accurate at long times; see Sec. 4). Energy deposition into a 128^3 voxel grid uses fixed-point integer `atomicAdd` ($\times 100$ per eV), since WGSL atomics are integer-only.

2.3 Ported models and deliberate divergences

We port Born ionisation (5 shells), Champion tabulated / screened-Rutherford elastic, Sanche vibrational excitation (9 modes), and the Karamitros 9-reaction IRT chemistry (6 species, Smoluchowski totally-diffusion-controlled and Onsager-screened partially-diffusion-controlled types) [11, 12]. Two deliberate departures from the `DNA_Opt2` defaults are documented (Table 1): we use the Emfietzoglou excitation model [4]—the model used by the Geant4-DNA constructor `G4EmDNAPhysics_option4`, which is unrelated to the similarly-named EM Standard list `G4EmStandardPhysics_option4`—rather than Born, because it yields the correct initial $G(H) = 0.33$; and two tuning scalars

Table 1: Deliberate divergences from Geant4-DNA DNA_Opt2, with the measured cost of each (experiment IDs reference § Numbers).

Item	DNA_Opt2	This work	Cost
Excitation	Born	Emfietzoglou	σ_{exc} $2.55\times$ (E6b)
σ_{exc} scale	—	$\times 0.5$	net $1.27\times$ (E6c)
Recomb. boost	—	$\times 2.0$	no physical basis
Chemistry pool	single	per-primary	$\Delta G(\text{H}_2)$ (E10f)
Dose reduction	fp64	fp32 atomics	MC-equivalent, not bit-exact

Table 2: Cross-section validation vs. G4EMLOW 8.8. Peak ratio is WebGPU/Geant4 at the worst single point; medians of the relative deviation are $\sim 10^{-3}$.

Channel	Model	Peak ratio	Exp.
Ionisation σ_{ion}	Born (5 shells)	0.9987	E1
Excitation σ_{exc}	Emfietzoglou (5 lev.)	0.9970	E2
Elastic σ_{el}	Champion / scr.-Ruth.	0.9751	E3
Vibrational σ_{vib}	Sanche (9 modes)	1.0000	E4

(SIGMA_EXC_SCALE= 0.5, RECOMB_BOOST= 2.0) partially close the chemistry yields. RECOMB_BOOST is an acknowledged fudge without a physical basis and is slated for removal.

2.4 WGS� constraints

WGS� forbids recursion, dynamic dispatch and dynamic allocation, and allows atomics only on u32. Consequently: all physics is inlined into `main()`; secondaries are deferred to a buffer + wavefront rather than recursed; dose uses fixed-point integers; and all memory is pre-allocated.

2.5 Validation protocol

The repository follows a six-level falsifiable ladder. Every experiment emits a JSON artifact carrying the git SHA, a named seed, WGS� shader hashes, a pass bar, and per-row observations; failed experiments are committed with `status:"fail"` and a diagnosis rather than rerun. All numbers below are drawn from this artifact set. The reference configuration is $N = 4096$ primaries at 10 keV on an Apple M2 Pro (apple/metal-3 adapter). A reproducibility caveat applies: fp32 atomic reductions are not order-deterministic across GPU vendors, so cross-hardware results are statistically equivalent within Monte Carlo noise but not bit-exact.

3 Results

3.1 Cross sections (Level 1)

All 9 cross-section channels match the G4EMLOW reference (Table 2). Sanche vibrational is bit-exact; the largest deviation is the Champion elastic peak ratio at 0.975.

3.2 Track structure (Level 2)

The continuous-slowing-down (CSDA) range at 10 keV is 2714.4 nm vs. Geant4’s 2747.5 nm, i.e. $0.988\times$ (3.59σ), with energy conservation at 100.0% (E5). Across the eight ESTAR energies the

[Figure placeholder: CSDA range ratio vs. energy, pre/post joint-fix]

Plot from E5b (pre) and E5d (post) artifacts: WebGPU/Geant4 CSDA ratio at 100, 300, 500, 1000, 3000, 5000, 10000, 20000 eV; two curves (pre-fix 0.587→0.992, post-fix 0.736→0.996); the post curve sits above the pre curve, gap largest at low E.

Figure 2: Energy-dependent CSDA closure under the joint fix (8/8 improved).

Table 3: Track-structure metrics vs. Geant4 11.4.1 at 10 keV, $N = 4096$.

Metric	This work	Geant4	Ratio (exp.)
CSDA range	2714.4 nm	2747.5 nm	0.988 (E5)
Energy cons.	100.0 %	99.99 %	1.000 (E5)
Ions / primary	371.9	509.2	0.730 (E7)
W -value	26.89 eV	21.4 eV [†]	1.257 (E5c)
Sec. KE/primary	143.4	144.9	1.010 (E8)

[†] ICRU 31 low-LET liquid water reference.

deficit is energy-dependent and was closed monotonically by the joint fix: the 100 eV ratio rose from $0.587\times$ to $0.736\times$ and all eight energies improved, with the lift inversely proportional to the original deficit—the expected signature of the excitation cross-section correction (E5b, E5d; Fig. 2).

The mean free path is 5–11 % short of Geant4 across six energy bins (median 0.941, E6). The full-cascade ionisation count is 371.9 per primary vs. Geant4’s 509.2—a 27 % deficit at 263σ (E7)—and the implicit W -value is 26.89 eV vs. ICRU 31’s 21.4 eV ([8], +25.7 %, E5c). Both are the same physics: the intentional excitation inflation channels energy away from ionisation. The secondary-electron kinetic-energy spectrum at creation matches to 1.0 % overall, with 7/8 log-bins within 0.1–3.1 % (E8). Table 3 summarises.

3.3 Chemistry (Levels 3–4)

Against Geant4’s `chem6` at matched 10 keV, the G -values at 1 μs (post joint-fix, E10j) are: $G(\text{OH})$ $0.90\times$, $G(\text{e}_{\text{aq}}^-)$ $0.82\times$, $G(\text{H})$ $1.10\times$, $G(\text{H}_2)$ $0.86\times$, $G(\text{H}_2\text{O}_2)$ $0.69\times$ (Fig. 3). The five-species RMS deviation at 0.1 ps is 19.0 % (down from a 30.3 % baseline, E10i). A non-trivial physics result emerges from the protocol: $G(\text{e}_{\text{aq}}^-)$ is non-monotonic between 1 and 3 keV, a 12.5 % dip significant at 126σ by primary-bootstrap (E10b), which `chem6` independently reproduces (E10d)—i.e. it is real track-end / spur-structure physics, not an artifact.

3.4 DNA damage (Level 5)

On a 21×21 B-DNA fibre grid (3.89 Mbp target) the indirect/direct single-strand-break ratio is 2.46 ($\text{SSB}_{\text{dir}} = 26$, $\text{SSB}_{\text{ind}} = 64$, $\text{DSB} = 9$; post joint-fix, E13c), within PARTRAC’s reported 2–3 band [5]. Absolute per-Da yields are geometry-dependent because the concentrated fibre grid samples the track core rather than a uniform bulk; this is a stated geometric caveat, not a scoring error.

[Figure placeholder: $G(t)$ for five species, WebGPU vs. chem6]

Plot from E10c/E10j artifacts: $G(OH, e_{aq}^-, H, H_2, H_2O_2)$ vs. time on a log axis $0.1\text{ ps} \rightarrow 1\text{ }\mu\text{s}$; solid = WebGPU, dashed = chem6.

Figure 3: Radiolysis G -value time evolution against Geant4 chem6.

[Figure placeholder: Speedup bars + throughput vs. N]

From E15/E15b/E15c/E15d/E16: (a) bar chart $455\times$ (ST), $280\times$ (MT-8), $40\times$ (fusion), $1.48\times$ (end-to-end); (b) throughput vs. N across 8 energies.

Figure 4: Performance summary.

3.5 Performance (Level 6)

Phase A+B tracking is **455** $\times$ faster than Geant4 11.4.1 single-thread (635 ms vs. a 289.1 s median over 3 trials, E15b) and **280** $\times$ vs. an 8-thread build (178.0 s, E15c). Peak Phase A throughput is 538 947 primaries/s at $N = 16384$ (E15). A within-WebGPU comparison of the fused kernel against a modelled per-step dispatch gives a **40** $\times$ kernel-fusion factor (E16). The end-to-end pre-DNA pipeline is only 1.48 $\times$ faster, because the IRT chemistry runs on CPU and dominates the wall-clock (194 s of 194.6 s); this motivates Sec. 4.

4 The inter-track chemistry limitation

The largest remaining gap vs. chem6 is an *inter-track* effect: per-primary IRT partitioning misses cross-primary reactions that, when included, add $\Delta G(H_2) = +0.149$ at $1\text{ }\mu\text{s}$ —96 % of the residual implementation gap (E10f). We asked whether this is recoverable browser-native, and found it is not, by two independent routes.

Cross-primary IRT is $O(N^2)$ for a point source. The IRT reaction horizon is the diffusion length over the full $1\text{ }\mu\text{s}$ window, $R_{\text{cut}} = 1.45 + 2\sqrt{8D_{\text{max}}t_{\text{max}}} \approx 551\text{ nm}$, not the few-nm contact scale. At 10 keV the radicals occupy a $\sim 5\text{ }\mu\text{m}$ blob, so a 551 nm spatial hash spans most of it and prunes almost nothing; over 50 % of accepted cross-primary reactions involve pairs more than 100 nm apart (E10k). The single-pool scan is $\sim 10^{12}$ reaction-time samples (~ 70 hr in-browser) and the heap is ~ 2 GB.

GPU step-by-step (SBS) chemistry is encounter-radius-bounded. The GPU radiolysis codes use a step-by-step Brownian-motion model in which a reaction fires when two reactants fall within a Smoluchowski reaction radius (gMicroMC [14]; MPEXS-DNA adds a during-step reaction probability at small time steps [13]). We implemented this in WGS and, as a refinement, a Green’s-function (Brownian-bridge) during-step reaction probability [3] from the step-by-step radiation-chemistry literature, and measured its convergence against the analytic Smoluchowski value for a single pair. The per-step cost is ~ 110 ms at $N = 4096$ (E10L), and the IRT-parity budget is ~ 1750 steps. But accuracy requires a time step that resolves the encounter radius σ , $\Delta t \lesssim \sigma^2/(2D) \approx 0.02\text{ ns}$, *independent of pair separation or density* (E10Q; Fig. 5)—so $\sim 50\,000$

[Figure placeholder: Bridge convergence ratio vs. Δt at several separations]
From E10Q: SBS/analytic reaction-probability ratio vs. Δt (4, 1, 0.25, 0.05 ns) for $r_0 = 1, 10, 30, 100$ nm; the four curves overlap, showing the Δt requirement is set by σ , not separation.

Figure 5: Kill criterion: the SBS time-step bound is separation-independent.

steps are needed regardless. Compaction lowers per-step cost but not step count (E10P). No SBS schedule, full or hybrid, beats the IRT worker at the required accuracy on laptop WebGPU.

Both routes converge on the same conclusion: the inter-track physics requires a *native* runtime (e.g. Node/Deno + `wgpu-native`), where RAM is cheap (unblocking single-pool cross-primary IRT) or many cores parallelise the step count. The browser remains the validation and demonstration surface.

5 Discussion

Browser-native execution buys accessibility and reproducibility that the CUDA/OpenCL engines cannot: the simulation and its validation harness run on any modern GPU with no install, which is valuable for teaching, sharing, and independent re-checking of every artifact. The cost is the WebGPU feature envelope (integer-only atomics, no recursion) and—per Sec. 4—an inter-track chemistry ceiling that pushes the last increment of fidelity off-browser. The cascade-ionisation deficit and the `RECOMB_BOOST` fudge are open items with identified, falsifiable next steps (super-excitation autoionisation; an Onsager escape-probability treatment to replace the fudge). Relative to the CUDA codes `gMicroMC` and `MPEXS-DNA`, our speedups come from kernel fusion on a single consumer GPU (`MPEXS-DNA`, for reference, reports up to $2900\times$ over single-core Geant4-DNA on an NVIDIA TITAN V [13]); extending the same methodology to the EM Standard list `G4EmStandardPhysics_option4` for macroscopic photon dose is a natural, if substantially larger, direction.

6 Conclusion

A browser-native, kernel-fused WebGPU re-implementation reproduces Geant4-DNA track structure to within a fraction of a percent on cross sections and $0.988\times$ on CSDA range, at $455\times/280\times$ the speed of single-/multi-threaded Geant4, with an honest, artifact-backed account of its limits—most notably the quantified result that the inter-track chemistry contribution is not browser-native and requires a native runtime.

Data and code availability

Source, the full validation ladder, and every JSON artifact are at <https://github.com/abgnydn/webgpu-dna>; a live build runs at <https://webgpudna.com>. A versioned Zenodo deposit accompanies release v0.4.0 (DOI to be inserted). Each experiment reruns via `npm run experiments - <id>`.

References

- [1] S. Agostinelli et al. Geant4—a simulation toolkit. *Nuclear Instruments and Methods in Physics Research A*, 506(3):250–303, 2003.
- [2] M. A. Bernal et al. Track structure modeling in liquid water: A review of the Geant4-DNA very low energy extension of the Geant4 Monte Carlo simulation toolkit. *Physica Medica*, 31(8):861–874, 2015. VERIFY exact pages.
- [3] P. Clifford, N. J. B. Green, M. J. Oldfield, M. J. Pilling, and S. M. Pimblott. Stochastic models of multi-species kinetics in radiation-induced spurs. *Journal of the Chemical Society, Faraday Transactions 1*, 82, 1986. VERIFY exact volume/pages; this is the bridge/Green’s-function reaction-probability source, NOT gMicroMC.
- [4] D. Emfietzoglou, F. A. Cucinotta, and H. Nikjoo. A complete dielectric response model for liquid water: A solution of the Bethe ridge problem. *Radiation Research*, 2005. VERIFY author list, exact title, volume/pages.
- [5] W. Friedland, M. Dingfelder, P. Kunderát, and P. Jacob. Track structures, DNA targets and radiation effects in the biophysical Monte Carlo simulation code PARTRAC. *Mutation Research*, 711(1–2):28–40, 2011. VERIFY exact pages.
- [6] GGEMS contributors. GGEMS: GPU GEant4-based Monte Carlo simulations. <https://ggems.fr>. VERIFY: cite the canonical GGEMS paper (e.g. GGEMS-Brachy, Lemaître/Bert et al.) instead of the website if available.
- [7] S. Hissoiny, B. Ozell, H. Bouchard, and P. Després. GPUMCD: A new GPU-oriented Monte Carlo dose calculation platform. *Medical Physics*, 38(1):754–764, 2011.
- [8] ICRU. Average energy required to produce an ion pair. Technical Report Report 31, International Commission on Radiation Units and Measurements, 1979.
- [9] S. Incerti et al. The Geant4-DNA project. *International Journal of Modeling, Simulation, and Scientific Computing*, 1(2):157–178, 2010. VERIFY exact volume/pages.
- [10] S. Incerti et al. Geant4-DNA example applications for track structure simulations in liquid water: A report from the Geant4-DNA project. *Medical Physics*, 45(8):e722–e739, 2018. VERIFY exact pages.
- [11] M. Karamitros et al. Modeling radiation chemistry in the Geant4 toolkit. *Progress in Nuclear Science and Technology*, 2:503–508, 2011. VERIFY volume/pages.
- [12] M. Karamitros et al. Diffusion-controlled reactions modeling in Geant4-DNA. *Journal of Computational Physics*, 274:841–882, 2014. VERIFY exact pages.
- [13] S. Okada, K. Murakami, S. Incerti, K. Amako, and T. Sasaki. MPEXS-DNA, a new GPU-based Monte Carlo simulator for track structures and radiation chemistry at subcellular scale. *Medical Physics*, 46(3):1483–1500, 2019. VERIFY author order/list and end page.
- [14] M.-Y. Tsai, Z. Tian, N. Qin, H. Yan, Y. Lai, S.-H. Hung, Y. Chi, and X. Jia. A new open-source GPU-based microscopic Monte Carlo simulation tool for the calculations of DNA damages caused by ionizing radiation—Part I: Core algorithm and validation. *Medical Physics*, 47(4), 2020. VERIFY exact pages and author order.