

Class 5-Linear Algebra

Dot Product

Let u and v be any two vectors in \mathbb{R}^n .

Dot Product

Let u and v be any two vectors in \mathbb{R}^n . That is., $u = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix}$ and $v = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$.

Dot Product

Let u and v be any two vectors in \mathbb{R}^n . That is., $u = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix}$ and $v = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$.

Then the dot product of u and v is,

Dot Product

Let u and v be any two vectors in \mathbb{R}^n . That is., $u = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix}$ and $v = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$.

Then the dot product of u and v is,

$$u.v = u^T v = \begin{bmatrix} u_1 & u_2 & \dots & u_n \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$$

Dot Product

Let u and v be any two vectors in \mathbb{R}^n . That is., $u = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix}$ and $v = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix}$.

Then the dot product of u and v is,

$$\begin{aligned} u \cdot v &= u^T v = \begin{bmatrix} u_1 & u_2 & \dots & u_n \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} \\ &= u_1 v_1 + u_2 v_2 + \dots + u_n v_n. \end{aligned}$$

Example

Compute $u \cdot v$ and $v \cdot u$ for $u = \begin{bmatrix} 2 \\ -5 \\ 1 \end{bmatrix}$ and $v = \begin{bmatrix} 3 \\ 2 \\ -3 \end{bmatrix}$

Properties of dot product

Let u , v and w be vectors in \mathbb{R}^n , and c be a scalar. Then

- ① $(u + v) \cdot w = u \cdot w + v \cdot w$
- ② $(cu) \cdot v = c(u \cdot v) = u \cdot (cv)$
- ③ $u \cdot v = v \cdot u$
- ④ $u \cdot u \geq 0$ and $u \cdot u = 0$ if and only if $u = 0$.

Inner Product on \mathbb{R}^n

An **inner product** on \mathbb{R}^n is a function

$$\langle \cdot, \cdot \rangle : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$$

that satisfies the following properties for all $u, v, w \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$:

Inner Product on \mathbb{R}^n

An **inner product** on \mathbb{R}^n is a function

$$\langle \cdot, \cdot \rangle : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$$

that satisfies the following properties for all $u, v, w \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$:

① **Linearity in the first argument:**

$$\langle \alpha u + v, w \rangle = \alpha \langle u, w \rangle + \langle v, w \rangle$$

Inner Product on \mathbb{R}^n

An **inner product** on \mathbb{R}^n is a function

$$\langle \cdot, \cdot \rangle : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$$

that satisfies the following properties for all $u, v, w \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$:

① **Linearity in the first argument:**

$$\langle \alpha u + v, w \rangle = \alpha \langle u, w \rangle + \langle v, w \rangle$$

② **Symmetry:**

$$\langle u, v \rangle = \langle v, u \rangle$$

Inner Product on \mathbb{R}^n

An **inner product** on \mathbb{R}^n is a function

$$\langle \cdot, \cdot \rangle : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$$

that satisfies the following properties for all $u, v, w \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$:

❶ **Linearity in the first argument:**

$$\langle \alpha u + v, w \rangle = \alpha \langle u, w \rangle + \langle v, w \rangle$$

❷ **Symmetry:**

$$\langle u, v \rangle = \langle v, u \rangle$$

❸ **Positive-definiteness:**

$$\langle v, v \rangle \geq 0, \quad \text{and} \quad \langle v, v \rangle = 0 \iff v = 0$$

Example: Dot Product

Dot Product as an Inner Product

For $u = (u_1, u_2, \dots, u_n)$ and $v = (v_1, v_2, \dots, v_n)$ in \mathbb{R}^n , define

$$\langle u, v \rangle = u \cdot v = \sum_{i=1}^n u_i v_i$$

Example: Dot Product

Dot Product as an Inner Product

For $u = (u_1, u_2, \dots, u_n)$ and $v = (v_1, v_2, \dots, v_n)$ in \mathbb{R}^n , define

$$\langle u, v \rangle = u \cdot v = \sum_{i=1}^n u_i v_i$$

Dot product is an example of inner product on \mathbb{R}^n .

Example: Inner Product via Positive Definite Matrix

Let

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}, \quad u = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}, \quad v = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}.$$

Example: Inner Product via Positive Definite Matrix

Let

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}, \quad u = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}, \quad v = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}.$$

Define

$$\langle u, v \rangle_A = u^T A v = 2u_1 v_1 + 3u_2 v_2.$$

Example: Inner Product via Positive Definite Matrix

Let

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}, \quad u = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}, \quad v = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}.$$

Define

$$\langle u, v \rangle_A = u^T A v = 2u_1 v_1 + 3u_2 v_2.$$

Verification:

Example: Inner Product via Positive Definite Matrix

Let

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}, \quad u = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}, \quad v = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}.$$

Define

$$\langle u, v \rangle_A = u^T A v = 2u_1 v_1 + 3u_2 v_2.$$

Verification:

- Linearity: follows from matrix multiplication.

Example: Inner Product via Positive Definite Matrix

Let

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}, \quad u = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}, \quad v = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}.$$

Define

$$\langle u, v \rangle_A = u^T A v = 2u_1 v_1 + 3u_2 v_2.$$

Verification:

- Linearity: follows from matrix multiplication.
- Symmetry: A is symmetric, so $\langle u, v \rangle_A = \langle v, u \rangle_A$.

Example: Inner Product via Positive Definite Matrix

Let

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}, \quad u = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}, \quad v = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}.$$

Define

$$\langle u, v \rangle_A = u^T A v = 2u_1 v_1 + 3u_2 v_2.$$

Verification:

- Linearity: follows from matrix multiplication.
- Symmetry: A is symmetric, so $\langle u, v \rangle_A = \langle v, u \rangle_A$.
- Positive-definite: $\langle u, u \rangle_A = 2u_1^2 + 3u_2^2 > 0$ for $u \neq 0$.

Example: Inner Product via Positive Definite Matrix

Let

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}, \quad u = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}, \quad v = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}.$$

Define

$$\langle u, v \rangle_A = u^T A v = 2u_1 v_1 + 3u_2 v_2.$$

Verification:

- Linearity: follows from matrix multiplication.
- Symmetry: A is symmetric, so $\langle u, v \rangle_A = \langle v, u \rangle_A$.
- Positive-definite: $\langle u, u \rangle_A = 2u_1^2 + 3u_2^2 > 0$ for $u \neq 0$.

Note: We will mostly consider dot product from now on.

Norm on \mathbb{R}^n

A **norm** on \mathbb{R}^n is a function

$$\| \cdot \| : \mathbb{R}^n \rightarrow \mathbb{R}$$

that assigns to each vector $v \in \mathbb{R}^n$ a nonnegative real number, and satisfies the following properties for all $u, v \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$:

Norm on \mathbb{R}^n

A **norm** on \mathbb{R}^n is a function

$$\| \cdot \| : \mathbb{R}^n \rightarrow \mathbb{R}$$

that assigns to each vector $v \in \mathbb{R}^n$ a nonnegative real number, and satisfies the following properties for all $u, v \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$:

① **Positivity:**

$$\|v\| \geq 0, \quad \|v\| = 0 \iff v = 0$$

Norm on \mathbb{R}^n

A **norm** on \mathbb{R}^n is a function

$$\| \cdot \| : \mathbb{R}^n \rightarrow \mathbb{R}$$

that assigns to each vector $v \in \mathbb{R}^n$ a nonnegative real number, and satisfies the following properties for all $u, v \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$:

① **Positivity:**

$$\|v\| \geq 0, \quad \|v\| = 0 \iff v = 0$$

② **Homogeneity (absolute scalability):**

$$\|\alpha v\| = |\alpha| \|v\|$$

Norm on \mathbb{R}^n

A **norm** on \mathbb{R}^n is a function

$$\| \cdot \| : \mathbb{R}^n \rightarrow \mathbb{R}$$

that assigns to each vector $v \in \mathbb{R}^n$ a nonnegative real number, and satisfies the following properties for all $u, v \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$:

① **Positivity:**

$$\|v\| \geq 0, \quad \|v\| = 0 \iff v = 0$$

② **Homogeneity (absolute scalability):**

$$\|\alpha v\| = |\alpha| \|v\|$$

③ **Triangle inequality:**

$$\|u + v\| \leq \|u\| + \|v\|$$

The length(or norm) of a vector

The length (or norm) of v in \mathbb{R}^n is the non-negative scalar $\|v\|$ defined by

$$\|v\| = \sqrt{v \cdot v} = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$$

The length(or norm) of a vector

The length (or norm) of v in \mathbb{R}^n is the non-negative scalar $\|v\|$ defined by

$$\|v\| = \sqrt{v \cdot v} = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$$

Can you write formula for norm in

The length(or norm) of a vector

The length (or norm) of v in \mathbb{R}^n is the non-negative scalar $\|v\|$ defined by

$$\|v\| = \sqrt{v \cdot v} = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$$

Can you write formula for norm in

- ① Length in \mathbb{R}^2
- ② Length in \mathbb{R}^3

Unit vector

A vector whose length is 1 is called the unit vector.

Example

$$v = \begin{bmatrix} \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \end{bmatrix}$$

Normalizing a vector

If we divide a non-zero v vector by its norm $\|v\|$, then the resulting vector is unit vector in the direction of v .

Example

$$v = \begin{bmatrix} 1 \\ -2 \\ 2 \\ 0 \end{bmatrix}$$