

Class- 6 and 7: Matrix Decomposition

Gram-Schmidt process and the Singular Value Decomposition

August 26, 2025

Orthogonal vectors

Two vectors u and v in \mathbb{R}^n are **orthogonal** if $u \cdot v = 0$

Orthogonal vectors

Two vectors u and v in \mathbb{R}^n are **orthogonal** if $u \cdot v = 0$

1. Check whether $u = \begin{bmatrix} \frac{4}{3} \\ -1 \\ \frac{2}{3} \end{bmatrix}$ and $v = \begin{bmatrix} 5 \\ 6 \\ -1 \end{bmatrix}$ are orthogonal vectors.

2. Determine which pair of vectors are orthogonal

a $a = \begin{bmatrix} 8 \\ -5 \end{bmatrix}$ and $b = \begin{bmatrix} -2 \\ -3 \end{bmatrix}$

2. Determine which pair of vectors are orthogonal

a $a = \begin{bmatrix} 8 \\ -5 \end{bmatrix}$ and $b = \begin{bmatrix} -2 \\ -3 \end{bmatrix}$

b $y = \begin{bmatrix} 3 \\ 2 \\ -5 \\ 0 \end{bmatrix}$ and $z = \begin{bmatrix} -4 \\ 1 \\ -2 \\ 6 \end{bmatrix}$

Orthogonal sets

A set of vectors $\{u_1, u_2, \dots, u_p\}$ in \mathbb{R}^n are orthogonal if each pair of distinct vectors in the set are orthogonal, that is $u_i \cdot u_j = 0$ for $i \neq j$.

Orthogonal sets

A set of vectors $\{u_1, u_2, \dots, u_p\}$ in \mathbb{R}^n are orthogonal if each pair of distinct vectors in the set are orthogonal, that is $u_i \cdot u_j = 0$ for $i \neq j$.

Example

Check if $u_1 = \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix}$, $u_2 = \begin{bmatrix} -1 \\ 2 \\ 1 \end{bmatrix}$ and $u_3 = \begin{bmatrix} -\frac{1}{2} \\ -2 \\ \frac{7}{2} \end{bmatrix}$ are orthogonal.

Orthonormal vectors

Definition

Two vectors u and v in \mathbb{R}^n are orthonormal if $u \cdot v = 0$ and $\|u\| = \|v\| = 1$.

Orthonormal vectors

Definition

Two vectors u and v in \mathbb{R}^n are orthonormal if $u \cdot v = 0$ and $\|u\| = \|v\| = 1$.

Example

Check if $u = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$ and $v = \begin{bmatrix} \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$ are orthonormal.

Orthonormal sets

A set of vectors $\{u_1, u_2, \dots, u_p\}$ in \mathbb{R}^n are orthonormal if each pair of distinct vectors in the set are orthogonal and $\|u_i\| = 1$ for all $1 \leq i \leq p$.

Orthonormal sets

A set of vectors $\{u_1, u_2, \dots, u_p\}$ in \mathbb{R}^n are orthonormal if each pair of distinct vectors in the set are orthogonal and $\|u_i\| = 1$ for all $1 \leq i \leq p$.

Example

Show that $\{v_1, v_2, v_3\}$ is an orthonormal set in \mathbb{R}^3 , where $v_1 = \begin{bmatrix} \frac{3}{\sqrt{11}} \\ \frac{1}{\sqrt{11}} \\ \frac{1}{\sqrt{11}} \end{bmatrix}$,

$$v_2 = \begin{bmatrix} \frac{-1}{\sqrt{6}} \\ \frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix} \text{ and } v_3 = \begin{bmatrix} \frac{-1}{\sqrt{66}} \\ \frac{-4}{\sqrt{66}} \\ \frac{7}{\sqrt{66}} \end{bmatrix}$$

Theorem

*An $m \times n$ matrix U has orthonormal column if and only if $U^T U = I$. If a matrix has orthonormal columns, then the matrix is **orthogonal matrix**.*

Theorem

An $m \times n$ matrix U has orthonormal column if and only if $U^T U = I$. If a matrix has orthonormal columns, then the matrix is **orthogonal matrix**.

Example

Check if U is orthogonal matrix, where $U = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{2}{3} \\ \frac{1}{\sqrt{2}} & -\frac{2}{3} \\ 0 & \frac{1}{3} \end{bmatrix}$

The Gram-Schmidt Process

The Gram-Schmidt Process is a simple algorithm for producing an orthogonal or orthonormal set in \mathbb{R}^n .

Gram-Schmidt orthonormalization process

Consider the vectors as columns of the matrix A . That is,

$$A = [a_1 \mid a_2 \mid \cdots \mid a_n] .$$

Gram-Schmidt orthonormalization process

Consider the vectors as columns of the matrix A . That is,

$$A = [a_1 \mid a_2 \mid \cdots \mid a_n].$$

Then,

$$\mathbf{u}_1 = \mathbf{a}_1, \quad \mathbf{e}_1 = \frac{\mathbf{u}_1}{\|\mathbf{u}_1\|},$$

$$\mathbf{u}_2 = \mathbf{a}_2 - (\mathbf{a}_2 \cdot \mathbf{e}_1)\mathbf{e}_1, \quad \mathbf{e}_2 = \frac{\mathbf{u}_2}{\|\mathbf{u}_2\|},$$

$$\mathbf{u}_3 = \mathbf{a}_3 - (\mathbf{a}_3 \cdot \mathbf{e}_1)\mathbf{e}_1 - (\mathbf{a}_3 \cdot \mathbf{e}_2)\mathbf{e}_2, \quad \mathbf{e}_3 = \frac{\mathbf{u}_3}{\|\mathbf{u}_3\|},$$

$$\vdots$$

$$\mathbf{u}_{k+1} = \mathbf{a}_{k+1} - (\mathbf{a}_{k+1} \cdot \mathbf{e}_1)\mathbf{e}_1 - \cdots - (\mathbf{a}_{k+1} \cdot \mathbf{e}_k)\mathbf{e}_k, \quad \mathbf{e}_{k+1} = \frac{\mathbf{u}_{k+1}}{\|\mathbf{u}_{k+1}\|}.$$

Eigenvalues and Eigenvectors: Definition

A real number λ is an *eigenvalue* of an $n \times n$ square matrix A if there exists a nonzero vector $\mathbf{x} \in \mathbb{R}^n$ such that

Eigenvalues and Eigenvectors: Definition

A real number λ is an *eigenvalue* of an $n \times n$ square matrix A if there exists a nonzero vector $\mathbf{x} \in \mathbb{R}^n$ such that

$$A\mathbf{x} = \lambda \mathbf{x}.$$

Eigenvalues and Eigenvectors: Definition

A real number λ is an *eigenvalue* of an $n \times n$ square matrix A if there exists a nonzero vector $\mathbf{x} \in \mathbb{R}^n$ such that

$$A\mathbf{x} = \lambda \mathbf{x}.$$

The vector \mathbf{x} is then called an *eigenvector* corresponding to λ .

Eigenvalues and Eigenvectors: Definition

A real number λ is an *eigenvalue* of an $n \times n$ square matrix A if there exists a nonzero vector $\mathbf{x} \in \mathbb{R}^n$ such that

$$A\mathbf{x} = \lambda \mathbf{x}.$$

The vector \mathbf{x} is then called an *eigenvector* corresponding to λ .

For example, consider $\begin{bmatrix} 2 & 0 & 2 \\ 0 & 2 & 0 \\ 2 & 0 & 2 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} =$

Eigenvalues and Eigenvectors: Definition

A real number λ is an *eigenvalue* of an $n \times n$ square matrix A if there exists a nonzero vector $\mathbf{x} \in \mathbb{R}^n$ such that

$$A\mathbf{x} = \lambda \mathbf{x}.$$

The vector \mathbf{x} is then called an *eigenvector* corresponding to λ .

For example, consider
$$\begin{bmatrix} 2 & 0 & 2 \\ 0 & 2 & 0 \\ 2 & 0 & 2 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix} =$$

Eigenvalues and Eigenvectors: Definition

A real number λ is an *eigenvalue* of an $n \times n$ square matrix A if there exists a nonzero vector $\mathbf{x} \in \mathbb{R}^n$ such that

$$A\mathbf{x} = \lambda \mathbf{x}.$$

The vector \mathbf{x} is then called an *eigenvector* corresponding to λ .

For example, consider
$$\begin{bmatrix} 2 & 0 & 2 \\ 0 & 2 & 0 \\ 2 & 0 & 2 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix} = 2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}.$$

Eigenvalues and Eigenvectors: Definition

A real number λ is an *eigenvalue* of an $n \times n$ square matrix A if there exists a nonzero vector $\mathbf{x} \in \mathbb{R}^n$ such that

$$A\mathbf{x} = \lambda \mathbf{x}.$$

The vector \mathbf{x} is then called an *eigenvector* corresponding to λ .

For example, consider
$$\begin{bmatrix} 2 & 0 & 2 \\ 0 & 2 & 0 \\ 2 & 0 & 2 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix} = 2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}.$$

Therefore, $\lambda = 2$ is an eigenvalue corresponding to the eigenvector $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$.

Example

Example

Equivalently, λ is an eigenvalue iff $\det(A - \lambda I) = 0$.

Example

Equivalently, λ is an eigenvalue iff $\det(A - \lambda I) = 0$.

Find the eigenvalues of $A = \begin{bmatrix} 2 & 0 & 2 \\ 0 & 2 & 0 \\ 2 & 0 & 2 \end{bmatrix}$.

Example

Equivalently, λ is an eigenvalue iff $\det(A - \lambda I) = 0$.

Find the eigenvalues of $A = \begin{bmatrix} 2 & 0 & 2 \\ 0 & 2 & 0 \\ 2 & 0 & 2 \end{bmatrix}$.

Compute the characteristic polynomial and equate it to zero:

$$\det(A - \lambda I) = \det \begin{bmatrix} 2 - \lambda & 0 & 2 \\ 0 & 2 - \lambda & 0 \\ 2 & 0 & 2 - \lambda \end{bmatrix} = 0.$$

Example

Equivalently, λ is an eigenvalue iff $\det(A - \lambda I) = 0$.

Find the eigenvalues of $A = \begin{bmatrix} 2 & 0 & 2 \\ 0 & 2 & 0 \\ 2 & 0 & 2 \end{bmatrix}$.

Compute the characteristic polynomial and equate it to zero:

$$\det(A - \lambda I) = \det \begin{bmatrix} 2 - \lambda & 0 & 2 \\ 0 & 2 - \lambda & 0 \\ 2 & 0 & 2 - \lambda \end{bmatrix} = 0.$$

so the eigenvalues are $\lambda \in \{0, 2, 4\}$.

Definition of singular values

For rectangular (or square) matrices, *singular values* play a role analogous to eigenvalues.

Definition of singular values

For rectangular (or square) matrices, *singular values* play a role analogous to eigenvalues.

Definition: Let A be an $m \times n$ matrix. A scalar λ is called a *singular value* of A if there exist nonzero vectors $\mathbf{u} \in \mathbb{R}^n$ and $\mathbf{v} \in \mathbb{R}^m$ such that

$$A\mathbf{v} = \lambda \mathbf{u} \quad \text{and} \quad A^T \mathbf{u} = \lambda \mathbf{v}.$$

Definition of singular values

For rectangular (or square) matrices, *singular values* play a role analogous to eigenvalues.

Definition: Let A be an $m \times n$ matrix. A scalar λ is called a *singular value* of A if there exist nonzero vectors $\mathbf{u} \in \mathbb{R}^n$ and $\mathbf{v} \in \mathbb{R}^m$ such that

$$A\mathbf{v} = \lambda \mathbf{u} \quad \text{and} \quad A^T \mathbf{u} = \lambda \mathbf{v}.$$

The vectors \mathbf{u} and \mathbf{v} are a *singular vector pair* corresponding to λ .

Definition of singular values

For rectangular (or square) matrices, *singular values* play a role analogous to eigenvalues.

Definition: Let A be an $m \times n$ matrix. A scalar λ is called a *singular value* of A if there exist nonzero vectors $\mathbf{u} \in \mathbb{R}^n$ and $\mathbf{v} \in \mathbb{R}^m$ such that

$$A\mathbf{v} = \lambda \mathbf{u} \quad \text{and} \quad A^T \mathbf{u} = \lambda \mathbf{v}.$$

The vectors \mathbf{u} and \mathbf{v} are a *singular vector pair* corresponding to λ .

Equivalently: the singular values are the positive square roots of the eigenvalues of $A^T A$ or AA^T .

Finding singular values: Example (i)

Let $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$. Then

$$A^T A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}^T \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix}.$$

Finding singular values: Example (i)

Let $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$. Then

$$A^T A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}^T \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix}.$$

The eigenvalues of $\begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix}$ are 9 and 1.

Finding singular values: Example (i)

Let $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$. Then

$$A^T A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}^T \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix}.$$

The eigenvalues of $\begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix}$ are 9 and 1.

Therefore the singular values of A are 3 and 1.

Finding singular values: Example (ii)

Let $A = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \end{bmatrix}$. Then

$$AA^T = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 2 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 3 & 6 \\ 6 & 12 \end{bmatrix}.$$

Finding singular values: Example (ii)

Let $A = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \end{bmatrix}$. Then

$$AA^T = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 2 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 3 & 6 \\ 6 & 12 \end{bmatrix}.$$

Characteristic polynomial: $\det \begin{bmatrix} 3 - \lambda & 6 \\ 6 & 12 - \lambda \end{bmatrix} = \lambda(\lambda - 15).$

Finding singular values: Example (ii)

Let $A = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \end{bmatrix}$. Then

$$AA^T = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 2 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 3 & 6 \\ 6 & 12 \end{bmatrix}.$$

Characteristic polynomial: $\det \begin{bmatrix} 3 - \lambda & 6 \\ 6 & 12 - \lambda \end{bmatrix} = \lambda(\lambda - 15)$.

Eigenvalues: 15 and 0.

Finding singular values: Example (ii)

Let $A = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \end{bmatrix}$. Then

$$AA^T = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 2 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 3 & 6 \\ 6 & 12 \end{bmatrix}.$$

Characteristic polynomial: $\det \begin{bmatrix} 3 - \lambda & 6 \\ 6 & 12 - \lambda \end{bmatrix} = \lambda(\lambda - 15)$.

Eigenvalues: 15 and 0.

Singular values: $\sqrt{15}$ and 0.

The Singular Value Decomposition (SVD)

Theorem (SVD)

Let A be an $m \times n$ matrix and let $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_r > 0$ be the nonzero singular values. Then

$$A = U \Sigma V^T,$$

where U is $m \times m$ orthogonal, V is $n \times n$ orthogonal, and

$$\Sigma = \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix}, \quad D = \text{diag}(\sigma_1, \dots, \sigma_r).$$

The Singular Value Decomposition (SVD)

Theorem (SVD)

Let A be an $m \times n$ matrix and let $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_r > 0$ be the nonzero singular values. Then

$$A = U \Sigma V^T,$$

where U is $m \times m$ orthogonal, V is $n \times n$ orthogonal, and

$$\Sigma = \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix}, \quad D = \text{diag}(\sigma_1, \dots, \sigma_r).$$

Columns of U are orthonormal eigenvectors of AA^T (left singular vectors).

The Singular Value Decomposition (SVD)

Theorem (SVD)

Let A be an $m \times n$ matrix and let $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_r > 0$ be the nonzero singular values. Then

$$A = U \Sigma V^T,$$

where U is $m \times m$ orthogonal, V is $n \times n$ orthogonal, and

$$\Sigma = \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix}, \quad D = \text{diag}(\sigma_1, \dots, \sigma_r).$$

Columns of U are orthonormal eigenvectors of AA^T (left singular vectors). Columns of V are orthonormal eigenvectors of $A^T A$ (right singular vectors).

Working procedure for computing an SVD

- 1 Determine an orthonormal set of eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$ corresponding to eigenvalues $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($n - r$ times).

Working procedure for computing an SVD

- 1 Determine an orthonormal set of eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$ corresponding to eigenvalues $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($n - r$ times).
- 2 Form $V = [\mathbf{v}_1 \ \dots \ \mathbf{v}_n]$.

Working procedure for computing an SVD

- 1 Determine an orthonormal set of eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$ corresponding to eigenvalues $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($n - r$ times).
- 2 Form $V = [\mathbf{v}_1 \cdots \mathbf{v}_n]$.
- 3 For $1 \leq i \leq r$, set $\mathbf{u}_i := \frac{A\mathbf{v}_i}{\|A\mathbf{v}_i\|}$.

Working procedure for computing an SVD

- 1 Determine an orthonormal set of eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$ corresponding to eigenvalues $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($n - r$ times).
- 2 Form $V = [\mathbf{v}_1 \ \dots \ \mathbf{v}_n]$.
- 3 For $1 \leq i \leq r$, set $\mathbf{u}_i := \frac{A\mathbf{v}_i}{\|A\mathbf{v}_i\|}$.
- 4 Choose $\mathbf{u}_{r+1}, \dots, \mathbf{u}_m$ to complete $\{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ to an orthonormal basis of \mathbb{R}^m .

Working procedure for computing an SVD

- 1 Determine an orthonormal set of eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$ corresponding to eigenvalues $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($n - r$ times).
- 2 Form $V = [\mathbf{v}_1 \cdots \mathbf{v}_n]$.
- 3 For $1 \leq i \leq r$, set $\mathbf{u}_i := \frac{A\mathbf{v}_i}{\|A\mathbf{v}_i\|}$.
- 4 Choose $\mathbf{u}_{r+1}, \dots, \mathbf{u}_m$ to complete $\{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ to an orthonormal basis of \mathbb{R}^m .
- 5 Form $U = [\mathbf{u}_1 \cdots \mathbf{u}_m]$.

Working procedure for computing an SVD

- 1 Determine an orthonormal set of eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$ of $A^T A$ corresponding to eigenvalues $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($n - r$ times).
- 2 Form $V = [\mathbf{v}_1 \cdots \mathbf{v}_n]$.
- 3 For $1 \leq i \leq r$, set $\mathbf{u}_i := \frac{A\mathbf{v}_i}{\|A\mathbf{v}_i\|}$.
- 4 Choose $\mathbf{u}_{r+1}, \dots, \mathbf{u}_m$ to complete $\{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ to an orthonormal basis of \mathbb{R}^m .
- 5 Form $U = [\mathbf{u}_1 \cdots \mathbf{u}_m]$.
- 6 Let Σ have diagonal entries $\sigma_i = \sqrt{\lambda_i}$ for $1 \leq i \leq r$, zeros otherwise. Then $A = U\Sigma V^T$.

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \leq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = A^T A$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($n - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \leq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = A^T A$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($n - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.
- 2 Find the orthonormal set of eigenvectors of L corresponding to each eigenvalue (taken in decreasing order): $\mathbf{v}_1, \dots, \mathbf{v}_n$.

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \leq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = A^T A$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($n - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.
- 2 Find the orthonormal set of eigenvectors of L corresponding to each eigenvalue (taken in decreasing order): $\mathbf{v}_1, \dots, \mathbf{v}_n$.
- 3 Form $V = [\mathbf{v}_1 \dots \mathbf{v}_n]$.

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \leq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = A^T A$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($n - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.
- 2 Find the orthonormal set of eigenvectors of L corresponding to each eigenvalue (taken in decreasing order): $\mathbf{v}_1, \dots, \mathbf{v}_n$.
- 3 Form $V = [\mathbf{v}_1 \dots \mathbf{v}_n]$.
- 4 For $1 \leq i \leq r$, set $\mathbf{u}_i := \frac{A\mathbf{v}_i}{\|A\mathbf{v}_i\|}$.

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \leq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = A^T A$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($n - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.
- 2 Find the orthonormal set of eigenvectors of L corresponding to each eigenvalue (taken in decreasing order): $\mathbf{v}_1, \dots, \mathbf{v}_n$.
- 3 Form $V = [\mathbf{v}_1 \dots \mathbf{v}_n]$.
- 4 For $1 \leq i \leq r$, set $\mathbf{u}_i := \frac{A\mathbf{v}_i}{\|A\mathbf{v}_i\|}$.
- 5 Choose $\mathbf{u}_{r+1}, \dots, \mathbf{u}_m$ to complete $\{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ to an orthonormal vectors in \mathbb{R}^m .

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \leq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = A^T A$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($n - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.
- 2 Find the orthonormal set of eigenvectors of L corresponding to each eigenvalue (taken in decreasing order): $\mathbf{v}_1, \dots, \mathbf{v}_n$.
- 3 Form $V = [\mathbf{v}_1 \dots \mathbf{v}_n]$.
- 4 For $1 \leq i \leq r$, set $\mathbf{u}_i := \frac{A\mathbf{v}_i}{\|A\mathbf{v}_i\|}$.
- 5 Choose $\mathbf{u}_{r+1}, \dots, \mathbf{u}_m$ to complete $\{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ to an orthonormal vectors in \mathbb{R}^m .
- 6 Form $U = [\mathbf{u}_1 \dots \mathbf{u}_m]$.

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \leq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = A^T A$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($n - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.
- 2 Find the orthonormal set of eigenvectors of L corresponding to each eigenvalue (taken in decreasing order): $\mathbf{v}_1, \dots, \mathbf{v}_n$.
- 3 Form $V = [\mathbf{v}_1 \dots \mathbf{v}_n]$.
- 4 For $1 \leq i \leq r$, set $\mathbf{u}_i := \frac{A\mathbf{v}_i}{\|A\mathbf{v}_i\|}$.
- 5 Choose $\mathbf{u}_{r+1}, \dots, \mathbf{u}_m$ to complete $\{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ to an orthonormal vectors in \mathbb{R}^m .
- 6 Form $U = [\mathbf{u}_1 \dots \mathbf{u}_m]$.
- 7 Let Σ have diagonal entries $\sigma_i = \sqrt{\lambda_i}$ for $1 \leq i \leq r$, zeros otherwise. Then $A = U\Sigma V^T$.

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \geq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = AA^T$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($m - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \geq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = AA^T$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($m - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.
- 2 Find the orthonormal set of eigenvectors of L corresponding to each eigenvalue (taken in decreasing order): $\mathbf{u}_1, \dots, \mathbf{u}_m$.

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \geq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = AA^T$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($m - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.
- 2 Find the orthonormal set of eigenvectors of L corresponding to each eigenvalue (taken in decreasing order): $\mathbf{u}_1, \dots, \mathbf{u}_m$.
- 3 Form $U = [\mathbf{u}_1 \dots \mathbf{u}_m]$.

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \geq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = AA^T$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($m - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.
- 2 Find the orthonormal set of eigenvectors of L corresponding to each eigenvalue (taken in decreasing order): $\mathbf{u}_1, \dots, \mathbf{u}_m$.
- 3 Form $U = [\mathbf{u}_1 \dots \mathbf{u}_m]$.
- 4 For $1 \leq i \leq r$, set $\mathbf{v}_i := \frac{A^T \mathbf{u}_i}{\|A^T \mathbf{u}_i\|}$.

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \geq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = AA^T$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($m - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.
- 2 Find the orthonormal set of eigenvectors of L corresponding to each eigenvalue (taken in decreasing order): $\mathbf{u}_1, \dots, \mathbf{u}_m$.
- 3 Form $U = [\mathbf{u}_1 \dots \mathbf{u}_m]$.
- 4 For $1 \leq i \leq r$, set $\mathbf{v}_i := \frac{A^T \mathbf{u}_i}{\|A^T \mathbf{u}_i\|}$.
- 5 Choose $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$ to complete $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ to an orthonormal vectors in \mathbb{R}^n .

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \geq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = AA^T$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($m - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.
- 2 Find the orthonormal set of eigenvectors of L corresponding to each eigenvalue (taken in decreasing order): $\mathbf{u}_1, \dots, \mathbf{u}_m$.
- 3 Form $U = [\mathbf{u}_1 \dots \mathbf{u}_m]$.
- 4 For $1 \leq i \leq r$, set $\mathbf{v}_i := \frac{A^T \mathbf{u}_i}{\|A^T \mathbf{u}_i\|}$.
- 5 Choose $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$ to complete $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ to an orthonormal vectors in \mathbb{R}^n .
- 6 Form $V = [\mathbf{v}_1 \dots \mathbf{v}_n]$.

Working procedure for computing an SVD

Suppose A is $m \times n$ matrix with $m \geq n$

- 1 Determine the largest order matrix between $A^T A$ and AA^T which is $L = AA^T$ in this case. Find the eigenvalues L and arrange in decreasing order: $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $0, \dots, 0$ ($m - r$) times. Singular values are the nonnegative square root of these eigenvalues say $\sigma_1 \geq \dots \geq \sigma_r > 0$ and possibly 0 as well.
- 2 Find the orthonormal set of eigenvectors of L corresponding to each eigenvalue (taken in decreasing order): $\mathbf{u}_1, \dots, \mathbf{u}_m$.
- 3 Form $U = [\mathbf{u}_1 \dots \mathbf{u}_m]$.
- 4 For $1 \leq i \leq r$, set $\mathbf{v}_i := \frac{A^T \mathbf{u}_i}{\|A^T \mathbf{u}_i\|}$.
- 5 Choose $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$ to complete $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ to an orthonormal vectors in \mathbb{R}^n .
- 6 Form $V = [\mathbf{v}_1 \dots \mathbf{v}_n]$.
- 7 Let Σ have diagonal entries $\sigma_i = \sqrt{\lambda_i}$ for $1 \leq i \leq r$, zeros otherwise. Then $A = U\Sigma V^T$.