

# Graph Theory-Class 4

A **directed graph or digraph**  $D$  consists of


- ▶ a finite nonempty set  $V$  of points together with,

A **directed graph or digraph**  $D$  consists of

- ▶ a finite nonempty set  $V$  of points together with,
- ▶ a prescribed collection  $E$  of ordered pairs of distinct points.


A **directed graph or digraph**  $D$  consists of

- ▶ a finite nonempty set  $V$  of points together with,
- ▶ a prescribed collection  $E$  of ordered pairs of distinct points.



A **directed graph or digraph**  $D$  consists of


- ▶ a finite nonempty set  $V$  of points together with,
- ▶ a prescribed collection  $E$  of ordered pairs of distinct points.



The elements of  $E$  are **directed lines** or **arcs**.

A **directed graph or digraph**  $D$  consists of

- ▶ a finite nonempty set  $V$  of points together with,
- ▶ a prescribed collection  $E$  of ordered pairs of distinct points.




The elements of  $E$  are **directed lines** or **arcs**.

In above graph  $E = \{(v_1, v_2), (v_2, v_1), (v_1, v_3)\}$ .

A **weighted graph** is a graph where each edge is assigned a numerical value, called a weight.

A **weighted graph** is a graph where each edge is assigned a numerical value, called a weight.





# Finding shortest path from a given point to all other points

Let  $G$  be a directed graph (digraph) with  $p$  vertices.

# Finding shortest path from a given point to all other points

Let  $G$  be a directed graph (digraph) with  $p$  vertices.

The **distance matrix**  $D = (d_{ij})$  associated with  $G$  is defined as follows:

# Finding shortest path from a given point to all other points

Let  $G$  be a directed graph (digraph) with  $p$  vertices.

The **distance matrix**  $D = (d_{ij})$  associated with  $G$  is defined as follows:

- ▶  $d_{ij} = 0$ , if  $i = j$

# Finding shortest path from a given point to all other points

Let  $G$  be a directed graph (digraph) with  $p$  vertices.

The **distance matrix**  $D = (d_{ij})$  associated with  $G$  is defined as follows:

- ▶  $d_{ij} = 0$ , if  $i = j$
- ▶  $d_{ij} = \infty$ , if there is no edge from  $v_i$  to  $v_j$

# Finding shortest path from a given point to all other points

Let  $G$  be a directed graph (digraph) with  $p$  vertices.

The **distance matrix**  $D = (d_{ij})$  associated with  $G$  is defined as follows:

- ▶  $d_{ij} = 0$ , if  $i = j$
- ▶  $d_{ij} = \infty$ , if there is no edge from  $v_i$  to  $v_j$
- ▶  $d_{ij} = \text{weight/distance of the edge from } v_i \text{ to } v_j$ , if such an edge exists.

**PLEASE REFER TO GRAPH THEORY NOTES FOR THE Dijkstra's algorithm.**