

Graph Theory-Class 2

Basic Definitions

Basic Definitions

A graph in which every pair of points are adjacent is called

Basic Definitions

A graph in which every pair of points are adjacent is called **Complete graph**.

Basic Definitions

A graph in which every pair of points are adjacent is called **Complete graph**.

A complete graph with p vertices is denoted by K_p .

Basic Definitions

A graph in which every pair of points are adjacent is called **Complete graph**.

A complete graph with p vertices is denoted by K_p .

Result:

Basic Definitions

A graph in which every pair of points are adjacent is called **Complete graph**.

A complete graph with p vertices is denoted by K_p .

Result: A complete graph with p vertices has $\frac{p(p-1)}{2}$.

Basic Definitions

A graph in which every pair of points are adjacent is called **Complete graph**.

A complete graph with p vertices is denoted by K_p .

Result: A complete graph with p vertices has $\frac{p(p-1)}{2}$. **Hint:** Use degree sum formula.


Definitions

Definitions

- ▶ **Length of a walk:** The number of occurrences of edges in the walk.

Definitions


- **Length of a walk:** The number of occurrences of edges in the walk.



1. $s \ a \ t \ e \ u \ f \ v \ d \ t \ a \ s$ -

Definitions


- **Length of a walk:** The number of occurrences of edges in the walk.



1. $s \ a \ t \ e \ u \ f \ v \ d \ t \ a \ s - 5.$

Definitions


- **Length of a walk:** The number of occurrences of edges in the walk.



1. $s a t e u f v d t a s - 5.$
2. $u e t a s b u c w -$

Definitions


- **Length of a walk:** The number of occurrences of edges in the walk.



1. $s a t e u f v d t a s - 5.$
2. $u e t a s b u c w - 4.$

Definitions

- **Length of a walk:** The number of occurrences of edges in the walk.




1. $s a t e u f v d t a s - 5.$

2. $u e t a s b u c w - 4.$

- **Distance between two vertices:** The distance between vertices u and v in a graph G is the length of a shortest path joining them.

Definitions

- **Length of a walk:** The number of occurrences of edges in the walk.



1. $s a t e u f v d t a s - 5$.
2. $u e t a s b u c w - 4$.

- **Distance between two vertices:** The distance between vertices u and v in a graph G is the length of a shortest path joining them. If no such path exists, the distance is defined to be ∞ .

Connectivity

Connectivity

Connected Vertices: Two vertices u and v are said to be *connected* if there exists a path between them in G .

Connectivity


Connected Vertices: Two vertices u and v are said to be *connected* if there exists a path between them in G .

Connected Graph: A graph is *connected* if every pair of vertices is connected by a path.

Connectivity

Connected Vertices: Two vertices u and v are said to be *connected* if there exists a path between them in G .



Connected Graph: A graph is *connected* if every pair of vertices is connected by a path.



Connectivity

Connected Vertices: Two vertices u and v are said to be *connected* if there exists a path between them in G .


Connected Graph: A graph is *connected* if every pair of vertices is connected by a path.





Subgraph: Let $G(V, E)$ be a graph. A graph H is called a subgraph of G if

Subgraph: Let $G(V, E)$ be a graph. A graph H is called a subgraph of G if $V(H) \subset V(G)$ and $E(H) \subset E(G)$.




Subgraph: Let $G(V, E)$ be a graph. A graph H is called a subgraph of G if $V(H) \subset V(G)$ and $E(H) \subset E(G)$.



Subgraph: Let $G(V, E)$ be a graph. A graph H is called a subgraph of G if $V(H) \subset V(G)$ and $E(H) \subset E(G)$.



Subgraph: Let $G(V, E)$ be a graph. A graph H is called a subgraph of G if $V(H) \subset V(G)$ and $E(H) \subset E(G)$.




Supergraph: If H is a subgraph of G , then G is called supergraph of H .

Supergraph: If H is a subgraph of G , then G is called supergraph of H .

Spanning subgraph: A subgraph H of G such that $V(H) = V(G)$.



Supergraph: If H is a subgraph of G , then G is called supergraph of H .

Spanning subgraph: A subgraph H of G such that $V(H) = V(G)$.




Supergraph: If H is a subgraph of G , then G is called supergraph of H .

Spanning subgraph: A subgraph H of G such that $V(H) = V(G)$.




Spanning Subgraphs of C_3




the graph C_3

Spanning Subgraphs of C_3




the graph C_3


Spanning Subgraphs of C_3




the graph C_3




Spanning Subgraphs of C_3



the graph C_3




Spanning Subgraphs of C_3




the graph C_3




Spanning Subgraphs of C_3




the graph C_3




Spanning Subgraphs of C_3




the graph C_3




Spanning Subgraphs of C_3




the graph C_3



Spanning Subgraphs of C_3



the graph C_3



Spanning subgraphs of C_3

Different types of subgraphs

Different types of subgraphs

Removal of a point v : Let G be a graph and $v \in V(G)$, the subgraph $H = G - v$ consists of all points of G except v and all lines of G except those lines incident with v .

Different types of subgraphs

Removal of a point v : Let G be a graph and $v \in V(G)$, the subgraph $H = G - v$ consists of all points of G except v and all lines of G except those lines incident with v .

Removal of an edge x : Let G be a graph and $x \in E(G)$, the subgraph $H = G - x$ is a spanning subgraph of G with all edges except x .

Different types of subgraphs

Removal of a point v : Let G be a graph and $v \in V(G)$, the subgraph $H = G - v$ consists of all points of G except v and all lines of G except those lines incident with v .

Removal of an edge x : Let G be a graph and $x \in E(G)$, the subgraph $H = G - x$ is a spanning subgraph of G with all edges except x .

Maximal subgraph A subgraph H of G is said to be maximal with respect to a property P if

- i. H satisfies property P
- ii. if H' is another subgraph of G which is a supergraph of H then H' does not satisfy property P .

Different types of subgraphs

Removal of a point v : Let G be a graph and $v \in V(G)$, the subgraph $H = G - v$ consists of all points of G except v and all lines of G except those lines incident with v .

Removal of an edge x : Let G be a graph and $x \in E(G)$, the subgraph $H = G - x$ is a spanning subgraph of G with all edges except x .

Maximal subgraph A subgraph H of G is said to be maximal with respect to a property P if

- i. H satisfies property P
- ii. if H' is another subgraph of G which is a supergraph of H then H' does not satisfy property P . **OR**
 H is not a subgraph of any supergraph with property P .

Different types of subgraphs

Removal of a point v : Let G be a graph and $v \in V(G)$, the subgraph $H = G - v$ consists of all points of G except v and all lines of G except those lines incident with v .

Removal of an edge x : Let G be a graph and $x \in E(G)$, the subgraph $H = G - x$ is a spanning subgraph of G with all edges except x .

Maximal subgraph A subgraph H of G is said to be maximal with respect to a property P if

- i. H satisfies property P
- ii. if H' is another subgraph of G which is a supergraph of H then H' does not satisfy property P . **OR**
 H is not a subgraph of any supergraph with property P .

We will see an example now.

Different types of subgraphs

Removal of a point v : Let G be a graph and $v \in V(G)$, the subgraph $H = G - v$ consists of all points of G except v and all lines of G except those lines incident with v .

Removal of an edge x : Let G be a graph and $x \in E(G)$, the subgraph $H = G - x$ is a spanning subgraph of G with all edges except x .


Maximal subgraph A subgraph H of G is said to be maximal with respect to a property P if

- i. H satisfies property P
- ii. if H' is another subgraph of G which is a supergraph of H then H' does not satisfy property P . **OR**
 H is not a subgraph of any supergraph with property P .


We will see an example now.

Acyclic graph: A graph with no cycles.

An example




An example




Let property P : Subgraph has no cycles.

An example




Let property P : Subgraph has no cycles. Draw a maximal subgraph with property P .


An example




Let property P : Subgraph has no cycles. Draw a maximal subgraph with property P .



An example



Let property P : Subgraph has no cycles. Draw a maximal subgraph with property P .



Maximal Acyclic Subgraph of C_3

Induced subgraph Let G be a graph and $S \subset V(G)$, the induced subgraph with respect to set S , denoted by $\langle S \rangle$ is the maximal subgraph of G with point set S .

Induced subgraph Let G be a graph and $S \subset V(G)$, the induced subgraph with respect to set S , denoted by $\langle S \rangle$ is the maximal subgraph of G with point set S .


In otherwords, if G is a graph, then a subgraph $H = \langle S \rangle$ is induced subgraph if

- ▶ $V(H) = S$ and

Induced subgraph Let G be a graph and $S \subset V(G)$, the induced subgraph with respect to set S , denoted by $\langle S \rangle$ is the maximal subgraph of G with point set S .


In otherwords, if G is a graph, then a subgraph $H = \langle S \rangle$ is induced subgraph if

- ▶ $V(H) = S$ and
- ▶ H is not a subgraph of any supergraph H' with vertex set S .




Graph G

Let $S = v_1, v_4, v_5$. Draw induced subgraph of G with set S .



Graph G

Let $S = v_1, v_4, v_5$. Draw induced subgraph of G with set S .




Task

Task

- ▶ We know that two vertices u and v in a graph G are connected if _____.

Task

- ▶ We know that two vertices u and v in a graph G are connected if _____.
- ▶ Based on connectedness of vertices can you separate/divide the vertex set V of the following graph?



An equivalence relation on vertex set

Let $G(V, E)$ be a graph. Define a relation on $V(G)$ as follows:
 $u \sim v$ (u is related to v) if u and v are connected.

An equivalence relation on vertex set

Let $G(V, E)$ be a graph. Define a relation on $V(G)$ as follows:
 $u \sim v$ (u is related to v) if u and v are connected.

- ▶ This relation is an **equivalence relation** on the vertex set $V(G)$.

An equivalence relation on vertex set

Let $G(V, E)$ be a graph. Define a relation on $V(G)$ as follows:
 $u \sim v$ (u is related to v) if u and v are connected.

- ▶ This relation is an **equivalence relation** on the vertex set $V(G)$.
- ▶ Thus, there is a partition of $V(G)$ into non empty subsets V_1, V_2, \dots, V_k such that two vertices u and v are connected if and only if both u and v belongs to the same set V_i .

An equivalence relation on vertex set

Let $G(V, E)$ be a graph. Define a relation on $V(G)$ as follows:
 $u \sim v$ (u is related to v) if u and v are connected.

- ▶ This relation is an **equivalence relation** on the vertex set $V(G)$.
- ▶ Thus, there is a partition of $V(G)$ into non empty subsets V_1, V_2, \dots, V_k such that two vertices u and v are connected if and only if both u and v belongs to the same set V_i .
- ▶ The induced subgraphs $\langle V_1 \rangle, \langle V_2 \rangle, \dots, \langle V_k \rangle$ are called the (connected) components of G .

An equivalence relation on vertex set

Let $G(V, E)$ be a graph. Define a relation on $V(G)$ as follows:
 $u \sim v$ (u is related to v) if u and v are connected.

- ▶ This relation is an **equivalence relation** on the vertex set $V(G)$.
- ▶ Thus, there is a partition of $V(G)$ into non empty subsets V_1, V_2, \dots, V_k such that two vertices u and v are connected if and only if both u and v belongs to the same set V_i .
- ▶ The induced subgraphs $\langle V_1 \rangle, \langle V_2 \rangle, \dots, \langle V_k \rangle$ are called the (connected) components of G .
- ▶ If G has exactly one component, G is connected; otherwise G is disconnected.

An equivalence relation on vertex set

Let $G(V, E)$ be a graph. Define a relation on $V(G)$ as follows:
 $u \sim v$ (u is related to v) if u and v are connected.

- ▶ This relation is an **equivalence relation** on the vertex set $V(G)$.
- ▶ Thus, there is a partition of $V(G)$ into non empty subsets V_1, V_2, \dots, V_k such that two vertices u and v are connected if and only if both u and v belongs to the same set V_i .
- ▶ The induced subgraphs $\langle V_1 \rangle, \langle V_2 \rangle, \dots, \langle V_k \rangle$ are called the (connected) components of G .
- ▶ If G has exactly one component, G is connected; otherwise G is disconnected.
- ▶ We denote the number of components of G by $\omega(G)$.