

# Toward Particle ID in Granular Hadron Calorimeters

Andrea De Vita<sup>1,2</sup>, Tommaso Dorigo<sup>1,6</sup>, Abhishek<sup>5</sup>, Max Aehle<sup>4</sup>, Nicholas R. Gauger<sup>4</sup>, Muhammad Awais<sup>2,6</sup>,

Jan Kieseler<sup>3</sup>, Ralf Keidel<sup>4</sup>, Tobias Kortus<sup>4</sup>, Enrico Lupi<sup>1,2</sup>, Xuan-Tung Nguyen<sup>1,4</sup>,

Fredrik Sandin<sup>6</sup>, Alexander Schilling<sup>4</sup>, Joseph Willmore<sup>1</sup>

<sup>1</sup> National Institute for Nuclear Physics, Padova, Italy


<sup>2</sup> University of Padova, Italy

<sup>3</sup> Karlsruhe Institute of Technology, Germany

<sup>4</sup> University of Kaiserslautern-Landau, Germany

<sup>5</sup> National Institute of Science Education and Research, India

<sup>6</sup> Luleå University of Technology, Sweden



4th Mode Workshop - Valencia (Spain)

Poster Session - 24th September 2024



## Introduction

This study investigates whether **high-granularity** hadronic calorimeters can differentiate between protons, charged pions, and kaons by analyzing detailed **energy deposition patterns**, with promising preliminary results from Geant4 simulations.

## Simulation Setup

Material: PbWO<sub>4</sub>  
Segmentation: 100x100x100 cells  
Cell size: 3x3x12 mm<sup>3</sup>


$p, \pi^+, K^+$   
 $E = 100 \text{ GeV}$



Projection of the energy distribution in the XY and ZY planes.


## Machine Learning Strategy

Our work proposes the use of **XGBoost Boosted Decision Trees (BDTs)** to analyse descriptive features for each event. The approach includes a preprocessing step that generates variables for each event, which are then input into the machine learning algorithm. Hyperparameter optimization is conducted using **GridSearch**, exploring different configurations, including the choice of booster and tree method type.




## Meaningful Shower Features

To study particle interactions, identifying the **primary interaction vertex** is crucial, as it reveals key information about the particle. Detector segmentation, particularly longitudinal, enables detailed analysis of the shower's energy profile. A **moving window algorithm** helps locate the primary vertex near an energy peak. Further studies can focus on the energy around the vertex, its relationship to secondary vertices, and **shower dimensions**, including average size and asymmetries from non-interacting secondary particles.



Shower's energy profile along the beam axis.



Distribution of the radius of the shower for  $p, \pi^+$  and  $K^+$ .

## Performance

After preprocessing and generating a set of descriptive variables for each event, BDTs were trained using around 40,000 samples per particle, characterised by **49 features**. The training process employed 5-fold cross-validation to assess the architecture's performance. The results for two classification tasks are presented: on the left is the confusion matrix for binary classification of protons vs pions, achieving **63% accuracy**; on the right, the confusion matrix shows the classification of protons, charged pions, and charged kaons, with an **accuracy of 45%**.

	Predicted	
	proton	pion
True proton	7013	2966
True pion	4483	5450

	Predicted		
	proton	pion	kaon
True proton	5585	1887	2506
True pion	3216	4032	2686
True kaon	3788	2630	3550

## Future Perspectives

Future studies will examine **how cell size affects the performance** of the particle identification algorithm, balancing cost and benefit. Moreover, the **behaviour at different energies** will be studied. Our study will also combine different machine learning algorithms in order to exploit the 3D shower pattern, for example using **BDTs with CNNs**. Additionally, new materials and geometries will be investigated to highlight specific hadron properties.

## References

- N Akchurin, et al., On the differences between high-energy proton and pion showers and their signals in a non-compensating calorimeter, Nuclear Instruments and Methods in Physics Research Section A, [[https://doi.org/10.1016/S0168-9002\(98\)00021-7](https://doi.org/10.1016/S0168-9002(98)00021-7)]
- Related works: Enrico Lupi's and Xuan-Tung Nguyen's posters (4th Mode Workshop)