

# SVM Dual Problem

He He

CDS, NYU

Feb 23, 2021

## SVM as a Quadratic Program

- The SVM optimization problem is equivalent to

$$\begin{aligned} \text{minimize} \quad & \frac{1}{2} \|w\|^2 + \frac{c}{n} \sum_{i=1}^n \xi_i \\ \text{subject to} \quad & -\xi_i \leq 0 \quad \text{for } i = 1, \dots, n \\ & (1 - y_i [w^T x_i + b]) - \xi_i \leq 0 \quad \text{for } i = 1, \dots, n \end{aligned}$$

- Differentiable objective function
- $n + d + 1$  unknowns and  $2n$  affine constraints.
- A **quadratic program** that can be solved by any off-the-shelf QP solver.
- Let's learn more by examining the dual.

## Why Do We Care About the Dual?

# The Lagrangian

The general [inequality-constrained] optimization problem is:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m\end{array}$$

## Definition

The **Lagrangian** for this optimization problem is

$$L(x, \lambda) = f_0(x) + \sum_{i=1}^m \lambda_i f_i(x).$$

- $\lambda_i$ 's are called **Lagrange multipliers** (also called the **dual variables**).
- Weighted sum of the objective and constraint functions
- Hard constraints  $\rightarrow$  soft constraints

# Lagrange Dual Function

## Definition

The **Lagrange dual function** is

$$g(\lambda) = \inf_x L(x, \lambda) = \inf_x \left( f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) \right)$$

- $g(\lambda)$  is **concave** (why?)
- **Lower bound property**: if  $\lambda \succeq 0$ ,  $g(\lambda) \leq p^*$  where  $p^*$  is the optimal value of the optimization problem.
- $g(\lambda)$  can be  $-\infty$  (uninformative lower bound)

1.  $g$  is concave because it is the infimum of affine functions. Note that we are not assuming convexity of  $f_0$ .
2. Note that the proof is straightforward:  $\sum_i \lambda_i f_i(x)$  is always negative.
3. For example when  $L(x, \lambda)$  is affine in  $x$ .
4. We can consider  $g(\lambda)$  as a parametrized lower bound that depends on  $\lambda$ . So we might want to find  $\lambda$  that gives us the best lower bound, which motivates the dual problem.

# The Primal and the Dual

- For any **primal form** optimization problem,

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m,\end{array}$$


there is a recipe for constructing a corresponding **Lagrangian dual problem**:

$$\begin{array}{ll}\text{maximize} & g(\lambda) \\ \text{subject to} & \lambda_i \geq 0, \quad i = 1, \dots, m,\end{array}$$

- The dual problem is always a convex optimization problem.
- The dual variables often have interesting and relevant interpretations.
- The dual variables provide certificate for optimality.

## Weak Duality


We always have **weak duality**:  $p^* \geq d^*$ .



Plot courtesy of Brett Bernstein.

## Strong Duality

For some problems, we have **strong duality**:  $p^* = d^*$ .



For convex problems, strong duality is fairly typical.

Plot courtesy of Brett Bernstein.



- Assume strong duality. Let  $x^*$  be primal optimal and  $\lambda^*$  be dual optimal. Then:

$$\begin{aligned} f_0(x^*) &= g(\lambda^*) = \inf_x L(x, \lambda^*) \quad (\text{strong duality and definition}) \\ &\leq L(x^*, \lambda^*) \\ &= f_0(x^*) + \sum_{i=1}^m \lambda_i^* f_i(x^*) \\ &\leq f_0(x^*). \end{aligned}$$

Each term in sum  $\sum_{i=1}^m \lambda_i^* f_i(x^*)$  must actually be 0. That is

$$\lambda_i > 0 \implies f_i(x^*) = 0 \quad \text{and} \quad f_i(x^*) < 0 \implies \lambda_i = 0 \quad \forall i$$

This condition is known as **complementary slackness**.

## The SVM Dual Problem

$$\begin{aligned} \text{minimize} \quad & \frac{1}{2} \|w\|^2 + \frac{c}{n} \sum_{i=1}^n \xi_i \\ \text{subject to} \quad & -\xi_i \leq 0 \quad \text{for } i = 1, \dots, n \\ & (1 - y_i [w^T x_i + b]) - \xi_i \leq 0 \quad \text{for } i = 1, \dots, n \end{aligned}$$

Lagrange Multiplier	Constraint
$\lambda_i$	$-\xi_i \leq 0$
$\alpha_i$	$(1 - y_i [w^T x_i + b]) - \xi_i \leq 0$

$$L(w, b, \xi, \alpha, \lambda) = \frac{1}{2} \|w\|^2 + \frac{c}{n} \sum_{i=1}^n \xi_i + \sum_{i=1}^n \alpha_i (1 - y_i [w^T x_i + b] - \xi_i) + \sum_{i=1}^n \lambda_i (-\xi_i)$$

Dual optimum value:  $d^* = \sup_{\alpha, \lambda \succeq 0} \inf_{w, b, \xi} L(w, b, \xi, \alpha, \lambda)$

1. What are the primal and dual variables?

## Strong Duality by Slater's Constraint Qualification

The SVM optimization problem:

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \|w\|^2 + \frac{c}{n} \sum_{i=1}^n \xi_i \\ &\text{subject to} && -\xi_i \leq 0 \text{ for } i = 1, \dots, n \\ &&& (1 - y_i [w^T x_i + b]) - \xi_i \leq 0 \text{ for } i = 1, \dots, n \end{aligned}$$

Slater's constraint qualification:

- Convex problem + affine constraints  $\implies$  strong duality iff problem is feasible
- Do we have a feasible point?
- For SVM, we have **strong duality**.

1. Constraints are satisfied by  $w = b = 0$  and  $\xi_i = 1$  for  $i = 1, \dots, n$

## SVM Dual Function: First Order Conditions

Lagrange dual function is the inf over primal variables of  $L$ :

$$g(\alpha, \lambda) = \inf_{w, b, \xi} L(w, b, \xi, \alpha, \lambda)$$
$$= \inf_{w, b, \xi} \left[ \frac{1}{2} w^T w + \sum_{i=1}^n \xi_i \left( \frac{c}{n} - \alpha_i - \lambda_i \right) + \sum_{i=1}^n \alpha_i (1 - y_i [w^T x_i + b]) \right]$$

$$\partial_w L = 0 \iff w - \sum_{i=1}^n \alpha_i y_i x_i = 0 \iff \boxed{w = \sum_{i=1}^n \alpha_i y_i x_i}$$

$$\partial_b L = 0 \iff - \sum_{i=1}^n \alpha_i y_i = 0 \iff \boxed{\sum_{i=1}^n \alpha_i y_i = 0}$$

$$\partial_{\xi_i} L = 0 \iff \frac{c}{n} - \alpha_i - \lambda_i = 0 \iff \boxed{\alpha_i + \lambda_i = \frac{c}{n}}$$

1. How do we solve the minimization problem?
2. Is it convex? [yes, quadratic term]

- Substituting these conditions back into  $L$ , the second term disappears.
- First and third terms become

$$\begin{aligned}\frac{1}{2}w^T w &= \frac{1}{2} \sum_{i,j=1}^n \alpha_i \alpha_j y_i y_j x_i^T x_j \\ \sum_{i=1}^n \alpha_i (1 - y_i [w^T x_i + b]) &= \sum_{i=1}^n \alpha_i - \sum_{i,j=1}^n \alpha_i \alpha_j y_i y_j x_j^T x_i - b \underbrace{\sum_{i=1}^n \alpha_i y_i}_{=0}.\end{aligned}$$

- Putting it together, the dual function is

$$g(\alpha, \lambda) = \begin{cases} \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i,j=1}^n \alpha_i \alpha_j y_i y_j x_j^T x_i & \begin{matrix} \sum_{i=1}^n \alpha_i y_i = 0 \\ \alpha_i + \lambda_i = \frac{\epsilon}{n}, \text{ all } i \end{matrix} \\ -\infty & \text{otherwise.} \end{cases}$$

- The **dual function** is

$$g(\alpha, \lambda) = \begin{cases} \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i,j=1}^n \alpha_i \alpha_j y_i y_j x_j^T x_i & \begin{array}{l} \sum_{i=1}^n \alpha_i y_i = 0 \\ \alpha_i + \lambda_i = \frac{c}{n}, \text{ all } i \end{array} \\ -\infty & \text{otherwise.} \end{cases}$$

- The **dual problem** is  $\sup_{\alpha, \lambda \succeq 0} g(\alpha, \lambda)$ :

$$\begin{aligned} \sup_{\alpha, \lambda} \quad & \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i,j=1}^n \alpha_i \alpha_j y_i y_j x_j^T x_i \\ \text{s.t.} \quad & \sum_{i=1}^n \alpha_i y_i = 0 \\ & \alpha_i + \lambda_i = \frac{c}{n} \quad \alpha_i, \lambda_i \geq 0, \quad i = 1, \dots, n \end{aligned}$$

## Insights from the Dual Problem



# KKT Conditions

For **convex** problems, if **Slater's condition** is satisfied, then **KKT conditions** provide **necessary and sufficient** conditions for the optimal solution.

- Primal feasibility:  $f_i(x) \leq 0 \quad \forall i$
- Dual feasibility:  $\lambda \succeq 0$
- Complementary slackness:  $\lambda_i f_i(x) = 0$
- First-order condition:

$$\frac{\partial}{\partial x} L(x, \lambda) = 0$$

1.  $x$  needs to be a stationary point of the Lagrangian.

## The SVM Dual Solution

- We found the SVM dual problem can be written as:

$$\begin{aligned} \sup_{\alpha} \quad & \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i,j=1}^n \alpha_i \alpha_j y_i y_j x_j^T x_i \\ \text{s.t.} \quad & \sum_{i=1}^n \alpha_i y_i = 0 \\ & \alpha_i \in \left[0, \frac{c}{n}\right] \quad i = 1, \dots, n. \end{aligned}$$

- Given solution  $\alpha^*$  to dual, primal solution is  $w^* = \sum_{i=1}^n \alpha_i^* y_i x_i$ .
- The solution is in the space spanned by the inputs.
- Note  $\alpha_i^* \in [0, \frac{c}{n}]$ . So  $c$  controls max weight on each example. (**Robustness!**)
  - What's the relation between  $c$  and regularization?

1. If  $c$  is small, the solution is not sensitive to any single example—strong regularization. We can also see this in the primal problem: small  $c$  corresponds to larger coefficients for the regularization term.

## Complementary Slackness Conditions

- Recall our primal constraints and Lagrange multipliers:

Lagrange Multiplier	Constraint
$\lambda_i$	$-\xi_i \leq 0$
$\alpha_i$	$(1 - y_i f(x_i)) - \xi_i \leq 0$

- Recall first order condition  $\nabla_{\xi_i} L = 0$  gave us  $\lambda_i^* = \frac{c}{n} - \alpha_i^*$ .
- By strong duality, we must have **complementary slackness**:

$$\alpha_i^* (1 - y_i f^*(x_i) - \xi_i^*) = 0$$

$$\lambda_i^* \xi_i^* = \left( \frac{c}{n} - \alpha_i^* \right) \xi_i^* = 0$$

## Consequences of Complementary Slackness

By strong duality, we must have **complementary slackness**.

$$\alpha_i^* (1 - y_i f^*(x_i) - \xi_i^*) = 0$$
$$\left(\frac{c}{n} - \alpha_i^*\right) \xi_i^* = 0$$

Recall “**slack variable**”  $\xi_i^* = \max(0, 1 - y_i f^*(x_i))$  is the hinge loss on  $(x_i, y_i)$ .

- If  $y_i f^*(x_i) > 1$  then the margin loss is  $\xi_i^* = 0$ , and we get  $\alpha_i^* = 0$ .
- If  $y_i f^*(x_i) < 1$  then the margin loss is  $\xi_i^* > 0$ , so  $\alpha_i^* = \frac{c}{n}$ .
- If  $\alpha_i^* = 0$ , then  $\xi_i^* = 0$ , which implies no loss, so  $y_i f^*(x) \geq 1$ .
- If  $\alpha_i^* \in (0, \frac{c}{n})$ , then  $\xi_i^* = 0$ , which implies  $1 - y_i f^*(x_i) = 0$ .

## Complementary Slackness Results: Summary

If  $\alpha^*$  is a solution to the dual problem, then primal solution is

$$w^* = \sum_{i=1}^n \alpha_i^* y_i x_i \quad \text{where } \alpha_i^* \in [0, \frac{c}{n}].$$

Relation between margin and example weights ( $\alpha_i$ 's):

$$\alpha_i^* = 0 \implies y_i f^*(x_i) \geq 1$$

$$\alpha_i^* \in \left(0, \frac{c}{n}\right) \implies y_i f^*(x_i) = 1$$

$$\alpha_i^* = \frac{c}{n} \implies y_i f^*(x_i) \leq 1$$

$$y_i f^*(x_i) < 1 \implies \alpha_i^* = \frac{c}{n}$$

$$y_i f^*(x_i) = 1 \implies \alpha_i^* \in \left[0, \frac{c}{n}\right]$$

$$y_i f^*(x_i) > 1 \implies \alpha_i^* = 0$$

- If  $\alpha^*$  is a solution to the dual problem, then primal solution is

$$w^* = \sum_{i=1}^n \alpha_i^* y_i x_i$$

with  $\alpha_i^* \in [0, \frac{c}{n}]$ .

- The  $x_i$ 's corresponding to  $\alpha_i^* > 0$  are called **support vectors**.
- Few margin errors or “on the margin” examples  $\implies$  [sparsity in input examples](#).

## The Bias Term: $b$

- For our SVM primal, the complementary slackness conditions are:

$$\alpha_i^* (1 - y_i [x_i^T w^* + b] - \xi_i^*) = 0 \quad (1)$$

$$\lambda_i^* \xi_i^* = \left( \frac{c}{n} - \alpha_i^* \right) \xi_i^* = 0 \quad (2)$$

- Suppose there's an  $i$  such that  $\alpha_i^* \in (0, \frac{c}{n})$ .

- (2) implies  $\xi_i^* = 0$ .

- (1) implies

$$y_i [x_i^T w^* + b^*] = 1$$

$$\iff x_i^T w^* + b^* = y_i \text{ (use } y_i \in \{-1, 1\})$$

$$\iff \boxed{b^* = y_i - x_i^T w^*}$$

## The Bias Term: $b$

- We get the same  $b^*$  for any choice of  $i$  with  $\alpha_i^* \in (0, \frac{c}{n})$

$$b^* = y_i - x_i^T w^*$$

- With numerical error, more robust to average over all eligible  $i$ 's:

$$b^* = \text{mean} \left\{ y_i - x_i^T w^* \mid \alpha_i^* \in \left( 0, \frac{c}{n} \right) \right\}.$$

- If there are no  $\alpha_i^* \in (0, \frac{c}{n})$ ?
  - Then we have a **degenerate SVM training problem**<sup>1</sup> ( $w^* = 0$ ).

---

<sup>1</sup>See Rifkin et al.'s "A Note on Support Vector Machine Degeneracy", an MIT AI Lab Technical Report.



## Teaser for Kernelization

## Dual Problem: Dependence on $x$ through inner products

- SVM Dual Problem:

$$\begin{aligned} \sup_{\alpha} \quad & \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i,j=1}^n \alpha_i \alpha_j y_i y_j x_j^T x_i \\ \text{s.t.} \quad & \sum_{i=1}^n \alpha_i y_i = 0 \\ & \alpha_i \in \left[0, \frac{C}{n}\right] \quad i = 1, \dots, n. \end{aligned}$$

- Note that all dependence on inputs  $x_i$  and  $x_j$  is through their inner product:  $\langle x_j, x_i \rangle = x_j^T x_i$ .
- We can replace  $x_j^T x_i$  by other products...
- This is a “kernelized” objective function.