



# **CS 228 : Logic in Computer Science**

Krishna. S

# FO without equality

---

Let us focus on FO without “=”. Recall that “=” is always interpreted as equality.

## Herbrand Theorem

Let  $\Gamma = \{\varphi_1, \varphi_2, \dots\}$  be a set of equality-free sentences in Skolem Normal Form. Then  $\Gamma$  is satisfiable iff  $\Gamma$  has a Herbrand model.

If  $\Gamma$  has a Herbrand model, clearly  $\Gamma$  is satisfiable. The converse needs a proof.

# The converse

---

Assume  $\Gamma$  is satisfiable. Let  $\tau_H$  be the Herbrand signature for  $\Gamma$ .

- ▶ Let  $\mathcal{A}$  be a  $\tau_H$  structure such that  $\mathcal{A} \models \Gamma$ . ( $U^{\mathcal{A}}$  need not be the Herbrand universe)
- ▶ Let  $\mathcal{B}$  be a Herbrand structure over  $\tau_H$ . ( $U^{\mathcal{B}}$  is the Herbrand universe)

# The converse

---

Assume  $\Gamma$  is satisfiable. Let  $\tau_H$  be the Herbrand signature for  $\Gamma$ .

- ▶ Let  $\mathcal{A}$  be a  $\tau_H$  structure such that  $\mathcal{A} \models \Gamma$ . ( $U^{\mathcal{A}}$  need not be the Herbrand universe)
- ▶ Let  $\mathcal{B}$  be a Herbrand structure over  $\tau_H$ . ( $U^{\mathcal{B}}$  is the Herbrand universe)
- ▶ We need to create a Herbrand model for  $\Gamma$

# The converse

---

Assume  $\Gamma$  is satisfiable. Let  $\tau_H$  be the Herbrand signature for  $\Gamma$ .

- ▶ Let  $\mathcal{A}$  be a  $\tau_H$  structure such that  $\mathcal{A} \models \Gamma$ . ( $U^{\mathcal{A}}$  need not be the Herbrand universe)
- ▶ Let  $\mathcal{B}$  be a Herbrand structure over  $\tau_H$ . ( $U^{\mathcal{B}}$  is the Herbrand universe)
- ▶ We need to create a Herbrand model for  $\Gamma$
- ▶ Try “merging”  $\mathcal{A}$  and  $\mathcal{B}$  to obtain a Herbrand model  $M$  for  $\Gamma$ .

# The converse

---

Assume  $\Gamma$  is satisfiable. Let  $\tau_H$  be the Herbrand signature for  $\Gamma$ .

- ▶ Let  $\mathcal{A}$  be a  $\tau_H$  structure such that  $\mathcal{A} \models \Gamma$ . ( $U^{\mathcal{A}}$  need not be the Herbrand universe)
- ▶ Let  $\mathcal{B}$  be a Herbrand structure over  $\tau_H$ . ( $U^{\mathcal{B}}$  is the Herbrand universe)
- ▶ We need to create a Herbrand model for  $\Gamma$
- ▶ Try “merging”  $\mathcal{A}$  and  $\mathcal{B}$  to obtain a Herbrand model  $M$  for  $\Gamma$ .
  - ▶ Define  $M$  so that its universe is the Herbrand universe over  $\tau_H$ .
  - ▶ Let  $M$  interpret functions and constants like  $\mathcal{B}$  (both have the same Herbrand universe)
  - ▶ How to interpret the relations from  $\tau_H$  (same as those of  $\tau$ )?

# The converse

---

Assume  $\Gamma$  is satisfiable. Let  $\tau_H$  be the Herbrand signature for  $\Gamma$ .

- ▶ Let  $\mathcal{A}$  be a  $\tau_H$  structure such that  $\mathcal{A} \models \Gamma$ . ( $U^{\mathcal{A}}$  need not be the Herbrand universe)
- ▶ Let  $\mathcal{B}$  be a Herbrand structure over  $\tau_H$ . ( $U^{\mathcal{B}}$  is the Herbrand universe)
- ▶ We need to create a Herbrand model for  $\Gamma$
- ▶ Try “merging”  $\mathcal{A}$  and  $\mathcal{B}$  to obtain a Herbrand model  $M$  for  $\Gamma$ .
  - ▶ Define  $M$  so that its universe is the Herbrand universe over  $\tau_H$ .
  - ▶ Let  $M$  interpret functions and constants like  $\mathcal{B}$  (both have the same Herbrand universe)
  - ▶ How to interpret the relations from  $\tau_H$  (same as those of  $\tau$ )?
  - ▶ Let  $M$  interpret relations like  $\mathcal{A}$  (not obvious, their universes are not the same.)

# Building the Herbrand Model $M$

---

- ▶ Let  $R$  be an  $n$ -ary relation in  $\tau_H$  (hence in  $\tau$ ).
- ▶ For each  $n$ -tuple  $(t_1, \dots, t_n)$  with  $t_i$  coming from the Herbrand universe  $H(\Gamma)$ , we must say whether  $(t_1, \dots, t_n) \in R^M$  or not
- ▶ Each  $t_i \in H(\Gamma)$  is a ground term in  $\tau_H$  (so variable free).
- ▶ Since  $\mathcal{A}$  is a structure over  $\tau_H$ , if  $t \in H(\Gamma)$  is a ground term from  $\tau_H$ ,  $\mathcal{A}$  interprets  $t$  as an element of  $U^{\mathcal{A}}$ .
- ▶ For each  $n$ -tuple  $(t_1, \dots, t_n)$ ,  $t_i \in H(\Gamma)$ , we know whether  $(t_1, \dots, t_n) \in R^{\mathcal{A}}$  or not
- ▶ Define  $R^M = R^{\mathcal{A}}$ .
- ▶ Prove that if  $\mathcal{A} \models \varphi$  for any  $\varphi \in \Gamma$ , then  $M \models \varphi$ .
- ▶ The proof is by induction on the number of quantifiers in  $\varphi$ . Recall that each  $\varphi$  is in Skolem Normal Form.



# Base case : $\varphi$ has 0 quantifiers

---

$\mathcal{A} \models \varphi$  iff  $M \models \varphi$ . Do structural induction on  $\varphi$ .

- ▶ Assume  $\varphi$  is an atomic formula. Then  $\varphi$  is  $R(t_1, \dots, t_n)$  where  $R$  is an  $n$ -ary relation from  $\tau_H$ , and  $t_1, \dots, t_n$  are all terms from  $H(\Gamma)$ .
- ▶ By the construction of  $M$ ,  $R^M = R^{\mathcal{A}}$ .
- ▶ Hence  $M \models \varphi$  iff  $\mathcal{A} \models \varphi$ .
- ▶ Same reasoning holds for  $\varphi_1 \wedge \varphi_2$ ,  $\varphi_1 \vee \varphi_2$  and  $\neg \varphi$ .
- ▶ Hence,  $\mathcal{A} \models \varphi$  iff  $M \models \varphi$ .

# Post Inductive Hypothesis

---

Assume that for any  $\psi \in \Gamma$  with  $\leq k - 1$  quantifiers, if  $\mathcal{A} \models \psi$ , then  $M \models \psi$ . Let  $\varphi \in \Gamma$  have  $k$  quantifiers,  $\varphi = \forall x_1 \forall x_2 \dots \forall x_k \zeta(x_1, \dots, x_k)$  where  $\zeta$  is quantifier free.

# Post Inductive Hypothesis

---

Assume that for any  $\psi \in \Gamma$  with  $\leq k - 1$  quantifiers, if  $\mathcal{A} \models \psi$ , then  $M \models \psi$ . Let  $\varphi \in \Gamma$  have  $k$  quantifiers,  $\varphi = \forall x_1 \forall x_2 \dots \forall x_k \zeta(x_1, \dots, x_k)$  where  $\zeta$  is quantifier free.

- ▶ Let  $\kappa(x_1) = \forall x_2 \dots \forall x_k \zeta(x_1, \dots, x_k)$ , and  $\varphi = \forall x_1 \kappa(x_1)$ .

# Post Inductive Hypothesis

---

Assume that for any  $\psi \in \Gamma$  with  $\leq k - 1$  quantifiers, if  $\mathcal{A} \models \psi$ , then  $M \models \psi$ . Let  $\varphi \in \Gamma$  have  $k$  quantifiers,  $\varphi = \forall x_1 \forall x_2 \dots \forall x_k \zeta(x_1, \dots, x_k)$  where  $\zeta$  is quantifier free.

- ▶ Let  $\kappa(x_1) = \forall x_2 \dots \forall x_k \zeta(x_1, \dots, x_k)$ , and  $\varphi = \forall x_1 \kappa(x_1)$ .
- ▶  $\mathcal{A} \models \varphi$  implies  $\mathcal{A} \models \forall x_1 \kappa(x_1)$ . That is,  $\mathcal{A} \models \kappa(a)$  for any  $a \in U^{\mathcal{A}}$ .

# Post Inductive Hypothesis

---

Assume that for any  $\psi \in \Gamma$  with  $\leq k - 1$  quantifiers, if  $\mathcal{A} \models \psi$ , then  $M \models \psi$ . Let  $\varphi \in \Gamma$  have  $k$  quantifiers,  $\varphi = \forall x_1 \forall x_2 \dots \forall x_k \zeta(x_1, \dots, x_k)$  where  $\zeta$  is quantifier free.

- ▶ Let  $\kappa(x_1) = \forall x_2 \dots \forall x_k \zeta(x_1, \dots, x_k)$ , and  $\varphi = \forall x_1 \kappa(x_1)$ .
- ▶  $\mathcal{A} \models \varphi$  implies  $\mathcal{A} \models \forall x_1 \kappa(x_1)$ . That is,  $\mathcal{A} \models \kappa(a)$  for any  $a \in U^{\mathcal{A}}$ .
- ▶ Since  $\mathcal{A}$  is a structure over  $\tau_H$ , if  $t \in H(\Gamma)$  is a ground term from  $\tau_H$ ,  $\mathcal{A}$  interprets  $t$  as an element of  $U^{\mathcal{A}}$ .
- ▶ Thus,  $\mathcal{A} \models \kappa(t)$  for any  $t \in H(\Gamma)$ .

# Post Inductive Hypothesis

---

Assume that for any  $\psi \in \Gamma$  with  $\leq k - 1$  quantifiers, if  $\mathcal{A} \models \psi$ , then  $M \models \psi$ . Let  $\varphi \in \Gamma$  have  $k$  quantifiers,  $\varphi = \forall x_1 \forall x_2 \dots \forall x_k \zeta(x_1, \dots, x_k)$  where  $\zeta$  is quantifier free.

- ▶ Let  $\kappa(x_1) = \forall x_2 \dots \forall x_k \zeta(x_1, \dots, x_k)$ , and  $\varphi = \forall x_1 \kappa(x_1)$ .
- ▶  $\mathcal{A} \models \varphi$  implies  $\mathcal{A} \models \forall x_1 \kappa(x_1)$ . That is,  $\mathcal{A} \models \kappa(a)$  for any  $a \in U^{\mathcal{A}}$ .
- ▶ Since  $\mathcal{A}$  is a structure over  $\tau_H$ , if  $t \in H(\Gamma)$  is a ground term from  $\tau_H$ ,  $\mathcal{A}$  interprets  $t$  as an element of  $U^{\mathcal{A}}$ .
- ▶ Thus,  $\mathcal{A} \models \kappa(t)$  for any  $t \in H(\Gamma)$ .
- ▶ By induction hypothesis,  $M \models \kappa(t)$  for any  $t \in H(\Gamma)$ .
- ▶ Since  $H(\Gamma)$  is the universe of  $M$ ,  $M \models \forall x_1 \kappa(x_1)$ . That is,  $M \models \varphi$ .

# Equality

---

$$\varphi = \forall x[(f(x) \neq x) \wedge (f(f(x)) = x)].$$

# Equality

---

$$\varphi = \forall x[(f(x) \neq x) \wedge (f(f(x)) = x)].$$

- ▶ Herbrand universe over  $\tau_H = \{c, f\}$  is  $\{c, f(c), f(f(c)), \dots\}$  all distinct
- ▶ Then  $f(f(c)) \neq c$ , and a Herbrand structure cannot satisfy  $\varphi$



# Equality

---

$$\varphi = \forall x[(f(x) \neq x) \wedge (f(f(x)) = x)].$$

- ▶ Herbrand universe over  $\tau_H = \{c, f\}$  is  $\{c, f(c), f(f(c)), \dots\}$  all distinct
- ▶ Then  $f(f(c)) \neq c$ , and a Herbrand structure cannot satisfy  $\varphi$
- ▶ However,  $\varphi$  is satisfiable. Define a structure  $\mathcal{A} = (\{0, 1\}, f^{\mathcal{A}}(0) = 1, f^{\mathcal{A}}(1) = 0)$ ,  $\mathcal{A} \models \varphi$
- ▶ For formulae which have equality, Herbrand's Theorem does not apply directly
- ▶ If  $\varphi$  has equality, convert it to an equisatisfiable sentence without equality and apply Herbrand

# Equality

---

$$\varphi = \forall x[(f(x) \neq x) \wedge (f(f(x)) = x)].$$

- ▶ Herbrand universe over  $\tau_H = \{c, f\}$  is  $\{c, f(c), f(f(c)), \dots\}$  all distinct
- ▶ Then  $f(f(c)) \neq c$ , and a Herbrand structure cannot satisfy  $\varphi$
- ▶ However,  $\varphi$  is satisfiable. Define a structure  $\mathcal{A} = (\{0, 1\}, f^{\mathcal{A}}(0) = 1, f^{\mathcal{A}}(1) = 0), \mathcal{A} \models \varphi$
- ▶ For formulae which have equality, Herbrand's Theorem does not apply directly
- ▶ If  $\varphi$  has equality, convert it to an equisatisfiable sentence without equality and apply Herbrand

Let  $\varphi$  be in Skolem normal form with equality. Then  $\varphi$  is satisfiable iff there is an equisatisfiable formula  $\psi$  in Skolem normal form without equality which has a Herbrand model.