

# Exponential Distribution

Abhijith Asok, Chris Hilger, Liam Loscalzo, Katherine Wang

# Point Estimation: Methods

We compared the following point estimators:

- Maximum Likelihood Estimator
- Unbiased correction for the MLE
- Second Method of Moment Estimator

##add criteria for comparing estimators!

# Maximum Likelihood Estimator

```
##code here
```

# Unbiased correction for the MLE

```
##code here
```

## Second Method of Moment Estimator

```
##code here
```

# Confidence Intervals: Methods

We compared the following confidence intervals:

- Wald-based Confidence Interval
- Gamma-based Confidence Interval
- Score-based Confidence Interval
- Bootstrap Confidence Interval

##add criteria for comparing CIs!!

# Wald Confidence Interval

```
wald_ci <- function(N, rate, alpha = 0.05){  
  x <- rexp(N, rate = rate)  
  x_bar <- mean(x)  
  se <- sd(x)/sqrt(N)  
  ci <- x_bar + c(-1, 1)*qnorm(1 - (alpha / 2))  
  return(ci)  
}
```

## Gamma Confidence Interval

If  $X_1, X_2, \dots, X_n \stackrel{\text{iid}}{\sim} \text{Exponential}(\lambda)$ , then

$\sum_{i=1}^n x_i \sim \text{Gamma}(n, \lambda) \implies \lambda \bar{x} \sim \text{Gamma}(n, n)$ . Let  $g_y$  be the  $y$ th percentile of this distribution. Then we can say:

$$1 - \alpha = P(g_{\alpha/2} \leq \lambda \bar{x} \leq g_{1-\alpha/2}) = P(g_{\alpha/2}/\bar{x} \leq \lambda \leq g_{1-\alpha/2}/\bar{x})$$

And therefore, a  $(1 - \alpha)\%$  confidence interval for  $\lambda$  is

$$(g_{\alpha/2}/\bar{x}, g_{1-\alpha/2}/\bar{x})$$

```
gamma_ci <- function(N, rate, alpha = 0.05){  
  # Inspiration: https://math.stackexchange.com/questions/1288  
  x <- rexp(N, rate = rate)  
  x_bar <- mean(x)  
  ci_rate <- qgamma(c(alpha / 2, 1 - (alpha / 2)), N, N) / x_bar  
  ci_mean <- 1 / ci_rate  
  return(c(ci_mean[2], ci_mean[1]))  
}
```



# Score Confidence Interval

```
score_ci <- function(N, rate, alpha = 0.05) {  
  x <- rexp(N, rate = rate)  
  xbar <- mean(x)  
  ci_lambda <- (1/xbar)*(1 + c(-1, 1) * qnorm(1 - alpha / 2) / xbar)  
  ci_mean <- 1 / ci_lambda  
  return(c(ci_mean[2], ci_mean[1]))  
}
```


## Bootstrap Confidence Interval

```
bootstrap_ci <- function(N, rate, alpha = 0.05){  
  # Function to calculate bootstrap CI  
  x <- rexp(N, rate = rate)  
  x_bar <- mean(x)  
  # Number of bootstrap samples  
  nb <- 1000  
  # Take bootstrap samples  
  bootstrap_samples <- sample(x, N * nb, replace = TRUE) %>%  
    matrix(nrow = N, ncol = nb)  
  # Get means of columns  
  means <- colMeans(bootstrap_samples)  
  # Get deltas ( $x^* - x$ )  
  deltas <- means - x_bar  
  deltas <- sort(deltas)  
  # Calculate CIs  
  ci <- x_bar - quantile(deltas, probs = c(alpha / 2, 1 - (alpha / 2)))  
  return(c(ci[2], ci[1]))  
}
```

## Other

```
## Parsed with column specification:
## cols(
##   n = col_integer(),
##   bs = col_double(),
##   gamma = col_double(),
##   wald = col_double(),
##   score = col_double()
## )

## Warning: Ignoring unknown parameters: se
```



# Summary of Findings and Recommendations

```
summary(cars)
```

##	speed	dist
##	Min. : 4.0	Min. : 2.00
##	1st Qu.:12.0	1st Qu.: 26.00
##	Median :15.0	Median : 36.00
##	Mean :15.4	Mean : 42.98
##	3rd Qu.:19.0	3rd Qu.: 56.00
##	Max. :25.0	Max. :120.00