

UNIT - 2.

Relations

Subset of Cartesian product of the n sets.

$$R_1 = \{(a, 1), (1, 2)\}$$


$$R_2 = \{(b, 1) (a, 1)\}$$

Cartesian product

$$A = \{a, b, c\}$$

$$B = \{1, 2\}$$

$$A \times B = \{(a, 1) (a, 2) (b, 1) (b, 2) (c, 1) (c, 2)\}$$



* Representing Reln

Matrix

Digraph.

$$\text{Eg: } A = \{1, 2, 3\}$$

$$B = \{1, 2\}$$

$$R = \{(2, 1) (3, 1) (3, 2)\}$$

$$\begin{matrix} & 1 & 2 \\ 1 & \begin{bmatrix} 0 & 0 \end{bmatrix} \\ 2 & \begin{bmatrix} 1 & 0 \end{bmatrix} \\ 3 & \begin{bmatrix} 1 & 1 \end{bmatrix} \end{matrix}$$

If ordered pair = 1.

Not = 0.

Ques $A = \{a_1, a_2, a_3\}$

$$B = \{b_1, b_2, b_3, b_4, b_5\}$$

$$R = \{(a_1, b_2) (a_2, b_2) (a_2, b_3)\}$$

$$\begin{matrix} & b_1 & b_2 & b_3 & b_4 & b_5 \\ a_1 & \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \end{bmatrix} \\ a_2 & \begin{bmatrix} 1 & 0 & 1 & 1 & 0 \end{bmatrix} \\ a_3 & \begin{bmatrix} 1 & 0 & 1 & 0 & 1 \end{bmatrix} \end{matrix}$$

... }



* $R = \{(1,1)(1,3)(2,1)(2,3)(2,4)(3,1)(3,2)\}$

$$(4,1) \quad (4,3)\}$$

* Representing Reln:-

Matrix Diagram

$A \times B$


a b

$$\begin{bmatrix} 1 & 1 & 0 \\ 2 & 1 & 1 \\ 3 & 0 & 1 \end{bmatrix}$$

$$\begin{array}{ccc} 1 & \nearrow a & \\ & \nearrow b & \\ 2 & & \\ & \searrow a & \\ & \searrow b & \\ 3 & & \end{array}$$

$R_1 = \{(1,a)(2,b)$

$(2a)(3,b)\}$



* Types of Relations

→ Reflexive :-

A reln R on set A is called reflexive if $(a,a) \in R$ for every element $a \in A$

e.g.: $A = \{1, 2, 3\}$

$R_1 = \{(1,1)(2,2)(3,3)\}$

Ques
 $A = \{a, b, c\}$

$R = \{(1,3), (1,4), (2,2), (2,1), (2,3), (3,1), (3,1), (4,1), (4,3)\}$

i) $R_1 = A \times A$

ii) $R_2 = \emptyset$

iii) $R_3 = \emptyset$

iv) $R_4 = \{(a,a)(b,b)(c,c)(a,b)(b,c)\}$

v) $R_5 = \{(a,b)(b,a)(a,a)(b,b)\}$

vi) $R_6 = \{(a,b)(b,a)(a,a)(b,b)\}$

vii) $R_7 = \{(a,b)(b,a)(a,a)(b,b)\}$

viii) $R_8 = \{(a,b)(b,a)(a,a)(b,b)\}$

ix) $R_9 = \{(a,b)(b,a)(a,a)(b,b)\}$

x) $R_{10} = \{(a,b)(b,a)(a,a)(b,b)\}$

xi) $R_{11} = \{(a,b)(b,a)(a,a)(b,b)\}$

xii) $R_{12} = \{(a,b)(b,a)(a,a)(b,b)\}$

xiii) $R_{13} = \{(a,b)(b,a)(a,a)(b,b)\}$

xiv) $R_{14} = \{(a,b)(b,a)(a,a)(b,b)\}$

xv) $R_{15} = \{(a,b)(b,a)(a,a)(b,b)\}$

xvi) $R_{16} = \{(a,b)(b,a)(a,a)(b,b)\}$

xvii) $R_{17} = \{(a,b)(b,a)(a,a)(b,b)\}$

xviii) $R_{18} = \{(a,b)(b,a)(a,a)(b,b)\}$

xix) $R_{19} = \{(a,b)(b,a)(a,a)(b,b)\}$

xx) $R_{20} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxi) $R_{21} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxii) $R_{22} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxiii) $R_{23} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxiv) $R_{24} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxv) $R_{25} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxvi) $R_{26} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxvii) $R_{27} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxviii) $R_{28} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxix) $R_{29} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxx) $R_{30} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxxi) $R_{31} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxxii) $R_{32} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxxiii) $R_{33} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxxiv) $R_{34} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxxv) $R_{35} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxxvi) $R_{36} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxxvii) $R_{37} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxxviii) $R_{38} = \{(a,b)(b,a)(a,a)(b,b)\}$

xxxix) $R_{39} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl) $R_{40} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl1) $R_{41} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl2) $R_{42} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl3) $R_{43} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl4) $R_{44} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl5) $R_{45} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl6) $R_{46} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl7) $R_{47} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl8) $R_{48} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl9) $R_{49} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl10) $R_{50} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl11) $R_{51} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl12) $R_{52} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl13) $R_{53} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl14) $R_{54} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl15) $R_{55} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl16) $R_{56} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl17) $R_{57} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl18) $R_{58} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl19) $R_{59} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl20) $R_{60} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl21) $R_{61} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl22) $R_{62} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl23) $R_{63} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl24) $R_{64} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl25) $R_{65} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl26) $R_{66} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl27) $R_{67} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl28) $R_{68} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl29) $R_{69} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl30) $R_{70} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl31) $R_{71} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl32) $R_{72} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl33) $R_{73} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl34) $R_{74} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl35) $R_{75} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl36) $R_{76} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl37) $R_{77} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl38) $R_{78} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl39) $R_{79} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl40) $R_{80} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl41) $R_{81} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl42) $R_{82} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl43) $R_{83} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl44) $R_{84} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl45) $R_{85} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl46) $R_{86} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl47) $R_{87} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl48) $R_{88} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl49) $R_{89} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl50) $R_{90} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl51) $R_{91} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl52) $R_{92} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl53) $R_{93} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl54) $R_{94} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl55) $R_{95} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl56) $R_{96} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl57) $R_{97} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl58) $R_{98} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl59) $R_{99} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl60) $R_{100} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl61) $R_{101} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl62) $R_{102} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl63) $R_{103} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl64) $R_{104} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl65) $R_{105} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl66) $R_{106} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl67) $R_{107} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl68) $R_{108} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl69) $R_{109} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl70) $R_{110} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl71) $R_{111} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl72) $R_{112} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl73) $R_{113} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl74) $R_{114} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl75) $R_{115} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl76) $R_{116} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl77) $R_{117} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl78) $R_{118} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl79) $R_{119} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl80) $R_{120} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl81) $R_{121} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl82) $R_{122} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl83) $R_{123} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl84) $R_{124} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl85) $R_{125} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl86) $R_{126} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl87) $R_{127} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl88) $R_{128} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl89) $R_{129} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl90) $R_{130} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl91) $R_{131} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl92) $R_{132} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl93) $R_{133} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl94) $R_{134} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl95) $R_{135} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl96) $R_{136} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl97) $R_{137} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl98) $R_{138} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl99) $R_{139} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl100) $R_{140} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl101) $R_{141} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl102) $R_{142} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl103) $R_{143} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl104) $R_{144} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl105) $R_{145} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl106) $R_{146} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl107) $R_{147} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl108) $R_{148} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl109) $R_{149} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl110) $R_{150} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl111) $R_{151} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl112) $R_{152} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl113) $R_{153} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl114) $R_{154} = \{(a,b)(b,a)(a,a)(b,b)\}$

xl115) $R_{155} = \{(a,b)(b,a)(a,a)(b,b)\}$

Ques

$$A = \{a, b, c\}$$

- i) $R_1 = \emptyset$ ← Everything present
- ii) $R_2 = A \times A = \{(a, a), (b, b), (c, c)\}$
- iii) $R_3 = \{(a, a), (b, b), (c, c), (a, b), (b, a), (a, c), (c, a)\}$
- iv) $R_4 = \{(a, b), (b, a), (a, c), (c, a)\}$
- v) $R_5 = \{(a, b), (b, c), (a, c)\}$

→ Symmetric:

A reln R on set is said to be symmetric if $\forall a, b \in A$

If $(a, b) \in R$ then $(b, a) \in R$

→ Asymmetric:

A reln R on set A is asymmetric if $\forall a, b \in A$

If $(a, b) \in R$ then $(b, a) \notin R$.

→ Transitive

A reln R on set A is said to be transitive if $\forall a, b, c \in A$, $(a, b) \in R$ & $(b, c) \in R$ then $(a, c) \in R$.
If \exists a and b then must \exists
it only (a, b) is there but (b, a) is not there
then say it is trans.

- Ques
- i) $\{(a, b), (b, a)\}$
 - ii) $\{(b, c), (c, b), (b, b)\}$
 - iii) $\{(a, a), (b, b), (c, c)\}$
 - iv) \emptyset ← Nothing to compare
 - v) $\{(a, b), (b, c), (a, c)\}$
 - vi) $\{(a, b), (b, a), (a, c)\}$

Ques

$$R_1 = \{(a, b), (b, a), (a, a), (b, b)\}$$

→ Antisymmetric:

A reln R on set A is said to be antisymmetric if $\forall a, b \in A$. If $(a, b) \in R$ and $(b, a) \in R$ then $a = b$

$\rightarrow (b, a) \notin R$ if $a \neq b$.

* Sym. pairs not allowed diagonal pairs allowed.

\downarrow

$$(a, a) \quad b \quad c$$

\downarrow

$$\begin{array}{c} i) \{(a, a), (b, b), (c, c)\} \\ ii) R_3 = \{(a, b)\} \quad \leftarrow \text{No reln starting with } b \end{array}$$

Ques
 $A = \{(1, 2, 3, 4, 5)\}$ Ans

- i) $\{(2, 2), (2, 3), (2, 4), (3, 2), (3, 3), (3, 4)\}$ Not Anti-symmetric
 Not Reflexive Not Asymmetric
 Not Symmetric Not Transitive.
 Not Invertive

ii) $\{(1, 1), (1, 2), (2, 1), (2, 2), (3, 3), (4, 4)\}$ Reflexive
Symmetric
Transitive.

iii) $\{(2, 4), (4, 2)\}$ Reflexive
Symmetric
Transitive.

iv) $\{(2, 1), (2, 2), (3, 3), (4, 4)\}$ Not transitive (2, 2) not precur


v) $\{(1, 1), (1, 2), (2, 1), (2, 2), (3, 3), (4, 4)\}$ Symmetric, Inv.

vi) $\{(1, 1), (1, 2), (2, 1), (2, 2), (3, 1), (3, 2), (3, 3), (4, 1), (4, 2)\}$ Invertive, Inv., Any

vii) $\{(1, 1), (2, 2), (3, 3), (4, 4)\}$ Reflexive, Symmetric, Anti-symmetric, Transitive.

viii) $\{(1, 1), (1, 2), (2, 1), (2, 2), (3, 1), (3, 2), (3, 3), (4, 1), (4, 2)\}$ Reflexive, Symmetric, Anti-symmetric, Transitive.


ix) $\{(1, 1), (1, 2), (2, 1), (2, 2), (3, 1), (3, 2), (3, 3), (4, 1), (4, 2)\}$ Invertive.



Ques

Ques Draw:
 $R = \{(1, 1), (1, 3), (2, 1), (2, 3), (3, 1), (3, 2), (4, 1)\}$

Set {1, 2, 3, 4}



Ques

$R = \{(a, a), (a, b), (b, b), (b, c), (c, a), (c, b)\}$ Symmetric X

Reflexive

$$\begin{matrix} & a & b & c \\ a & 1 & 1 & 0 \\ b & 0 & 1 & 1 \\ c & 1 & 1 & 1 \end{matrix} \quad \text{Matrix representation.}$$

* Partial Order Reln::

A is a PO Reln on set A if it is Reflexive, Anti-symmetric and Transitive

Ques $A = \{1, 2, 3\}$

i) $R_1 = \emptyset$ Not reflexive

ii)

$R_2 = \{(1, 1), (2, 2), (3, 3)\}$ Reflexive, symmetric, anti-symmetric

iii)

$R_3 = \{(1, 1), (2, 2), (3, 3), (1, 3), (2, 3)\}$ Reflexive, symmetric, anti-symmetric, transitive

iv)

$R_4 = A \times A$ Reflexive, symmetric, anti-symmetric, transitive

x) $R = \{(1, 1), (1, 3), (2, 1), (2, 2), (3, 1), (3, 3), (4, 1)\}$

xi)

* Equivalence Relation: to be equivalent

A new form A is said to be
achiral, asymmetric and chiral

$$\text{Key leave } \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

| | $\mathcal{E} \left[\begin{smallmatrix} 1 & 0 & 1 \\ 1 & 1 & 1 \end{smallmatrix} \right]$ | $\mathcal{E} \left[\begin{smallmatrix} 1 & 0 & 1 \\ 1 & 1 & 1 \end{smallmatrix} \right]$ | Not Ref |
|-----------------------|---|---|--------------------|
| $R = f(a, a) f(a, c)$ | $R = f(f(a, b))$ | $f(a, b)$ | POD or λ . |

~~D~~ $\frac{d}{dx} \left(b_1 b_2 \right) = b_1' b_2 + b_1 b_2'$

TV
KOMV

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix} \quad \begin{bmatrix} 0 & 1 \end{bmatrix}$$

| | | |
|---------|----------|---------|
| Not ref | Not sym. | Not ref |
| POR X | POR X | POR X |

Eq x Eq x Eq

$$S_{\text{new}} = \{1, 2, 3, 4\}$$

$$R_2 = \{ (1,2), (1,3), (1,4), (2,3), (2,4), (3,4) \}$$

$$R_3 = \{ (1,1) \}$$

卷之三

100

EQUIVALENCE CLASS

To bind the equ. class and parallel

The equ. class of x is denoted by $[x]$ or $[x]_R$
 where $[x] = \{y \mid xRy\}$ and R is a relation.

$$A = \{1, 2, 3, 4, 5\}$$

$$R = \{ (1,1), (2,2), (3,3), (4,4), (5,5), (1,2), (2,1), (4,5), (5,4) \}$$

11 = {1, 2} One is with one and two

$$|3\rangle = \{3\} \quad \left. \begin{array}{c} \\ \\ \end{array} \right\} \quad P_1 \quad P_2 \quad P_3$$

151 = {4, 53}]

$$\{ P_1 \cup P_2 \cup P_3 = A \\ P_1 \cap P_2 \cap P_3 = \emptyset \}$$

—

* Closure: Rehearsal 11:00 AM - 1:00 PM Kell.

all pairs (a,a)

$R = \{0, 1, 2, 3\}$

→ Address

$$R = \{(0,0)(3,3)(5,2)(1,1)(1,2) \\ (2,0)(2,2)(3,1)\}$$

Cornell Page

→ Symmetric Closure
The SC of R is R together with all pairs (b, a)

such (ab) is in R .

$$\text{Ex: } A = \{(0, 1, 2, 3)\} \\ R = \{(0, 1), (1, 2), (1, 3), (2, 0), (2, 2), (3, 0)\}$$

$$R^S = \{(0, 1)(1, 0), (1, 1)(2, 2), (2, 1)(2, 0), (0, 2), \\ (2, 2)(3, 0), (0, 3)\}$$


→ added.

• Transitive closure:
→ Warshall's Algorithm

$$\text{Ex: } A = \{1, 2, 3, 4\} \\ R = \{(1, 2), (2, 1), (2, 3), (3, 4)\}$$

$$\text{Ques } A = \{a, b, c, d\}$$

$$R = \{(a, d), (b, a), (b, c), (c, a)\}$$



Column = 2] where we
Row = 2] are getting 1 first.

$$C \times R = (2, 2)$$

$$W_2 = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{Column} = 1, 2 \\ \text{Row} = 4$$

$$\text{Rows} = 1, 2, 3$$

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} = (1, 1)(1, 2)(1, 3) \\ (2, 1)(2, 2)(2, 3)$$

$$W_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad \text{Column} = \emptyset \\ \text{Rows} = a, c, d$$

$$W_3 = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{Column} = 1, 2, 3 \\ \text{Row} = \emptyset$$

$$W_4 = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{Column} = 1, 2, 3 \\ \text{Row} = \emptyset$$

4 changes
in A:

$$R^S = \{(1, 1)(1, 2)(1, 3)(1, 4), (2, 1)(2, 2)(2, 3) \\ (2, 4)(3, 4)\}$$

→ This is Transitive.

$$M_R = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} = W_0 \quad \text{With 0's only}$$

Ques

$$A = \{a, b, c, d\}$$

$$R = \{(a, d), (b, a), (b, c), (c, a)\}$$

$$M_R = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{bmatrix} = W_0$$

$$\text{Column} = 1, 2 \\ \text{Row} = 1, 2$$

$$C \times R = (4, 4)$$

$$W_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad \text{Column} = \emptyset \\ \text{Row} = a, b, c, d$$

$$W_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad \text{Column} = \emptyset \\ \text{Row} = a, b, c, d$$

Ques
Find τ_C using Trans M^m .

Ex:

$$M_R = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \end{bmatrix}$$

* Hasse Diagram | POSET Diagram
→ If a reln is partial order then there only we can draw the HD of it.

$$A = \{1, 2, 3, 4\}$$

$$R = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,3), (2,4), (3,3), (3,4), (4,4)\}.$$

$$M_R^{(2)} = M_R \circ M_R$$

$$= \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \end{bmatrix}$$

$$\epsilon = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix}$$

$$M_R^{(3)} = M_R^{(2)} \circ M_R$$

$$= \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix}$$

$$M_R^* = M_R^{(2)} \vee M^{(3)}$$

$$= \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix} \vee \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix} : \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix}$$

$$In \downarrow ing \text{ order. } \begin{array}{c} 1 \\ 2 \\ 3 \\ 4 \end{array}$$

$$3 \text{ is R to } 3$$

$$4 \text{ is R to } 4$$

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

Ques

$$A = \{1, 2, 3, 6\}$$


$$R = \{(1, 1) (1, 2) (1, 3) (1, 8) (2, 2) (2, 6)$$

$$(3, 3) (3, 6) (6, 6)\}$$

Ans

$$\{1, 2, 3, 4, 6, 12\}$$



$$\{1, 2, 3, 4, 6, 9\}$$

Ans
Remove Reflexive Pairs

1 - 2 - 6 Remove from this


not from this

1 - 3 - 6 i.e. not from refi pair

Ques $\{1, 2, 3, 5, 6, 10, 15, 30\}$ 

We can draw

this directly

Ques $\{1, 2, 3, 6, 9, 18\}$ Ans (A) $\{1, 2, 3, 4, 5\}, \leq$
 $1 \leq 2 \quad 2 \leq 2 \quad / / / \text{ by for } 3, 4, 5.$

 $1 \leq 2 \quad 2 \leq 2 \quad / / / \text{ by for } 3, 4, 5.$

→ Marital separation - from
partner

TOP element

* Maximal Element: - in not related to -

is in poset
other element.

* Minimal Element:
No element is related to it.

Eg:

Op only, in quest

UB: {a, b, c}

LB: {d, e, f}

L →

U →

$$\Rightarrow B = \{c, f, d\}$$

$$E_{\alpha_1} = \{5, 10, 2, 4\}$$

no longer in use.

* Upper Bound and Lower Bound.

* UB

[A] is a POSET if $a \in A$ is called upper bound of subset B ($B \subseteq A$) if $x \geq a$, $\forall x \in B$

Representation of POSEI

UNIT - 3

Ques: $\{D_o, D_i\}$ convert to Hanoi then conclude.




domain

codomain

5. It is a special type of reln
6. \rightarrow A funcⁿ F which maps from A to B ($F: A \rightarrow C_B$)
7. when every element of domain must have a mapping
8. \rightarrow domain is also called as pre-image
9. \rightarrow For every pre-image we should get unique pre-image (In reln we can avoid some elements of A but in funcⁿ we should use all elements of set A)
10. \rightarrow Range is subset of co-domain
11. Range is nothing but elements of funcⁿ
12. \rightarrow also called as image.

Domain — A $B \rightarrow$ codomain



A

2

25

X

B

p

q

r

2

3

X

- Ques $f(x) = x^2$ $R \rightarrow R$ ✓ $f(x) = \sqrt{x}$ $R \rightarrow R$ \times {For -ve elements} no mapping.

30

3

* Types of functions :

→ One to One (Injective)

→ Onto (Surjective)

→ One to One and Onto (Bijective).

* One to One :

A function f is said to be one to one if
 $f(a) = f(b) \Rightarrow a = b$ $\forall a, b$ in domain of f .


Distinct element → Distinct element

of A of B.



e.g.: $\{a, b, c, d\} \rightarrow \{1, 2, 3, 4, 5\}$

$$\begin{array}{l} a \rightarrow 1 \\ b \rightarrow 2 \\ c \rightarrow 3 \\ d \rightarrow 4 \end{array}$$

If it is one to one and onto.



* Bijective:



Nothing.

Not a function.

Ques (i) $f(x) = x^2$ x for \perp and \rightarrow (Point → Int)

$$z(i) f(x) = x+1 \quad R \rightarrow R \quad \checkmark$$

* Onto: $F: X \rightarrow Y$ \rightarrow Capitalized.

Every image in Y must have a pre-image in X

Eg: $\begin{cases} x \rightarrow 1 \\ y \rightarrow 2 \\ z \rightarrow 3 \end{cases} \rightarrow$ Every element has
 one pre-image.

* Composition of Function

$$f: A \rightarrow B$$

$$g: B \rightarrow C$$

Composition of f and g
or $g \circ f$ by $f \circ g$

Ex:

$$\begin{aligned}f(x) &= 2x + 3 \\g(x) &= 3x + 2\end{aligned}$$

$$\begin{aligned}(f \circ g)x &= f(g(x)) \\&= f(3x+2) = 2(3x+2) + 3 \\&= 6x + 7.\end{aligned}$$

$$\begin{aligned}(g \circ f)x &= g(2x+3) \\&= 3(2x+3) + 3 \\&= 6x + 12.\end{aligned}$$

$$(g \circ f)x \rightarrow g(f(x))$$

*


Inverse of a function: If $\text{set } f^{-1} : B \rightarrow A$ given fun "f": $A \rightarrow B$ if it is called inverse of a function in a function.

$$\text{Ex: } \begin{array}{ccc} A & \xrightarrow{\quad} & B \\ a & \xrightarrow{1} & 2 \\ b & \xrightarrow{2} & 3 \\ c & \xrightarrow{3} & 4 \\ d & \xrightarrow{4} & \end{array}$$

$$\begin{array}{ccc} B & \xrightarrow{\quad} & A \\ 2 & \xrightarrow{1} & a \\ 3 & \xrightarrow{2} & b \\ 4 & \xrightarrow{3} & c \\ \end{array}$$

Not a function

$\therefore B$ is not inverse of A .



*

Discrete Numeric fun ": Let N be the set of natural nos. and R be the set of Real nos. Then fun "f": $N \rightarrow R$ is called DNF.

In practice we write DNF as a sequences like $(f(0), f(1), f(2), \dots)$

The sequences can be written like this,

$$a = a_0, a_1, a_2, \dots, a_k, \dots$$

Ex: Let a be DNF.

$$a_n = 7n^2 + 1, n \geq 0$$

$$\begin{aligned} a_0 &= 7(0)^2 + 1 = 8 \\ a_1 &= 7(1)^2 + 1 = 8 \\ a_2 &= 7(2)^2 = 29 \end{aligned}$$

* DNF can be defined by using diff. formulas

$$a_n = \begin{cases} 2+n & 0 \leq n \leq 5 \\ 2-n & n > 5 \text{ and } n \text{ is odd.} \\ n/2 & n > 5 \text{ and } n \text{ is even} \end{cases}$$

Eg:

$$\begin{aligned} A &= \{a, b, c\} & f(a) &= 2 \\ B &= \{1, 2, 3\} & f(b) &= 3 \\ & & f(c) &= 1 \end{aligned}$$

*

Generating fun ":

* If G_f is a way of encoding an infinite seq. of nos. (a_m) by treating them as a coeff. of power series

Def ": Let a_0, a_1, \dots, a_m be a series of real nos. denoted by $\{a_m\}$ then a series in power of n such

$$\text{that } g(x) = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \dots \in \mathbb{R}$$

$$\text{or } \left\{ g(x) = \sum_{n=0}^{\infty} a_n x^n \right\}$$

It is used to solve the counting problems and recurrence relⁿ.

Ques Find gf for following series
 $1, -1, 2, -2, \dots$

$$g(x) = a_0 + a_1 x + a_2 x^2 + \dots$$

$$a_0 = 1$$

$$a_1 = -1$$

$$a_2 = 2$$

$$g(x) = 1 - x + x^2 - x^3 + \dots$$

$$\Rightarrow g(x) = \sum_{n=0}^{\infty} (-1)^n (x)^n.$$

*

Find Order Linear Recurrence Relⁿ:

$$\begin{aligned} & 2^0, 2^1, 2^3, 2^4, \dots \\ & = 2^0 x + 2^1 x \\ & = 2^0 + 2^1 x + 2^2 x^2 + \dots \\ & = \sum_{n=0}^{\infty} 2^n x^n. \end{aligned}$$

$$\text{Rec Rel}^n : a_n = c_1 a_{n-1} + c_2 a_{n-2} + \dots + c_k a_{n-k}$$

↓ A form at

$$x^n = c_1 x^{n-1} + c_2 x^{n-2} + \dots + c_k x^{n-k}$$

$$x^n = c_1 x^{n-1} + c_2 x^{n-2} + \dots + c_k x^{n-k}$$

* Recurrence Relⁿ

- It is a recursively defined sequence
- A Relⁿ is an eqn that recursively defines seq.
- where next term depends on prev. term.
- Ex: Fibonacci series.

Find first 4 terms.
 i) $a_0 = 2a_{n-1} + n$, $a_1 = 1$

$$a_2 = 2a_{n-1} + 2$$

$$= 2a_1 + 2$$

$$= 4 + 2$$

$$a_3 = 2a_2 + 3$$

$$= 8 + 3$$

$$= 11.$$

$$a_4 = 2a_3 + 4$$

$$= 2(11) + 4$$

$$= 26.$$

$$a_1 = a_0 + 2 = 3$$

$$a_2 = a_1 + 2 = 5$$

$$a_3 = a_2 + 2 = 7.$$

$\Rightarrow a_{n+1} = 2a_n + a_0$ \rightarrow

Convert eqn to L format.

Divide it by power value of L

Find the value of A (here A is constant)

Find values of constant a and A in

Substitute the values of a and A in

$$\{a_n = A(L^n)\}$$

$$at \quad n=0$$

$$a_0 = A(3)^0 + B(2)^0$$

10

$$1 = A + B \quad \text{--- (1)}$$

$$at \quad n=1$$

$$a_1 = A(3)^1 + B(2)^1$$

$$J = 3A + 2B$$

from (1)

$$J = 3(J - B) + 2B$$

$$J = 3 - 3B + 2B$$

$$\Rightarrow B = 2.$$

$$\therefore A = -J.$$

* 2nd Order Linear Homogeneous Rec. Reln.

$$\text{Final eqn} = a_n = -J(3)^n + 2.(2)^n.$$

→ Q we have

$$B_{n+2} + C a_{n+1} + D a_n = 0.$$

Convert to L format

Divide it by lowest L term

Find the roots e.g. $L^2 + L + 1$ \rightarrow λ_1 and λ_2

→ Bew Solve $a_n = A(\lambda_1)^n + B(\lambda_2)^n$ \rightarrow

Find A and B and put in

$$a_n = S a_{n-1} - 6 a_{n-2}, \quad a_0 = 1, \quad a_1 = 1.$$

\Rightarrow

$$① \quad a_n - 5a_{n-1} + 6a_{n-2} = 0$$

$$② \quad \lambda^2 - 5\lambda + 6 = 0$$

$$\lambda_1 = -3 \quad \lambda_2 = -2 \quad \lambda_1 = 3 \quad \lambda_2 = 2$$

$$a_n = A.(3)^n + B(2)^n$$

at $n=0$

10

$$a_0 = A(3)^0 + B(2)^0$$

1

$$1 = A + B \quad \text{--- (1)}$$

at $n=1$

$$a_1 = A(3)^1 + B(2)^1$$

2

$$J = 3A + 2B$$

3

$$J = 3(J - B) + 2B$$

4

$$J = 3 - 3B + 2B$$

5

$$\Rightarrow B = 2.$$

$$\therefore A = -J.$$

* Binary sequence

$$\tau_m = \tau(m-1) + I(m-1)$$

$$\underbrace{\begin{array}{|c|c|c|c|c|} \hline 0 & & & & \\ \hline \end{array}}_m \quad \tau(n) = 2\tau(m-1)$$

$$\underbrace{\begin{array}{|c|c|c|c|c|} \hline 1 & & & & \\ \hline \end{array}}_{m-1} \quad \tau(1) = 2 \text{ ways}$$


$$\underbrace{\begin{array}{|c|c|c|c|c|} \hline 1 & & & & \\ \hline \end{array}}_{(m-1)} \quad (\text{ways})$$

∴ When $\tau(2) = 2\tau(1)$


= 4 ways.

Ques Binary seq without consecutive 0's

Ques 2:



Ques 3:



only R and ↑

Take only 1 or 2
steps.

10

15

20