

Pattern Recognition
Course Code: CSE313/423
by
Dr Sudarshan Nandy

Module -II

Statistical Pattern Recognition:

Bayesian Decision Theory, Classifiers, Normal density and discriminant functions

Basic of probability

- Probability denotes “chance”
- Laws that governs the chance of occurrence of an event is maintained in the theory of the probability
- How much chance of an event -----> is known as probability.
- Random Experiment: when an occurrence of the event can not be predicted.
- Outcome: The result of a random experiment
- Event: Phenomenon which occurs in a random experiment

Basic of probability

- **Mutually Exclusive**: if two or more events are not occurring simultaneously or together then events are mutually exclusive. They occurs one at a time.
Example: events A , B, C, D are mutually exclusive, only if A, or B, or C or D can occur.
- **Exhustive**: it is the a group of events from a random experiment from which at least one event has occured.
- **Equally Likely/ Equaly Probable** :If after taking into consideration all relevant evidance, none of them can be expected in preference to another.

Basic of probability

- Probability:

if a random experiment has n possible outcomes, which are mutually exclusive, exhaustive and equally likely and m of these are favourable to an event A then the probability of A is $P(A)$ and is calculated as follows:

$P(A) = m/n = \text{Number of outcomes favourable to the Event} / \text{total number of mutually exclusive event.}$

Basic of probability

- What is the process of calculation?

Step1 : count the number of outcomes from an experiment where the outcomes must satisfy: a. Mutually Exclusive b. Exhaustive c. Equally likely ----->(n)

Step2 : Count the numbers of Event in favour of the Event for which you are calculating the probability.----->(m)

Step3 : Divide Step2 by Step1 which gives the probability of the event.

- But the calculation is always $0 < P(A) < 1$

Basic of probability

- **Example 1: Two coins are tossed. Find the probability of getting both heads or both tails.**
 1. Events are ---> HH, HT, TH and TT and hence $n=4$
 2. Both head or tail means HH or TT then $m = 2$
 3. So the probability : $P(HH) \text{ or } P(TT) = 2/4 = 0.50$
- **Example 2: Two dice with faces marked 1,2,3,4,5,6 are thrown simultaneously and the points on the dice are multiplied together. Find the probability that the product is 12.**
 1. for two dice if thrown simultaneously then $n = 6*6 = 36$
 2. now the products will be 12 for the following event- (2,6), (3,4), (4,3), (6,2)
Hence , $m = 4$
 3. So, $P = 4/36 = 1/9$

Basic of probability

- **A bag contains 6 white balls and 4 black balls. One ball is drawn. What is the probability that it is white?**
 1. There are 6 favourable event possible, hence $n = 10$
 2. The number of outcomes is, $m = 6$
 3. Hence, $P(W) = 6/10$

Basic of probability

- **Multiplication Rule:**

Probability is to find the chance of an event.

- In an **independent event**, two events does not effect each other.

Ex: rolling a six sided normal die for 5 and 3

Event is independent .

So, $P(5) = 1/6$ and $P(3) = 1/6$

- Now, to understand the probability of the 5 and 3 , we need to go through the multiplication rule in the probability:

So, $P(A \text{ and } B) = P(A \cap B) = P(AB) = P(A) * P(B)$

Hence, $P(5 \text{ and } 3) = 1/6 * 1/6 = 1/36$

- **P(AB)** is also known as a compound probability. This is actually means probability of occurance of A as well as B envent.

Basic of probability

- The **Dependent events** on other side are those where occurrence of an event effects the another event.
- A famous example of this is the drawing the king and then queen from the deck of cards. In this experiment the queen is drawn without putting the king back.
- So, $P(\text{KING}) = \text{no of king} / \text{total number of cards} = 4/52$
 $P(\text{QUEEN}) = \text{no of queen} / (\text{total number of cards} - \text{King}) = 4/51$
- Now the probability of any dependent events is caculated as follows:

$$P(A \cap B) = P(AB) = P(A) * P(B|A)$$

Basic of probability

- So,

$$\begin{aligned} P(\text{King and Queen}) &= P(\text{King}) * P(\text{Queen} | \text{King}) \\ &= 4/52 * 4/51 = 16/2652 \end{aligned}$$

- Hence the Conditional probability where we consider that the probability of occurrence A, assuming that B has already occurred and it is denoted as $P(A | B)$.

$$P(A|B) = \frac{P(AB)}{P(B)}$$

$$P(B|A) = \frac{P(AB)}{P(A)}$$

Basic of probability

- Given $P(A) = 1/2$, $P(B) = 1/3$ and $P(AB) = 1/4$ then state whether the events A and B are (i) mutually exclusive, (ii) exhaustive (III) Equally Likely/ Equally Probable (iv) independent.

Ans:

- i) No, here $P(AB) = 1/4$ and $P(AB) \neq 0$

II) So, here $P(A+B) = P(A) + P(B) - P(AB) = 7/12$ and if $P(A+B) \neq 1$ then the events are not exhaustive.

III) no, it is not equally likely because $P(A)$ and $P(B)$ are different.

IV) Here, $P(AB) \neq P(A) \cdot P(B)$

So, A and B are not independent.

Basic of probability

- IF TWO EVENTS A AND B ARE MUTUALLY EXCLUSIVE THEN THE PROBABILITY OF OCCURANCE:

$$P(A \text{ or } B) = P(A) + P(B)$$

- **Proof:** suppose there are n outcomes from a random experiment. Events occur for this experiments are mutually exclusive. So,

Favourable to Event A is = m_1 ;

Favourable to Event B is = m_2 ;

Hence, $P(A) = m_1/n$

and $P(B) = m_2/n$

So, number of case favourable to either A or B = (m_1+m_2)

- Hence, the probability for either A or B is = $P(A \text{ or } B) = P(A \cup B) = P(A+B)$

$$P(A+B) = (m_1+m_2) / n = (m_1/n) + (m_2/n) = P(A)+P(B)$$

- Hence the probability of the occurrence of any events from series of events which are mutually exclusive, is actually

$$P(A_1+A_2+A_3+.....)=P(A_1)+P(A_2)+.....+P(A_k)$$

Basic of probability

- Given that $P(A) = 3/8$ and $P(B) = 5/3$ and $P(A+B) = 3/4$ find $P(A|B)$ and $P(B|A)$. Are A and B independent?

Bayes theorem

- In probability theory and statistics, Bayes' theorem (alternatively Bayes' law or Bayes' rule), named after Reverend Thomas Bayes
- It basically describe the probability of an Event, and the knowledge on the condition related to the event is known.
- This is mathematically expressed as follows:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

- Where, $P(A|B)$ is a conditional probability. This is the calculation of probability A , considering the B event has already happend. $P(A)$ and $P(B)$ is the probability of A and B event.

Bayes theorem

Proof:

$$P(A|B) = \frac{P(A \cap B)}{P(B)} \quad \text{if } P(B) \neq 0$$

and

$$P(B|A) = \frac{P(B \cap A)}{P(A)} \quad \text{if } P(A) \neq 0$$

Now,

$$P(A \cap B) = P(B \cap A)$$

$$\Rightarrow P(A|B)P(B) = P(B|A)P(A)$$

$$\Rightarrow P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

Naive Bayes classifier

- This is based on Bayes theorem.
- Naive Bayes classifiers assume that the value of a particular feature is independent of the value of any other feature, given the class variable.
- For example, a fruit may be considered to be an apple if it is red, round, and about 10 cm in diameter. A naive Bayes classifier considers each of these features to contribute independently to the probability that this fruit is an apple, regardless of any possible correlations between the color, roundness, and diameter features.
- Naive Bayes classifiers can be trained very efficiently in a supervised learning environment.
- Work well in the complex environment.
- It is work very well even with a small dataset.

Naive Bayes classifier

- According to the Bayes theorem:

$$P(A|B) = \frac{P(B|A) * P(A)}{P(B)}$$

- The features of the patterns is representing a class which is represented as follows:

$$L = (x_1, x_2, x_3, \dots, x_n)$$

- These features are independent from each other and can be treated as a list of independent event. It is also representing the instance and hence probability or the chance of K^{th} class, denoted as L_K , is represented by all the attributes or features in the pattern:

$$P(L_k | x_1, x_2, x_3, \dots)$$

Naive Bayes classifier

- According to the Baye's Theorem, the conditional probability can be decomposed as:

$$P(L_k | X) = \frac{P(L_k) * P(X | L_k)}{P(X)}$$

- Now consider that the all features

$$(x_1, x_2, x_3, \dots, x_n) \in X$$

are mutually exclusive and conditional on the category L_k . Under this assumption,

$$P(L_k | x_1, x_2, x_3, \dots) = P(x_i | L_k)$$

Naive Bayes classifier

- So, now the Baye's theorem can be re-written as:

$$P(L_k | x_1, x_2, x_3, \dots) = \frac{P(L_k) P(x_i | L_k)}{P(x_1) P(x_2) P(x_3) \dots P(x_n)}$$

•
•
•
•
•
•
•
•
•
•
•
•
•
•
•

$$= \frac{P(L_k) \prod_{i=1}^n P(x_i | L_k)}{P(x_1) P(x_2) P(x_3) \dots P(x_n)}$$

$$= \frac{P(L_{k=\text{yes}}) \prod_{i=1}^n P(x_i | L_{k=\text{yes}})}{P(x_1) P(x_2) P(x_3) \dots P(x_n)}$$

- Now as the denominator is constant hence, it is canceled and then it is always

$$P(L_k | x_1, x_2, x_3, \dots) \propto P(L_k) \prod_{i=1}^n P(x_i | L_k)$$

Naive Bayes classifier

- Now, the Naive bayes classifier comes with a decision rule which is based on the maximum probable likelihood and it is represented as follows:

$$D = \operatorname{argmax}_k P(L_k) \prod_{i=1}^n P(x_i | L_k)$$

Naive Bayes classifier

| SI No | A1=Outlook | A2=Temperature | A3=Humidity | A4=Windy | Target=Play Golf |
|-------|------------|----------------|-------------|----------|------------------|
| P0 | Rainy | Hot | High | False | No |
| P1 | Rainy | Hot | High | True | No |
| P2 | Overcast | Hot | High | False | Yes |
| P3 | Sunny | Mild | High | False | Yes |
| P4 | Sunny | Cool | Normal | False | Yes |
| P5 | Sunny | Cool | Normal | True | No |
| P6 | Overcast | Cool | Normal | True | Yes |
| 7 | Rainy | Mild | High | False | No |
| 8 | Rainy | Cool | Normal | False | Yes |
| 9 | Sunny | Mild | Normal | False | Yes |
| 10 | Rainy | Mild | Normal | True | Yes |
| 11 | Overcast | Mild | High | True | Yes |
| 12 | Overcast | Hot | Normal | False | Yes |
| 13 | Sunny | Mild | High | True | No |

Naive Bayes classifier

Outlook

| | Yes | No | P(yes) | P(no) |
|--------------|-----|----|--------|-------|
| Sunny | 2 | 3 | 2/9 | 3/5 |
| Overcast | 4 | 0 | 4/9 | 0/5 |
| Rainy | 3 | 2 | 3/9 | 2/5 |
| Total | 9 | 5 | 100% | 100% |

Temperature

| | Yes | No | P(yes) | P(no) |
|--------------|-----|----|--------|-------|
| Hot | 2 | 2 | 2/9 | 2/5 |
| Mild | 4 | 2 | 4/9 | 2/5 |
| Cool | 3 | 1 | 3/9 | 1/5 |
| Total | 9 | 5 | 100% | 100% |

Humidity

| | Yes | No | P(yes) | P(no) |
|--------------|-----|----|--------|-------|
| High | 3 | 4 | 3/9 | 4/5 |
| Normal | 6 | 1 | 6/9 | 1/5 |
| Total | 9 | 5 | 100% | 100% |

Wind

| | Yes | No | P(yes) | P(no) |
|--------------|-----|----|--------|-------|
| False | 6 | 2 | 6/9 | 2/5 |
| True | 3 | 3 | 3/9 | 3/5 |
| Total | 9 | 5 | 100% | 100% |

| Play | | P(Yes)/P(No) |
|--------------|----|--------------|
| Yes | 9 | 9/14 |
| No | 5 | 5/14 |
| Total | 14 | 100% |

Naive Bayes classifier

- Problem = $P(\text{yes}|\text{today}=(\text{outlook}_{\text{Sunny}}, \text{Temp}_{\text{hot}}, \text{Humid}_{\text{normal}}, \text{Wind}_{\text{false}}))$
- Solution ->
$$P(\text{yes}|\text{today}=(\text{outlook}_{\text{Sunny}}, \text{Temp}_{\text{hot}}, \text{Humid}_{\text{normal}}, \text{Wind}_{\text{false}})) = P(\text{yes}) * P(\text{Sunny}|\text{yes}) * P(\text{hot}|\text{yes}) * P(\text{normal}|\text{yes}) * P(\text{False}|\text{yes})$$
- $P(\text{yes}|\text{today}) = (9/14) * (2/9 * 2/9 * 6/9 * 6/9) = 0.0141$
- $P(\text{no}|\text{today}) = (5/14) * (3/5 * 2/5 * 1/5 * 2/5) = 0.0069$
- $P(\text{yes}|\text{today}) + P(\text{no}|\text{today}) = 1$
- $P(\text{yes}|\text{today}) = 0.0141 / (0.0141 + 0.0069) = 0.671$
- $P(\text{no}|\text{today}) = 0.0069 / (0.0069 + 0.0141) = 0.323$
- $D = \text{argmax}(P(\text{yes}|\text{today}), P(\text{no}|\text{today})) = \text{argmax}(0.671, 0.323) = P(\text{yes}|\text{today}) \Rightarrow 0.671$

Naive Bayes classifier

- Problem = $P(\text{yes} | \text{today} = (\text{outlook}_{\text{Rainy}}, \text{Temp}_{\text{hot}}, \text{Humid}_{\text{high}}, \text{Wind}_{\text{true}}))$

Code

```
• import numpy as np # linear algebra
• import pandas as pd # data processing, CSV file I/O (e.g. pd.read_csv)
• import matplotlib.pyplot as plt
• import seaborn as sns
• from sklearn.model_selection import KFold,train_test_split,cross_val_score
•
• from sklearn.model_selection import train_test_split
•
• from sklearn.naive_bayes import GaussianNB
• from sklearn.metrics import make_scorer, accuracy_score,precision_score
•
• df=pd.read_csv("../root/Desktop/Class/pattern/code/iris.csv")
• print(df.head())
•
• df.drop(columns="Id",inplace=True)
•
•
• print(df.head())
• corr_matrix=df.corr()
• fig,ax=plt.subplots(figsize=(15,10))
• ax = sns.heatmap(corr_matrix,
•                 annot=True,
•                 linewidths=0.7,
•                 fmt=".3f",
•                 cmap="Accent_r")
• X=df.iloc[:,0:4].values
• y=df.iloc[:,4].values
•
•
• X_train,X_test,y_train,y_test=train_test_split(X,y,test_size=0.5,random_state=42)
• gaussian = GaussianNB()
• gaussian.fit(X_train, y_train)
• Y_pred = gaussian.predict(X_test)
• accuracy_nb=round(accuracy_score(y_test,Y_pred)* 100, 2)
• #acc_gaussian = round(gaussian.score(X_train, y_train) * 100, 2)
•
• accuracy = accuracy_score(y_test,Y_pred)
•
• print("accuracy_Naive Bayes: %.3f %accuracy)
```