

MTH301: Analysis I

Abhimanyu Sethia

190023

sethia@iitk.ac.in

Assignment 5

Question 9

Q9 $I = (\mathbb{R} \setminus \mathbb{Q}) \cap [0, 1]$

$Q = \mathbb{Q} \cap [0, 1]$

To prove:

(i) There is continuous map from I onto Q

(ii) There does not exist a continuous map from $[0, 1]$ onto Q

Proof of (i):

Consider the partition of I into countably many nonempty pieces, given by-

$$I_0 = (\frac{1}{2}, 1) \cap I$$

$$I_1 = (\frac{1}{4}, \frac{1}{2}) \cap I$$

$$\text{In general, } I_k = (\frac{1}{2^{k+1}}, \frac{1}{2^k}) \cap I$$

Note that, each I_k is open in I and $I = \bigcup_{k=0}^{\infty} I_k$.

Let Q be enumerated as $\{q_0, q_1, q_2, \dots\}$.

For any $x \in I$, $x \in I_k$ for exactly one k . Thereby, we define $f(x) = q_k$.

Since each I_k is non-empty, the surjectiveness of f follows.

We have seen in the lecture that proving f is continuous is equivalent to proving that for any open set O in Q , the set $f^{-1}(O)$ is open in I .

Now, consider any open set $O \subset Q$. So we can write, $O = \bigcup_{q_k \in O} \{q_k\}$

Therefore, $f^{-1}(O) = \bigcup_{q_k \in O} f^{-1}(q_k) = \bigcup_{q_k \in O} I_k$.

Since all I_k are open in I , union of open sets in I will also be open in I .

Therefore, by the theorem quoted above (from lecture), f is continuous.

Hence, proved. ■

Proof of (ii):

To prove (ii), we will use the following lemma:

Lemma: A non-empty connected subset of \mathbb{Q} must be singleton.

The lemma's proof is provided in Appendix 1.

Now, let there be a function $f : [0, 1] \rightarrow Q$ s.t. f is continuous.

Since $[0, 1]$ is connected, $f([0, 1])$ will also be connected. (from lecture)

Since $f([0, 1])$ is the range, it must be a subset of the co-domain i.e. Q . Also, $Q \subseteq \mathbb{Q}$.

Therefore, $f([0, 1])$ is a connected subset of \mathbb{Q} .

Hence, by our lemma, $f([0, 1])$ is a singleton. But Q is not a singleton. So, range is not same as the co-domain. Hence, f is not onto.

Therefore, we have proved that any continuous function from $[0, 1]$ to Q cannot be surjective. In other words, there cannot be a continuous map from $[0, 1]$ onto Q .

Hence, proved. ■

Appendix 1: Proof of lemma

Let X be a nonempty connected subset of \mathbb{Q} . We have to show that X is singleton. We prove by contradiction. Hence, assume that X is not singleton.

Therefore, $\exists a, b \in X$ s.t. $a < b$

Since irrationals are dense in rationals, $\exists c \in \mathbb{R} \setminus \mathbb{Q}$ s.t. $a < c < b$.

The sets $(c, +\infty)$ and $(-\infty, c)$ are open (disjoint) subsets of X and $X = (c, +\infty) \cup (-\infty, c)$.

Therefore, X is disconnected, which is a contradiction.

Hence, our assumption was wrong. That is, X is singleton.

Hence, proved. ■