

CRYSTALLINE REPRESENTATIONS AND WACH MODULES IN THE RELATIVE CASE

ABHINANDAN

Abstract. We study the notion of Wach modules in relative setting, generalizing the arithmetic case. Over an unramified base, for a p -adic representation admitting such structure, we examine the relationship between its relative Wach module and filtered (φ, ∂) -module. Moreover, we show that such a representation is crystalline (in the sense of Faltings-Brinon), and one can recover its filtered (φ, ∂) -module from the relative Wach module. Conversely, for low Hodge-Tate weights $[0, p-2]$, we construct relative Wach modules from free relative Fontaine-Laffaille modules (in the sense of Faltings).

Contents

1	Introduction	2
1.1	The arithmetic case	2
1.2	The relative case	3
1.3	Relative Fontaine-Laffaille modules	6
1.4	Setup and notations	6
2	p-adic Hodge theory	9
2.1	The de Rham period ring	9
2.2	The crystalline period ring	10
2.3	p -adic representations	12
3	(φ, Γ)-modules and crystalline coordinates	15
3.1	(φ, Γ) -modules	15
3.2	Crystalline coordinates	19
3.3	Cyclotomic embedding	20
3.4	Fat period rings	22
4	Finite height representations	23
4.1	The arithmetic case	23
4.2	The relative case	25
4.3	Statement of the main result	28
4.4	From (φ, Γ) -modules to (φ, ∂) -modules	33
4.5	Proof of Theorem 4.25	42
4.6	One-dimensional representations	47
5	Relative Fontaine-Laffaille modules	50
5.1	Some ideals of $\mathbf{A}_{R, \varpi}^+$ and $\mathbf{A}_{R, \varpi}^{\text{PD}}$	51
5.2	Equivalence of categories	54
5.3	Wach modules from Fontaine-Laffaille data	57
	References	64

1. Introduction

The theory of Wach modules for p -adic crystalline representations of the absolute Galois group of a finite unramified extension of \mathbb{Q}_p was introduced in the paper of Fontaine [Fon90]. This notion was further developed by Wach [Wac96; Wac97] and Berger [Ber04]. Over the years, this theory has found many applications, for example, to the Iwasawa theory of crystalline representations in [Ben00; BB08], and in the study of the p -adic local Langlands program [BB10]. Wach modules were also among one of the motivations for Scholze's idea of q -deformations [Sch17], which in turn paved the way for the theory of prisms and prismatic cohomology of Bhatt and Scholze developed in [BS19].

Our goal in this article is to upgrade the notion of Wach modules to the relative case by which we mean certain étale algebras over a formal torus (see §1.4 for precise setup). But before examining the relative case, let us recall the relation between Wach modules and crystalline representations in the arithmetic case.

1.1. The arithmetic case. Let p be a fixed prime number and let κ denote a finite field of characteristic p ; set $O_F = W(\kappa)$ to be the ring of p -typical Witt vectors with coefficients in κ and $F = \text{Fr}(O_F)$. Let \overline{F} denote a fixed algebraic closure of F , $\mathbb{C}_p := \widehat{\overline{F}}$ the p -adic completion, and $G_F = \text{Gal}(\overline{F}/F)$ the absolute Galois group of F . Further, let $F_\infty = \bigcup_n F(\mu_{p^n})$ with $\Gamma_F := \text{Gal}(F_\infty/F)$ and $H_F := \text{Gal}(\overline{F}/F_\infty)$. Finally, let \mathbb{C}_p^b denote the tilt of \mathbb{C}_p .

1.1.1. (φ, Γ_F) -modules. Using a certain period ring $\mathbf{A} \subset W(\mathbb{C}_p^b)$ stable under the Frobenius on Witt vectors and the G_F -action (see §3.1 for precise definition), Fontaine functorially attached to any \mathbb{Z}_p -representation T of G_F (i.e. finitely generated \mathbb{Z}_p -modules equipped with a linear and continuous G_F -action), the module $\mathbf{D}(T) = (\mathbf{A} \otimes_{\mathbb{Z}_p} T)^{H_F}$ over the two dimensional local ring $\mathbf{A}_F = \mathbf{A}^{H_F}$. The module $\mathbf{D}(T)$ is equipped with a (induced from \mathbf{A}) Frobenius-semilinear operator φ such that the image of φ generates $\mathbf{D}(T)$, i.e. $\mathbf{D}(T)$ is étale. Moreover, $\mathbf{D}(T)$ is equipped with a continuous and semilinear action of Γ_F and if T is free the \mathbf{A}_F -rank of $\mathbf{D}(T)$ equals the \mathbb{Z}_p -rank of T . In [Fon90] Fontaine established an equivalence of categories between \mathbb{Z}_p -representations of G_F and étale (φ, Γ_F) -modules over \mathbf{A}_F . Furthermore, this construction naturally extends to p -adic representations of G_F . Namely, using the period ring $\mathbf{B} = \mathbf{A}[\frac{1}{p}]$, Fontaine functorially attached to any p -adic representation V of G_F an étale (φ, Γ_F) -module $\mathbf{D}(V) = (\mathbf{B} \otimes_{\mathbb{Q}_p} V)^{H_F}$ over $\mathbf{B}_F = \mathbf{B}^{H_F}$ (i.e. there exists a \mathbb{Z}_p -lattice $T \subset V$ such that $\mathbf{D}(T)$ is an étale (φ, Γ_F) -module over \mathbf{A}_F). Moreover, he showed that this induces an equivalence between p -adic representations of G_F and étale (φ, Γ_F) -modules over \mathbf{B}_F .

1.1.2. Crystalline representations of G_F . Using another period ring \mathbf{B}_{cris} also equipped with a Frobenius and continuous G_F -action (see §2.2 for precise definition), Fontaine functorially attached to any p -adic representation V of G_F an F -vector space $\mathbf{D}_{\text{cris}}(V) = (\mathbf{B}_{\text{cris}} \otimes_{\mathbb{Q}_p} V)^{G_F}$. The F -vector space $\mathbf{D}_{\text{cris}}(V)$ is a filtered φ -module, i.e. it is equipped with a (induced from \mathbf{B}_{cris}) Frobenius-semilinear operator φ and a filtration. In case $\dim_F \mathbf{D}_{\text{cris}}(V) = \dim_{\mathbb{Q}_p} V$, such a representation is said to be crystalline (the terminology *crystalline* comes from the fact that for a smooth proper scheme X/O_F and $i \in \mathbb{N}$ the p -adic étale cohomology group of the generic fiber $V_i = H_{\text{ét}}^i(X_{\overline{F}}, \mathbb{Q}_p)$ is crystalline as a G_F -representation and the crystalline cohomology group of the special fiber $H_{\text{cris}}^i(X_\kappa/F)$ is naturally isomorphic to $\mathbf{D}_{\text{cris}}(V_i)$). Restricting the functor \mathbf{D}_{cris} to the subcategory of crystalline representations, in [Fon82] Fontaine observed that the associated filtered φ -modules are weakly admissible (a property relating the endomorphism φ and filtration on $\mathbf{D}_{\text{cris}}(V)$ in a non-trivial manner). In fact, in [CF00] Colmez and Fontaine showed that crystalline representations of G_F are equivalent to weakly admissible filtered φ -modules.

1.1.3. Arithmetic Wach modules. From the discussion above, it is a natural question to ask: Does there exist some direct relation between the étale (φ, Γ) -module of a crystalline representation and its associated weakly admissible filtered φ -module? For a fixed representation, this question could be rephrased in terms of comparing certain elements of the period rings \mathbf{B} and \mathbf{B}_{cris} . However, the rings \mathbf{B} and \mathbf{B}_{cris} are not comparable. So to answer this question, Fontaine considered a smaller period ring $\mathbf{B}^+ \subset \mathbf{B}$ stable under Frobenius and G_F -action and such that $\mathbf{B}^+ \hookrightarrow \mathbf{B}_{\text{cris}}$ stable under Frobenius and G_F -action. Using \mathbf{B}^+ he defined: a p -adic representation V of G_F is said to be of finite height if the associated (φ, Γ_F) -module $\mathbf{D}(V)$ admits a (φ, Γ_F) -stable lattice over the subring $\mathbf{B}_F^+ = (\mathbf{B}^+)^{H_F} \subset \mathbf{B}_F$ (see §4.1 for precise definitions).

In [Fon90] Fontaine conjectured that for a crystalline representation V of G_F there exist lattices inside $\mathbf{D}(V)$ over which the action of Γ_F admits a simpler form. More precisely, finite height and crystalline representations of G_F are related as follows:

Theorem 1.1 ([Wac96, Wach], [Col99, Colmez], [Ber04, Berger]). *Let V be a p -adic representation of G_F . Then V is crystalline if and only if it is of finite height and there exists $r \in \mathbb{Z}$ and a free \mathbf{B}_F^+ -submodule $N \subset \mathbf{D}(V)$ of rank $= \dim_{\mathbb{Q}_p} V$, stable under the action of Γ_F and such that Γ_F acts trivially over $(N/\pi N)(-r)$.*

Here $(-r)$ denotes the Tate twist. Note that in the situation of Theorem 1.1, the module N is not unique. A functorial construction was given by Berger in [Ber04], i.e. to any p -adic crystalline representation V of G_F he attached a canonical \mathbf{B}_F^+ -submodule $\mathbf{N}(V) \subset \mathbf{D}(V)$ which he called the Wach module of V . Moreover, Berger established an equivalence of categories between crystalline representations of G_F and Wach modules over \mathbf{B}_F^+ . Furthermore, Berger obtained an integral version of his result by considering the period ring $\mathbf{A}^+ = \mathbf{A} \cap \mathbf{B}^+ \subset \mathbf{B}$ stable under Frobenius and G_F -action. He showed that for a crystalline representation V of G_F , there exists a bijection between G_F -stable \mathbb{Z}_p -lattices $T \subset V$ and integral Wach modules $\mathbf{N}(T) \subset \mathbf{N}(V)$ where $\mathbf{N}(T)$ is defined over the integral subring $\mathbf{A}_F^+ = (\mathbf{A}^+)^{H_F}$. Finally, given $\mathbf{N}(V)$ one can canonically recover the other linear algebraic object attached to V , i.e. $\mathbf{D}_{\text{cris}}(V)$ (see [Ber04, Propositions II.2.1 & III.4.4]).

1.2. The relative case. The motivation for defining Wach modules in the relative case and exploring its relation with $\mathcal{O}\mathbf{D}_{\text{cris}}(V)$ (see §2 for notations) comes from the hope of computing Galois cohomology of p -adic representations using syntomic complexes with coefficients in $\mathcal{O}\mathbf{D}_{\text{cris}}(V)$. Using syntomic complexes and techniques from the theory of (φ, Γ) -modules, this was done for the trivial representation by Colmez and Nizioł [CN17]. A generalization of these complexes to non-trivial coefficients can be found in [Abh22] and [Abh21, Chapter 5].

In this article, we are interested in the p -adic Hodge theory of an étale algebra over a formal torus defined over O_F . More precisely, let $d \in \mathbb{N}$ and $X = (X_1, X_2, \dots, X_d)$ be some indeterminates, $O_F\{X, X^{-1}\}$ the p -adic completion of a d -dimensional torus over O_F and let R denote the p -adic completion of an étale algebra over $O_F\{X, X^{-1}\}$ with non-empty and geometrically integral special fiber. Next, let G_R denote the étale fundamental group of $R[\frac{1}{p}]$ and Γ_R the Galois group of the cyclotomic tower over R and $H_R = \text{Ker}(G_R \rightarrow \Gamma_R)$ (see §3.1 for precise definitions). In the relative setting, on one hand Brinon has developed the theory of crystalline representations of G_R [Bri08], while on the other hand Andreatta, Brinon and Iovita have developed the theory of (φ, Γ_R) -modules in [And06; AB08; AI08].

Remark 1.2. Note that in Theorem 1.1 it is important to restrict to an unramified extension F/\mathbb{Q}_p . For ramified extensions, such a statement does not hold in general. Therefore, in the relative setting we consider an analogue of “unramified extension of \mathbb{Q}_p ” (indeed, by removing the geometric coordinates one obtains $R = O_F$).

1.2.1. (φ, Γ_R) -modules. Analogous to the arithmetic case, we have relative period rings $\mathbf{A} \subset \mathbf{B} \supset \mathbf{B}^+$ and $\mathbf{A}^+ = \mathbf{A} \cap \mathbf{B}^+ \subset \mathbf{B}$ (see §3.1 for precise definition) equipped with Frobenius and a continuous action of G_R . Let V be a p -adic representation of G_R , then one can functorially attach to V a projective and étale (φ, Γ_R) -module $\mathbf{D}(V) = (\mathbf{B} \otimes_{\mathbb{Q}_p} V)^{H_R}$ over $\mathbf{B}_R = \mathbf{B}^{H_R}$ of rank $= \dim_{\mathbb{Q}_p} V$ equipped with a Frobenius-semilinear operator φ and a semilinear and continuous action of Γ_R . This induces an equivalence of categories between p -adic representations of G_R and étale (φ, Γ_R) -modules over \mathbf{B}_R . Similarly, using the period ring \mathbf{A} one can functorially attach to any \mathbb{Z}_p -representation T of G_R an étale (φ, Γ_R) -module $\mathbf{D}(T) = (\mathbf{A} \otimes_{\mathbb{Z}_p} T)^{H_R}$ over the period ring $\mathbf{A}_R = \mathbf{A}^{H_R}$. Again, this induces an equivalence between \mathbb{Z}_p -representations of G_R and étale (φ, Γ_R) -modules over \mathbf{A}_R .

1.2.2. Relative Wach modules. Using the period ring \mathbf{A}^+ we set $\mathbf{D}^+(T) = (\mathbf{A}^+ \otimes_{\mathbb{Z}_p} T)^{H_R}$, which is a (φ, Γ_R) -module over $\mathbf{A}_R^+ = (\mathbf{A}^+)^{H_R}$ and let $q = \frac{\varphi(\pi)}{\pi}$, where π is the usual element in Fontaine's constructions (see §2.1 for notations). Note that for a finite free \mathbb{Z}_p -representation T of G_R the \mathbf{A}_R -module $\mathbf{D}(T)$ is finite projective, however it is not known whether $\mathbf{D}^+(T)$ is projective. So, we introduce the following definition:

Definition 1.3. A *positive finite q -height* representation is a p -adic representation V of G_R admitting a \mathbb{Z}_p -lattice $T \subset V$ such that there exists a finite projective \mathbf{A}_R^+ -submodule $\mathbf{N}(T) \subset \mathbf{D}^+(T)$ of rank $= \dim_{\mathbb{Q}_p} V$ satisfying the following conditions:

- (i) $\mathbf{N}(T)$ is stable under the action of φ and Γ_R and $\mathbf{A}_R \otimes_{\mathbf{A}_R^+} \mathbf{N}(T) \xrightarrow{\sim} \mathbf{D}(T)$;
- (ii) The \mathbf{A}_R^+ -module $\mathbf{N}(T)/\varphi^*(\mathbf{N}(T))$ is killed by q^s for some $s \in \mathbb{N}$;
- (iii) The action of Γ_R is trivial on $\mathbf{N}(T)/\pi \mathbf{N}(T)$;
- (iv) There exists $R' \subset \overline{R}$ finite étale over R such that the $\mathbf{A}_{R'}^+$ -module $\mathbf{A}_{R'}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ is free.

The module $\mathbf{N}(T)$ is a *Wach module* associated to T and we set $\mathbf{N}(V) := \mathbf{N}(T)[\frac{1}{p}]$ which satisfies analogous properties. The *height* of V is the smallest $s \in \mathbb{N}$ satisfying (ii) above.

Remark 1.4. (i) A finite q -height representation is twist of a positive one by some power of the p -adic cyclotomic character (see Definition 4.9 for details). The terminology “positive” refers to the fact that the Wach module $\mathbf{N}(T)$ is stable under the Frobenius-semilinear operator φ . It is motivated by the fact (and as we will show) that V is positive crystalline (see Theorem 1.6).

- (ii) In the arithmetic case, i.e. $R = O_F$, the notion of finite height representations in Theorem 1.1 and finite q -height representations in Definition 1.3 are related. In fact, in the arithmetic case using Definition 1.3 one obtains the functorial Wach module of Berger mentioned above (see [Ber04, Proposition II.1.1]).

1.2.3. Crystalline representations of G_R . Using the period ring $\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ Brinon functorially attaches to any p -adic representation V of G_R an $R[\frac{1}{p}]$ -module

$$\mathcal{O}\mathbf{D}_{\text{cris}}(V) := (\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} V)^{G_R}.$$

The module $\mathcal{O}\mathbf{D}_{\text{cris}}(V)$ is called a filtered (φ, ∂) -module, i.e. it is equipped with a filtration, a Frobenius-semilinear endomorphism φ and a quasi-nilpotent integrable connection ∂ satisfying Griffiths transversality with respect to the filtration (see §2.3 for precise definitions). The representation V is said to be crystalline if the natural map is an isomorphism

$$\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{R[1/p]} \mathcal{O}\mathbf{D}_{\text{cris}}(V) \xrightarrow{\sim} \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} V,$$

compatible with Frobenius, filtration, connection and the action of G_R on each side. Moreover, Brinon also defined the notion of weak admissibility in the relative case and showed that $\mathcal{OD}_{\text{cris}}(V)$ is weakly admissible for crystalline representations (see [Bri08, Chapitre 8] for more details).

Notation 1.5. We use period rings such as $\mathcal{OB}_{\text{cris}}(\overline{R})$ which is a modified version of Fontaine's relative period ring $\mathbf{B}_{\text{cris}}(\overline{R})$ (see §2.2 for details). The notation \mathcal{O} here indicates that apart from Frobenius, filtration and G_R -action, we have a connection over $\mathcal{OB}_{\text{cris}}(\overline{R})$ and we will call such rings fat relative period rings. However, note that in [Bri08] Brinon denotes these rings as $B_{\text{cris}}(R)$ and $B_{\text{cris}}^{\nabla}(R)$, respectively. Similarly, we will use the notation $\mathcal{OD}_{\text{cris}}(V)$ and $\mathbf{D}_{\text{cris}}(V)$ for modules instead of Brinon's $D_{\text{cris}}(V)$ and $D_{\text{cris}}^{\nabla}(V)$, respectively. We hope it is not too confusing for the reader.

1.2.4. Main result. Our aim is to show that for positive finite q -height representations, the \mathbf{B}_R^+ -module $\mathbf{N}(V)$ and the $R[\frac{1}{p}]$ -module $\mathcal{OD}_{\text{cris}}(V)$ are related in a precise manner and the latter can be recovered from the former. To relate these objects we consider the ring $R[\varpi]$ where $\varpi = \zeta_p - 1$ for a primitive p -th root of unity ζ_p (take $\varpi = \zeta_{p^2} - 1$ if $p = 2$ for a primitive p^2 -th root of unity ζ_{p^2}), and using this ring we construct a fat relative period ring $\mathcal{OA}_{R,\varpi}^{\text{PD}} \subset \mathcal{OB}_{\text{cris}}(\overline{R})$ equipped with compatible Frobenius, filtration, connection and the action of Γ_R (see §4.3 for precise definitions). The main result of this article is as follows:

Theorem 1.6 (see Theorem 4.25). *Let V be a positive finite q -height representation of G_R , then*

- (i) *V is a positive crystalline representation.*
- (ii) *Let $M := (\mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))^{\Gamma_R}$, then after extending scalars to $\mathcal{OA}_{R,\varpi}^{\text{PD}}$ and inverting p , we obtain a natural isomorphism*

$$\mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_R M[\frac{1}{p}] \xrightarrow{\sim} \mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V),$$

compatible with Frobenius, filtration, connection and the action of Γ_R on each side.

- (iii) *We have an isomorphism of $R[\frac{1}{p}]$ -modules*

$$\mathcal{OD}_{\text{cris}}(V) \xleftarrow{\sim} (\mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))^{\Gamma_R}[\frac{1}{p}],$$

compatible with Frobenius, filtration, and connection on each side. Therefore, we obtain a comparison isomorphism

$$\mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V) \xrightarrow{\sim} \mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_R \mathcal{OD}_{\text{cris}}(V),$$

compatible with Frobenius, filtration, connection and the action of Γ_R on each side.

Let us mention the idea of the proof. In case $\mathbf{N}(T)$ is free, we proceed in two steps: First, we describe a process (see Proposition 4.28 for details) by which we can recover a submodule of $\mathcal{OD}_{\text{cris}}(V)$ starting with the Wach module $\mathbf{N}(T)$, establishing a comparison over $\mathcal{OA}_{R,\varpi}^{\text{PD}}$ between the submodule obtained and the Wach module. Next, the claims made in the theorem are shown by exploiting properties of Wach modules and the comparison obtained in the first step. In the first step, one can take two approaches to obtain generators of the promised submodule of $\mathcal{OD}_{\text{cris}}(V)$: either by taking Γ_R -fixed points of $\mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ (by successively approximating for Γ_R -action on a basis of $\mathbf{N}(T)$); or by taking elements killed by differential operators defined using topological generators of Γ_R (see Lemma 4.41 for details). In this paper, we take the latter approach whereas the former approach is detailed in [Abh21, Chapter 3]. In

the general case when $\mathbf{N}(T)$ is projective, using property (iv) in Definition 1.3 one can pass to an extension $\mathbf{A}_R^+ \subset \mathbf{A}_{R'}^+$ to obtain a free Wach module, then use the preceding argument and finally apply Galois descent to obtain the theorem (see Proposition 4.28 for details). Finally, we also show that all one-dimensional crystalline representations are of finite q -height and for such representations one can directly compare $\mathcal{O}\mathbf{D}_{\text{cris}}(V)$ and the Wach module $\mathbf{N}(V)$.

1.3. Relative Fontaine-Laffaille modules. After obtaining Theorem 1.6 above, it is natural to wonder if a converse statement could be true, i.e. starting with a lattice $T \subset V$ of a crystalline representation G_R , is it possible to construct the Wach module $\mathbf{N}(T)$? In the arithmetic setting, for p -adic crystalline representations of G_F , this was shown to be true by Wach [Wac96], and the statement was refined by Berger [Ber04]. In the relative case, the picture is quite encouraging when we restrict to Hodge-Tate length $\leq p - 2$ (also see Remark 1.9).

For a p -adic crystalline representation of G_F with Hodge-Tate length $\leq p - 1$, there exists a canonical O_F -lattice inside $\mathbf{D}_{\text{cris}}(V)$ called the Fontaine-Laffaille module defined in [FL82]. In this case, Wach constructed Wach modules out of Fontaine-Laffaille data in [Wac97]. In the relative setting, Faltings studied relative Fontaine-Laffaille modules in [Fal89] and used them to functorially recover \mathbb{Z}_p -lattices inside crystalline representations of G_R . Recently, for free relative Fontaine-Laffaille modules of filtration length $\leq p - 2$, adapting techniques from Wach's computations, Tsuji has constructed generalized representations of G_R over $\mathbf{A}_{\text{inf}}(\overline{R})$ (see [Tsu20]). In fact, it is possible to show that starting with a free relative Fontaine-Laffaille module, one can obtain a free relative Wach module over \mathbf{A}_R^+ .

Theorem 1.7 (see Theorem 5.5). *Let M be a free relative Fontaine-Laffaille module over R of level $[0, p - 2]$, and let $T_{\text{cris}}(M)$ denote the associated \mathbb{Z}_p -representation of G_R . Then, the p -adic representation $V_{\text{cris}}(M) := \mathbb{Q}_p \otimes_{\mathbb{Z}_p} T_{\text{cris}}(M)$ is a positive finite q -height representation.*

Twisting the representation thus obtained by powers of the cyclotomic character, generalizes the statement to all free Fontaine-Laffaille modules with filtration length $\leq p - 2$.

The proof of the theorem crucially exploits the computation of Fontaine [Fon94], Wach [Wac97] and Tsuji [Tsu20]. It follows in three steps: First, starting with a Fontaine-Laffaille module, we obtain an $\mathbf{A}_{R, \varpi}^{\text{PD}}$ -module using formal properties of crystalline site for maps $\theta : \mathbf{A}_{R, \varpi}^{\text{PD}} \twoheadrightarrow R$ and $\theta_R : \mathcal{O}\mathbf{A}_{R, \varpi}^{\text{PD}} \twoheadrightarrow R$ (see §5.3.1 for details). Next, we exploit equivalence of categories obtained in Theorem 5.21 by extending scalars along the maps $\mathbf{A}_{R, \varpi}^{\text{PD}} \twoheadrightarrow \mathbf{A}_{R, \varpi}^{\text{PD}}/I^{(p-1)}\mathbf{A}_{R, \varpi}^{\text{PD}} \xleftarrow{\sim} \mathbf{A}_{R, \varpi}^+/I^{(p-1)}\mathbf{A}_{R, \varpi}^+ \leftarrow \mathbf{A}_{R, \varpi}^+$ (see Proposition 5.12 for explanations). This gives us an $\mathbf{A}_{R, \varpi}^+$ -module with precise description of the Frobenius and the action of Γ_R . Finally, we descend over to the ring \mathbf{A}_R^+ by exploiting the Frobenius and Γ_R -action, thus obtaining a Wach module over \mathbf{A}_R^+ and proving the theorem (see §5.3.2).

Remark 1.8. In a recent work, Morrow and Tsuji have developed a theory of coefficients for integral p -adic Hodge theory in [MT20]. Extending scalars of relative Wach modules along $O_F[[\pi]] \rightarrow \mathbf{A}_{\text{inf}}(O_{\overline{F}})$ would yield generalized representations over $\mathbf{A}_{\text{inf}}^{\square}(R)$ in the sense of Morrow-Tsuji.

Remark 1.9. Recent developments in the theory of prismatic crystals [BS21; DLMS22; GR22], indicate that to obtain a full converse statement, i.e. to construct Wach modules from lattices inside crystalline representations, one needs to generalize Definition 1.3 slightly. This is a work in progress and we will report further on this line of investigation in future.

1.4. Setup and notations. In this section we will describe the setup for the rest of the text and fix some notations.

Convention. We will work under the convention that $0 \in \mathbb{N}$, the set of natural numbers.

Let p be a fixed prime number, κ a finite field of characteristic p , $W := W(\kappa)$ the ring of p -typical Witt vectors with coefficients in κ and $F := W[\frac{1}{p}]$, the fraction field of W . In

particular, F is an unramified extension of \mathbb{Q}_p with ring of integers $O_F = W$. Let \bar{F} be a fixed algebraic closure of F so that its residue field, denoted as $\bar{\kappa}$, is an algebraic closure of κ . Further, we denote by $G_F = \text{Gal}(\bar{F}/F)$, the absolute Galois group of F .

Let $Z = (Z_1, \dots, Z_s)$ denote a set of indeterminates and $\mathbf{k} = (k_1, \dots, k_s) \in \mathbb{N}^s$ be a multi-index, then we write $Z^{\mathbf{k}} := Z_1^{k_1} \cdots Z_s^{k_s}$. For $\mathbf{k} \rightarrow +\infty$ we will mean that $\sum k_i \rightarrow +\infty$. Now for a topological algebra Λ we define

$$\Lambda\{Z\} := \left\{ \sum_{\mathbf{k} \in \mathbb{N}^s} a_{\mathbf{k}} Z^{\mathbf{k}}, \text{ where } a_{\mathbf{k}} \in \Lambda \text{ and } a_{\mathbf{k}} \rightarrow 0 \text{ as } \mathbf{k} \rightarrow +\infty \right\}.$$

We fix $d \in \mathbb{N}$ and let $X = (X_1, X_2, \dots, X_d)$ be some indeterminates. Let R be the p -adic completion of an étale algebra over $O_F\{X, X^{-1}\}$ with non-empty geometrically integral special fiber. In particular, we have a presentation

$$R = O_F\{X, X^{-1}\}\{Z_1, \dots, Z_s\}/(Q_1, \dots, Q_s),$$

where $Q_i(Z_1, \dots, Z_s) \in O_F\{X, X^{-1}\}[Z_1, \dots, Z_s]$ for $1 \leq i \leq s$ are multivariate polynomials such that $\det(\frac{\partial Q_i}{\partial Z_j})_{1 \leq i, j \leq s}$ is invertible in R . The algebra $R[\frac{1}{p}]$ is the relative analogue of “finite unramified extension of \mathbb{Q}_p ” (indeed, by removing the geometric coordinates we will obtain $R[\frac{1}{p}] = F$).

Remark 1.10. Note that Theorem 1.1 serves as our main motivation for the theory developed in this article. The assumptions we put on R generalizes the fact that “ F is unramified over \mathbb{Q}_p ”.

The p -adic Hodge theory over R is the study of p -adic representations of the étale fundamental group of $R[\frac{1}{p}]$, which we introduce next. We fix an algebraic closure $\overline{\text{Fr}}(\bar{R})$ of $\text{Fr}(R)$ containing \bar{F} . Let \bar{R} denote the union of finite R -subalgebras $S \subset \overline{\text{Fr}}(\bar{R})$, such that $S[\frac{1}{p}]$ is étale over $R[\frac{1}{p}]$. Let $\bar{\eta}$ denote a geometric point of the generic fiber $\text{Spec } R[\frac{1}{p}]$ and let $G_R := \pi_1^{\text{ét}}(\text{Spec } R[\frac{1}{p}], \bar{\eta})$ denote the étale fundamental group. By [Gro63, Exposé V, §8], we can write this étale fundamental group as the Galois group (of the fraction field of $\bar{R}[\frac{1}{p}]$ over the fraction field of $R[\frac{1}{p}]$)

$$G_R = \pi_1^{\text{ét}}(\text{Spec } R[\frac{1}{p}], \bar{\eta}) = \text{Gal}(\bar{R}[\frac{1}{p}]/R[\frac{1}{p}]).$$

For $n \in \mathbb{N}$, let $F_n := F(\mu_{p^n})$. From now onwards, we will fix some $m \in \mathbb{N}_{\geq 1}$ (take $m \in \mathbb{N}_{\geq 2}$ if $p = 2$) and set $K := F_m$, with its ring of integers O_K . The element $\varpi = \zeta_{p^m} - 1 \in O_K$ is a uniformizer of K , and its minimal polynomial $P_{\varpi}(X) = \frac{(1+X)^{p^m} - 1}{(1+X)^{p^{m-1}} - 1}$ is an Eisenstein polynomial in $W[X]$ of degree $e := [K : F] = p^{m-1}(p-1)$. Finally, for $S = R[\varpi] = O_K \otimes_{O_F} R$ we have that $R[\varpi]$ is totally ramified at the prime ideal $(p) \subset R[\varpi]$. And similar to above, we obtain Galois groups $G_K \triangleleft G_F$ and $G_S \triangleleft G_R$ respectively, such that $G_R/G_S = G_F/G_K = \text{Gal}(K/F)$. Finally, we have that R and $R[\varpi]$ are *small* algebras in the sense of Faltings (see [Fal88, §II 1(a)]).

For $k \in \mathbb{N}$, let Ω_R^k denote the p -adic completion of module of k -differentials of R relative to \mathbb{Z} . Then, we have

$$\Omega_R^1 = \bigoplus_{i=1}^d R d \log X_i, \text{ and } \Omega_R^k = \bigwedge_R^k \Omega_R^1.$$

We also have that $R/pR \xrightarrow{\sim} S/\varpi S$ and for all $n \in \mathbb{N}$, $R/p^n R$ is a smooth $\mathbb{Z}/p^n \mathbb{Z}$ -algebra. Finally, we have a unique lift $\varphi : R \rightarrow R$ of the absolute Frobenius $x \mapsto x^p$ over R/pR such that $\varphi(X_i) = X_i^p$, for all $1 \leq i \leq d$ (in general, a lift of Frobenius modulo p need not be unique, see [Bri08, p.9]).

Convention. Let A be a ring and $I \subsetneq A$ an ideal. We say that an A -module M is I -adically complete if and only if $M \xrightarrow{\sim} \varprojlim_n M/I^n M$.

Acknowledgements. The work presented here was part of my doctoral thesis. I would like to thank my supervisor Denis Benois for encouraging me to think about this question, having many helpful discussions, reading an earlier version of the article and suggesting improvements. I would also like to thank Takeshi Tsuji for pointing out connections between this work and his results in [Tsu20] and helpful discussions concerning contents of §5 which helped improve the exposition, in particular, the proof of Theorem 5.5. Moreover, I would like to thank Lorenzo Ramero for discussions around Remark 3.3 and suggesting improvements in writing and Nicola Mazzari for suggesting several improvements in the exposition. Finally, many thanks to the referee for carefully reading the article and suggesting many improvements, in particular, proofs of Lemma 3.9 and Lemma 4.41 could be shortened.

2. p -adic Hodge theory

In this section we will recall some constructions and results in relative p -adic Hodge theory developed in [Bri08], albeit in a simpler setting compared to Brinon's book. As we will be using different notations compared to Brinon, we will make most of the definitions explicit.

We are interested in exploring the relationship between p -adic crystalline representations and finite height representations of G_R . This will be detailed in §4 and §5. To carry out some computations in the aforementioned sections, we will need to extend our base field (hence the base ring) by adjoining some p -power roots of unity (see the field K and the ring $S = R[\varpi]$ in §1.4). As a consequence, we will also require the corresponding period rings defined for such rings. However, in §2.1, §2.2 & §2.3 we will only recall results from [Bri08] by fixing our base as R . As we shall see the period rings will only depend on \overline{R} and we have $\overline{S} = \overline{R} \subset \overline{\text{Fr}(R)} = \overline{\text{Fr}(S)}$, therefore fixing our base as R is sufficient (see [Bri08] for general constructions).

2.1. The de Rham period ring. We will recall definitions and properties of the relative version of Fontaine's period ring \mathbf{B}_{dR} (see [Fon94] for classical case).

2.1.1. The ring $\mathbb{C}^+(\overline{R})$ and its tilt. Let \mathbb{C}_p denote the p -adic completion of \overline{F} . Recall that \overline{R} is the union of finite R -subalgebras $S \subset \overline{\text{Fr}(R)} = \overline{\text{Fr}(R[\varpi])}$, such that $S[\frac{1}{p}]$ is étale over $R[\frac{1}{p}]$. Let $\mathbb{C}^+(\overline{R})$ denote the p -adic completion of \overline{R} and $\mathbb{C}(\overline{R}) = \mathbb{C}^+(\overline{R})[\frac{1}{p}]$. We define the tilt $\mathbb{C}^+(\overline{R})^\flat := \lim_{x \mapsto x^p} \mathbb{C}^+(\overline{R})/p = \lim_{x \mapsto x^p} \overline{R}/p$ and equip it with the inverse limit topology (where we equip \overline{R}/p with the discrete topology) and let $\mathbb{C}(\overline{R})^\flat = \mathbb{C}^+(\overline{R})^\flat[\frac{1}{p}]$ for $p^\flat := (p, p^{1/p}, p^{1/p^2}, \dots) \in \mathbb{C}^+(\overline{R})^\flat$ and equipped with the coarsest ring topology such that $\mathbb{C}^+(\overline{R})$ is an open subring. Note that an element $x \in \mathbb{C}(\overline{R})^\flat$ can be described as a sequence $(x_n)_{n \in \mathbb{N}}$, with $x_n \in \mathbb{C}(\overline{R})$ and $x_{n+1}^p = x_n$ for all $n \in \mathbb{N}$. These rings admit a continuous G_R -action for the topology described.

We will fix some choices of compatible p -power roots which will appear in the sequel. Let $\varepsilon := (1, \zeta_p, \zeta_{p^2}, \dots) \in \mathbb{C}_p^\flat$, $X_i^\flat := (X_i, X_i^{1/p}, X_i^{1/p^2}, \dots) \in \mathbb{C}(\overline{R})^\flat$ for $1 \leq i \leq d$. We set $\mathbf{A}_{\text{inf}}(\overline{R}) := W(\mathbb{C}^+(\overline{R})^\flat)$, the ring of p -typical Witt vectors with coefficients in $\mathbb{C}^+(\overline{R})^\flat$ equipped with weak topology (see [AI08, §2.10]). The absolute Frobenius on $\mathbb{C}^+(\overline{R})^\flat$ lifts to an endomorphism $\varphi : \mathbf{A}_{\text{inf}}(\overline{R}) \rightarrow \mathbf{A}_{\text{inf}}(\overline{R})$ and the G_R -action extends to $\mathbf{A}_{\text{inf}}(\overline{R})$ such that the action is continuous for the weak topology. For $x \in \mathbb{C}^+(\overline{R})^\flat$, let $[x] = (x, 0, 0, \dots) \in \mathbf{A}_{\text{inf}}(\overline{R})$ denote its Teichmüller representative. So we have $[\varepsilon] \in \mathbf{A}_{\text{inf}}(\overline{R})$ with $g[\varepsilon] = [\varepsilon]^{\chi(g)}$ for $g \in G_R$ and $\chi : G_R \rightarrow \mathbb{Z}_p^\times$ the p -adic cyclotomic character and $\varphi([\varepsilon]) = [\varepsilon]^p$. Now any element $x \in \mathbf{A}_{\text{inf}}(\overline{R})$ can be uniquely written as $x = \sum_{k \in \mathbb{N}} p^k [x_k]$ for $x_k \in \mathbb{C}^+(\overline{R})^\flat$. So we set $\pi := [\varepsilon] - 1$, $\pi_1 := \varphi^{-1}(\pi) = [\varepsilon^{1/p}] - 1$, and $\xi := \frac{\pi}{\pi_1}$. Clearly we have $g(\pi) = (1 + \pi)^{\chi(g)} - 1$ for $g \in G_R$ and $\varphi(\pi) = (1 + \pi)^p - 1$.

2.1.2. Definition of $\mathcal{O}\mathbf{B}_{\text{dR}}(\overline{R})$. We have Fontaine's θ -map defined as $\theta : \mathbf{A}_{\text{inf}}(\overline{R}) \rightarrow \mathbb{C}^+(\overline{R})$ sending $\sum_{k \in \mathbb{N}} p^k [x_k] \mapsto \sum_{k \in \mathbb{N}} p^k x_k^\sharp$, it is a G_R -equivariant surjective ring homomorphism whose kernel is principal and generated by, for example, $p - [p^\flat]$ or ξ (see [Fon82, Proposition 2.4 (ii)]). By \mathbb{Q}_p -linearity, the map θ can be extended to $\theta : \mathbf{A}_{\text{inf}}(\overline{R})[\frac{1}{p}] \rightarrow \mathbb{C}(\overline{R})$ and we define

$$\mathbf{B}_{\text{dR}}^+(\overline{R}) := \lim_n \mathbf{A}_{\text{inf}}(\overline{R})[\frac{1}{p}] / (\text{Ker } \theta)^n,$$

as the $(\text{Ker } \theta)$ -adic completion of $\mathbf{A}_{\text{inf}}(\overline{R})[\frac{1}{p}]$. The ring $\mathbf{B}_{\text{dR}}^+(\overline{R})$ is an F -algebra and admits a G_R -action. The map θ further extends to a G_R -equivariant surjective ring homomorphism $\theta : \mathbf{B}_{\text{dR}}^+(\overline{R}) \rightarrow \mathbb{C}(\overline{R})$ with $\text{Ker } \theta = t\mathbf{B}_{\text{dR}}^+(\overline{R})$, where $t := \log[\varepsilon] = \log(1 + \pi) = \sum_{k \in \mathbb{N}} (-1)^k \frac{\pi^{k+1}}{k+1} \in \mathbf{B}_{\text{dR}}^+(\overline{R})$ such that $g \in G_R$ acts by $g(t) = \chi(g)t$. By functoriality of the construction of $\mathbf{B}_{\text{dR}}^+(\overline{R})$, the homomorphism $O_{\overline{F}} \rightarrow \overline{R}$ induces an injection $\mathbf{B}_{\text{dR}}^+(O_{\overline{F}}) \rightarrow \mathbf{B}_{\text{dR}}^+(\overline{R})$. The ring $\mathbf{B}_{\text{dR}}^+(\overline{R})$ is t -torsion free, so we set $\mathbf{B}_{\text{dR}}(\overline{R}) := \mathbf{B}_{\text{dR}}^+(\overline{R})[\frac{1}{t}]$. The G_R -action extends to $\mathbf{B}_{\text{dR}}(\overline{R})$ and the ring

$\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ admits a natural G_R -stable filtration given as $\mathrm{Fil}^r \mathbf{B}_{\mathrm{dR}}(\overline{R}) := t^r \mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ for $r \in \mathbb{Z}$ and we equip $\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ with the induced filtration (see [Bri08, §5.1] for details).

We can extend the map $\theta : \mathbf{A}_{\mathrm{inf}}(\overline{R}) \rightarrow \mathbb{C}^+(\overline{R})$ by R -linearity to obtain a ring homomorphism $\theta_R : R \otimes_{\mathbb{Z}} \mathbf{A}_{\mathrm{inf}}(\overline{R}) \rightarrow \mathbb{C}^+(\overline{R})$. Let $\mathcal{O}\mathbf{A}_{\mathrm{inf}}(\overline{R})$ denote the $\theta_R^{-1}(p\mathbb{C}^+(\overline{R}))$ -adic completion of $R \otimes_{\mathbb{Z}} \mathbf{A}_{\mathrm{inf}}(\overline{R})$ (the ideal $\theta_R^{-1}(p\mathbb{C}^+(\overline{R}))$ is generated by p and $\mathrm{Ker} \theta_R$), then θ_R extends to a surjective homomorphism $\theta_R : \mathcal{O}\mathbf{A}_{\mathrm{inf}}(\overline{R})[\frac{1}{p}] \rightarrow \mathbb{C}(\overline{R})$. Define

$$\mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R}) := \lim_n \mathcal{O}\mathbf{A}_{\mathrm{inf}}(\overline{R})[\frac{1}{p}] / (\mathrm{Ker} \theta_R)^n,$$

as the $(\mathrm{Ker} \theta_R)$ -adic completion of $\mathcal{O}\mathbf{A}_{\mathrm{inf}}(\overline{R})[\frac{1}{p}]$. The ring $\mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ is an $R[\frac{1}{p}]$ -algebra and admits a G_R -action. The homomorphism θ_R extends to a G_R -equivariant surjective homomorphism $\theta_R : \mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R}) \rightarrow \mathbb{C}(\overline{R})$. The ring $\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ is t -torsion free and we set $\mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R}) := \mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R})[\frac{1}{t}]$. Moreover, the G_R -action extends to $\mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R})$.

2.1.3. Structure and properties of $\mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R})$. A more explicit description of the ring $\mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ can be given. Note that $X_i \otimes 1 - 1 \otimes [X_i^p] \in \mathrm{Ker} \theta_R \subset R \otimes_{\mathbb{Z}} \mathbf{A}_{\mathrm{inf}}(\overline{R})$ for $1 \leq i \leq d$ and let z_i denote its image in $\mathcal{O}\mathbf{A}_{\mathrm{inf}}(\overline{R}) \subset \mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$. Since $\mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ is complete for the $(\mathrm{Ker} \theta_R)$ -adic topology, the homomorphism $\mathbf{B}_{\mathrm{dR}}^+(\overline{R}) \rightarrow \mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ extends to a homomorphism

$$\begin{aligned} f : \mathbf{B}_{\mathrm{dR}}^+(\overline{R})[[T_1, \dots, T_d]] &\longrightarrow \mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R}) \\ T_i &\longmapsto z_i, \quad \text{for } 1 \leq i \leq d. \end{aligned}$$

In fact, f is an isomorphism and $\mathrm{Ker} \theta_R = (t, z_1, \dots, z_d) \subset \mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$. Therefore, one can identify $\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ with a subring of $\mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$. There is a natural G_R -stable filtration on $\mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ given by $\mathrm{Fil}^r \mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R}) = (\mathrm{Ker} \theta_R)^r$ for $r \in \mathbb{N}$. We set $\mathrm{Fil}^0 \mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R}) := \sum_{n=0}^{+\infty} t^{-n} \mathrm{Fil}^n \mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R}) = \mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R})[\frac{z_1}{t}, \dots, \frac{z_d}{t}]$ and $\mathrm{Fil}^r \mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R}) := t^r \mathrm{Fil}^0 \mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R})$ for $r \in \mathbb{Z}$, satisfying the same conditions. Moreover, the induced filtrations on $\mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$, $\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ and $\mathbf{B}_{\mathrm{dR}}(\overline{R})$ match with the ones defined before. Finally, we have $(\mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R}))^{G_R} = R[\frac{1}{p}]$ (see [Bri08, §5.2] for details).

We can equip the rings $\mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ and $\mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R})$ with a connection. Let N_i denote the unique $(\mathrm{Ker} \theta_R)$ -adically continuous, $\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ -linear derivation on $\mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ given as $N_i(z_j) = \delta_{ij} X_j$ for $1 \leq i, j \leq d$, where δ_{ij} denotes the Kronecker delta symbol. Furthermore, the derivation N_i extends to a $\mathbf{B}_{\mathrm{dR}}(\overline{R})$ -linear derivation on $\mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R})$, since $N_i(t) = 0$. Define a connection

$$\begin{aligned} \partial : \mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R}) &\longrightarrow \mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R}) \otimes_{R[\frac{1}{p}]} \Omega_R^1[\frac{1}{p}] \\ x &\longmapsto \sum_{i=1}^d N_i(x) \otimes d \log X_i. \end{aligned}$$

The connection ∂ is G_R -equivariant and satisfies Griffiths transversality with respect to the filtration, i.e. $\partial(\mathrm{Fil}^r \mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R})) \subset \mathrm{Fil}^{r-1} \mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R}) \otimes_{R[\frac{1}{p}]} \Omega_R^1[\frac{1}{p}]$. Its restriction to $R[\frac{1}{p}]$

is the canonical differential operator. Moreover, we have $(\mathcal{O}\mathbf{B}_{\mathrm{dR}}^+(\overline{R}))^{\partial=0} = \mathbf{B}_{\mathrm{dR}}^+(\overline{R})$ and $(\mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R}))^{\partial=0} = \mathbf{B}_{\mathrm{dR}}(\overline{R})$ (see [Bri08, §5.3] for details).

2.2. The crystalline period ring. In this section, we will recall the definition and properties of crystalline period rings following [Bri08]. Note that Brinon defines these rings under a certain assumption on his base ring (see condition (BR) on [Bri08, p. 9]) which is always true in our setting.

2.2.1. Definition of $\mathcal{OB}_{\text{cris}}(\bar{R})$. Let us consider the map $\theta : \mathbf{A}_{\text{inf}}(\bar{R}) \rightarrow \mathbb{C}^+(\bar{R})$ whose kernel is a principal ideal generated by ξ or $p - [p^b]$. Let us denote $x^{[k]} := \frac{x^k}{k!}$ for $x \in \text{Ker } \theta \subset \mathbf{A}_{\text{inf}}(\bar{R})$ and $k \in \mathbb{N}$. The divided power envelope of $\mathbf{A}_{\text{inf}}(\bar{R})$ with respect to $\text{Ker } \theta$ is given as $\mathbf{A}_{\text{inf}}(\bar{R})[x^{[k]}, x \in \text{Ker } \theta]_{k \in \mathbb{N}} = \mathbf{A}_{\text{inf}}(\bar{R})[\xi^{[k]}]_{k \in \mathbb{N}}$. We define

$$\mathbf{A}_{\text{cris}}(\bar{R}) := p\text{-adic completion of } \mathbf{A}_{\text{inf}}(\bar{R})[\xi^{[k]}]_{k \in \mathbb{N}}.$$

This is a $W(\kappa)$ -algebra equipped with a continuous action of G_R . The ring $\mathbf{A}_{\text{cris}}(\bar{R})$ is p -torsion free and the Frobenius on $\mathbf{A}_{\text{inf}}(\bar{R})$ extends to $\mathbf{A}_{\text{cris}}(\bar{R})$. The homomorphism θ in §2.1.2 extends to a surjective homomorphism $\theta : \mathbf{A}_{\text{cris}}(\bar{R}) \rightarrow \mathbb{C}^+(\bar{R})$. Also, we have $t = \log(1 + \pi) \in \text{Ker } \theta \subset \mathbf{A}_{\text{cris}}(\bar{R})$ and the Frobenius φ on this element is given as $\varphi(t) = pt$. Moreover, $\text{Ker } \theta \subset \mathbf{A}_{\text{cris}}(\bar{R})$ is a divided power ideal. Further, the ring $\mathbf{A}_{\text{cris}}(\bar{R})$ is t -torsion free, so we set $\varphi(\frac{1}{t}) = \frac{1}{pt}$ and define $\mathbf{B}_{\text{cris}}^+(\bar{R}) := \mathbf{A}_{\text{cris}}(\bar{R})[\frac{1}{p}]$ and $\mathbf{B}_{\text{cris}}(\bar{R}) := \mathbf{B}_{\text{cris}}^+(\bar{R})[\frac{1}{t}]$. These are F -algebras, equipped with a continuous action of G_R and the Frobenius φ (see [Bri08, §6.1 and §6.2] for details).

Next, let us consider the map $\theta_R : R \otimes_{\mathbb{Z}} \mathbf{A}_{\text{inf}}(\bar{R}) \rightarrow \mathbb{C}^+(\bar{R})$. The kernel of this map is an ideal generated by $\{1 \otimes \xi, z_1, \dots, z_d\}$, where $z_i = X_i \otimes 1 - 1 \otimes [X_i^p]$ for $1 \leq i \leq d$. The divided power envelope of $R \otimes_{\mathbb{Z}} \mathbf{A}_{\text{inf}}(\bar{R})$ with respect to $\text{Ker } \theta_R$ is given as $R \otimes_{\mathbb{Z}} \mathbf{A}_{\text{inf}}(\bar{R})[x^{[k]}, x \in \text{Ker } \theta_R]_{k \in \mathbb{N}}$. We define

$$\mathcal{OA}_{\text{cris}}(\bar{R}) := p\text{-adic completion of } R \otimes_{\mathbb{Z}} \mathbf{A}_{\text{inf}}(\bar{R})[x^{[k]}, x \in \text{Ker } \theta_R]_{k \in \mathbb{N}}.$$

This is an R -algebra equipped with a continuous action of G_R . Taking the diagonal action of the Frobenius on $R \otimes_{\mathbb{Z}} \mathbf{A}_{\text{inf}}(\bar{R})$ it can be shown that the Frobenius extends to $\mathcal{OA}_{\text{cris}}(\bar{R})$ and we denote this extension again by φ . The homomorphism θ_R from §2.1 extends to surjective homomorphism $\theta_R : \mathcal{OA}_{\text{cris}}(\bar{R}) \rightarrow \mathbb{C}^+(\bar{R})$ (see [Bri08, p. 64] for details).

2.2.2. Structure and properties of $\mathcal{OB}_{\text{cris}}(\bar{R})$. Let $T = (T_1, \dots, T_d)$ be some indeterminates as in §2.1.3 and let us denote by $\mathbf{A}_{\text{cris}}(\bar{R})\langle T \rangle^\wedge$ the p -adic completion of the divided power polynomial algebra in indeterminates T and coefficients in $\mathbf{A}_{\text{cris}}(\bar{R})$. Then we obtain an isomorphism of $\mathbf{A}_{\text{cris}}(\bar{R})$ -algebras (see [Bri08, Proposition 6.1.5])

$$f_{\text{cris}} : \mathbf{A}_{\text{cris}}(\bar{R})\langle T \rangle^\wedge \longrightarrow \mathcal{OA}_{\text{cris}}(\bar{R}) \quad (2.1)$$

$$T_i \longmapsto z_i \quad \text{for } 1 \leq i \leq d. \quad (2.2)$$

The ring $\mathcal{OA}_{\text{cris}}(\bar{R})$ is p -torsion free as well as t -torsion free, so we set $\mathcal{OB}_{\text{cris}}^+(\bar{R}) := \mathcal{OA}_{\text{cris}}(\bar{R})[\frac{1}{p}]$ and $\mathcal{OB}_{\text{cris}}(\bar{R}) := \mathcal{OB}_{\text{cris}}^+(\bar{R})[\frac{1}{t}]$. These $R[\frac{1}{p}]$ -algebras are equipped with a continuous action of G_R and the action of Frobenius extends to these rings and we denote this extension again by φ (see [Bri08, §6.1 and §6.2] for details).

Note that there exist natural morphisms of rings $\mathbf{A}_{\text{cris}}(\bar{R}) \rightarrow \mathbf{B}_{\text{dR}}^+(\bar{R})$ and $\mathcal{OA}_{\text{cris}}(\bar{R}) \rightarrow \mathcal{OB}_{\text{dR}}^+(\bar{R})$. So we obtain induced homomorphisms $\mathbf{B}_{\text{cris}}^+(\bar{R}) \rightarrow \mathbf{B}_{\text{dR}}^+(\bar{R})$, $\mathcal{OB}_{\text{cris}}^+(\bar{R}) \rightarrow \mathcal{OB}_{\text{dR}}^+(\bar{R})$, $\mathbf{B}_{\text{cris}}(\bar{R}) \rightarrow \mathbf{B}_{\text{dR}}(\bar{R})$ and $\mathcal{OB}_{\text{cris}}(\bar{R}) \rightarrow \mathcal{OB}_{\text{dR}}(\bar{R})$, which are injective and G_R -equivariant. Using this, we get induced filtrations on crystalline period rings as $\text{Fil}^r \mathbf{B}_{\text{cris}}(\bar{R}) := \mathbf{B}_{\text{cris}}(\bar{R}) \cap \text{Fil}^r \mathbf{B}_{\text{dR}}(\bar{R})$ and $\text{Fil}^r \mathcal{OB}_{\text{cris}}(\bar{R}) := \mathcal{OB}_{\text{cris}}(\bar{R}) \cap \text{Fil}^r \mathcal{OB}_{\text{dR}}(\bar{R})$ for $r \in \mathbb{Z}$ (see [Bri08, §6.2] for details).

Next, we will consider a connection on $\mathcal{OB}_{\text{cris}}(\bar{R})$ induced from the connection on $\mathcal{OB}_{\text{dR}}(\bar{R})$. For $n \in \mathbb{N}$, we have $\partial(z_i^{[n]}) = z_i^{[n-1]} \otimes dX_i$ for $1 \leq i \leq d$, so we get that for any $x \in \mathcal{OA}_{\text{cris}}(\bar{R}) = \mathbf{A}_{\text{cris}}(\bar{R})\langle T \rangle^\wedge$, we have $\partial(x) \in \mathcal{OA}_{\text{cris}}(\bar{R}) \otimes_R \Omega_R^1$. This gives us an induced connection

$$\partial : \mathcal{OB}_{\text{cris}}(\bar{R}) \longrightarrow \mathcal{OB}_{\text{cris}}(\bar{R}) \otimes_{R[\frac{1}{p}]} \Omega_R^1[\frac{1}{p}].$$

The connection ∂ is G_R -equivariant and satisfies Griffiths transversality with respect to the filtration, since the same is true over $\mathcal{OB}_{\text{dR}}(\bar{R})$. Its restriction to $R[\frac{1}{p}]$ is the canonical differential operator. Moreover, taking horizontal sections we get $(\mathcal{OA}_{\text{cris}}^+(\bar{R}))^{\partial=0} = \mathbf{A}_{\text{cris}}(\bar{R})$,

$(\mathcal{O}\mathbf{B}_{\text{cris}}^+(\overline{R}))^{\partial=0} = \mathbf{B}_{\text{cris}}^+(\overline{R})$ and $(\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}))^{\partial=0} = \mathbf{B}_{\text{cris}}(\overline{R})$. We equip $\Omega_R^1[\frac{1}{p}]$ with the unique Frobenius-linear map φ satisfying $\varphi(dx) = d\varphi(x)$ for $x \in R$. Then, over $\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ the Frobenius operator commutes with the connection, i.e. $\varphi\partial = \partial\varphi$ (see [Bri08, Proposition 6.2.5]). Furthermore, we have $(\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}))^{G_R} = R[\frac{1}{p}]$. Finally, the natural map $R[\frac{1}{p}] \rightarrow \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ is faithfully flat (see [Bri08, §6.2 and §6.3] for details).

2.3. p -adic representations. In this section we will recall results on linear algebra data associated to p -adic de Rham and crystalline representations of the Galois group G_R . We will use the G_R -regularity of a topological \mathbb{Q}_p -algebra B in the sense of [Bri08, p. 106]. If V is a p -adic representation of G_R , we set

$$\mathbf{D}_B(V) := (B \otimes_{\mathbb{Q}_p} V)^{G_R}.$$

This is a B^{G_R} -module and we have a natural morphism of B -modules, functorial in V

$$\begin{aligned} \alpha_B(V) : B \otimes_{B^{G_R}} \mathbf{D}_B(V) &\longrightarrow B \otimes_{\mathbb{Q}_p} V \\ b \otimes d &\longmapsto bd. \end{aligned}$$

The representation V is said to be *B -admissible* if α_B is an isomorphism.

2.3.1. Unramified representations. Let R^{ur} denote the union of finite étale R -subalgebras $S \subset \overline{R}$, and let \widehat{R}^{ur} denote its p -adic completion. It is an R -subalgebra of $\mathbb{C}(\overline{R})$ equipped with a continuous action of G_R . Further, we have $(\widehat{R}^{\text{ur}}[\frac{1}{p}])^{G_R} = R[\frac{1}{p}]$ and $\widehat{R}^{\text{ur}}[\frac{1}{p}]$ is G_R -regular. Let us set $G_R^{\text{ur}} := \text{Gal}(R^{\text{ur}}/R)$ which is a quotient of G_R . A p -adic representation $\rho : G_R \rightarrow \text{GL}(V)$ is said to be *unramified*, if ρ factorizes through $G_R \rightarrow G_R^{\text{ur}}$.

Let V be a p -adic representation of G_R and we set

$$\mathbf{D}_{\text{ur}}(V) := (\widehat{R}^{\text{ur}}[\frac{1}{p}] \otimes_{\mathbb{Q}_p} V)^{G_R},$$

which is an $R[\frac{1}{p}]$ -module and we say that V is unramified if and only if V is $\widehat{R}^{\text{ur}}[\frac{1}{p}]$ -admissible (see [Bri08, §8.1]).

Remark 2.1. Let V be an h -dimensional p -adic representation of G_R and $T \subset V$ a \mathbb{Z}_p -lattice stable under the action of G_R such that the action is trivial modulo p . Consider the associated continuous cocycle $f : G_R^{\text{ur}} \rightarrow \text{GL}_h(\widehat{R}^{\text{ur}})$ describing the action of G_R^{ur} over $\widehat{R}^{\text{ur}} \otimes_{\mathbb{Z}_p} T$. Since V is unramified, f is cohomologous to the trivial cocycle and from [Bri08, proof of Proposition 8.1.2], there exists $b \in 1 + p \cdot \text{Mat}(h, \widehat{R}^{\text{ur}})$ such that f is given as $g \mapsto f(g) = g(b)b^{-1}$ for $g \in G_R$. In this case, we say that f is *trivialised* by $b \in 1 + p \cdot \text{Mat}(h, \widehat{R}^{\text{ur}})$.

2.3.2. de Rham representations. Note that $\mathcal{O}\mathbf{B}_{\text{dR}}(\overline{R})$ is a G_R -regular $R[\frac{1}{p}]$ -algebra. We set

$$\mathcal{O}\mathbf{D}_{\text{dR}}(V) := (\mathcal{O}\mathbf{B}_{\text{dR}}(\overline{R}) \otimes_{\mathbb{Q}_p} V)^{G_R}.$$

The representation V is said to be de Rham if it is $\mathcal{O}\mathbf{B}_{\text{dR}}(\overline{R})$ -admissible. The $R[\frac{1}{p}]$ -module $\mathcal{O}\mathbf{D}_{\text{dR}}(V)$ is equipped with a decreasing, separated and exhaustive filtration induced from the filtration on $\mathcal{O}\mathbf{B}_{\text{dR}}(\overline{R}) \otimes_{\mathbb{Q}_p} V$ where we consider the G_R -stable filtration on $\mathcal{O}\mathbf{B}_{\text{dR}}(\overline{R})$ from §2.1.3. Moreover, the module $\mathcal{O}\mathbf{D}_{\text{dR}}(V)$ is equipped with an integrable connection, induced from the G_R -equivariant integrable connection

$$\begin{aligned} \partial : V \otimes_{\mathbb{Q}_p} \mathcal{O}\mathbf{B}_{\text{dR}}(\overline{R}) &\longrightarrow V \otimes_{\mathbb{Q}_p} \mathcal{O}\mathbf{B}_{\text{dR}}(\overline{R}) \otimes_{R[\frac{1}{p}]} \Omega_R^1[\frac{1}{p}] \\ v \otimes b &\longmapsto v \otimes \partial(b). \end{aligned}$$

We denote the induced connection on $\mathcal{O}\mathbf{D}_{\mathrm{dR}}(V)$ again by ∂ . Since the connection ∂ on $\mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R})$ satisfies Griffiths transversality, the same is true for $\mathcal{O}\mathbf{D}_{\mathrm{dR}}(V)$, i.e. $\partial(\mathrm{Fil}^r \mathcal{O}\mathbf{D}_{\mathrm{dR}}(V)) \subset \mathrm{Fil}^{r-1} \mathcal{O}\mathbf{D}_{\mathrm{dR}}(V) \otimes_{R[\frac{1}{p}]} \Omega_R^1[\frac{1}{p}]$. Further, $\mathcal{O}\mathbf{D}_{\mathrm{dR}}(V)$ is projective of rank $\leq \dim(V)$ over $(\mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R}))^{G_R} = R[\frac{1}{p}]$. If V is de Rham then for all $r \in \mathbb{Z}$, the $R[\frac{1}{p}]$ -modules $\mathrm{Fil}^r \mathcal{O}\mathbf{D}_{\mathrm{dR}}(V)$ and $\mathrm{gr}^r \mathcal{O}\mathbf{D}_{\mathrm{dR}}(V)$ are projective of finite type and for such a representation the collection of integers r_i for $1 \leq i \leq \dim_{\mathbb{Q}_p}(V)$ such that $\mathrm{gr}^{-r_i} \mathcal{O}\mathbf{D}_{\mathrm{dR}}(V) \neq 0$ are called *Hodge-Tate weights* of V . Moreover, we say that V is positive if and only if $r_i \leq 0$ for all $1 \leq i \leq \dim_{\mathbb{Q}_p}(V)$ (see [Bri08, §8.3] for details).

2.3.3. Crystalline representations. Note that $\mathcal{O}\mathbf{B}_{\mathrm{cris}}(\overline{R})$ is a G_R -regular $R[\frac{1}{p}]$ -algebra. We set

$$\mathcal{O}\mathbf{D}_{\mathrm{cris}}(V) := (\mathcal{O}\mathbf{B}_{\mathrm{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} V)^{G_R}.$$

The representation V is said to be crystalline if it is $\mathcal{O}\mathbf{B}_{\mathrm{cris}}(\overline{R})$ -admissible and we denote the category of all crystalline representations of G_R by $\mathrm{Rep}_{\mathbb{Q}_p}^{\mathrm{cris}}(G_R)$. The $R[\frac{1}{p}]$ -module $\mathcal{O}\mathbf{D}_{\mathrm{cris}}(V)$ is equipped with a Frobenius-semilinear operator φ induced from the Frobenius on $\mathcal{O}\mathbf{B}_{\mathrm{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} V$, where we consider the G_R -equivariant Frobenius on $\mathcal{O}\mathbf{B}_{\mathrm{cris}}(\overline{R})$. Further, $\mathcal{O}\mathbf{D}_{\mathrm{cris}}(V)$ is an $R[\frac{1}{p}]$ -submodule of $\mathcal{O}\mathbf{D}_{\mathrm{dR}}(V)$, and we equip the former with induced filtration and connection which satisfies Griffiths transversality with respect to the filtration. Additionally, we have $\partial\varphi = \varphi\partial$ over $\mathcal{O}\mathbf{D}_{\mathrm{cris}}(V)$ (see [Bri08, §8.3] for details).

The $R[\frac{1}{p}]$ -module $\mathcal{O}\mathbf{D}_{\mathrm{cris}}(V)$ is projective of rank $\leq \dim(V)$. If V is crystalline, then the $R[\frac{1}{p}]$ -linear homomorphism $1 \otimes \varphi : R[\frac{1}{p}] \otimes_{R[\frac{1}{p}], \varphi} \mathcal{O}\mathbf{D}_{\mathrm{cris}}(V) \rightarrow \mathcal{O}\mathbf{D}_{\mathrm{cris}}(V)$ is an isomorphism and $\mathcal{O}\mathbf{D}_{\mathrm{cris}}(V)$ is called a filtered (φ, ∂) -module. The inclusion $\mathcal{O}\mathbf{B}_{\mathrm{cris}}(\overline{R}) \hookrightarrow \mathcal{O}\mathbf{B}_{\mathrm{dR}}(\overline{R})$ induces the inclusion $\mathcal{O}\mathbf{D}_{\mathrm{cris}}(V) \hookrightarrow \mathcal{O}\mathbf{D}_{\mathrm{dR}}(V)$. Let V be a non-trivial de Rham representation of G_R , then the inclusion $\mathcal{O}\mathbf{D}_{\mathrm{cris}}(V) \hookrightarrow \mathcal{O}\mathbf{D}_{\mathrm{dR}}(V) \neq 0$ is surjective if and only if V is crystalline (see [Bri08, §8.2 and §8.3] for details).

In conclusion, we have a functor

$$\mathcal{O}\mathbf{D}_{\mathrm{cris}} : \mathrm{Rep}_{\mathbb{Q}_p}^{\mathcal{O}\mathrm{cris}}(G_R) \longrightarrow \text{filtered } (\varphi, \partial)\text{-modules over } R[\frac{1}{p}].$$

Objects in the essential image are called *admissible* filtered (φ, ∂) -modules and the functor induces an equivalence of categories with the essential image (see [Bri08, Théorèmes 8.4.2, 8.5.1]).

Remark 2.2. In the arithmetic case, the essential image of $\mathbf{D}_{\mathrm{cris}}$, i.e. admissible filtered φ -modules can be described more explicitly. In particular, using certain invariants attached to filtered φ -modules one considers the full subcategory of *weakly admissible* filtered φ -modules and it is a result of Colmez and Fontaine that weakly admissible filtered φ -modules are admissible (in the sense above, see [CF00, Théorème A]). In the relative case, Brinon gave a definition of weakly admissible filtered (φ, ∂) -modules (see [Bri08, p. 136]). However, the notion is not completely satisfactory as one does not obtain an equivalence between admissible and weakly admissible filtered (φ, ∂) -modules (see [Moo18, Theorem 1.3]).

2.3.4. One dimensional de Rham and crystalline representations. In the 1-dimensional case, it is possible to classify all crystalline representations:

Proposition 2.3 ([Bri08, Propositions 8.4.1, 8.6.1]). *Let $\eta : G_R \rightarrow \mathbb{Z}_p^\times$ be a continuous character.*

- (i) *η is de Rham if and only if we can write $\eta = \eta_f \eta_{\mathrm{ur}} \chi^n$ where η_f is a finite character, η_{ur} is an unramified character taking values in $1 + p\mathbb{Z}_p$ (therefore trivialized $\alpha \in 1 + pR^{\mathrm{ur}}$, see Remark 2.1) and χ is the p -adic cyclotomic character and $n \in \mathbb{Z}$.*

- (ii) η is crystalline if and only if we can write $\eta = \eta_{\text{f}} \eta_{\text{ur}} \chi^n$ where η_{f} is a finite unramified character, η_{ur} is an unramified character taking values in $1 + p\mathbb{Z}_p$ (therefore trivialized by some $\alpha \in 1 + p\widehat{R^{\text{ur}}}$, see Remark 2.1) and χ is the p -adic cyclotomic character and $n \in \mathbb{Z}$.

In particular, a 1-dimensional de Rham representation is potentially crystalline.

- (iii) Let $V = \mathbb{Q}_p(\eta)$ be a one-dimensional crystalline representation. Then there exists a finite étale extension $R \rightarrow R'$ such that the $R'[\frac{1}{p}]$ -module $R'[\frac{1}{p}] \otimes_{R[\frac{1}{p}]} \mathcal{O}\mathbf{D}_{\text{cris}}(V)$ is free. In particular, if η_{f} is trivial then $\mathcal{O}\mathbf{D}_{\text{cris}}(V)$ is a free $R[\frac{1}{p}]$ -module of rank 1.

3. (φ, Γ) -modules and crystalline coordinates

We will keep the setting and notations of §2. In particular, we have that F is a finite unramified extension of \mathbb{Q}_p and $K = F(\mu_{p^m})$ for a fixed $m \in \mathbb{N}_{\geq 1}$ (fix $m \in \mathbb{N}_{\geq 2}$ if $p = 2$). Recall that R is étale over $O_F\{X, X^{-1}\}$ and we have multivariate polynomials $Q_i(Z_1, \dots, Z_s) \in O_F\{X, X^{-1}\}[Z_1, \dots, Z_s]$ for $1 \leq i \leq s$ such that $\det(\frac{\partial Q_i}{\partial Z_j})$ is invertible in R . In particular, the ring $O_F\{X, X^{-1}\}$ provides a system of coordinates for R .

3.1. (φ, Γ) -modules. In this section, we briefly recall the theory of relative (φ, Γ) -modules from [And06; AB08; AI08].

Let $F_n = F(\mu_{p^n})$ for $n \in \mathbb{N}$ and $F_\infty = \cup_n F_n$. We take R_n to be the integral closure of $R \otimes_{O_F[X^{\pm 1}]} O_{F_n}[X_1^{p^{-n}}, \dots, X_d^{p^{-n}}]$ inside $\overline{R}[\frac{1}{p}]$ and set $R_\infty := \cup_{n \geq m} R_n$ noting that $F_\infty \subset R_\infty[\frac{1}{p}]$. From §2.1.2 recall that $\mathbb{C}(\overline{R}) = \mathbb{C}^+(\overline{R})[\frac{1}{p}]$ and $\mathbb{C}(\overline{R})^b$ denotes its tilt. The ring $\mathbb{C}(\overline{R})^b$ is perfect of characteristic p and we set $\mathbf{A}_{\overline{R}} := W(\mathbb{C}(\overline{R})^b)$, the ring of p -typical Witt vectors with coefficients in $\mathbb{C}(\overline{R})^b$ and endowed with the weak topology (see [AI08, §2.10]). The absolute Frobenius over $\mathbb{C}(\overline{R})^b$ lifts to an endomorphism $\varphi : \mathbf{A}_{\overline{R}} \rightarrow \mathbf{A}_{\overline{R}}$, which we again call the Frobenius. The action of G_R on $\mathbb{C}(\overline{R})^b$ extends to a continuous action on $\mathbf{A}_{\overline{R}}$ commuting with the Frobenius. The inclusion $\overline{F} \subset \overline{R}[\frac{1}{p}]$ induces inclusions $\mathbb{C}_p^b \subset \mathbb{C}(\overline{R})^b$ and $\mathbf{A}_{\overline{F}} \subset \mathbf{A}_{\overline{R}}$. Recall that we set $\mathbf{A}_{\text{inf}}(\overline{R}) := W(\mathbb{C}^+(\overline{R})^b)$. The inclusion $O_{\overline{F}} \subset \overline{R}$ induces inclusions $O_{\mathbb{C}_p}^b \subset \mathbb{C}^+(\overline{R})^b$ and $\mathbf{A}_{\text{inf}}(O_{\overline{F}}) \subset \mathbf{A}_{\text{inf}}(\overline{R})$.

3.1.1. The group Γ_R . The ring $R_\infty[\frac{1}{p}]$ is a Galois extension of $R[\frac{1}{p}]$ with Galois group $\Gamma_R := \text{Gal}(R_\infty[\frac{1}{p}]/R[\frac{1}{p}])$ isomorphic to the semidirect product of Γ_F and Γ'_R , where $\Gamma_F = \text{Gal}(F_\infty/F)$ and $\Gamma'_R = \text{Gal}(R_\infty[\frac{1}{p}]/F_\infty R[\frac{1}{p}])$. In particular, we have an exact sequence

$$1 \longrightarrow \Gamma'_R \longrightarrow \Gamma_R \longrightarrow \Gamma_F \longrightarrow 1, \quad (3.1)$$

where (see [Bri08, p. 9] and [And06, §2.4])

$$\begin{aligned} \Gamma'_R &= \text{Gal}(R_\infty[\frac{1}{p}]/F_\infty R[\frac{1}{p}]) \xrightarrow{\sim} \mathbb{Z}_p^d, \\ \chi : \Gamma_F &= \text{Gal}(F_\infty/F) \xrightarrow{\sim} \mathbb{Z}_p^\times. \end{aligned}$$

The group Γ_F can be viewed as a subgroup of Γ_R , i.e. we can take a section of the projection map in (3.1) such that for $\gamma \in \Gamma_F$ and $g \in \Gamma'_R$, we have $\gamma g \gamma^{-1} = g^{\chi(\gamma)}$. So we can choose topological generators $\{\gamma, \gamma_1, \dots, \gamma_d\}$ of Γ_R such that

$$\begin{aligned} \gamma(\varepsilon) &= \varepsilon^{\chi(\gamma)}, \quad \gamma_i(\varepsilon) = \varepsilon && \text{for } 1 \leq i \leq d, \\ \gamma_i(X_j^b) &= \varepsilon X_i^b, \quad \gamma_i(X_j^b) = X_j^b && \text{for } i \neq j \text{ and } 1 \leq j \leq d, \end{aligned}$$

and that $\gamma_0 = \gamma^e$ with $\chi(\gamma_0) = \exp(p^m)$, is a topological generator of $\Gamma_K = \text{Gal}(K_\infty/K)$, where $K_\infty = F_\infty$ and $e = [K : F]$. It follows that $\{\gamma_1, \dots, \gamma_d\}$ are topological generators of Γ'_R , γ is a lift of a topological generator of Γ_F , and γ_0 is a topological generator of Γ_K . In particular,

$$\chi : \Gamma_K = \text{Gal}(F_\infty/K) \xrightarrow{\sim} 1 + p^m \mathbb{Z}_p.$$

3.1.2. Setup. In [FW79b; FW79a; Win83], using the field-of-norms functor, Fontaine and Wintenberger constructed a non-archimedean complete discrete valuation field $\mathbf{E}_K \subset \widehat{K}_\infty^b$ of characteristic p with residue field κ and admitting a continuous action of Γ_K (notation is a bit unfortunate as \mathbf{E}_K depends only on K_∞). Utilizing the isomorphism of Galois groups

$\text{Gal}(\overline{F}/K_\infty) \xrightarrow{\sim} \text{Gal}(\mathbf{E}_K^{\text{sep}}/\mathbf{E}_K)$ (also see tilting correspondence in [Sch12] for a modern treatment), Fontaine classified mod- p representations of G_K in terms of étale (φ, Γ_K) -modules over \mathbf{E}_K . By some technical considerations one can then lift this to the classification of \mathbb{Z}_p -representations of G_F in terms of étale (φ, Γ_K) -modules over a certain two dimensional regular local ring $\mathbf{A}_K \subset W(\widehat{K}_\infty^\flat)$ (see [Fon90] for details).

We have an analogous theory in the relative setting, to describe which we need to consider generically étale algebras over finite extensions of R in the cyclotomic tower R_∞/R . More precisely, let $S \subset \overline{R}$ be a finite R_n -algebra with $S[\frac{1}{p}]$ étale over $R_n[\frac{1}{p}]$. For $k \geq n$ denote by S_k the integral closure of $S \otimes_{R_n} R_k$ in $\overline{R}[\frac{1}{p}]$ and set $S_\infty := \cup_{k \geq n} S_k$. We have that S_∞ is a normal R_∞ -algebra and an integral domain as a subring of \overline{R} . As in the case of R , for S we define $G_S := \text{Gal}(\overline{R}[\frac{1}{p}]/S[\frac{1}{p}])$, $\Gamma_S := \text{Gal}(S_\infty[\frac{1}{p}]/S[\frac{1}{p}])$ and $H_S := \text{Ker}(G_S \rightarrow \Gamma_S)$. Again, Γ_S is isomorphic to the semidirect product of Γ_{F_n} and Γ'_S , where $\Gamma'_S = \text{Gal}(S_\infty[\frac{1}{p}]/F_\infty S[\frac{1}{p}])$ is a finite index subgroup of $\Gamma'_R \xrightarrow{\sim} \mathbb{Z}_p^d$.

3.1.3. Rings in characteristic p . In the relative setting, Andreatta in [And06] constructed an analogue of the subfield $\mathbf{E}_K \subset \widehat{K}_\infty^\flat$, i.e. to any S as above, he associated a ring $\mathbf{E}_S \subset \text{Fr } \widehat{S}_\infty^\flat$ functorial in S_∞ . Let us recall his definition: Let \mathbf{E}_F^+ denote the valuation ring of \mathbf{E}_F and we have $\pi \in W(\widehat{F}_\infty^\flat)$ such that its reduction modulo p , denoted as $\bar{\pi} = \varepsilon - 1$, is a uniformizer of \mathbf{E}_F^+ . Depending on S , let $\delta \in \mathbb{Q} \cap [0, 1]$ small enough and $N \in \mathbb{N}$ large enough (see [And06, Definition 4.2] for precise formulations of δ and N), and define the ring

$$\mathbf{E}_S^+ := \{(a_0, \dots, a_k, \dots) \in \widehat{S}_\infty^\flat, \text{ such that } a_k \in S_k/p^\delta S_k \text{ for all } k \geq N\}.$$

The ring \mathbf{E}_S^+ is finite and torsion free as an \mathbf{E}_R^+ -module. It is a reduced Noetherian ring which is $\bar{\pi}$ -adically complete. By construction, it is endowed with a $\bar{\pi}$ -adically continuous action of Γ_S and a Frobenius endomorphism φ , commuting with each other and compatible with respective structures on \widehat{S}_∞^\flat . Moreover, \mathbf{E}_S^+ is a normal extension of \mathbf{E}_R^+ , étale after inverting $\bar{\pi}$ and of degree equal to the generic degree of $R_m \subset S$. Further, the set of elements $\{\bar{\pi}, X_1^\flat, \dots, X_d^\flat\}$ form an absolute p -basis of \mathbf{E}_R^+ (see [And06, Proposition 4.5, Corollaries 5.3 & 5.4]). The ring \widehat{S}_∞^\flat coincides with the $\bar{\pi}$ -adic completion of the perfect closure of \mathbf{E}_S^+ and the extension $\mathbf{E}_S^+ \rightarrow \widehat{S}_\infty^\flat$ is faithfully flat. Finally, set $\mathbf{E}_S := \mathbf{E}_S^+[\frac{1}{\bar{\pi}}]$.

Definition 3.1. Define $\mathbf{E}^+ := \cup_S \mathbf{E}_S^+$, where the union runs over R_n -subalgebras $S \subset \overline{R}$ for some $n \in \mathbb{N}$ such that S is normal and finite as an R_n -module and $S[\frac{1}{p}]$ is étale over $R_n[\frac{1}{p}]$. Also, we set $\mathbf{E} := \mathbf{E}^+[\frac{1}{\bar{\pi}}]$. These rings are $\bar{\pi}$ -adically complete and equipped with a Frobenius and a continuous G_R -action.

Remark 3.2. From [AI08, Proposition 2.9], we have $(\mathbb{C}^+(\overline{R}))^{H_R} = \widehat{R}_\infty$, $(\mathbb{C}^+(\overline{R})^\flat)^{H_R} = \widehat{R}_\infty^\flat$, $(\mathbb{C}(\overline{R})^\flat)^{H_R} = \widehat{R}_\infty^\flat[\frac{1}{\bar{\pi}}]$, $(\mathbf{E}^+)^{H_R} = \mathbf{E}_R^+$ and $\mathbf{E}^{H_R} = \mathbf{E}_R$.

Remark 3.3. We will describe $\mathbb{C}^+(\overline{R})^\flat$ as the ring of power-bounded elements inside $\mathbb{C}(\overline{R})^\flat$ (for the spectral norm). Recall that \overline{R} is the union of finite R -subalgebras $S \subset \text{Fr}(\overline{R})$ such that $S[\frac{1}{p}]$ is étale over $R[\frac{1}{p}]$. Since \overline{R} is an integral domain and p -adically separated, i.e. $\cap_{k \in \mathbb{N}} p^k \overline{R} = 0$, we obtain that the filtration by powers of the ideal $p\overline{R} \subset \overline{R}$ induces a sub-multiplicative norm (see [BGR84, §1.3.3, Proposition 1]) which extends to $\overline{R}[\frac{1}{p}]$. A further “smoothing” of the aforementioned norm yields a power-multiplicative norm on $\overline{R}[\frac{1}{p}]$ (see [BGR84, §1.3.2]) which we call the *spectral norm* on $\overline{R}[\frac{1}{p}]$. Let C denote the completion of $\overline{R}[\frac{1}{p}]$ for the spectral norm and C° its power-bounded elements.

Next, one can show that under the spectral norm the power-bounded elements (or equivalently, the closed unit ball) of $\overline{R}[\frac{1}{p}]$ written as $(\overline{R}[\frac{1}{p}])^\circ$ is exactly \overline{R} . Indeed, we have the obvious inclusion $\overline{R} \subset (\overline{R}[\frac{1}{p}])^\circ$ and for the converse taking $x \in (\overline{R}[\frac{1}{p}])^\circ$, one can reduce the claim to

a finite R -subalgebra $S \subset \overline{R}$ integrally closed in $\overline{R}[\frac{1}{p}]$ and such that $x \in S[\frac{1}{p}]$. Then it easily follows that $S = (S[\frac{1}{p}])^\circ = S[\frac{1}{p}] \cap (\overline{R}[\frac{1}{p}])^\circ \subset \overline{R}[\frac{1}{p}]$. So we obtain that the topology induced by the spectral norm is equivalent to the p -adic topology on $\overline{R}[\frac{1}{p}]$, therefore $C = \mathbb{C}(\overline{R})$ and $C^\circ = \mathbb{C}^+(\overline{R})$ and $(\mathbb{C}(\overline{R}), \mathbb{C}^+(\overline{R}))$ is a uniform adic Banach \mathbb{Q}_p -algebra (see [KL15, Definitions 2.4.1 and 2.8.1]).

Finally, by the perfectoid correspondence of uniform adic Banach algebras in [KL15, Theorem 3.6.5], we obtain that $(\mathbb{C}(\overline{R})^\flat, \mathbb{C}^+(\overline{R})^\flat)$ is a uniform adic Banach \mathbb{F}_p -algebra such that the topology induced by the spectral norm (arising from the sub-multiplicative norm induced by the ideal $p^\flat \mathbb{C}^+(\overline{R})^\flat \subset \mathbb{C}^+(\overline{R})^\flat$) is equivalent to the topology on $(\mathbb{C}(\overline{R})^\flat, \mathbb{C}^+(\overline{R})^\flat)$ described in §2.1.2. Finally, since $\mathbb{C}^+(\overline{R})$ is the ring of power-bounded elements in $\mathbb{C}(\overline{R})$ we obtain that the its tilt $\mathbb{C}^+(\overline{R})^\flat$ is the ring of power-bounded elements in $\mathbb{C}(\overline{R})^\flat$.

Remark 3.4. Let us denote the natural valuation on \mathbb{C}_p^\flat by v^\flat . Then one can show that $v^\flat(\overline{\pi}) = \frac{p}{p-1} > 0$, i.e. $\overline{\pi}$ is not invertible in $O_{\mathbb{C}_p}^\flat$. Since $O_{\mathbb{C}_p}^\flat = \mathbb{C}_p^\flat \cap \mathbb{C}^+(\overline{R})^\flat \subset \mathbb{C}(\overline{R})^\flat$, we obtain that $\overline{\pi}$ is not invertible in $\mathbb{C}^+(\overline{R})^\flat$. Moreover, as $\mathbb{C}^+(\overline{R})^\flat$ is the ring of power-bounded elements in $\mathbb{C}^+(\overline{R})^\flat$ (see Remark 3.3) we conclude that $\mathbf{E}^+ = \mathbf{E} \cap \mathbb{C}^+(\overline{R})^\flat \subset \mathbb{C}(\overline{R})^\flat$.

3.1.4. Rings in characteristic 0. We have liftings of the rings discussed above to characteristic 0. In other words, there exists a Noetherian regular domain $\mathbf{A}_R \subset W(\widehat{R}_\infty^\flat[\frac{1}{\pi}])$, complete for the weak topology and endowed with a continuous action of Γ_R and a Frobenius such that $\mathbf{A}_R/p\mathbf{A}_R = \mathbf{E}_R$. Moreover, \mathbf{A}_R contains a subring \mathbf{A}_R^+ lifting \mathbf{E}_R^+ complete for the weak topology with $\pi, [X_1^\flat], \dots, [X_d^\flat] \in \mathbf{A}_R^+$ (see [And06, Appendix C]). Furthermore, for S as in Definition 3.1 let \mathbf{A}_S denote the unique finite étale \mathbf{A}_R -algebra lifting the finite étale extension $\mathbf{E}_R \subset \mathbf{E}_S$. It is a Noetherian regular domain, complete for the weak topology and endowed with a continuous action of Γ_S and a Frobenius, lifting the ones defined on \mathbf{E}_S . Moreover, it contains a subring \mathbf{A}_S^+ lifting \mathbf{E}_S^+ so that the former is complete for the weak topology. In characteristic 0, we set $\mathbf{B}_{\overline{R}} := \mathbf{A}_{\overline{R}}[\frac{1}{p}] = \bigcup_{j \in \mathbb{N}} p^{-j} \mathbf{A}_{\overline{R}}$ equipped with the direct limit topology (see [And06, §7] for details).

Definition 3.5. Define $\mathbf{A} :=$ completion of $\bigcup_S \mathbf{A}_S \subset \mathbf{A}_{\overline{R}}$ for the p -adic topology, where the union runs over all R_n -subalgebras $S \subset \overline{R}$ as in Definition 3.1. Equip \mathbf{A} with the weak topology induced by the inclusion $\mathbf{A} \subset \mathbf{A}_{\overline{R}}$. Moreover, we set $\mathbf{A}^+ := \mathbf{A} \cap \mathbf{A}_{\text{inf}}(\overline{R})$, $\mathbf{B}^+ := \mathbf{A}^+[\frac{1}{p}]$ and $\mathbf{B} := \mathbf{A}[\frac{1}{p}]$ equipped with induced weak topology. These rings are stable under φ and admit a continuous G_R -action.

Remark 3.6. In Definition 3.5 one can take the base ring as $R[\varpi]$ instead of R to obtain period rings $\mathbf{A}_{\varpi}^+ \subset \mathbf{A}_{\varpi}$ (instead of $\mathbf{A}^+ \subset \mathbf{A}$). In particular, one has that $\pi_m = \varphi^{-m}(\pi) \in \mathbf{A}_{\varpi}^+$ and it easily follows that $\mathbf{A}^+ \subset \mathbf{A}_{\varpi}^+ \subset \mathbf{A}_{\text{inf}}(\overline{R})$ compatible with Frobenius and G_R -action.

Remark 3.7. (i) It follows from definitions that $p\mathbf{A}^+ = p\mathbf{A} \cap \mathbf{A}_{\text{inf}}(\overline{R}) = \mathbf{A} \cap p\mathbf{A}_{\text{inf}}(\overline{R}) = p(\mathbf{A} \cap \mathbf{A}_{\text{inf}}(\overline{R}))$. Therefore, from Remark 3.4 it easily follows that $\mathbf{A}^+/p\mathbf{A}^+ = \mathbf{E}^+$.

(ii) From [AI08, Lemma 2.11] we have $\mathbf{A}^{H_R} = \mathbf{A}_R$ and $(\mathbf{A}^+)^{H_R} = \mathbf{A}_R^+$.

3.1.5. Some lemmas on matrices. Let us note some results which will be useful in the proof of Proposition 4.11.

Lemma 3.8. *Let $h \in \mathbb{N}$ and matrices $Y \in \text{Mat}(h, \mathbf{E})$ and $X, Z, W \in \text{Mat}(h, \mathbf{E}^+)$ such that $\varphi(Y) = XYZ + W$, then $Y \in \text{Mat}(h, \mathbf{E}^+)$.*

Proof. From Remark 3.4 we have $\mathbf{E}^+ = \mathbf{E} \cap \mathbb{C}^+(\overline{R})^\flat$. So it is enough to show that $Y \in \text{Mat}(h, \mathbb{C}^+(\overline{R})^\flat)$. Recall that we have $\mathbb{C}(\overline{R})^\flat = \mathbb{C}^+(\overline{R})^\flat[\frac{1}{p^\flat}]$. Therefore, for some smallest $k \in \mathbb{N}$, we can write $Y = \frac{1}{(p^\flat)^k} Y_1$ with $Y_1 \in \text{Mat}(h, \mathbb{C}^+(\overline{R})^\flat)$. Now, applying φ we get that $\varphi(\frac{1}{(p^\flat)^k} Y_1) =$

$\frac{1}{(p^b)^k}XY_1Z+W$, which can be rewritten as $\frac{(p_1^b)^k}{(p^b)^k}Y_1 = \varphi^{-1}(XY_1Z + (p^b)^k W)$, where $p_1^b = \varphi^{-1}(p^b)$. In the last equality, note that the expression on the left $\frac{(p_1^b)^k}{(p^b)^k}Y_1 \in \text{Mat}(h, \mathbb{C}(\overline{R})^b)$, whereas the expression on the right $\varphi^{-1}(XY_1Z + (p^b)^k W) \in \text{Mat}(h, \varphi^{-1}(\mathbb{C}^+(\overline{R})^b)) = \text{Mat}(h, \mathbb{C}^+(\overline{R})^b)$ since $\mathbb{C}^+(\overline{R})^b$ is perfect. So we obtain that $\frac{(p_1^b)^k}{(p^b)^k}Y_1 \in \text{Mat}(h, \mathbb{C}^+(\overline{R})^b)$, i.e. $Y = \frac{1}{(p^b)^k}Y_1 \in \text{Mat}(h, \frac{1}{(p_1^b)^k}\mathbb{C}^+(\overline{R})^b)$. Next, we write $Y = \frac{1}{(p_1^b)^k}Y_2$ with $Y_2 \in \text{Mat}(h, \mathbb{C}^+(\overline{R})^b)$. Again, applying φ and arguing as above, one gets $Y \in \text{Mat}(h, \frac{1}{(p_2^b)^k}\mathbb{C}^+(\overline{R})^b)$, where $p_2^b = \varphi^{-2}(p^b)$. Now, it easily follows by induction on $n \in \mathbb{N}$ that $Y \in \text{Mat}(h, \frac{1}{(p_n^b)^k}\mathbb{C}^+(\overline{R})^b)$, where $p_n^b = \varphi^{-n}(p^b)$. Therefore, $Y \in \text{Mat}(h, \cap_{n \in \mathbb{N}} \frac{1}{(p_n^b)^k}\mathbb{C}^+(\overline{R})^b) \subset \text{Mat}(h, \mathbb{C}(\overline{R})^b)$. But since $\mathbb{C}^+(\overline{R})^b$ is the ring of power-bounded elements in $\mathbb{C}(\overline{R})^b$, we obtain that $\cap_{n \in \mathbb{N}} \frac{1}{(p_n^b)^k}\mathbb{C}^+(\overline{R})^b = \mathbb{C}^+(\overline{R})^b$. Hence, we get $Y \in \text{Mat}(h, \mathbb{C}^+(\overline{R})^b)$ as desired. ■

Lemma 3.9. *Let $h \in \mathbb{N}$ and matrices $Y \in \text{Mat}(h, \mathbf{A})$ and $X, Z, W \in \text{Mat}(h, \mathbf{A}^+)$ such that $\varphi(Y) = XYZ + W$, then $Y \in \text{Mat}(h, \mathbf{A}^+)$.*

Proof. Reducing the equation modulo p we have $\varphi(\overline{Y}) = \overline{X}\overline{Y}\overline{Z} + \overline{W}$, with $\overline{Y} \in \text{Mat}(h, \mathbf{E})$ and $\overline{X}, \overline{Z}, \overline{W} \in \text{Mat}(h, \mathbf{E}^+)$. Therefore, from Lemma 3.8 we obtain that $\overline{Y} \in \text{Mat}(h, \mathbf{E}^+)$. As we have $\mathbf{A}^+/p\mathbf{A}^+ = \mathbf{E}^+$ (see Remark 3.7 (ii)), let $V_0 \in \text{Mat}(h, \mathbf{A}^+)$ such that $\overline{Y} = \overline{V}_0$ and $\varphi(\overline{V}_0) = \overline{X}\overline{V}_0\overline{Z} + \overline{W}$. So we can write $Y = V_0 + pY_1$ with $Y_1 \in \text{Mat}(h, \mathbf{A})$, and obtain that $\varphi(V_0 + pY_1) = X(V_0 + pY_1)Z + W$. Simplifying the latter expression, we have $\varphi(V_0) - (XV_0Z + W) = p(XY_1Z - \varphi(Y_1))$. Since $\varphi(V_0) - (XV_0Z + W) \in \text{Mat}(h, p\mathbf{A}^+)$, we conclude that $\varphi(Y_1) - XY_1Z \in \text{Mat}(h, \mathbf{A}^+)$. In other words, we have an equality $\varphi(Y_1) = XY_1Z + W_1$ with $Y_1 \in \text{Mat}(h, \mathbf{A})$ and $X, Z, W_1 \in \text{Mat}(h, \mathbf{A}^+)$. Repeating the argument as above, we get that $\overline{Y}_1 \in \text{Mat}(h, \mathbf{E}^+)$ and we can take a lift to write $Y_1 = V_1 + pY_2$ with $V_1 \in \text{Mat}(h, \mathbf{A}^+)$ and $Y_2 \in \text{Mat}(h, \mathbf{A})$. This gives us that $Y = V_0 + pV_1 + p^2Y_2$. Now, it easily follows by induction on $n \in \mathbb{N}$ that $Y = V_0 + pV_1 + \dots + p^{n-1}V_{n-1} + p^nY_n$ with $V_i \in \text{Mat}(h, \mathbf{A}^+)$ for $0 \leq i \leq n-1$ and $Y_n \in \text{Mat}(h, \mathbf{A})$. Letting $n \rightarrow +\infty$ and noting that \mathbf{A}^+ is p -adically complete, we obtain that $Y \in \text{Mat}(h, \mathbf{A}^+)$ as desired. ■

3.1.6. Étale (φ, Γ_R) -modules.

Definition 3.10. A (φ, Γ_R) -module D over \mathbf{A}_R is a finitely generated module equipped with

- (i) A semilinear action of Γ_R , continuous for the weak topology,
- (ii) A Γ_R -equivariant Frobenius-semilinear endomorphism φ .

We say that D is *étale* if the natural map $1 \otimes \varphi : \mathbf{A}_R \otimes_{\mathbf{A}_R, \varphi} D \rightarrow D$ is an isomorphism of \mathbf{A}_R -modules.

Denote by $(\varphi, \Gamma_R)\text{-Mod}_{\mathbf{A}_R}^{\text{ét}}$ the category of étale (φ, Γ_R) -modules over \mathbf{A}_R with morphisms between objects being continuous, (φ, Γ_R) -equivariant morphisms of \mathbf{A}_R -modules. Next, denote by $\text{Rep}_{\mathbb{Z}_p}(G_R)$ the category of finitely generated \mathbb{Z}_p -modules equipped with a linear and continuous action of G_R , with morphisms between objects being continuous and G_R -equivariant morphisms of \mathbb{Z}_p -modules.

Let T be a \mathbb{Z}_p -representation of G_R . The \mathbf{A}_R -module $\mathbf{D}(T) := (\mathbf{A} \otimes_{\mathbb{Z}_p} T)^{H_R}$ is equipped with a semilinear operator φ and a continuous (for the weak topology) and semilinear action of Γ_R , commuting with each other. Moreover, $\mathbf{D}(T)$ is an étale (φ, Γ_R) -module. Furthermore, if T is free of finite rank, then $\mathbf{D}(T)$ is a projective module of rank $= \text{rk}_{\mathbb{Z}_p} T$ (see [And06, Theorem 7.11]). The functor

$$\mathbf{D} : \text{Rep}_{\mathbb{Z}_p}(G_R) \longrightarrow (\varphi, \Gamma_R)\text{-Mod}_{\mathbf{A}_R}^{\text{ét}}, \quad (3.2)$$

induces an equivalence of categories (see [And06, Theorem 7.11] and [AB08, Théorème 4.35]), and the natural map $\mathbf{A} \otimes_{\mathbf{A}_R} \mathbf{D}(T) \xrightarrow{\sim} \mathbf{A} \otimes_{\mathbb{Z}_p} T$ is an isomorphism of \mathbf{A} -modules compatible with Frobenius and the action of G_R on each side.

3.2. Crystalline coordinates. In this section we will introduce certain “coordinate” rings. As we shall see in the next section, these rings are related to period rings appearing in §2 and §3.1.

Let r_ϖ^+ and r_ϖ denote the algebras $O_F[[X_0]]$ and $O_F[[X_0]]\{X_0^{-1}\}$. Sending X_0 to ϖ induces a surjective homomorphism $r_\varpi^+ \twoheadrightarrow O_K$. Let $R_{\varpi, \square}^+$ denote the completion of $O_F[X_0, X, X^{-1}]$ for the (p, X_0) -adic topology. Sending X_0 to ϖ induces a surjective homomorphism $R_{\varpi, \square}^+ \twoheadrightarrow O_K\{X, X^{-1}\}$, whose kernel is generated by $P = P_\varpi(X_0)$. This provides a closed embedding of $\mathrm{Spf} O_K\{X, X^{-1}\}$ into a formal scheme $\mathrm{Spf} R_{\varpi, \square}^+$, which is smooth over O_F . Recall that R is étale over $O_F\{X, X^{-1}\}$ and we have multivariate polynomials $Q_i(Z_1, \dots, Z_s) \in O_F\{X, X^{-1}\}[Z_1, \dots, Z_s]$ for $1 \leq i \leq s$ such that $\det(\frac{\partial Q_i}{\partial Z_j})$ is invertible in R . So we can set R_ϖ^+ to be the quotient by (Q_1, \dots, Q_s) of the completion of $R_{\varpi, \square}^+[Z_1, \dots, Z_s]$ for (p, X_0) -adic topology. Again, we have that $\det(\frac{\partial Q_i}{\partial Z_j})$ is invertible in R_ϖ^+ (since $R \twoheadrightarrow R_\varpi^+$). Hence, R_ϖ^+ is étale over $R_{\varpi, \square}^+$ and smooth over O_F . Sending X_0 to ϖ induces a surjective homomorphism $R_\varpi^+ \twoheadrightarrow R[\varpi]$ whose kernel is generated by $P = P_\varpi(X_0)$. This can be summarized by the commutative diagram

$$\begin{array}{ccccc}
 \mathrm{Spf} R[\varpi] & \xrightarrow{\quad} & & \xrightarrow{\quad} & \mathrm{Spf} R_\varpi^+ \\
 \downarrow & \searrow & & \swarrow & \downarrow \\
 & & \mathrm{Spf} R & & \\
 & & \downarrow & & \\
 & & \mathrm{Spf} O_F\{X, X^{-1}\} & & \\
 \swarrow & & & & \searrow \\
 \mathrm{Spf} O_K\{X, X^{-1}\} & \xrightarrow{\quad} & & \xrightarrow{\quad} & \mathrm{Spf} R_{\varpi, \square}^+
 \end{array}$$

where the vertical arrows are étale extensions and the horizontal maps are obtained by sending $X_0 \mapsto \varpi$, and the rest are natural maps. Finally, we set $R_\varpi = p$ -adic completion of $R_\varpi^+[\frac{1}{X_0}]$.

Next, since $P \equiv X_0^e \pmod{p}$, we have $R_\varpi^+[\frac{P^k}{k!}]_{k \in \mathbb{N}} = R_\varpi^+[\frac{X_0^k}{[k/e]!}]_{k \in \mathbb{N}}$. So, we set $R_\varpi^{\mathrm{PD}} := p$ -adic completion of $R_\varpi^+[\frac{P^k}{k!}]_{k \in \mathbb{N}}$. In summary, we have a diagram of formal schemes where the horizontal arrows are closed embeddings into formal schemes smooth over O_F , obtained by sending $X_0 \mapsto \varpi$ on the level of algebras,

$$\begin{array}{ccc}
 & \mathrm{Spf} R_\varpi^{\mathrm{PD}} & \\
 \swarrow & & \searrow \\
 \mathrm{Spf} R[\varpi] & \xrightarrow{\quad} & \mathrm{Spf} R_\varpi^+ \\
 \downarrow & & \downarrow \\
 \mathrm{Spf} O_K\{X, X^{-1}\} & \xrightarrow{\quad} & \mathrm{Spf} R_{\varpi, \square}^+ \\
 \downarrow & & \downarrow \\
 \mathrm{Spf} O_K & \xrightarrow{\quad} & \mathrm{Spf} r_\varpi^+ \\
 \downarrow & \swarrow & \\
 \mathrm{Spf} O_F & &
 \end{array}$$

Recall that P generates the kernel of the surjective map $R_{\varpi}^+ \rightarrow R[\varpi]$ and divided powers of P generate the kernel of the surjective map $R_{\varpi}^{\text{PD}} \rightarrow R[\varpi]$.

Definition 3.11. Endow the ring R_{ϖ}^{PD} with a filtration by divided power ideals as

$$\text{Fil}^k R_{\varpi}^{\text{PD}} = (P^{[n]}, n \geq k) \subset R_{\varpi}^{\text{PD}} \text{ for } k \in \mathbb{N}.$$

In other words, the filtration on R_{ϖ}^{PD} is given by divided powers of the kernel of $R_{\varpi}^{\text{PD}} \rightarrow R[\varpi]$. Furthermore, the ring R_{ϖ}^+ is endowed with the induced filtration

$$\text{Fil}^k R_{\varpi}^+ := R_{\varpi}^+ \cap \text{Fil}^k R_{\varpi}^{\text{PD}} = P^k R_{\varpi}^+ \text{ for } k \in \mathbb{N},$$

where the last equality follows since P generates the kernel of $R_{\varpi}^+ \rightarrow R[\varpi]$.

3.3. Cyclotomic embedding. In this section, we will describe the relationship between R_{ϖ}^{\star} for $\star \in \{+, \text{PD}\}$ and the period rings discussed in §2 and §3.1. We start by defining the (cyclotomic) Frobenius endomorphism on the former rings. Over $R_{\varpi, \square}^+$ define a lift of the absolute Frobenius on $R_{\varpi, \square}^+/p$ by

$$\begin{aligned} \varphi : R_{\varpi, \square}^+ &\longrightarrow R_{\varpi, \square}^+ \\ X_0 &\longmapsto (1 + X_0)^p - 1 \\ X_i &\longmapsto X_i^p, \text{ for } 1 \leq i \leq d, \end{aligned}$$

which we will call the (cyclotomic) Frobenius. Clearly, $\varphi(x) - x^p \in pR_{\varpi, \square}^+$ for $x \in R_{\varpi, \square}^+$. Using the implicit function theorem for topological rings [CN17, Proposition 2.1], we can extend the Frobenius homomorphism to $\varphi : R_{\varpi}^+ \rightarrow R_{\varpi}^+$. By continuity, the Frobenius endomorphism φ admits unique extensions $\varphi : R_{\varpi}^{\text{PD}} \rightarrow R_{\varpi}^{\text{PD}}$ and $\varphi : R_{\varpi} \rightarrow R_{\varpi}$.

3.3.1. The rings $\mathbf{A}_{R, \varpi}^{\star}$. We will describe the (cyclotomic) embeddings of R_{ϖ}^{\star} into various period rings discussed in §2 and §3.1. Define an embedding

$$\begin{aligned} \iota_{\text{cycl}} : R_{\varpi, \square}^+ &\longrightarrow \mathbf{A}_{\text{inf}}(\overline{R}) \\ X_0 &\longmapsto \pi_m = \varphi^{-m}(\pi), \\ X_i &\longmapsto [X_i^{\flat}], \text{ for } 1 \leq i \leq d. \end{aligned}$$

Lemma 3.12. *The map ι_{cycl} has a unique extension to an embedding $R_{\varpi}^+ \rightarrow \mathbf{A}_{\text{inf}}(\overline{R})$ such that $\theta \circ \iota_{\text{cycl}}$ is the projection $R_{\varpi}^+ \rightarrow R[\varpi]$.*

Proof. We can use the implicit function theorem [CN17, Proposition 2.1] to extend the embedding to $\iota_{\text{cycl}} : R_{\varpi}^+ \rightarrow \mathbf{A}_{\text{inf}}(\overline{R})$. Next, from definitions we already have that $\theta \circ \iota_{\text{cycl}} : R_{\varpi, \square}^+ \rightarrow O_K\{X, X^{-1}\}$ coincides with the canonical projection and R_{ϖ}^+ is étale over $R_{\varpi, \square}^+$, hence the second claim follows. ■

This embedding commutes with Frobenius on either side, i.e. $\iota_{\text{cycl}} \circ \varphi = \varphi \circ \iota_{\text{cycl}}$. By continuity, the morphism ι_{cycl} extends to embeddings $\iota_{\text{cycl}} : R_{\varpi}^{\text{PD}} \rightarrow \mathbf{A}_{\text{cris}}(\overline{R})$ and $\iota_{\text{cycl}} : R_{\varpi} \rightarrow \mathbf{A}_{\overline{R}}$. Denote by $\mathbf{A}_{R, \varpi}^+$ and $\mathbf{A}_{R, \varpi}$ the image in $\mathbf{A}_{\overline{R}}$ of R_{ϖ}^+ and R_{ϖ} respectively, under the map ι_{cycl} . Similarly, let $\mathbf{A}_{R, \varpi}^{\text{PD}} := \iota_{\text{cycl}}(R_{\varpi}^{\text{PD}}) \subset \mathbf{A}_{\text{cris}}(\overline{R})$. These rings are stable under the action of Γ_R (see [CN17, §2.5.3]). Moreover, these embeddings induce a filtration on $\mathbf{A}_{R, \varpi}^{\star}$ for $\star \in \{+, \text{PD}\}$ and $r \in \mathbb{Z}$ (use Definition 3.11).

Remark 3.13. Note that we write $\mathbf{A}_{R, \varpi}^+$ and so on instead of slightly cumbersome notation $\mathbf{A}_{R[\varpi]}^+$ or simpler notation \mathbf{A}_S^+ for $S = R[\varpi]$, in order to emphasize the choice of root of unity in the definition.

We note a simple lemma that will be useful later.

Lemma 3.14. $\frac{t}{\pi}$ is a unit in $\mathbf{A}_{F,\varpi}^{\text{PD}} \subset \mathbf{A}_{R,\varpi}^{\text{PD}}$.

Proof. We can write the fraction

$$\frac{t}{\pi} = \frac{\log(1+\pi)}{\pi} = \sum_{k \geq 0} (-1)^k \frac{\pi^k}{k+1}.$$

Formally, we can write

$$\frac{\pi}{t} = \frac{\pi}{\log(1+\pi)} = 1 + b_1\pi + b_2\pi^2 + b_3\pi^3 + \dots,$$

where $v_p(b_k) \geq -\frac{k}{p-1}$ for all $k \geq 1$. Since $\pi = (1 + \pi_m)^{p^m} - 1$, we get that $\pi \in (p, \pi_m^{p^m})\mathbf{A}_{F,\varpi}^+$ (as $m \geq 1$). By induction over k , we can easily conclude that $\pi^k \in (p, \pi_m^{p^m})^k \mathbf{A}_{F,\varpi}^{\text{PD}}$. Using this, we can re-express the series $\sum_k b_k \pi^k$ as a power series in π_m , written as $\sum_i c_i \pi_m^i$. We need to check that this re-expressed series converges in $\mathbf{A}_{F,\varpi}^{\text{PD}}$. To do this, we collect the terms with coefficients having the smallest p -adic valuation for each power of $\pi_m^{p^m}$ in the re-expressed series. For $k \geq 1$, b_k has the smallest p -adic valuation among the coefficients of $\pi_m^{p^m k}$, and therefore it has the least p -adic valuation among coefficients of π_m^i for $p^m k \leq i < p^m(k+1)$. We write the collection of these terms as

$$\sum_{k \geq 1} (-1)^{k+1} b_k \pi_m^{p^m k} = \sum_{k \geq 1} (-1)^{k+1} b_k \left\lfloor \frac{p^m k}{e} \right\rfloor! \frac{\pi_m^{p^m k}}{[p^m k/e]!}, \quad (3.3)$$

and by the preceding discussion it is sufficient to show that these coefficients go to 0 as $k \rightarrow +\infty$. Moreover, for (3.3) it would suffice to check the estimate for $k = (p-1)j$ as $j \rightarrow +\infty$ (this gets rid of the floor function above). With the observation in Remark 3.15, we have

$$v_p\left(b_k \left\lfloor \frac{p^m k}{e} \right\rfloor!\right) = v_p(b_k) + v_p((pj)!) \geq -\frac{(p-1)j}{p-1} + \frac{pj - s_p(pj)}{p-1} = \frac{j - s_p(j)}{p-1} = v_p(j!),$$

which goes to $+\infty$ as $j \rightarrow +\infty$. Hence, $\frac{\pi}{t}$ converges in $\mathbf{A}_{F,\varpi}^{\text{PD}}$ and is an inverse to $\frac{t}{\pi}$. \blacksquare

The following elementary observation was used above,

Remark 3.15. Let $n \in \mathbb{N}$, so we can write $n = \sum_{i=0}^k n_i p^i$ for some $k \in \mathbb{N}$, where $0 \leq n_i \leq p-1$ for $0 \leq i \leq k$. Let us set $s_p(n) = \sum_{i=0}^k n_i$. Then we have

$$\begin{aligned} v_p(n!) &= \sum_{j \geq 1} \left\lfloor \frac{n}{p^j} \right\rfloor = \sum_{j \geq 0} \left\lfloor \frac{\sum_{i=0}^k n_i p^i}{p^j} \right\rfloor = \sum_{j=1}^k \sum_{i=j}^k n_i p^{i-j} \\ &= \sum_{i=1}^k n_i \sum_{j=1}^i p^j = \sum_{i=1}^k n_i \frac{p^i - 1}{p-1} = \frac{n - s_p(n)}{p-1}. \end{aligned}$$

Also, note that we have $s_p(pn) = s_p(n)$ for any $n \in \mathbb{N}$.

Lemma 3.16. Let $i \in \{0, 1, \dots, d\}$. Then $(\gamma_i - 1)\mathbf{A}_{R,\varpi}^\star \subset \pi\mathbf{A}_{R,\varpi}^\star$ for $\star \in \{+, \text{PD}\}$;

Proof. First, let $i = 0$. Then we have

$$\begin{aligned} (\gamma_0 - 1)\pi_m &= (1 + \pi_m)((1 + \pi_m)^{\chi(\gamma_0)-1} - 1) = (1 + \pi_m)((1 + \pi_m)^{p^m a} - 1) \\ &= (1 + \pi_m)((1 + \pi)^a - 1) = (1 + \pi_m)(a\pi + \frac{a(a-1)}{2!}\pi^2 + \frac{a(a-1)(a-2)}{3!}\pi^3 + \dots) = \pi x, \end{aligned}$$

for some $x \in \mathbf{A}_{F,\varpi}^+$, i.e. $(\gamma_0 - 1)\pi_m \in \pi\mathbf{A}_{F,\varpi}^+$. Then it follows that $(\gamma_0 - 1)\mathbf{A}_{F,\varpi}^\star \subset \pi\mathbf{A}_{F,\varpi}^\star$ for $\star \in \{+, \text{PD}\}$

Next, for $i \in \{1, \dots, d\}$ we have $(\gamma_i - 1)[X_i^b] = \pi[X_i^b] \in \pi\mathbf{A}_{R,\varpi}^+$ and $(\gamma_i - 1)([X_i^b]^{-1}) = -\pi(1 + \pi)^{-1}[X_i^b]^{-1} \in \pi\mathbf{A}_{R,\varpi}^+$. Therefore, we get the claim. \blacksquare

3.3.2. The ring \mathbf{A}_R^+ . The preceding discussion works well for $R[\varpi]$ where $\varpi = \zeta_{p^m} - 1$ for $m \in \mathbb{N}_{\geq 1}$ ($m \in \mathbb{N}_{\geq 2}$ if $p = 2$). For R one can repeat the construction above to obtain the period ring $\mathbf{A}_R^+ \subset \mathbf{A}_{R,\varpi}^+$ (the embedding $R_{\varpi}^+ \hookrightarrow \mathbf{A}_{\text{inf}}(\overline{R})$ for R sends $X_0 \mapsto \pi$). Moreover, restriction of the map θ gives us a surjective map $\theta : \mathbf{A}_R^+ \twoheadrightarrow R$ whose kernel is principal and generated by π (since $\theta \circ \iota_{\text{cycl}} = \text{id}$ on R). Next, over $\mathbf{A}_{R,\varpi}^+$ the filtration is given as $\text{Fil}^k \mathbf{A}_{R,\varpi}^+ = \xi^k \mathbf{A}_{R,\varpi}^+$, where $\xi = \frac{\pi}{\pi_1}$. However, $\xi \notin \mathbf{A}_R^+$. Therefore, we equip \mathbf{A}_R^+ with the induced filtration $\text{Fil}^k \mathbf{A}_R^+ = \mathbf{A}_R^+ \cap \text{Fil}^k \mathbf{A}_{R,\varpi}^+$. Then describing the filtration as kernel of the θ map, we obtain

Lemma 3.17. $\text{Fil}^k \mathbf{A}_R^+ = \pi^k \mathbf{A}_R^+$.

Remark 3.18. Let \mathbf{A}^+ be the ring from Definition 3.5 and \mathbf{A}_{ϖ}^+ be the ring defined in Remark 3.6. From the definitions it follows that $\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{A}^+ \xrightarrow{\sim} \mathbf{A}_{\varpi}^+$ compatible with Frobenius and G_R -action. Moreover, we have $\mathbf{A}_R^+ = (\mathbf{A}^+)^{H_R}$ and $\mathbf{A}_{R,\varpi}^+ = (\mathbf{A}_{\varpi}^+)^{H_{R,\varpi}}$ where $H_{R,\varpi} = H_R$. Now, if we equip $\mathbf{A}^+ \subset \mathbf{A}_{\varpi}^+ \subset \mathbf{A}_{\text{inf}}(\overline{R})$ with the induced filtration, then we see that the isomorphism $\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{A}^+ \xrightarrow{\sim} \mathbf{A}_{\varpi}^+$ is compatible with filtrations as well (where on the left we consider the tensor product filtration).

3.4. Fat period rings. In this section we will introduce an alternative construction of fat period rings. This will be helpful in constructing some auxiliary rings in the proof of Proposition 4.28. Let S and Λ be p -adically complete filtered O_F -algebras. Let $\iota : S \rightarrow \Lambda$ be a continuous injective morphism of filtered O_F -algebras and let $f : S \otimes \Lambda \rightarrow \Lambda$ be the morphism sending $x \otimes y \mapsto \iota(x)y$.

Definition 3.19. Define $S\Lambda$ to be the p -adic completion of the divided power envelope of $S \otimes \Lambda$ with respect to $\text{Ker } f$.

Now, let $S = R, R_{\varpi}^{\text{PD}}$, where over R we consider the trivial filtration, whereas over R_{ϖ}^{PD} we consider the filtration described in Definition 3.11. Then we have,

Remark 3.20. (i) The ring $S\Lambda$ is the p -adic completion of $S \otimes \Lambda$ adjoined $(x \otimes 1 - 1 \otimes \iota(x))^{[k]}$, for $x \in S$ and $n \in \mathbb{N}$ and $(V_i - 1)^{[k]}$ for $1 \leq i \leq d$ and $k \in \mathbb{N}$, where $V_i = \frac{X_i \otimes 1}{1 \otimes \iota(X_i)}$ for $1 \leq i \leq d$.

(ii) The morphism $f : S \otimes \Lambda \rightarrow \Lambda$ extends uniquely to a continuous morphism $f : S\Lambda \rightarrow \Lambda$.

(iii) There is a natural filtration over $S\Lambda$ where we define $\text{Fil}^r S\Lambda$ to be the topological closure of the ideal generated by the products of the form $x_1 x_2 \prod (V_i - 1)^{[k_i]}$, with $x_1 \in \text{Fil}^{r_1} S$, $x_2 \in \text{Fil}^{r_2} \Lambda$ and $r_1 + r_2 + \sum k_i \geq r$.

(iv) From [CN17, Lemma 2.36], we have that any element $x \in S\Lambda$ can be uniquely written as $x = \sum_{\mathbf{k} \in \mathbb{N}^d} x_{\mathbf{k}} (1 - V_1)^{[k_1]} \dots (1 - V_d)^{[k_d]}$ with $x_{\mathbf{k}} \in \Lambda$ for all $\mathbf{k} = (k_1, \dots, k_d) \in \mathbb{N}^d$ and $x_{\mathbf{k}} \rightarrow 0$ as $|\mathbf{k}| = \sum_{i=1}^d k_i \rightarrow +\infty$. Moreover, an element $x \in \text{Fil}^r S\Lambda$ if and only if $x_{\mathbf{k}} \in \text{Fil}^{r-|\mathbf{k}|} \Lambda$ for all $\mathbf{k} \in \mathbb{N}^d$.

4. Finite height representations

In this section we will study Wach modules and their relationship with crystalline modules for crystalline representations.

4.1. The arithmetic case. Recall that we have $G_F = \text{Gal}(\overline{F}/F)$ as the absolute Galois group of F , $\Gamma_F := \text{Gal}(F_\infty/F)$ and $H_F := \text{Gal}(\overline{F}/F_\infty)$, where $F_\infty = \cup_n F(\mu_{p^n})$. From the theory of (φ, Γ_F) -modules, we have a two dimensional local ring \mathbf{A}_F given as the p -adic completion of $O_F[[\pi]][\frac{1}{\pi}]$ and $\mathbf{B}_F := \mathbf{A}_F[\frac{1}{p}]$ is a complete discrete valuation field with uniformizer p and residue field $\kappa((\overline{\pi}))$, the field of Laurent series with uniformizer $\overline{\pi}$ (the reduction of π modulo p).

Next, we have certain subrings $\mathbf{A}_F^+ := O_F[[\pi]] \subset \mathbf{A}_F$ and $\mathbf{B}_F^+ = \mathbf{A}_F^+[\frac{1}{p}] \subset \mathbf{B}_F$, stable under the action of φ and Γ_F . Let V be a p -adic representation of G_F , then $\mathbf{D}^+(V) = (\mathbf{B}^+ \otimes_{\mathbb{Q}_p} V)^{H_F}$ is a free module over the principal domain \mathbf{B}_F^+ of rank $\leq \dim_{\mathbb{Q}_p} V$, equipped with a Frobenius-semilinear endomorphism φ and a continuous and semilinear action of Γ_F . Further, let $\mathbf{D}(V) = (\mathbf{B} \otimes_{\mathbb{Q}_p} V)^{H_F}$ be the associated (φ, Γ_F) -module which is a \mathbf{B}_F -vector space of dimension $= \dim_{\mathbb{Q}_p} V$, equipped with a Frobenius-semilinear endomorphism φ and a continuous and semilinear action of Γ_F . We have a \mathbf{B}_F^+ -linear inclusion $\mathbf{D}^+(V) \subset \mathbf{D}(V)$ compatible with the action of φ and Γ_F . We say that V is of *finite height* if $\mathbf{D}^+(V)$ is a \mathbf{B}_F^+ -lattice inside $\mathbf{D}(V)$.

Similarly, if $T \subset V$ is a free \mathbb{Z}_p -lattice, stable under the action of G_F , then $\mathbf{D}^+(T) = (\mathbf{A}^+ \otimes_{\mathbb{Z}_p} T)^{H_F}$ is a free \mathbf{A}_F^+ -module of rank $\leq \dim_{\mathbb{Q}_p} V$, stable under the action of φ and Γ_F (see [Fon90, §B.1.2]). Moreover, $\mathbf{D}(T) = (\mathbf{A} \otimes_{\mathbb{Z}_p} T)^{H_F}$ is a free \mathbf{A}_F -module of rank $= \dim_{\mathbb{Q}_p} V$ equipped with a Frobenius-semilinear operator φ and a continuous and semilinear action of Γ_F , and we have $\mathbf{D}^+(T) \subset \mathbf{D}(T)$.

Fontaine showed that V is of finite height if and only if there exists a finite free \mathbf{B}_F^+ -submodule of $\mathbf{D}(V)$ of rank $= \dim_{\mathbb{Q}_p} V$, stable under the operator φ (see [Fon90, §B.2.1] and [Col99, §III.2]). Moreover, if $T \subset V$ is a free \mathbb{Z}_p -lattice as above and V of finite height, then $\mathbf{D}^+(T)$ is a free \mathbf{A}_F^+ -module of rank $= \dim_{\mathbb{Q}_p} V$ such that $\mathbf{A}_F \otimes_{\mathbf{A}_F^+} \mathbf{D}^+(T) \xrightarrow{\sim} \mathbf{D}(T)$ (see [Fon90, Théorème B.1.4.2]).

For crystalline representations there exist submodules of $\mathbf{D}^+(V)$ admitting a simpler action of Γ_F . Finite height and crystalline representations of G_F are related by the following result:

Theorem 4.1 ([Wac96], [Col99], [Ber02]). *Let V be a p -adic representation of G_F . Then V is crystalline if and only if it is of finite height and there exists $r \in \mathbb{Z}$ and a \mathbf{B}_F^+ -submodule $N \subset \mathbf{D}^+(V)$ of rank $= \dim_{\mathbb{Q}_p} V$, stable under the action of Γ_F , such that Γ_F acts trivially over $(N/\pi N)(-r)$.*

In the situation of Theorem 4.1, the module N is not unique. A functorial construction was given by Berger:

Proposition 4.2 ([Ber04, Proposition II.1.1]). *Let V be a positive crystalline representation of G_F , i.e. all Hodge-Tate weights of V are ≤ 0 . Let $T \subset V$ be a free \mathbb{Z}_p -lattice, stable under the action of G_F . Then there exists a unique \mathbf{A}_F^+ -module $\mathbf{N}(T) \subset \mathbf{D}(T)$, which is free of rank $= \dim_{\mathbb{Q}_p} V$, stable under the action of φ and Γ_F , and the action of Γ_F is trivial over $\mathbf{N}(T)/\pi\mathbf{N}(T)$. Moreover, there exists $s \in \mathbb{N}$ such that $\pi^s \mathbf{D}^+(T) \subset \mathbf{N}(T)$. Finally, set $\mathbf{N}(V) := \mathbf{B}_F^+ \otimes_{\mathbf{A}_F^+} \mathbf{N}(T)$, then $\mathbf{N}(V)$ is a unique \mathbf{B}_F^+ -submodule of $\mathbf{D}^+(V)$ satisfying analogous properties.*

Notation 4.3. For an algebra S admitting an action of the Frobenius and an S -module M admitting a Frobenius-semilinear endomorphism $\varphi : M \rightarrow M$, we denote by $\varphi^*(M) \subset M$ the S -submodule generated by the image of φ .

Remark 4.4. (i) In Proposition 4.2 for positive crystalline representations, Berger applies Theorem 4.1 with $r = 0$ to define $\mathbf{N}(V) := \mathbf{D}^+(V) \cap N[\frac{1}{\varphi^{n-1}(q)}]_{n \geq 1}$, where $q = \frac{\varphi(\pi)}{\pi}$.

Using this one can take $\mathbf{N}(T) := \mathbf{N}(V) \cap \mathbf{D}(T)$ and it can be shown to satisfy the desired properties.

- (ii) Berger further showed that in the setup of Proposition 4.2, if we take s to be the maximum among the absolute values of Hodge-Tate weights of V , then $\mathbf{N}(T)/\varphi^*(\mathbf{N}(T))$ is killed by q^s and we have that $\pi^s \mathbf{A}^+ \otimes_{\mathbb{Z}_p} T \subset \mathbf{A}^+ \otimes_{\mathbf{A}_F^+} \mathbf{N}(T)$ (see [Ber04, Théorème III.3.1]). The former observation can be thought of as a finite q -height property of Wach modules. We will impose it as one of the main conditions for defining finite q -height representations in the relative case (see 4.9).

Definition 4.5. Let $a, b \in \mathbb{Z}$ with $b \geq a$. A *Wach module* with weights in the interval $[a, b]$ is a finite free \mathbf{A}_F^+ -module or a \mathbf{B}_F^+ -module N , equipped with a continuous and semilinear action of Γ_F such that the action of Γ_F is trivial on $N/\pi N$ and a Frobenius-semilinear operator $\varphi : N[\frac{1}{\pi}] \rightarrow N[\frac{1}{\varphi(\pi)}]$ commuting with the action of Γ_F , $\varphi(\pi^b N) \subset \pi^b N$ and $\pi^b N/\varphi^*(\pi^b N)$ is killed by q^{b-a} .

Remark 4.6. The definition of the functor \mathbf{N} can be extended to crystalline representations of arbitrary Hodge-Tate weights quite easily. Indeed, let $V \in \text{Rep}_{\mathbb{Q}_p}^{\text{cris}}(G_F)$ with Hodge-Tate weights in the interval $[a, b]$ and let $T \subset V$ a free \mathbb{Z}_p -lattice, stable under the action of G_F . Then $\mathbf{N}(T) = \pi^{-b} \mathbf{N}(T(-b)) \otimes_{\mathbb{Z}_p} \mathbb{Z}_p(b)$ is a Wach module over \mathbf{A}_F^+ with weights in the interval $[a, b]$.

As it turns out, one can recover the crystalline representation from a given Wach module:

Proposition 4.7 ([Ber04, Proposition III.4.2]). *The functor*

$$\begin{aligned} \mathbf{N} : \text{Rep}_{\mathbb{Q}_p}^{\text{cris}}(G_F) &\longrightarrow \text{Wach modules over } \mathbf{B}_F^+ \\ V &\longmapsto \mathbf{N}(V), \end{aligned}$$

establishes an equivalence of categories with a quasi-inverse given by $N \mapsto (\mathbf{B} \otimes_{\mathbf{B}_F^+} N)^{\varphi=1}$. These functors are compatible with tensor products, duality and preserve exact sequences. Moreover, for a crystalline representation V , the map $T \mapsto \mathbf{N}(T)$ induces a bijection between \mathbb{Z}_p -lattices inside V and Wach modules over \mathbf{A}_F^+ contained in $\mathbf{N}(V)$.

We have a natural filtration on Wach modules given as

$$\text{Fil}^k \mathbf{N}(V) = \{x \in \mathbf{N}(V) \text{ such that } \varphi(x) \in q^k \mathbf{N}(V)\} \text{ for } k \in \mathbb{Z}.$$

If V is positive crystalline, i.e. all its Hodge-Tate weights are ≤ 0 , then for $r \in \mathbb{N}$ we have

$$\text{Fil}^k \mathbf{N}(V(r)) = \text{Fil}^k \pi^{-r} \mathbf{N}(V)(r) = \pi^{-r} \text{Fil}^{k+r} \mathbf{N}(V)(r).$$

Using this filtration on $\mathbf{N}(V)$, one can also recover the other linear algebraic object associated to V , i.e. the filtered φ -module $\mathbf{D}_{\text{cris}}(V)$: Let $\mathbf{B}_{\text{rig}, F}^+ \subset F[[\pi]]$ denote the subring of convergent power series over the open unit disc. Then we have $\mathbf{D}_{\text{cris}}(V) \subset \mathbf{B}_{\text{rig}, F}^+ \otimes_{\mathbf{B}_F^+} \mathbf{N}(V)$ and this gives $\mathbf{D}_{\text{cris}}(V) = (\mathbf{B}_{\text{rig}, F}^+ \otimes_{\mathbf{B}_F^+} \mathbf{N}(V))^{\Gamma_F}$ (see [Ber04, Proposition II.2.1]). Moreover, the induced map

$$\mathbf{D}_{\text{cris}}(V) \longrightarrow (\mathbf{B}_{\text{rig}, F}^+ \otimes_{\mathbf{B}_F^+} \mathbf{N}(V)) / \pi (\mathbf{B}_{\text{rig}, F}^+ \otimes_{\mathbf{B}_F^+} \mathbf{N}(V)) = \mathbf{N}(V) / \pi \mathbf{N}(V),$$

is an isomorphism of filtered φ -modules (see [Ber04, Proposition III.4.4]).

4.2. The relative case. In this section, we will introduce the notion of relative Wach modules and study representations of finite height. Recall that we fixed $m \in \mathbb{N}_{\geq 1}$ (fix $m \in \mathbb{N}_{\geq 2}$ if $p = 2$) and we have $K = F_m = F(\zeta_{p^m})$. The element $\varpi = \zeta_{p^m} - 1$ is a uniformizer of K . We have $X = (X_1, \dots, X_d)$ a set of indeterminates and we defined R to be the p -adic completion of an étale algebra over $O_F[X, X^{-1}]$ having non-empty and geometrically integral special fiber and $R[\varpi] = O_K \otimes_{O_F} R$. For R and $R[\varpi]$, we can use the (φ, Γ) -module theory discussed in §3.1, as well as the constructions in §3.2 and §3.3.

Setting $q = \frac{\varphi(\pi)}{\pi}$ and using the formulation in Definition 4.5, we define relative Wach modules:

Definition 4.8. Let $a, b \in \mathbb{Z}$ with $b \geq a$. A *Wach module* over \mathbf{A}_R^+ (resp. \mathbf{B}_R^+) with weights in the interval $[a, b]$ is a finite projective \mathbf{A}_R^+ -module (resp. \mathbf{B}_R^+ -module) N , equipped with a continuous and semilinear action of Γ_R such that the action of Γ_R is trivial on $N/\pi N$. Further, there is a Frobenius-semilinear operator $\varphi : N[\frac{1}{\pi}] \rightarrow N[\frac{1}{\varphi(\pi)}]$ commuting with the action of Γ_R such that $\varphi(\pi^b N) \subset \pi^b N$ and $\pi^b N / \varphi^*(\pi^b N)$ is killed by q^{b-a} .

Let V be a p -adic representation of the Galois group G_R admitting a \mathbb{Z}_p -lattice $T \subset V$ stable under the action of G_R . Then we have the finitely generated \mathbf{A}_R^+ -module $\mathbf{D}^+(T) := (\mathbf{A}^+ \otimes_{\mathbb{Q}_p} T)^{H_R}$. We introduce the following definition:

Definition 4.9. A *positive finite q -height* representation is a p -adic representation V of G_R admitting a \mathbb{Z}_p -lattice $T \subset V$ such that there exists a finite projective \mathbf{A}_R^+ -submodule $\mathbf{N}(T) \subset \mathbf{D}^+(T)$ of rank $= \dim_{\mathbb{Q}_p} V$ satisfying the following conditions:

- (i) $\mathbf{N}(T)$ is stable under the action of φ and Γ_R , and $\mathbf{A}_R \otimes_{\mathbf{A}_R^+} \mathbf{N}(T) \xrightarrow{\sim} \mathbf{D}(T)$;
- (ii) The \mathbf{A}_R^+ -module $\mathbf{N}(T)/\varphi^*(\mathbf{N}(T))$ is killed by q^s for some $s \in \mathbb{N}$;
- (iii) The action of Γ_R is trivial on $\mathbf{N}(T)/\pi \mathbf{N}(T)$;
- (iv) There exists a $R' \subset \bar{R}$ finite étale over R such that the $\mathbf{A}_{R'}^+$ -module $\mathbf{A}_{R'}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ is free.

The module $\mathbf{N}(T)$ is a *Wach module* associated to T with weights in the interval $[-s, 0]$ and we set $\mathbf{N}(V) := \mathbf{N}(T)[\frac{1}{p}]$ satisfying properties analogous to (i)-(iv) above. The *height* of V is defined to be the smallest $s \in \mathbb{N}$ satisfying (ii) above.

For $r \in \mathbb{Z}$, we set $V(r) := V \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(r)$ and $T(r) := T \otimes_{\mathbb{Z}_p} \mathbb{Z}_p(r)$. We will call these twists as representations of *finite q -height* and define

$$\mathbf{N}(T(r)) := \frac{1}{\pi^r} \mathbf{N}(T)(r) \text{ and } \mathbf{N}(V(r)) := \frac{1}{\pi^r} \mathbf{N}(V)(r).$$

Since $\mathbf{N}(V)$ and $\mathbf{N}(T)$ are Wach modules with weights in the interval $[-s, 0]$, twisting by r gives us Wach modules in the sense of Definition 4.8 with weights in the interval $[r-s, r]$. We will say that *height* of $V(r) = \text{height of } V$.

Remark 4.10. (i) In the arithmetic case, i.e. $R = O_F$, the notion of finite height representations in Theorem 4.1 and finite q -height representations in Definition 4.9 are related. In fact, in the arithmetic case using Definition 4.9 one obtains the functorial object of Berger mentioned above (see [Ber04, Proposition II.1.1]).

- (ii) In Definition 4.9 conditions (i), (ii) and (iii) are motivated from the definition of finite height representations of G_F admitting a Wach module structure. The last condition, i.e. (iv) is inspired by Brinon's definition of weak admissibility in the relative case (see [Bri08, p. 136]).

- (iii) In Definition 4.9 following Remark 4.4 (i), one can first define Wach module for the representation V and then consider the module $\mathbf{N}(T) = \mathbf{N}(V) \cap \mathbf{D}(T)$ associated to T . However, it is not clear whether the latter module, defined in this fashion, is a projective \mathbf{A}_R^+ -module. Therefore, we impose the condition on $\mathbf{N}(T)$ to be projective, which is required in establishing several results in this section.

4.2.1. Some properties of Wach modules. Let us note some important properties of Wach modules associated to finite q -height representations

Proposition 4.11. *Let V be a positive finite q -height representation and $T \subset V$ a G_R -stable \mathbb{Z}_p -lattice. Then we have $\pi^s \mathbf{A}^+ \otimes_{\mathbb{Z}_p} T \subset \mathbf{A}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$, where $s \in \mathbb{N}$ is the height of the representation V .*

Proof. To show the claim, we can assume that $\mathbf{N}(T)$ is free by base changing to the period ring corresponding to the finite étale extension R' of R . Then $\mathbf{A}^+ \otimes_{\mathbf{A}_R^+} (\mathbf{A}_{R'}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)) = \mathbf{A}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ is free. Since the discussion of previous chapters hold for the p -adic completion of a finite étale extension of R (see [Bri08, Chapitre 2] and [AI08, §2] for more on this), base changing to R' is harmless. So with a slight abuse of notation, below we will replace R' obtained in this manner by R and assume $\mathbf{N}(T)$ to be free of rank $h = \dim_{\mathbb{Q}_p} V$ over \mathbf{A}_R^+ .

Note that by definition we have $\mathbf{N}(T) \subset \mathbf{D}^+(T) = (\mathbf{A}^+ \otimes_{\mathbb{Z}_p} T)^{H_R} \subset \mathbf{A}^+ \otimes_{\mathbb{Z}_p} T$. So let $A \in \text{Mat}(h, \mathbf{A}^+)$ be the matrix obtained by expressing a basis of $\mathbf{N}(T)$ in a chosen basis of T . Also, let $P \in \text{Mat}(h, \mathbf{A}_R^+)$ be the matrix of φ in the basis of $\mathbf{N}(T)$. Then we have $\varphi(A) = AP$ and therefore $\varphi(\pi^s A^{-1}) = (q^s P^{-1})(\pi^s A^{-1})$. The fact that $\mathbf{N}(T)/\varphi^*(\mathbf{N}(T))$ is killed by q^s implies that $q^s P^{-1} \in \text{Mat}(h, \mathbf{A}_R^+)$, therefore from Lemma 3.9 we obtain that $\pi^s A^{-1} \in \text{Mat}(h, \mathbf{A}^+)$. Hence, we conclude that $\pi^s \mathbf{A}^+ \otimes_{\mathbb{Z}_p} T \subset \mathbf{A}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$. ■

Corollary 4.12. *By taking H_R -invariants in Proposition 4.11 it follows that $\pi^s \mathbf{D}^+(T) \subset \mathbf{N}(T)$.*

Proposition 4.13. *Let V be a finite q -height representation G_R . The Wach module $\mathbf{N}(V)$ over \mathbf{B}_R^+ is unique. Same holds true for the \mathbf{A}_R^+ -module $\mathbf{N}(T)$.*

Proof. The argument carries over from the classical case [Ber04, p. 13]. First note that we can assume that V is positive, since by definition the uniqueness of Wach module for such a representation is equivalent to uniqueness for all its Tate twists. In this case, let N_1 and N_2 be two \mathbf{A}_R^+ -modules satisfying the conditions of Definition 4.9 (the proof stays the same for $\mathbf{N}(V)$). By symmetry, it is enough to show that $N_1 \subset N_2$. Since we have $\pi^s N_1 \subset \pi^s \mathbf{D}^+(T) \subset N_2$ (see Corollary 4.12) and N_2 is π -torsion free, therefore for any $x \in N_1$ there exists $k \leq s$ such that $\pi^k x \in N_2$ but $\pi^k x \notin \pi N_2$. Varying over all $x \in N_1 \setminus \pi N_1$, we can take $k \leq s$ to be the minimal integer such that $\pi^k N_1 \subset N_2$. Since $\pi^k x \in N_2$ and Γ_R acts trivially on $N_2/\pi N_2$, we have that $(\gamma_0 - 1)(\pi^k x) \in \pi N_2$. So we can write

$$(\gamma_0 - 1)(\pi^k x) = \gamma_0(\pi^k)(\gamma_0(x) - x) + (\gamma_0(\pi^k) - \pi^k)x.$$

Since Γ_R also acts trivially on $N_1/\pi N_1$ and $\pi^k N_1 \subset N_2$, we see that $\gamma_0(\pi^k)(\gamma_0(x) - x) \in \pi N_2$, therefore $(\gamma_0(\pi^k) - \pi^k)x \in \pi N_2$, which means that $(\chi(\gamma_0)^k - 1)\pi^k x \in \pi N_2$. But $\pi \nmid (\chi(\gamma_0)^k - 1)$ if $k \geq 1$, and $\pi^k x \notin \pi N_2$. Hence, we must have $k = 0$, i.e. $N_1 \subset N_2$. ■

The uniqueness of Wach modules helps us in establishing compatibility with usual operations:

Proposition 4.14. *Let V and V' be two finite q -height representations of G_R . Then we have that $\mathbf{N}(V \oplus V') = \mathbf{N}(V) \oplus \mathbf{N}(V')$ and $\mathbf{N}(V \otimes V') = \mathbf{N}(V) \otimes \mathbf{N}(V')$. Similar statements hold for $\mathbf{N}(T)$ and $\mathbf{N}(T')$.*

Proof. We note similar to previous lemma that it is enough to show the statement for V and V' such that both representations are positive. By uniqueness of Wach modules proved in Proposition 4.13, it is enough to show that direct sum and tensor product of finite q -height representations are again of finite q -height.

First, it is straightforward to see that $\mathbf{N}(T) \oplus \mathbf{N}(T') \subset \mathbf{D}^+(T \oplus T')$ is a projective \mathbf{A}_R^+ -module of rank $\mathrm{rk}_{\mathbb{Z}_p}(T \oplus T')$ such that $\mathbf{A}_R \otimes_{\mathbf{A}_R^+} (\mathbf{N}(T) \oplus \mathbf{N}(T')) \xrightarrow{\sim} \mathbf{D}(T) \oplus \mathbf{D}(T')$. Similarly, we have that $\mathbf{N}(T) \otimes \mathbf{N}(T') \subset \mathbf{D}^+(T \otimes T')$ is a projective \mathbf{A}_R^+ -module of rank $\mathrm{rk}_{\mathbb{Z}_p}(T \otimes T')$ such that $\mathbf{A}_R \otimes_{\mathbf{A}_R^+} (\mathbf{N}(T) \otimes \mathbf{N}(T')) \xrightarrow{\sim} \mathbf{D}(T) \otimes \mathbf{D}(T')$.

Next, let s and s' denote the height of representations V and V' respectively and let $i := \max(s, s')$. Then we see that $(\mathbf{N}(T) \oplus \mathbf{N}(T'))/\varphi^*(\mathbf{N}(T) \oplus \mathbf{N}(T'))$ is killed by q^i and $(\mathbf{N}(T) \otimes \mathbf{N}(T'))/\varphi^*(\mathbf{N}(T) \otimes \mathbf{N}(T'))$ is killed by $q^{s+s'}$. Further, Γ_R acts trivially modulo π on $\mathbf{N}(T) \oplus \mathbf{N}(T')$ and $\mathbf{N}(T) \otimes \mathbf{N}(T')$. This verifies conditions (i), (ii) and (iii) for these modules. For condition (iv), note that given any two finite étale extensions R' and R'' of R , there exists a finite étale extension S over R such that S is finite étale over R' as well as R'' . Hence, we get the claim. ■

Corollary 4.15. *Let V be a finite q -height representation of G_R . Then, for $k \in \mathbb{N}$ the representations $\mathrm{Sym}^k(V)$ and $\wedge^k V$ are of finite q -height.*

Proof. Note that the compatibility with tensor products in Proposition 4.14 is enough to establish the compatibility with symmetric powers and exterior powers because then we can set

$$\mathbf{N}(\mathrm{Sym}^k(T)) := \mathrm{Sym}^k(\mathbf{N}(T)), \text{ and } \mathbf{N}(\wedge^k T) := \wedge^k \mathbf{N}(T).$$

We have $\mathbf{N}(\mathrm{Sym}^k(T)) \subset \mathrm{Sym}^k(\mathbf{D}^+(T)) \subset \mathbf{D}^+(\mathrm{Sym}^k(T))$, since $\mathbf{A}^+ \otimes_{\mathbf{A}_R^+} \mathrm{Sym}^k(\mathbf{D}^+(T)) \subset \mathbf{A}^+ \otimes_{\mathbf{A}_R^+} \mathbf{D}^+(\mathrm{Sym}^k(T))$. Similarly, $\mathbf{N}(\wedge^k T) \subset \mathbf{D}^+(\wedge^k T)$. Rest of the assumptions of Definition 4.9 follows in a same manner as in the proof of Proposition 4.14. This establishes that $\mathrm{Sym}^k(V)$ and $\wedge^k V$ are finite q -height representations and gives us the corresponding Wach modules. ■

4.2.2. Filtration on Wach modules. There is a natural filtration on Wach modules associated to finite q -height representations. We will introduce this filtration next and prove a lemma concerning this filtration.

Definition 4.16. Let V be a positive finite q -height representation of G_R and $r \in \mathbb{N}$. Then there is a natural filtration on the associated Wach modules given as

$$\mathrm{Fil}^k \mathbf{N}(V(r)) := \{x \in \mathbf{N}(V(r)), \text{ such that } \varphi(x) \in q^k \mathbf{N}(V(r))\} \text{ for } k \in \mathbb{Z},$$

and we set $\mathrm{Fil}^k \mathbf{N}(T(r)) := \mathrm{Fil}^k \mathbf{N}(V(r)) \cap \mathbf{N}(T(r))$, where the intersection is taken inside $\mathbf{N}(V(r))$.

Lemma 4.17. *With notations as above, we have*

- (i) $\mathrm{Fil}^k \mathbf{N}(T(r)) = \{x \in \mathbf{N}(T(r)), \text{ such that } \varphi(x) \in q^k \mathbf{N}(T(r))\}$.
- (ii) $\mathrm{Fil}^k \mathbf{N}(V(r)) = \mathrm{Fil}^k \pi^{-r} \mathbf{N}(V)(r) = \pi^{-r} \mathrm{Fil}^{k+r} \mathbf{N}(V)(r)$ and similarly for $\mathrm{Fil}^k \mathbf{N}(T(r))$.

Proof. (i) For $k \leq 0$, the claim is obvious, so we assume that $k > 0$. Then we are reduced to showing that $q^k \mathbf{N}(V(r)) \cap \mathbf{N}(T(r)) = q^k \mathbf{N}(T(r))$.

To prove the latter claim, note that it is enough to work under the assumption that $\mathbf{N}(T(r))$ is free. Indeed, for any finite q -height representation $V(r)$, there exists a finite étale R -algebra R' such that $\mathbf{A}_{R'}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T(r))$ is free. Since $\mathbf{A}_{R'}^+$ is faithfully flat over \mathbf{A}_R^+ , the claim is equivalent to showing that $\mathbf{A}_{R'}^+ \otimes_{\mathbf{A}_R^+} (q^k \mathbf{N}(V) \cap \mathbf{N}(T)) = q^k \mathbf{A}_{R'}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$. But one can easily obtain that $\mathbf{A}_{R'}^+ \otimes_{\mathbf{A}_R^+} (q^k \mathbf{N}(V) \cap \mathbf{N}(T)) = (q^k \mathbf{A}_{R'}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)) \cap (\mathbf{A}_{R'}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))$.

$\mathbf{N}(T)$) (or see [Mat86, Theorem 7.4 (i)]) as submodules of $\mathbf{A}_{R'}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)$. So below we will assume that $\mathbf{N}(T(r))$ is free over \mathbf{A}_R^+ with a basis $\{f_1, \dots, f_h\}$, where $h = \dim_{\mathbb{Q}_p} V(r)$.

Let $x = \sum_{i=1}^h x_i f_i \in q^k \mathbf{N}(V(r)) \cap \mathbf{N}(T(r))$ with $x_i \in \mathbf{A}_R^+$. Since $\{f_1, \dots, f_h\}$ is also a \mathbf{B}_R^+ -basis of $\mathbf{N}(V(r))$, we can write $x = q^k \sum_{i=1}^h y_i f_i$ with $y_i \in \mathbf{B}_R^+$. Comparing the two expressions for x we obtain that $q^k y_i = x_i \in \mathbf{A}_R^+$, i.e. $y_i \in \mathbf{A}_R$ for $1 \leq i \leq h$. But this just means that $y_i \in \mathbf{B}_R^+ \cap \mathbf{A}_R = \mathbf{A}_R^+$, therefore $x_i = q^k y_i \in q^k \mathbf{A}_R^+$ for $1 \leq i \leq h$. Hence, $x \in q^k \mathbf{N}(T(r))$ as desired. The other inclusion is obvious. ■

- (ii) Note that the inclusion $\pi^{-r} \text{Fil}^{k+r} \mathbf{N}(V)(r) \subset \text{Fil}^k \pi^{-r} \mathbf{N}(V)(r)$ is obvious. To show the converse let $\pi^{-r} x \otimes \epsilon^{\otimes r} \in \text{Fil}^k \pi^{-r} \mathbf{N}(V)(r)$, with $x \in \mathbf{N}(V)$ and $\epsilon^{\otimes r}$ being a basis of $\mathbb{Q}_p(r)$. Then we have that $\varphi(\pi^{-r} x \otimes \epsilon^{\otimes r}) = q^{-r} \pi^{-r} \varphi(x) \otimes \epsilon^{\otimes r} \in q^k \pi^{-r} \mathbf{N}(V)(r)$. Therefore, we obtain that $\varphi(x) \in q^{k+r} \mathbf{N}(V)$, i.e. $x \in \text{Fil}^{k+r} \mathbf{N}(V)$. ■

Remark 4.18. For $V = \mathbb{Q}_p$ the filtration in Definition 4.16 coincides with the filtration in Lemma 3.17

Proof. We have $T = \mathbb{Z}_p$ and $\mathbf{N}(T) = \mathbf{A}_R^+$ and let $\varpi = \zeta_p - 1$ (let $\varpi = \zeta_{p^2} - 1$ if $p = 2$) in this proof. Since $\pi^k \mathbf{A}_R^+ \subset \text{Fil}^k \mathbf{N}(T)$ (where the term on right is the filtration in Definition 4.16), we only need to show that $\text{Fil}^k \mathbf{N}(T) \subset \pi^k \mathbf{A}_R^+ = \mathbf{A}_R^+ \cap \xi^k \mathbf{A}_{R,\varpi}^+$. Let $x \in \mathbf{A}_R^+$ such that $\varphi(x) = q^k y$ for some $y \in \mathbf{A}_R^+$. As we have $\mathbf{A}_R^+ \subset \mathbf{A}_{R,\varpi}$, we can also write $\varphi(x) = \varphi(\xi^k) y \in \varphi(\mathbf{A}_{R,\varpi}) \subset \mathbf{A}_R$, i.e. $y \in \varphi(\mathbf{A}_{R,\varpi}) \cap \mathbf{A}_R^+ = \varphi(\mathbf{A}_{R,\varpi}^+)$ (where the intersection is taken inside \mathbf{A}_R). Therefore, we obtain that $y = \varphi(z)$ for some $z \in \mathbf{A}_{R,\varpi}^+$. Since $\varphi : \mathbf{A}_{R,\varpi}^+ \rightarrow \mathbf{A}_{R,\varpi}^+$ is injective, we must have $x = \xi^k z \in \mathbf{A}_R^+ \cap \xi^k \mathbf{A}_{R,\varpi}^+$, as desired. ■

4.3. Statement of the main result. In this section, we will relate the notion of crystalline and finite q -height representations. As we will see, we can recover the $R[\frac{1}{p}]$ -module $\mathcal{O}\mathbf{D}_{\text{cris}}(V)$ from the \mathbf{A}_R^+ -module $\mathbf{N}(T)$ after passing to a larger period ring and inverting p . We begin by introducing this ring below.

Recall from §1.4 that we have F as a finite unramified extension of \mathbb{Q}_p with ring of integers O_F and we take $K = F(\zeta_{p^m})$ for a fixed $m \in \mathbb{N}_{\geq 1}$ (fix $m \in \mathbb{N}_{\geq 2}$ if $p = 2$). Note that the formulation of the results and proofs depend on m and it is necessary to have $m \geq 1$ ($m \geq 2$ if $p = 2$) for the discussion below to make sense.

4.3.1. The ring $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$. In this section, we will work with the ring $\mathbf{A}_{R,\varpi}^+$ defined in §3.3, equipped with an action of the Frobenius φ and a continuous action of Γ_R . Since we have a natural injection $\mathbf{A}_{R,\varpi}^+ \hookrightarrow \mathbf{A}_{\text{inf}}(\overline{R})$, we obtain a G_R -equivariant commutative diagram

$$\begin{array}{ccc} \mathbf{A}_{R,\varpi}^+ & \xrightarrow{\theta} & R[\varpi] \\ \downarrow & & \downarrow \\ \mathbf{A}_{\text{inf}}(\overline{R}) & \xrightarrow{\theta} & \mathbb{C}^+(\overline{R}). \end{array}$$

By R -linearity, extending scalars for the map θ above, we obtain a ring homomorphism

$$\theta_R : R \otimes_{\mathbb{Z}} \mathbf{A}_{R,\varpi}^+ \longrightarrow R[\varpi],$$

sending $X_i \otimes 1 \mapsto X_i$, $1 \otimes [X_i^b] \mapsto X_i$ for $1 \leq i \leq d$ and $1 \otimes \pi_m \mapsto \zeta_{p^m} - 1$. Note that we have inclusion of ideals $(\xi, X_i \otimes 1 - 1 \otimes [X_i^b], \text{ for } 1 \leq i \leq d) \subset \text{Ker } \theta_R \subset R \otimes_{\mathbb{Z}} \mathbf{A}_{R,\varpi}^+$, where $\xi = \frac{\pi}{\pi_1}$. We have $\mathbf{A}_{R,\varpi}^+ \subset \mathbf{A}_{\text{inf}}(\overline{R})$ and θ_R above is the restriction of $\theta_R : R \otimes_{\mathbb{Z}} \mathbf{A}_{\text{inf}}(\overline{R}) \rightarrow \mathbb{C}^+(\overline{R})$ (see §2.2.1). So similar to $\mathcal{O}\mathbf{A}_{\text{inf}}(\overline{R})$ in §2.1.3 and $\mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})$ in §2.2.2 we define the following rings:

Definition 4.19. (i) Define $\mathcal{O}\mathbf{A}_{R,\varpi}^+$ to be $\theta_R^{-1}(pR[\varpi])$ -adic completion of $R \otimes_{\mathbb{Z}} \mathbf{A}_{R,\varpi}^+$.
(ii) Let $x^{[n]} := x^n/n!$ for $x \in \text{Ker } \theta_R$. Define $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ to be the p -adic completion of the divided power envelope of $R \otimes_{\mathbb{Z}} \mathbf{A}_{R,\varpi}^+$ with respect to $\text{Ker } \theta_R$.
Note that we have $\mathcal{O}\mathbf{A}_{R,\varpi}^+ = \mathcal{O}\mathbf{A}_{\text{inf}}(\overline{R}) \cap \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \subset \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})$.

Taking the divided power envelope of θ_R/p^n , note that $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n \twoheadrightarrow \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})/p^n$. Since we have $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} = \lim_n \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n$ and $\mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R}) = \lim_n \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})/p^n$, and (projective) limit is left exact, it follows that for the p -adic completion of divided power envelope of θ_R , we have $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \subset \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})$. Now, over the ring $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ we can consider the induced action of Γ_R under which it is stable, and it admits a Frobenius endomorphism arising from the Frobenius on each component of the tensor product. In particular, from the diagram above we obtain a G_R -equivariant commutative diagram

$$\begin{array}{ccc} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} & \xrightarrow{\theta_R} & R[\varpi] \\ \downarrow & & \downarrow \\ \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R}) & \xrightarrow{\theta_R} & \mathbb{C}^+(\overline{R}). \end{array}$$

Note that the left vertical arrow is Frobenius-equivariant.

Next, we will give an alternative description of the ring $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$. Let $T = (T_1, \dots, T_d)$ denote a set of indeterminates and let $\mathbf{A}_{\text{cris}}(\overline{R})\langle T \rangle^\wedge$ denote the p -adic completion of the divided power polynomial algebra $\mathbf{A}_{\text{cris}}(\overline{R})\langle T \rangle = \mathbf{A}_{\text{cris}}(\overline{R})[T_i^{[n]}]$, $n \in \mathbb{N}$, $1 \leq i \leq d$. Recall from §2.2.2 that we have an isomorphism of rings

$$\begin{aligned} f_{\text{cris}} : \mathbf{A}_{\text{cris}}(\overline{R})\langle T \rangle^\wedge &\xrightarrow{\sim} \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R}) \\ T_i &\longmapsto X_i \otimes 1 - 1 \otimes [X_i^b], \text{ for } 1 \leq i \leq d. \end{aligned}$$

Now recall that $\mathbf{A}_{R,\varpi}^{\text{PD}}$ is the p -adic completion of the divided power envelope of the surjective map $\theta : \mathbf{A}_{R,\varpi}^+ \twoheadrightarrow R[\varpi]$ with respect to its kernel (see §3.2). Next, let $\mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge$ denote the p -adic completion of the divided power polynomial algebra $\mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle = \mathbf{A}_{R,\varpi}^{\text{PD}}[T_i^{[n]}]$, $n \in \mathbb{N}$, $1 \leq i \leq d$. Then via the isomorphism f^{PD} (see Lemma 4.20 below), we will show that the preimage of $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$, under f_{cris} is exactly $\mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge$. In other words,

Lemma 4.20. *The morphism of rings*

$$\begin{aligned} f^{\text{PD}} : \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge &\longrightarrow \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \\ T_i &\longmapsto X_i \otimes 1 - 1 \otimes [X_i^b], \text{ for } 1 \leq i \leq d, \end{aligned}$$

is an isomorphism.

Proof. The proof follows [Bri08, Proposition 6.1.5] closely.

Recall that we have a surjective ring homomorphism $\theta : \mathbf{A}_{R,\varpi}^{\text{PD}} \twoheadrightarrow R[\varpi]$, which is the restriction of the map $\theta : \mathbf{A}_{\text{cris}}(\overline{R}) \rightarrow \mathbb{C}^+(\overline{R})$ defined in §2.2. This can be extended in a unique manner into the homomorphism $\theta : \mathbf{A}_{\text{cris}}(\overline{R})\langle T \rangle^\wedge \rightarrow \mathbb{C}^+(\overline{R})$. Restriction of the latter map gives us $\theta : \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge \twoheadrightarrow R[\varpi]$ such that $\theta(T_i^{[n]}) = 0$ for $1 \leq i \leq d$ and $n \geq 1$.

First, we will show that the $\mathcal{O}_F\{X^{\pm 1}\}$ -algebra structure on $\mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge$ given by $X_i \mapsto [X_i^b] + T_i$, extends uniquely to an R -algebra structure. Let $\mathcal{A} := (\mathbf{E}_{R,\varpi}^+/\pi^{p-1}\mathbf{E}_{R,\varpi}^+)[T_1, \dots, T_d]/(T_1^p, \dots, T_d^p)$. We have a surjective map $\theta : \mathbf{A}_{R,\varpi}^+ \twoheadrightarrow R[\varpi]$ and its reduction modulo p is given as $\bar{\theta} : \mathbf{E}_{R,\varpi}^+ \twoheadrightarrow R[\varpi]/pR[\varpi]$. Since $\xi^p \equiv \pi^{p-1} \pmod{p}$, where $\xi = \frac{\pi}{\pi_1}$

is a generator of $\text{Ker } \theta \subset \mathbf{A}_{R,\varpi}^+$, we obtain that $\bar{\theta}$ factors as $\bar{\theta} : \mathbf{E}_{R,\varpi}^+/\bar{\pi}^{p-1}\mathbf{E}_{R,\varpi}^+ \twoheadrightarrow R[\varpi]/pR[\varpi]$. This can be extended to a map $\bar{\theta} : \mathcal{A} \twoheadrightarrow R[\varpi]/pR[\varpi]$ by setting $\bar{\theta}(T_i) = 0$ for $1 \leq i \leq d$. The kernel $\mathcal{I} = \text{Ker } \bar{\theta} \subset \mathcal{A}$ is generated by $\xi \equiv \bar{\pi}_1^{p-1} \pmod{p}$ and $\{T_i\}_{1 \leq i \leq d}$. Now from the natural inclusion $R/pR \hookrightarrow R[\varpi]/pR[\varpi]$ and the isomorphism $\mathcal{A}/\mathcal{I} \xrightarrow{\sim} R[\varpi]/pR[\varpi]$ via $\bar{\theta}$, we obtain a map $\bar{g} : R/pR \rightarrow \mathcal{A}/\mathcal{I}$ such that $\bar{g}(X_i) = X_i$, which is the image of $X_i^b \in \mathcal{A}$ under the map $\bar{\theta}$. So we obtain a commutative diagram

$$\begin{array}{ccc} \kappa[X^{\pm 1}] & \xrightarrow{\quad} & \mathcal{A} \\ \downarrow & \nearrow \bar{g} & \downarrow \\ R/pR & \xrightarrow{\quad} & \mathcal{A}/\mathcal{I} \end{array}$$

where the top horizontal arrow is the map $X_i \mapsto X_i^b + T_i$. Note that $\mathcal{I}^{(d+1)p} = 0$. Since R/pR is étale over $\kappa[X^{\pm 1}]$, there exists a unique lift of $\bar{g} : R/pR \rightarrow \mathcal{A}/\mathcal{I}$ to a homomorphism $\bar{g} : R/pR \rightarrow \mathcal{A}$ (which we again denote by \bar{g} by slight abuse of notations).

Further, by the description of divided power envelope in [Bri08, Proposition 6.1.1] we have that

$$\begin{aligned} \mathbf{A}_{R,\varpi}^+[Y_0, Y_1, \dots]/(pY_0 - \xi^p, pY_{n+1} - Y_n^p)_{n \geq 1} &\xrightarrow{\sim} \mathbf{A}_{R,\varpi}^{\text{PD}} \\ Y_n &\longmapsto \frac{\xi^{p^{n+1}}}{p^{n+1}}. \end{aligned}$$

Therefore,

$$(\mathbf{E}_{R,\varpi}^+/\bar{\pi}^{p-1}\mathbf{E}_{R,\varpi}^+)[Y_0, Y_1, \dots]/(Y_n^p)_{n \geq 1} \xrightarrow{\sim} \mathbf{A}_{R,\varpi}^{\text{PD}}/p\mathbf{A}_{R,\varpi}^{\text{PD}},$$

Similarly, we have

$$(\mathbf{A}_{R,\varpi}^{\text{PD}}[T_1, \dots, T_d])[T_{i,0}, T_{i,1}, \dots]/(pT_{i,0} - T_i^p, pT_{i,n+1} - T_{i,n}^p)_{1 \leq i \leq d, n \in \mathbb{N}} \xrightarrow{\sim} \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle.$$

Therefore,

$$(\mathbf{A}_{R,\varpi}^{\text{PD}}/p\mathbf{A}_{R,\varpi}^{\text{PD}})[T_1, \dots, T_d][T_{i,0}, T_{i,1}, \dots]/(T_i^p, T_{i,n}^p)_{1 \leq i \leq d, n \in \mathbb{N}} \xrightarrow{\sim} \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle/p\mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle.$$

In conclusion, we have

$$\mathcal{A}[Y_0, Y_1, \dots, T_{i,0}, T_{i,1}, \dots]/(Y_n^p, T_{i,n}^p)_{1 \leq i \leq d, n \in \mathbb{N}} \xrightarrow{\sim} \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle/p\mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle.$$

From the discussion above we obtain a natural map of $\kappa[X^{\pm 1}]$ -algebras by composition $\bar{g}_1 : R/pR \rightarrow \mathcal{A} \rightarrow \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle/p\mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle$.

Now let $n \in \mathbb{N}$, then modulo p^n we have a natural map $O_F\{X^{\pm 1}\}/p^n O_F\{X^{\pm 1}\} \rightarrow \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle/p^n \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle$. Again, since $R/p^n R$ is étale over $O_F\{X^{\pm 1}\}/p^n O_F\{X^{\pm 1}\}$, we have a unique lift of $\bar{g}_n : R/p^n R \rightarrow \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle/p^n \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle$ in the commutative diagram

$$\begin{array}{ccc} O_F\{X^{\pm 1}\}/p^n O_F\{X^{\pm 1}\} & \xrightarrow{\quad} & \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle/p^n \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle \\ \downarrow & \nearrow \bar{g}_n & \downarrow \\ R/p^n R & \xrightarrow{\quad} & \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle/p\mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle. \end{array}$$

Via this lifting, the following diagram commutes

$$\begin{array}{ccc} R/p^{n+1} R & \xrightarrow{\quad} & \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle/p^{n+1} \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle \\ \downarrow & & \downarrow \\ R/p^n R & \xrightarrow{\quad} & \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle/p^n \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle, \end{array}$$

where the vertical arrows are natural projection maps. From the universal property of inverse limit of the right side of the diagram, we obtain a natural map of $O_F\{X^{\pm 1}\}$ -algebras

$$g : R \longrightarrow \lim_n \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle / p^n \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle = \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge.$$

Now, let $\bar{\theta} : \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle / p \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle \rightarrow R[\varpi] / pR[\varpi]$ denote the reduction of θ modulo p . Recall that by construction, $\bar{\theta} \circ \bar{g}$ is the inclusion of R/pR in $R[\varpi] / pR[\varpi]$. Therefore, the reduction modulo p of $\theta \circ g$ and the natural inclusion $R \hookrightarrow R[\varpi]$ coincide. Since R is p -torsion free, arguing as above we obtain that for each $n \in \mathbb{N}$, the natural inclusion and $\theta \circ g$ coincide modulo p^n .

Next, by $\mathbf{A}_{R,\varpi}^+$ -linearity, g can be extended to a map $g : R \otimes_{O_F} \mathbf{A}_{R,\varpi}^+ \rightarrow \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge$. From the discussion above and the definition of θ_R , we have that θ_R coincides with the homomorphism $\theta \circ g : R \otimes_{O_F} \mathbf{A}_{R,\varpi}^+ \rightarrow R[\varpi]$. In particular, $g(\text{Ker } \theta_R) \subset \text{Ker } \theta \subset \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge$. Since $\text{Ker } \theta$ contains divided powers, the map g extends to a map

$$g : (R \otimes_{O_F} \mathbf{A}_{R,\varpi}^+)[x^{[n]}, x \in \text{Ker } \theta_R, n \in \mathbb{N}] \longrightarrow \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge.$$

Finally, since $\mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge$ is p -adically complete, g extends to a map $g : \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \rightarrow \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge$.

Now by uniqueness of $g : R \rightarrow \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge$, the composition

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \xrightarrow{g} \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge \xrightarrow{f^{\text{PD}}} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}},$$

coincides with the identity over $R \subset \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$. Since it also coincides with identity on the image of $\mathbf{A}_{R,\varpi}^+$ (by $\mathbf{A}_{R,\varpi}^+$ -linearity), we obtain that $f^{\text{PD}} \circ g = \text{id}$ over $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$. Similarly, the homomorphism $g \circ f^{\text{PD}}$ coincides with identity over $\mathbf{A}_{R,\varpi}^+$ as well as over $O_F\{X^{\pm 1}\}$ (since g lifts the map $O_F\{X^{\pm 1}\} \rightarrow \mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge$), therefore it is identity over $\mathbf{A}_{R,\varpi}^{\text{PD}}\langle T \rangle^\wedge$. This establishes that f^{PD} is an isomorphism of rings. \blacksquare

Remark 4.21. We can give an alternative construction of the ring $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$. Note that we have a ring homomorphism $\iota : R \rightarrow \mathbf{A}_{R,\varpi}^{\text{PD}}$, where $X_i \mapsto [X_i^\flat]$ for $1 \leq i \leq d$. As in Definition 3.19, we define a map $g : R \otimes_{\mathbb{Z}} \mathbf{A}_{R,\varpi}^{\text{PD}} \rightarrow \mathbf{A}_{R,\varpi}^{\text{PD}}$, where $x \otimes y \mapsto \iota(x)y$. We obtain that $\text{Ker } g = (X_i \otimes 1 - 1 \otimes [X_i^\flat], \text{ for } 1 \leq i \leq d) \subset \text{Ker } \theta_R \subset \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})$. Since $R \otimes_{\mathbb{Z}} \mathbf{A}_{R,\varpi}^{\text{PD}}$ already contains divided powers of ξ , from Definition 4.19 we obtain that the p -adic completion of the divided power envelope of $R \otimes_{\mathbb{Z}} \mathbf{A}_{R,\varpi}^{\text{PD}}$ with respect to $\text{Ker } g$ is the same as $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$.

There is a natural filtration over $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ by Γ_R -stable submodules:

Definition 4.22. Let $U_i := \frac{1 \otimes [X_i^\flat]}{X_i \otimes 1}$ for $1 \leq i \leq d$ and $r \in \mathbb{Z}$, define the filtration over $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ as

$$\text{Fil}^r \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} := \left\langle (a \otimes b) \prod_{i=1}^d (U_i - 1)^{[k_i]} \in \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}, \text{ such that } a \in R, b \in \text{Fil}^j \mathbf{A}_{R,\varpi}^{\text{PD}}, \text{ and } j + \sum_i k_i \geq r \right\rangle.$$

Remark 4.23. The filtration over $\mathbf{A}_{R,\varpi}^{\text{PD}}$ (via its identification with R_ϖ^{PD} , see §3.3 and Definition 3.11) coincides with the filtration induced from its embedding in $\mathbf{A}_{\text{cris}}(\overline{R})$. Indeed, in both cases we have $\text{Fil}^r \mathbf{A}_{R,\varpi}^{\text{PD}} = (\xi^{[k]}, k \leq r) \subset \mathbf{A}_{R,\varpi}^{\text{PD}}$ for $r \geq 0$, whereas $\text{Fil}^r \mathbf{A}_{R,\varpi}^{\text{PD}} = \mathbf{A}_{R,\varpi}^{\text{PD}}$ for $r < 0$. Next, the filtration on $\mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})$ is defined as the induced filtration from its embedding inside $\mathcal{O}\mathbf{B}_{\text{dR}}^+(\overline{R})$ and the filtration on the latter ring is given by powers of $\text{Ker } \theta_R$ (see §2.1 & 2.2 for definition and notation). The induced filtration over $\mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})$ is therefore given by divided powers of the ideal $\text{Ker } \theta_R \subset \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})$. Since the filtration over $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ in Definition 4.22 is again given by divided powers of the ideal $\text{Ker } \theta_R \subset \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$, we infer that this filtration coincides with the one induced by its embedding into $\mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})$.

Lemma 4.24. (i) *The action of $\Gamma_{R,\varpi}$ is trivial on $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}/\pi$, whereas $\Gamma_R/\Gamma_{R,\varpi}$ acts trivially over $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}/\pi_m$.*

(ii) *We have $(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}})^{\Gamma_R} = R$ and $(\text{Fil}^1 \mathcal{O}\mathbf{A}_R^{\text{PD}})^{\Gamma_R} = 0$.*

Proof. (i) The first part follows from the definition of $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ and the action of $\Gamma_{R,\varpi}$ on $\mathbf{A}_{R,\varpi}^{\text{PD}}$ (see Lemma 3.16). The second part follows from observing that $\Gamma_R/\Gamma_{R,\varpi} = \Gamma_F/\Gamma_K$ is a finite cyclic group of order $[K:F] = p^{m-1}(p-1)$, and a lift $g \in \Gamma_R$ of a generator of $\Gamma_R/\Gamma_{R,\varpi}$ acts as $g(\pi_m) = (1 + \pi_m)^{\chi(g)} - 1$.

(ii) This is straightforward, since $R \subset (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}})^{\Gamma_R} \subset (\mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R}))^{G_R} = R$ and $(\text{Fil}^1 \mathcal{O}\mathbf{A}_R^{\text{PD}})^{\Gamma_R} \subset (\text{Fil}^1 \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}))^{G_R} \subset (\text{Fil}^1 \mathcal{O}\mathbf{B}_{\text{dR}}(\overline{R}))^{G_R} = 0$ (for last equality see the proof of [Bri08, Proposition 5.2.12]). ■

Next we consider a connection over $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ induced by the connection on $\mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})$,

$$\partial : \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \longrightarrow \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes \Omega_R^1,$$

where we have $\partial(X_i \otimes 1 - 1 \otimes [X_i^b])^{[n]} = (X_i \otimes 1 - 1 \otimes [X_i^b])^{[n-1]} dX_i$. This connection over $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ satisfies Griffiths transversality with respect to the filtration since it does so over $\mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})$.

4.3.2. Main result.

Theorem 4.25. *Let V be a positive finite q -height representation of G_R , then*

- (i) *V is a positive crystalline representation.*
- (ii) *Let $M := (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))^{\Gamma_R}$, then after extending scalars to $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ and inverting p , we obtain a natural isomorphism*

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M[\frac{1}{p}] \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V),$$

compatible with Frobenius, filtration, connection and the action of Γ_R on each side.

- (iii) *We have an isomorphism of $R[\frac{1}{p}]$ -modules*

$$\mathcal{O}\mathbf{D}_{\text{cris}}(V) \xleftarrow{\sim} (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))^{\Gamma_R}[\frac{1}{p}],$$

compatible with Frobenius, filtration, and connection on each side. Therefore, we obtain a comparison isomorphism

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V) \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R \mathcal{O}\mathbf{D}_{\text{cris}}(V),$$

compatible with Frobenius, filtration, connection and the action of Γ_R on each side.

Remark 4.26. The statement of Theorem 4.25 can be seen an analogue of the result of Berger [Ber04, Proposition II.2.1] (see the discussion after Proposition 4.7).

Recall that from Definition 4.9 any finite q -height representation is a twist of a positive finite q -height representation by $\mathbb{Q}_p(r)$, for $r \in \mathbb{N}$. Since twist by $\mathbb{Q}_p(r)$ of crystalline representations are again crystalline, we obtain that:

Corollary 4.27. *All finite q -height representations of G_R are crystalline.*

The proof of Theorem 4.25 will proceed in two steps: First, we will describe a process by which we can recover a submodule of $\mathcal{OD}_{\text{cris}}(V)$ starting from the Wach module (see Proposition 4.28), here we establish the comparison displayed in (ii). Next, the remaining claims made in the theorem are shown by exploiting some properties of Wach modules and the comparison obtained in the first step.

In §4.6, we will explicitly state the structure of Wach module attached to a one-dimensional finite q -height representation and we will also show that all one-dimensional crystalline representations are of finite q -height and one can recover $\mathcal{OD}_{\text{cris}}(V)$ starting with the Wach module $\mathbf{N}(V)$. Combining this with the theorem above, we will obtain that the notion of crystalline representations and finite q -height representations coincide in dimension 1.

4.4. From (φ, Γ) -modules to (φ, ∂) -modules. The objective of this section is to prove the following:

Proposition 4.28. *Let V be an h -dimensional positive finite q -height representation of G_R , $T \subset V$ a \mathbb{Z}_p -lattice of rank h stable under the action of G_R and $\mathbf{N}(T)$ the associated Wach module. Then*

- (i) $M := (\mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))^{\Gamma_R}$ is a finitely generated R -module contained in $\mathcal{OD}_{\text{cris}}(V)$.
- (ii) $M[\frac{1}{p}]$ is a finitely generated projective $R[\frac{1}{p}]$ -module of rank h and the natural inclusion

$$\mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_R M[\frac{1}{p}] \longrightarrow \mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V),$$

is an isomorphism compatible with Frobenius, filtration, connection and the action of Γ_R .

- (iii) If $\mathbf{N}(T)$ is free over \mathbf{A}_R^+ then there exists a free R -module $M_0 \subset M$ such that $M_0[\frac{1}{p}] = M[\frac{1}{p}]$ are free modules of rank h over $R[\frac{1}{p}]$.

Proof. We will use the notation of Definition 4.9 without repeating them. The first claim is easy to establish. Since $H_R = \text{Gal}(\overline{R}[\frac{1}{p}]/R_\infty[\frac{1}{p}])$, therefore we can write

$$\begin{aligned} M &= (\mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))^{\Gamma_R} \subset (\mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{D}^+(T))^{\Gamma_R} \subset (\mathcal{OA}_{\text{cris}}(\overline{R})^{H_R} \otimes_{\mathbf{A}_R^+} \mathbf{D}^+(T))^{\Gamma_R} \\ &\subset (\mathcal{OA}_{\text{cris}}(\overline{R})^{H_R} \otimes_{\mathbf{A}_R^+} (\mathbf{A}^+ \otimes_{\mathbb{Z}_p} T)^{H_R})^{\Gamma_R} \subset (\mathcal{OA}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Z}_p} T)^{G_R} \subset \mathcal{OD}_{\text{cris}}(V). \end{aligned} \tag{4.1}$$

The module $(\mathcal{OA}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Z}_p} T)^{G_R}$ is finitely generated over R . Since R is Noetherian, M is finitely generated.

Independently, we have that $R[\frac{1}{p}]$ is Noetherian and $\mathcal{OD}_{\text{cris}}(V)$ is a finitely generated $R[\frac{1}{p}]$ -module, therefore $M[\frac{1}{p}] \subset \mathcal{OD}_{\text{cris}}(V)$ is finitely generated over $R[\frac{1}{p}]$. Moreover, the module $\mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ is equipped with an $\mathbf{A}_{R,\varpi}^{\text{PD}}$ -linear and integrable connection $\partial_N = \partial \otimes 1$, where ∂ is the connection on $\mathcal{OA}_{R,\varpi}^{\text{PD}}$ described after Lemma 4.24. Therefore, we can consider the induced connection on $M[\frac{1}{p}]$, which is integrable since it is integrable over $\mathcal{OA}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$. This connection is compatible with the one on $\mathcal{OD}_{\text{cris}}(V)$ since the connection over $\mathcal{OA}_{R,\varpi}^{\text{PD}}$ is induced from the connection over $\mathcal{OA}_{\text{cris}}(\overline{R})$. So by [Bri08, Proposition 7.1.2] we obtain that $M[\frac{1}{p}]$ must be projective of rank $\leq h$. Furthermore, the inclusion $M[\frac{1}{p}] \subset \mathcal{OD}_{\text{cris}}(V)$ is compatible with natural Frobenius on each module since all the inclusions in (4.1) are compatible with Frobenius.

Next, we will show that the rank of $M[\frac{1}{p}]$ as a projective $R[\frac{1}{p}]$ -module is exactly h . But first let us prove that it is enough to show that the rank is h after a finite étale extension of R . Let us consider R' to be a finite étale extension of R such that the corresponding scalar extension

$\mathbf{A}_{R'}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ is a free module of rank h (see Definition 4.9) and $R'[\frac{1}{p}]/R[\frac{1}{p}]$ is Galois. The discussion of previous chapters hold for R' (see [Bri08, Chapitre 2] and [AI08, §2] for more on this). In particular, for $R'[\varpi]$ we have rings $\mathbf{A}_{R'}^+$, $\mathbf{A}_{R',\varpi}^+$, $\mathbf{A}_{R',\varpi}^{\text{PD}}$ and $\mathcal{O}\mathbf{A}_{R',\varpi}^{\text{PD}}$. Let $R'_\infty[\frac{1}{p}]$ denote the cyclotomic tower over $R'[\frac{1}{p}]$ and

$$\Gamma_{R'} = \text{Gal}(R'_\infty[\frac{1}{p}]/R'[\frac{1}{p}]) \text{ and } H_{R'} = \text{Ker}(G_{R'} \rightarrow \Gamma_{R'}).$$

Similarly, we have Galois groups $\Gamma_{R'}$ and $H_{R'}$. Let

$$G' := \text{Gal}(R'_\infty[\frac{1}{p}]/R_\infty[\frac{1}{p}]) = \text{Gal}(R'[\varpi][\frac{1}{p}]/R[\varpi][\frac{1}{p}]) = \text{Gal}(R'[\frac{1}{p}]/R[\frac{1}{p}]),$$

then we have that $H_{R,\varpi}/H_{R',\varpi} = H_R/H_{R'} = G'$. So we obtain that

$$\mathbf{A}_R^+ = (\mathbf{A}^+)^{H_R} = ((\mathbf{A}^+)^{H_{R'}})^{H_R/H_{R'}} = (\mathbf{A}_{R'}^+)^{G'}.$$

Moreover, for the base ring $R[\varpi]$ (instead of R) one can consider the ring \mathbf{A}_ϖ^+ as in Remark 3.6. Then we have

$$\mathbf{A}_{R,\varpi}^+ = (\mathbf{A}_\varpi^+)^{H_{R,\varpi}} = ((\mathbf{A}_\varpi^+)^{H_{R',\varpi}})^{H_{R,\varpi}/H_{R',\varpi}} = (\mathbf{A}_{R',\varpi}^+)^{G'}.$$

From these equalities and the description of the action of Γ_R on $\xi = \frac{\pi}{\pi_1}$, it is clear that

$$\mathbf{A}_{R,\varpi}^{\text{PD}} = (\mathbf{A}_{R',\varpi}^{\text{PD}})^{G'}, \text{ and therefore } \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} = (\mathcal{O}\mathbf{A}_{R',\varpi}^{\text{PD}})^{G'}.$$

Now, since $\mathbf{N}(T)$ is projective and G' acts trivially on it, we obtain that

$$\begin{aligned} (\mathcal{O}\mathbf{A}_{R',\varpi}^{\text{PD}} \otimes_{\mathbf{A}_{R'}^+} (\mathbf{A}_{R'}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)))^{G'} &= \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T) \\ (\mathcal{O}\mathbf{A}_{R',\varpi}^{\text{PD}} \otimes_{R'} (R' \otimes_R M[\frac{1}{p}]))^{G'} &= \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M[\frac{1}{p}]. \end{aligned}$$

In particular, base changing to $\mathbf{A}_{R'}^+$ to obtain $\mathbf{N}(T)$ as a free module is harmless. For the convenience in notation, below we will replace R' obtained in this manner by R and assume $\mathbf{N}(T)$ to be free over \mathbf{A}_R^+ .

In order to show that the rank of $M[\frac{1}{p}]$ is at least h , we will find Γ_R -fixed elements of $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ corresponding to a basis of $\mathbf{N}(T)$, which are linearly independent elements of $M[\frac{1}{p}]$. To carry this out, first we will define several new rings following [Wac96, §B.1] and examine their relation with $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$. After extending scalars of $\mathbf{N}(T)$, we will define differential operators on the obtained module, corresponding to the topological generators of Γ_R . Next, for any element of $\mathbf{N}(T)$, we will write down a corresponding element killed by the differential operators which we will show is fixed by Γ_R .

Remark 4.29. Note that the Γ_R -fixed elements of $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ can be obtained by successive approximation as well. This computation was carried out in [Abh21, §3.2.3].

4.4.1. Auxiliary rings and modules. For $n \in \mathbb{N}$, let us define a p -adically complete ring

$$S_n^{\text{PD}} := \mathbf{A}_R^+ \left\{ \frac{\pi}{p^n}, \frac{\pi^2}{2!p^{2n}}, \dots, \frac{\pi^k}{k!p^{kn}}, \dots \right\}.$$

Let $I_n^{[i]}$ denote the ideal of S_n^{PD} generated by $\frac{\pi^k}{k!p^{kn}}$ for $k \geq i$ and we set

$$\widehat{S}_n^{\text{PD}} := \lim_i S_n^{\text{PD}} / I_n^{[i]}. \quad (4.2)$$

Note that $\widehat{S}_n^{\text{PD}}$ is p -adically complete as well. Further, note that we can write $\varphi(\pi) = (1 + \pi)^p - 1 = \pi^p + p\pi x$ for some $x \in \mathbf{A}_F^+$, therefore

$$\begin{aligned} \frac{\varphi(\pi^k)}{k!p^{kn}} &= \frac{(\pi^p + p\pi x)^k}{k!p^{kn}} = \frac{\sum_{i=0}^k \binom{k}{i} \pi^{pi} \cdot (p\pi x)^{k-i}}{k!p^{kn}} \\ &= \sum_{i=0}^k \frac{(k + (p-1)i)! p^{i(n(p-1)-p)}}{i!(k-i)!} \cdot \frac{\pi^{k+(p-1)i} x^{k-i}}{(k + (p-1)i)! p^{(k+(p-1)i)(n-1)}} \in \widehat{S}_{n-1}^{\text{PD}} \end{aligned}$$

Using this, the Frobenius operator on S can be extended to a map $\varphi : \widehat{S}_n^{\text{PD}} \rightarrow \widehat{S}_{n-1}^{\text{PD}}$, which we will again call Frobenius. The ring $\widehat{S}_n^{\text{PD}}$ readily admits a continuous action of Γ_R which commutes with the Frobenius.

Lemma 4.30. *The ring $\widehat{S}_0^{\text{PD}}$ is a subring of $\mathbf{A}_{R,\varpi}^{\text{PD}}$, and therefore $\varphi^n(\widehat{S}_n^{\text{PD}}) \subset \mathbf{A}_{R,\varpi}^{\text{PD}}$.*

Proof. The first claim is true because we have

$$\pi_1^p \equiv \pi \pmod{p\mathbf{A}_{F,\varpi}^+}, \text{ which gives } \pi_1^{p^i} \equiv \pi^{p^{i-1}} \pmod{p^i\mathbf{A}_{F,\varpi}^+}.$$

So for $k \geq p^i$ we can write

$$\frac{\pi^k}{k!} = \frac{\xi^k \pi_1^k}{k!} = \frac{\xi^k}{k!} \pi_1^{k-p^i} (\pi^{p^{i-1}} + p^i a) = p^i a \pi_1^{k-p^i} \frac{\xi^k}{k!} + p^{i-1} \pi_1^{p^{i-1}} \frac{(k + p^{i-1})!}{k!p^{i-1}} \frac{\xi^{k+p^{i-1}}}{(k + p^{i-1})!} \in p^{i-1} \mathbf{A}_{F,\varpi}^{\text{PD}},$$

for some $a \in \mathbf{A}_{F,\varpi}^+$. Therefore, we get that $I_0^{[p^i]} \subset p^{i-1} \mathbf{A}_{R,\varpi}^{\text{PD}}$ and hence $\widehat{S}_0^{\text{PD}} \subset \mathbf{A}_{R,\varpi}^{\text{PD}}$. The second claim is obvious. \blacksquare

In the relative setting, we need slightly larger rings. Let us consider the O_F -linear homomorphism of rings

$$\begin{aligned} \iota : R &\longrightarrow \widehat{S}_n^{\text{PD}} \\ X_j &\longmapsto [X_j^{\flat}] \text{ for } 1 \leq j \leq d. \end{aligned}$$

Using ι we can define an O_F -linear morphism of rings

$$\begin{aligned} f : R \otimes_{O_F} \widehat{S}_n^{\text{PD}} &\longrightarrow \widehat{S}_n^{\text{PD}} \\ a \otimes b &\longmapsto \iota(a)b. \end{aligned}$$

Let $\mathcal{O}\widehat{S}_n^{\text{PD}}$ denote the p -adic completion of the divided power envelope of $R \otimes_{O_F} \widehat{S}_n^{\text{PD}}$ with respect to $\text{Ker } f$. Further, the morphism f extends uniquely to a continuous morphism $f : \mathcal{O}\widehat{S}_n^{\text{PD}} \rightarrow \widehat{S}_n^{\text{PD}}$. Now, it easily follows from the discussion in §3.4 that the kernel of the morphism f is generated by divided powers of the ideal generated by $(1 - V_1, \dots, 1 - V_d)$, where $V_j = \frac{X_j \otimes 1}{1 \otimes [X_j^{\flat}]}$

for $1 \leq j \leq d$. The Frobenius operator extends to $\mathcal{O}\widehat{S}_n^{\text{PD}}$ as well as the continuous action of Γ_R . From the discussion above we have $\varphi^n(\widehat{S}_n^{\text{PD}}) \subset \widehat{S}_0^{\text{PD}} \subset \mathbf{A}_{R,\varpi}^{\text{PD}}$, and following the description of $\mathcal{O}\widehat{S}_0^{\text{PD}}$ in §3.4 and of $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ from Remark 4.21, we obtain that

$$\mathcal{O}\widehat{S}_0^{\text{PD}} \subset \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \text{ and } \varphi^n(\mathcal{O}\widehat{S}_n^{\text{PD}}) \subset \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}.$$

Moreover, we have a canonical inclusion of $\widehat{S}_n^{\text{PD}} \subset \mathcal{O}\widehat{S}_n^{\text{PD}}$ compatible with all the structures.

Now let us take $n \in \mathbb{N}_{\geq 1}$ and consider the ring $\mathcal{O}\widehat{S}_n^{\text{PD}}$ below. Set

$$J := \left(\frac{\pi}{p^n}, 1 - V_1, \dots, 1 - V_d \right) \subset \mathcal{O}\widehat{S}_n^{\text{PD}},$$

and its divided power

$$J^{[i]} := \left\langle \frac{\pi^{[k_0]}}{p^{nk_0}} \prod_{j=1}^d (1 - V_j)^{[k_j]}, \mathbf{k} = (k_0, k_1, \dots, k_d) \in \mathbb{N}^{d+1} \text{ such that } \sum_{j=0}^d k_j \geq i \right\rangle \subset \mathcal{O}\hat{S}_n^{\text{PD}}.$$

By construction of $\mathcal{O}\hat{S}_n^{\text{PD}}$, it is clear that a summation $\sum_{i \in \mathbb{N}} x_i a_i$ with $a_i \in J^{[i]}$ and $x_i \in \hat{S}_n^{\text{PD}}$ goes to 0 as $i \rightarrow +\infty$, converges in $\mathcal{O}\hat{S}_n^{\text{PD}}$. Moreover, every $x \in \mathcal{O}\hat{S}_n^{\text{PD}}$ has a presentation as $x = \sum_{\mathbf{k} \in \mathbb{N}^{d+1}} x_{\mathbf{k}} \frac{\pi^{[k_0]}}{p^{nk_0}} \prod_{j=1}^d (1 - V_j)^{[k_j]}$, where $x_{\mathbf{k}} \in \mathbf{A}_R^+$ goes to 0 as $|\mathbf{k}| = \sum_j k_j \rightarrow +\infty$.

Next, we set

$$\mathcal{O}N_n^{\text{PD}} := \mathcal{O}\hat{S}_n^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T).$$

Again, $\mathcal{O}N_n^{\text{PD}}$ is p -adically complete and it is equipped with a Frobenius-semilinear operator $\varphi : \mathcal{O}\hat{S}_n^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T) \rightarrow \mathcal{O}\hat{S}_{n-1}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ and a continuous and semilinear action of Γ_R . Now recall that we fixed $m \in \mathbb{N}_{\geq 1}$ (fix $m \in \mathbb{N}_{\geq 2}$ if $p = 2$) such that $K = F(\zeta_{p^m})$. So we take

$$M' := (\mathcal{O}N_m^{\text{PD}})^{\Gamma'_R} \text{ and } M'' := (M')^{\Gamma_F} = (\mathcal{O}N_m^{\text{PD}})^{\Gamma_R}.$$

Since we assumed $\mathbf{N}(T)$ to be free, therefore $\mathcal{O}N_m^{\text{PD}}$ is a free $\mathcal{O}\hat{S}_m^{\text{PD}}$ -module of rank h . As we have $\varphi^m(\mathcal{O}\hat{S}_m^{\text{PD}}) \subset \mathcal{O}\hat{S}_{R,\varpi}^{\text{PD}}$ so we get that $\varphi^m(M'') \subset (\mathcal{O}\hat{S}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))^{\Gamma_R} = M$. Therefore, to show that the $R[\frac{1}{p}]$ -rank of $M[\frac{1}{p}]$ is at least h , it is enough to show that for each $x \in \mathbf{N}(T)$ there exists unique $x'' \in M'' \subset \mathcal{O}N_m^{\text{PD}}$ fixed by Γ_R and $x \equiv x'' \pmod{J^{[1]}\mathcal{O}N_m^{\text{PD}}}$ (see Lemma 4.43).

4.4.2. Infinitesimal action of Γ_R . From §3.1 recall that we have $\{\gamma, \gamma_1, \dots, \gamma_d\}$ as a set of topological generators of Γ_R such that $\{\gamma_1, \dots, \gamma_d\}$ generate Γ'_R topologically, and γ is a lift of a topological generator of Γ_F where $\gamma^e = \gamma_0$ is a lift of a topological generator of Γ_K , $e = [K : F]$ and $\chi(\gamma_0) = \exp(p^m)$ where we fixed $m \in \mathbb{N}_{\geq 1}$ (fix $m \in \mathbb{N}_{\geq 2}$ if $p = 2$). Further, we have the identity $\gamma_0 \gamma_i = \gamma_i^{\chi(\gamma_0)} \gamma_0$ for $1 \leq i \leq d$. In this section we will study the infinitesimal action of Γ_R on the rings and modules constructed in previous section.

Lemma 4.31. *Let $k \in \mathbb{N}$, $n \geq m$ and $i \in \{0, 1, \dots, d\}$. Then $(\gamma_i - 1)(p^m, \pi)^k \hat{S}_n^{\text{PD}} \subset (p^m, \pi)^{k+1} \hat{S}_n^{\text{PD}}$.*

Proof. First, let $i = 0$. Recall that we have $\chi(\gamma_0) = \exp(p^m) = 1 + p^m a \in 1 + p^m \mathbb{Z}_p$. So we can write

$$\begin{aligned} (\gamma_0 - 1)\pi &= (1 + \pi)^{\chi(\gamma_0)} - (1 + \pi) \\ &= (\chi(\gamma_0)\pi + \frac{\chi(\gamma_0)(\chi(\gamma_0)-1)}{2!}\pi^2 + \frac{\chi(\gamma_0)(\chi(\gamma_0)-1)(\chi(\gamma_0)-2)}{3!}\pi^3 + \dots) - \pi \\ &= (\chi(\gamma_0)u - 1)\pi, \end{aligned}$$

for some $u = 1 + \pi x \in 1 + \pi \mathbf{A}_R^+$. Therefore, $\chi(\gamma_0)u - 1 = p^m a + \pi x + p^m a \pi x \in (p^m, \pi) \mathbf{A}_R^+$ which gives us that $(\gamma_0 - 1)\pi \in (p^m, \pi) \pi \mathbf{A}_R^+$. Now we have $(\gamma_0 - 1) \mathbf{A}_R^+ \subset \pi \mathbf{A}_R^+ \subset (p^m, \pi) \mathbf{A}_R^+$, so proceeding by induction on $k \geq 1$ and using the fact that $\gamma_0 - 1$ acts as a twisted derivation (i.e. $(\gamma_0 - 1)xy = (\gamma_0 - 1)x \cdot y + \gamma_0(x)(\gamma_0 - 1)y$ for $x, y \in \mathbf{A}_R^+$), we conclude that

$$(\gamma_0 - 1)(p^m, \pi)^k \mathbf{A}_R^+ \subset (p^m, \pi)^{k+1} \mathbf{A}_R^+.$$

Next, any $f \in \hat{S}_n^{\text{PD}}$ can be written as $f = \sum_{s \in \mathbb{N}} f_s \frac{\pi^s}{s! p^{ns}}$ such that $f_s \in \mathbf{A}_R^+$ goes to 0 as $s \rightarrow +\infty$. Clearly we have

$$(\gamma_0 - 1) \frac{\pi^s}{s! p^{ns}} = \frac{(\chi(\gamma_0)^s u^s - 1) \pi^s}{s! p^{ns}} \in (p^m, \pi) \frac{\pi^s}{s! p^{ns}} \hat{S}_n^{\text{PD}}.$$

Combining the discussion for \mathbf{A}_R^+ and $\frac{\pi^s}{s!p^{ns}}$, using induction on $k \geq 1$ and using the fact that $\gamma_0 - 1$ acts as a twisted derivation, we conclude that

$$(\gamma_0 - 1)(p^m, \pi)^k \widehat{S}_n^{\text{PD}} \subset (p^m, \pi)^{k+1} \widehat{S}_n^{\text{PD}}.$$

Finally, for $i \in \{1, \dots, d\}$ we have $(\gamma_i - 1)[X_i^\flat] = \pi[X_i^\flat] \in (p^m, \pi)\mathbf{A}_R^+$ and $(\gamma_i - 1)([X_i^\flat]^{-1}) = -\pi(1 + \pi)^{-1}[X_i^\flat]^{-1} \in (p^m, \pi)\mathbf{A}_R^+$. Again by induction on $k \geq 1$ and using the fact that $\gamma_i - 1$ acts as a twisted derivation, we get that

$$(\gamma_i - 1)(p^m, \pi)^k \mathbf{A}_R^+ \subset (p^m, \pi)^{k+1} \mathbf{A}_R^+.$$

Now any $f \in \widehat{S}_n^{\text{PD}}$ can be written as $f = \sum_{s \in \mathbb{N}} f_s \frac{\pi^s}{s!p^{ns}}$ such that $f_s \in \mathbf{A}_R^+$ goes to 0 as $s \rightarrow +\infty$, and γ_i acts trivially on π for $1 \leq i \leq d$, so we conclude that

$$(\gamma_i - 1)(p^m, \pi)^k \widehat{S}_n^{\text{PD}} \subset (p^m, \pi)^{k+1} \widehat{S}_n^{\text{PD}}.$$

■

Lemma 4.32. *For $n \geq m$ and $i \in \{0, 1, \dots, d\}$ the operators*

$$\nabla_i := \log \gamma_i = \sum_{k \in \mathbb{N}} (-1)^k \frac{(\gamma_i - 1)^{k+1}}{k+1},$$

converge as series of operators on $\widehat{S}_n^{\text{PD}}$.

Proof. From Lemma 4.31, we have that for $k \in \mathbb{N}$

$$(\gamma_i - 1)(p^m, \pi)^k \widehat{S}_n^{\text{PD}} \subset (p^m, \pi)^{k+1} \widehat{S}_n^{\text{PD}}.$$

Therefore, using the fact that $\gamma_i - 1$ acts as a twisted derivation (i.e. $(\gamma_i - 1)xy = (\gamma_i - 1)x \cdot y + \gamma_i(x)(\gamma_i - 1)y$ for $x, y \in \widehat{S}_n^{\text{PD}}$), we obtain that for $x \in \widehat{S}_n^{\text{PD}}$

$$(\gamma_i - 1)^{k+1}(x) \subset (p^m, \pi)^{k+1} \widehat{S}_n^{\text{PD}}. \quad (4.3)$$

Therefore, the following series converges in $\widehat{S}_n^{\text{PD}}$

$$\nabla_i(x) = \sum_{k \in \mathbb{N}} (-1)^k \frac{(\gamma_i - 1)^{k+1}(x)}{k+1}.$$

This allows us to conclude. ■

Remark 4.33. Note that Γ_R acts trivially modulo π on \mathbf{A}_R^+ . Therefore, we also get that it acts trivially modulo π over $\widehat{S}_n^{\text{PD}}$. Hence, for $0 \leq i \leq d$ we have $\nabla_i(\widehat{S}_n^{\text{PD}}) \subset \pi \widehat{S}_n^{\text{PD}} = t \widehat{S}_n^{\text{PD}}$, where the last equality follow from the fact that $\frac{t}{\pi}$ is a unit in $\widehat{S}_n^{\text{PD}}$ (see Lemma 4.35 below).

Remark 4.34. The operators ∇_i for $0 \leq i \leq d$, defined in Lemma 4.32, describe the action of the Lie algebra $\text{Lie } \Gamma_R$ on $\widehat{S}_n^{\text{PD}}$, i.e. ∇_i acts as a differential operator on $\widehat{S}_n^{\text{PD}}$.

Lemma 4.35. *$\frac{t}{\pi}$ is a unit in $\widehat{S}_n^{\text{PD}}$ for $n \geq m$.*

Proof. We can write the fraction

$$\frac{t}{\pi} = \frac{\log(1 + \pi)}{\pi} = \sum_{k \geq 0} (-1)^k \frac{\pi^k}{k+1}.$$

Formally, we can write

$$\frac{\pi}{t} = \frac{\pi}{\log(1 + \pi)} = b_0 + b_1 \pi + b_2 \pi^2 + b_3 \pi^3 + \dots,$$

where $b_0 = 1$ and $v_p(b_k) \geq -\frac{k}{p-1}$ for all $k \geq 1$. But rewriting the series as a power series in $\frac{\pi^k}{k!p^{nk}}$, we get that

$$\frac{\pi}{t} = \sum_{k \in \mathbb{N}} b_k k! p^{nk} \frac{\pi^k}{k! p^{nk}}.$$

The p -adic valuation of coefficients in the series above is given as

$$v_p(b_k k! p^{nk}) \geq \frac{-k}{p-1} + nk + v_p(k!) = \frac{p-2}{p-1} nk + v_p(k!),$$

which clearly goes to $+\infty$ as $k \rightarrow +\infty$. Hence, $\frac{\pi}{t}$ converges in $\widehat{S}_n^{\text{PD}}$ and is an inverse to $\frac{t}{\pi}$. \blacksquare

Now let us consider the ring $\mathcal{O}\widehat{S}_n^{\text{PD}}$ and divided power ideals

$$J^{[i]} := \left\langle \frac{\pi^{[k_0]}}{p^{nk_0}} \prod_{j=1}^d (1 - V_j)^{[k_j]}, \mathbf{k} = (k_0, k_1, \dots, k_d) \in \mathbb{N}^{d+1} \text{ such that } \sum_{j=0}^d k_j \geq i \right\rangle \subset \mathcal{O}\widehat{S}_n^{\text{PD}}.$$

Arguments similar to Lemmas 4.31 and 4.32 show that for $0 \leq i \leq d$ the series of operators $\nabla_i = \log \gamma_i = \sum_{k \in \mathbb{N}} (-1)^k \frac{\pi^{k+1}}{k+1}$ converge over $\mathcal{O}\widehat{S}_n^{\text{PD}}$. Moreover, from Remark 4.33 we obtain that for $0 \leq i \leq d$, we have $\nabla_i(\mathcal{O}\widehat{S}_n^{\text{PD}}) \subset t\mathcal{O}\widehat{S}_n^{\text{PD}}$. Also, it is easy to observe that we have $\nabla_0(t) = \log(\chi(\gamma_0))t = p^m t$ and $\nabla_i(V_i) = tV_i$ for $1 \leq i \leq d$. Finally, recall that $\gamma_i \gamma_j = \gamma_j \gamma_i$ for $1 \leq i, j \leq d$ and $\gamma_0 \gamma_i = \gamma_i^{\chi(\gamma_0)} \gamma_0$, therefore we conclude that

$$\begin{aligned} [\nabla_i, \nabla_j] &= 0, \\ [\nabla_i, \nabla_0] &= \log(\chi(\gamma_0)) \nabla_i = p^m \nabla_i. \end{aligned}$$

Now we will adapt the discussion above to scalar extension of Wach module $\mathbf{N}(T)$ to $\mathcal{O}\widehat{S}_n^{\text{PD}}$, i.e. for $\mathcal{O}N_n^{\text{PD}} := \mathcal{O}\widehat{S}_n^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$.

Lemma 4.36. *For $n \geq m$ and $i \in \{0, 1, \dots, d\}$ the operators*

$$\nabla_i = \log \gamma_i = \sum_{k \in \mathbb{N}} (-1)^{k+1} \frac{(\gamma_i - 1)^{k+1}}{k+1}$$

converge as series of operators on $\mathcal{O}N_n^{\text{PD}}$.

Proof. For $0 \leq i \leq d$, observe that $\gamma_i - 1$ acts as a twisted derivation, i.e. for $a \in \mathcal{O}\widehat{S}_n^{\text{PD}}$ and $x \in \mathbf{N}(T)$, we have

$$(\gamma_i - 1)(ax) = (\gamma_i - 1)a \cdot x + \gamma_i(a)(\gamma_i - 1)x.$$

The action of Γ_R is trivial on $\mathbf{N}(T)/\pi\mathbf{N}(T)$, so we can write $(\gamma_i - 1)x = \pi y$, for some $y \in \mathbf{N}(T)$, i.e. $(\gamma_i - 1)\mathcal{O}N_n^{\text{PD}} \subset (p^m, \pi)\mathcal{O}N_n^{\text{PD}}$. From the proof of Lemma 4.32 and (4.3) and induction over $k \geq 1$, it follows that

$$(\gamma_i - 1)(p^m, \pi)^k \mathcal{O}N_n^{\text{PD}} \subset (p^m, \pi)^{k+1} \mathcal{O}N_n^{\text{PD}}.$$

Next, using the fact that $\gamma_i - 1$ acts as a twisted derivation, we obtain that

$$(\gamma_i - 1)^{k+1}(ax) \subset (p^m, \pi)^{k+1} \mathcal{O}N_n^{\text{PD}}.$$

Therefore, the following series converges in $\mathcal{O}N_n^{\text{PD}}$

$$\nabla_i(ax) = \sum_{k \in \mathbb{N}} (-1)^k \frac{(\gamma_i - 1)^{k+1}(ax)}{k+1}.$$

This allows us to conclude. \blacksquare

Remark 4.37. Note that Γ_R acts trivially modulo π on $\mathcal{O}\widehat{S}_n^{\text{PD}}$ and $\mathbf{N}(T)$. Therefore, we also get that it acts trivially modulo π over $\mathcal{O}N_n^{\text{PD}}$. Hence, for $0 \leq i \leq d$ we have $\nabla_i(\mathcal{O}N_n^{\text{PD}}) \subset \pi\mathcal{O}N_n^{\text{PD}} = t\mathcal{O}N_n^{\text{PD}}$, where the last equality follows from the fact that $\frac{t}{\pi}$ is a unit in $\mathcal{O}\widehat{S}_n^{\text{PD}}$ (see Lemma 4.35).

Again, over $\mathcal{O}N_n^{\text{PD}}$ we have

$$\begin{aligned} [\nabla_i, \nabla_j] &= 0, \\ [\nabla_i, \nabla_0] &= \log(\chi(\gamma_0))\nabla_i = p^m\nabla_i, \end{aligned}$$

which enables us to define differential operators ∂_i over $\mathcal{O}N_n^{\text{PD}}$ using the formula

$$\partial_i = \begin{cases} -t^{-1}\nabla_0 & \text{for } i = 0, \\ t^{-1}V_i^{-1}\nabla_i & \text{for } 1 \leq i \leq d, \end{cases}$$

where $V_i = \frac{X_i \otimes 1}{1 \otimes [X_i^b]}$ for $1 \leq i \leq d$. Note that ∂_i is well defined since $\nabla_i(\mathcal{O}N_n^{\text{PD}}) \subset t\mathcal{O}N_n^{\text{PD}}$ (see Remark 4.37).

Lemma 4.38. *For $n \geq m$, the differential operators defined on $\mathcal{O}N_n^{\text{PD}}$ commute, i.e. $\partial_i \circ \partial_j = \partial_j \circ \partial_i$ for $0 \leq i, j \leq d$.*

Proof. From above we have $[\nabla_i, \nabla_j] = 0$ for $1 \leq i, j \leq d$, whereas $[\nabla_0, \nabla_i] = p^m\nabla_i$, for $1 \leq i \leq d$. So it follows that over $\mathcal{O}N_n^{\text{PD}}$ we have the composition of operators

$$t^2V_iV_j(\partial_i \circ \partial_j - \partial_j \circ \partial_i) = tV_i\partial_i \circ tV_j\partial_j - tV_j\partial_j \circ tV_i\partial_i = \nabla_i \circ \nabla_j - \nabla_j \circ \nabla_i = 0, \text{ for } 1 \leq i, j \leq d.$$

Next, for $1 \leq i \leq d$, we have

$$\begin{aligned} \nabla_0 \circ \nabla_i - \nabla_i \circ \nabla_0 &= -t\partial_0 \circ (tV_i\partial_i) + tV_i\partial_i \circ (t\partial_0) \\ &= -p^mtV_i\partial_i - t^2V_i\partial_0 \circ \partial_i + t^2V_i\partial_i \circ \partial_0 = p^m\nabla_i - t^2V_i(\partial_0 \circ \partial_i - \partial_i \circ \partial_0). \end{aligned}$$

In particular, $\partial_i \circ \partial_j - \partial_j \circ \partial_i = 0$ for $0 \leq i, j \leq d$ since $\mathcal{O}N_n^{\text{PD}}$ is t -torsion free. ■

For the rest of the section, let us now assume $n = m$.

Lemma 4.39. *Let $1 \leq i \leq d$ and $x \in \mathbf{N}(T)$, then we have that $\partial_i^k(x) \rightarrow 0$ in $\mathcal{O}N_m^{\text{PD}}$ as $k \rightarrow +\infty$.*

Proof. First, let us note that since $\partial_i(V_i) = 1$, $\partial_i(V_j) = 0$ for $j \neq i$ and $\partial_i(\pi) = 0$, so we have that $\partial_i^p(\mathcal{O}\widehat{S}_m^{\text{PD}}) \subset p\mathcal{O}\widehat{S}_m^{\text{PD}}$. Moreover, an easy computation shows that for $x \in \mathbf{N}(T)$ we have

$$\partial_i(\varphi(x)) = \frac{\nabla_i(\varphi(x))}{tV_i} = \frac{\varphi(\nabla_i(x))}{tV_i} = pV_i^{p-1}\varphi(\partial_i(x)) \in \mathcal{O}\widehat{S}_m^{\text{PD}} \otimes_{\varphi(\mathbf{A}_R^+)} \varphi(\mathbf{N}(T)),$$

where note that we have $\varphi(\partial_i(x)) \in \varphi(\mathcal{O}\widehat{S}_{m+1}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)) \subset \mathcal{O}\widehat{S}_m^{\text{PD}} \otimes_{\varphi(\mathbf{A}_R^+)} \varphi(\mathbf{N}(T))$ since $\partial_i(x)$ converges over $\mathcal{O}\widehat{S}_{m+1}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ by Lemma 4.36.

Next, from Definition 4.9 recall that we have $q^s\mathbf{N}(T) \subset \varphi^*(\mathbf{N}(T))$. Let us write $q^s x = \sum_{j=1}^h a_j \varphi(e_j)$ for $a_j \in \mathbf{A}_R^+$ and $\{e_1, \dots, e_h\}$ an \mathbf{A}_R^+ -basis of $\mathbf{N}(T)$. Then it follows that $\partial_i^p(q^s x) \in pq^s(\mathcal{O}\widehat{S}_m^{\text{PD}} \otimes_{\varphi(\mathbf{A}_R^+)} \varphi(\mathbf{N}(T)))$, therefore $\partial_i^p(x) \in p(\mathcal{O}\widehat{S}_m^{\text{PD}} \otimes_{\varphi(\mathbf{A}_R^+)} \varphi(\mathbf{N}(T)))$. By induction on k we see that $\partial_i^{pk}(x) \in p^k(\mathcal{O}\widehat{S}_m^{\text{PD}} \otimes_{\varphi(\mathbf{A}_R^+)} \varphi(\mathbf{N}(T))) \subset p^k\mathcal{O}N_m^{\text{PD}}$. Hence, the claim follows. ■

Remark 4.40. Note that one can recover the action of γ_i using the differential operator ∂_i . For $i \in \{1, \dots, d\}$ we have $\gamma_i = \exp(tV_i\partial_i)$, whereas for $i = 0$ we have $\gamma_0 = \exp(-t\partial_0)$.

From the remark above it is clear that for $0 \leq i \leq d$ and $x \in \mathcal{O}N_m^{\text{PD}}$ we have $\gamma_i(x) = x$ if and only if $\partial_i(x) = 0$.

Lemma 4.41. *For any $x \in \mathbf{N}(T)$ there exists a unique $x'' \in \mathcal{O}N_m^{\text{PD}}$ such that*

$$\begin{aligned} x'' &\equiv x \pmod{J^{[1]}\mathcal{O}N_m^{\text{PD}}}, \\ \gamma_i(x'') &= x'' \quad \text{for } 0 \leq i \leq d. \end{aligned}$$

In particular, $x'' \in M'' = (\mathcal{O}N_m^{\text{PD}})^{\Gamma_R}$.

Proof. For $x \in \mathbf{N}(T)$, we set

$$x' = \sum_{\mathbf{k} \in \mathbb{N}^d} \partial_1^{k_1} \circ \cdots \circ \partial_d^{k_d}(x)(1 - V_1)^{[k_1]} \cdots (1 - V_d)^{[k_d]} \in \mathcal{O}N_m^{\text{PD}}$$

The summation converges since for $1 \leq i \leq d$ we have that $\partial_0^{k_0} \circ \partial_1^{k_1} \circ \cdots \circ \partial_d^{k_d}(x) \rightarrow 0$ as $|\mathbf{k}| = \sum_{i=1}^d k_i \rightarrow +\infty$ from Lemma 4.39. Note that we have an isomorphism of rings $\widehat{S}_m^{\text{PD}} \xrightarrow{\sim} (\mathcal{O}\widehat{S}_m^{\text{PD}})^{\Gamma_{R'}}$ compatible with $\Gamma_R/\Gamma_{R'} = \Gamma_F$ -action. Therefore, by the description of $\widehat{S}_m^{\text{PD}}$ in (4.2) and since $x' \in (\mathcal{O}N_m^{\text{PD}})^{\Gamma_{R'}}$ we see that the following sum converges

$$x'' = \sum_{k_0 \in \mathbb{N}} \partial_0^{k_0}(x') \frac{t^{[k_0]}}{p^{mk_0}} \in \mathcal{O}N_m^{\text{PD}}.$$

Since the differential operators on $\mathcal{O}N_m^{\text{PD}}$ commute by Lemma 4.38, we get that

$$x'' = \sum_{\mathbf{k} \in \mathbb{N}^{d+1}} \partial_0^{k_0} \circ \partial_1^{k_1} \circ \cdots \circ \partial_d^{k_d}(x) \frac{t^{[k_0]}}{p^{mk_0}} (1 - V_1)^{[k_1]} \cdots (1 - V_d)^{[k_d]} \in \mathcal{O}N_m^{\text{PD}} \quad (4.4)$$

By the definition of x'' it is clear that $x'' \equiv x \pmod{J^{[1]}\mathcal{O}N_m^{\text{PD}}}$. Next, using the fact that $\partial_i \circ \partial_j = \partial_j \circ \partial_i$ for $0 \leq i, j \leq d$ (see Lemma 4.38) as well as $\partial_0(t) = -p^m$ and $\partial_i(V_i) = 1$ for $1 \leq i \leq d$, it is easy to deduce that $\partial_i(x'') = 0$ for $0 \leq i \leq d$. So by Remark 4.40, we get that $\gamma_i(x'') = x''$ for $0 \leq i \leq d$.

Uniqueness of x'' follows from Lemma 4.43. Finally, let $g \in \Gamma_F$ be a lift of a generator of the cyclic group Γ_F/Γ_K . Then we have that $g(x'') \in \mathcal{O}N_m^{\text{PD}}$ satisfies the conditions of the claim (since $(g - 1)x \in \pi\mathbf{N}(T) \subset J^{[1]}\mathcal{O}N_m^{\text{PD}}$). But by uniqueness, we obtain that $g(x'') = x''$, i.e. $x'' \in (\mathcal{O}N_m^{\text{PD}})^{\Gamma_R} = M''$. \blacksquare

Remark 4.42. Note that the lemma above can also be obtained by a “successive approximation” argument (see [Abh21, Lemmas 3.33 & 3.37]).

Following claim was used above:

Lemma 4.43. *For any $x \in \mathbf{N}(T)$ suppose there exists $x'' \in \mathcal{O}N_m^{\text{PD}}$ such that*

$$\begin{aligned} x'' &\equiv x \pmod{J^{[1]}\mathcal{O}N_m^{\text{PD}}}, \\ \gamma_i(x'') &= x'' \quad \text{for } 0 \leq i \leq d. \end{aligned}$$

Then x'' is unique.

Proof. Let $\{f_1, \dots, f_h\}$ denote an \mathbf{A}_R^+ -basis of $\mathbf{N}(T)$. Then $\{f_1, \dots, f_h\}$ is also an $\mathcal{O}\widehat{S}_m^{\text{PD}}$ -basis of $\mathcal{O}N_m^{\text{PD}}$. Now using the formula in (4.4), for all $1 \leq i \leq h$ let

$$f_i'' = \sum_{\mathbf{k} \in \mathbb{N}^{d+1}} \partial_0^{k_0} \circ \partial_1^{k_1} \circ \cdots \circ \partial_d^{k_d}(f_i) \frac{t^{[k_0]}}{p^{mk_0}} (1 - V_1)^{[k_1]} \cdots (1 - V_d)^{[k_d]} \in \mathcal{O}N_m^{\text{PD}}.$$

We want to show that $\{f_1'', \dots, f_h''\}$ also form an $\mathcal{O}\widehat{S}_m^{\text{PD}}$ -basis of $\mathcal{O}N_m^{\text{PD}}$. Let us write $f_i'' = f_i + \sum_{j=1}^h a_{ij} f_j$ with $a_{ij} \in J^{[1]}\mathcal{O}\widehat{S}_m^{\text{PD}}$ and let $A = id_h + (a_{ij}) \in \text{Mat}(h, \mathcal{O}\widehat{S}_m^{\text{PD}})$ denote the $h \times h$ matrix thus obtained. We have that $\det A = 1 + x$ with $x \in J^{[1]}\mathcal{O}\widehat{S}_m^{\text{PD}}$ and $1 - x + x^2 - x^3 + \cdots =$

$\sum_{n \in \mathbb{N}} (-1)^n n! x^{[n]}$ converges in $\mathcal{O}\hat{S}_m^{\text{PD}}$ as an inverse of $1 + x$, i.e. $\det A$ is invertible in $\mathcal{O}\hat{S}_m^{\text{PD}}$. Therefore, $\{f_1'', \dots, f_h''\}$ form a basis of $\mathcal{O}N_m^{\text{PD}}$.

Now for any $x \in \mathbf{N}(T)$, writing $x = \sum_{i=1}^h x_i f_i''$ and plugging into the formula (4.4) we obtain $x'' \in \mathcal{O}N_m^{\text{PD}}$ such that $x'' \equiv x \pmod{J^{[1]}\mathcal{O}N_m^{\text{PD}}}$ and $\gamma_j(x'') = x''$ for all $0 \leq j \leq d$. By linear independence of $\{f_1'', \dots, f_h''\}$ over $\mathcal{O}\hat{S}_m^{\text{PD}}$ we obtain that x'' is unique. \blacksquare

Remark 4.44. The uniqueness claim can also be established by a “successive approximation” argument (see [Abh21, p.63-p.65]).

Lemma 4.45. *We have $\mathcal{O}\hat{S}_m^{\text{PD}} \otimes_R M'' \xrightarrow{\sim} \mathcal{O}\hat{S}_m^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$.*

Proof. Let $\{f_1, \dots, f_h\}$ denote an \mathbf{A}_R^+ -basis of $\mathbf{N}(T)$. Then $\{f_1, \dots, f_h\}$ is also an $\mathcal{O}\hat{S}_m^{\text{PD}}$ -basis of $\mathcal{O}N_m^{\text{PD}}$. From the proof of Lemmas 4.41 & 4.43 we have $f_i'' \in M''$ for all $1 \leq i \leq h$, such that $\{f_1'', \dots, f_h''\}$ also form an $\mathcal{O}\hat{S}_m^{\text{PD}}$ -basis of $\mathcal{O}N_m^{\text{PD}}$. Therefore, $\mathcal{O}\hat{S}_m^{\text{PD}} \otimes_R M'' \xrightarrow{\sim} \mathcal{O}N_m^{\text{PD}}$. \blacksquare

4.4.3. Finishing the proof of Proposition 4.28. Recall that at the beginning of the proof we assumed $\mathbf{N}(T)$ to be free of rank h (after extension of scalars to $\mathbf{A}_{R'}^+$ which we again wrote as \mathbf{A}_R^+ by abusing notations), therefore $\mathcal{O}N_m^{\text{PD}}$ is free of rank h . Further, we have $M = (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))^{\Gamma_R}$ and since $M[\frac{1}{p}]$ is equipped with an integrable connection, it is projective of rank $\leq h$ (see the beginning of the proof). So applying Lemma 4.41 to a basis of $\mathbf{N}(T)$, we obtain that the rank of $M[\frac{1}{p}]$ as an $R[\frac{1}{p}]$ -module is exactly h .

Next, we want to show that the natural inclusion $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M[\frac{1}{p}] \hookrightarrow \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)$ is bijective. To show this claim, we require the following lemma:

Lemma 4.46. *We have $\varphi^*(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)) \xrightarrow{\sim} (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V))$.*

Proof. Recall that we are working under the assumption that $\mathbf{N}(V)$ is free and by definition of a positive finite q -height representation we have that the cokernel of the inclusion $\varphi^*(\mathbf{N}(V)) \rightarrow \mathbf{N}(V)$ is killed by q^s where $s \in \mathbb{N}$ is the height of the representation V . Extending scalars to $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$, we obtain that the cokernel of the inclusion $\varphi^*(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)) \rightarrow (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V))$ is killed by q^s . Now note that we have $q = \frac{\varphi(\pi)}{\pi} = p\varphi(\frac{\pi}{t})\frac{t}{\pi}$ where $\frac{t}{\pi}$ is a unit in $\mathbf{A}_{R,\varpi}^{\text{PD}}$ (see Lemma 3.14), i.e. p and q are associates in $\mathbf{A}_{R,\varpi}^{\text{PD}}$. Therefore, the cokernel of the inclusion in the claim is killed by p^s . But, p is invertible in $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}[\frac{1}{p}]$. Hence, we obtain that $\varphi^*(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)) \xrightarrow{\sim} (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V))$. \blacksquare

Since we assumed $\mathbf{N}(T)$ to be a free module, let $\{f_1, \dots, f_h\}$ be its \mathbf{A}_R^+ -basis. Let $P \in \text{Mat}(h, \mathbf{A}_R^+)$ denote the matrix for the action of Frobenius on $\mathbf{N}(T)$ in the chosen basis. In Lemma 4.46 we obtained that $\det P$ is invertible in $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}[\frac{1}{p}]$.

Now, recall that $\mathcal{O}N_m^{\text{PD}} = \mathcal{O}\hat{S}_m^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ and $M'' = (\mathcal{O}N_m^{\text{PD}})^{\Gamma_R}$. So we consider the following commutative diagram

$$\begin{array}{ccc} \mathcal{O}\hat{S}_m^{\text{PD}} \otimes_R M'' & \xrightarrow{\sim} & \mathcal{O}N_m^{\text{PD}} \\ \varphi^m \otimes \varphi^m \downarrow & & \downarrow \varphi^m \\ \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M & \longrightarrow & \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T), \end{array}$$

where the top horizontal arrow is bijective (see Lemma 4.45) and all other arrows are injective. We also have that $\{f_1, \dots, f_h\}$ is an $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ -basis of $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ as well as an $\mathcal{O}\hat{S}_m^{\text{PD}}$ -basis of $\mathcal{O}N_m^{\text{PD}}$. From Lemmas 4.41 & 4.45 and the discussion above, for $1 \leq i \leq h$ we have $f_i'' \in M''$ such that $f_i'' = f_i + \sum_{j=1}^h a_{ij} f_j$ for $a_{ij} \in J^{[1]}\mathcal{O}\hat{S}_m^{\text{PD}}$ and let $A := id_h + (a_{ij}) \in \text{Mat}(h, \mathcal{O}\hat{S}_m^{\text{PD}})$

denote the $h \times h$ matrix obtained in this manner. From the proof of Lemma 4.43 we have that $\det A$ is invertible in $\mathcal{O}\widehat{S}_m^{\text{PD}}$.

Now let $v_i = (\varphi^m \otimes \varphi^m) f_i'' = \varphi^m(f_i) + \sum_{j=1}^h \varphi^m(a_{ij}) \varphi^m(f_j) \in M$ and let M_0 be the free R -submodule of M generated by $\{v_1, \dots, v_h\}$. From the expression of $\{v_1, \dots, v_h\}$ in the basis of $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$, we get that the determinant of the inclusion $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M_0 \hookrightarrow \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ is given by $\varphi^m(\det A) \varphi^{m-1}(\det P) \varphi^{m-2}(\det P) \cdots \varphi(\det P)(\det P)$. Since $\det A$ is invertible in $\mathcal{O}\widehat{S}_m^{\text{PD}}$, we have that $\varphi^m(\det A)$ is invertible in $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ and from above we already have that $\det P$ is invertible in $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}[\frac{1}{p}]$. Therefore, the natural inclusions

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M_0[\frac{1}{p}] \longrightarrow \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M[\frac{1}{p}] \longrightarrow \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V),$$

are bijective. These inclusions are compatible with Frobenius, filtration, connection and the action of Γ_R on each side, which shows the second claim of Proposition 4.28.

Finally, note that above we assumed $\mathbf{N}(T)$ to be free of rank h , therefore we obtain a free R -submodule $M_0 \subset M$ such that

$$M_0[\frac{1}{p}] = (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M_0[\frac{1}{p}])^{\Gamma_R} \xrightarrow{\sim} (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M[\frac{1}{p}])^{\Gamma_R} = M[\frac{1}{p}],$$

which are free of rank h over $R[\frac{1}{p}]$. This shows the last claim of Proposition 4.28. In general, when $\mathbf{N}(T)$ is projective of rank h , we obtain that $M[\frac{1}{p}]$ is projective of rank h . This sums up our proof. \blacksquare

4.5. Proof of Theorem 4.25. Let $M = (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))^{\Gamma_R}$. From Proposition 4.28 we already have the isomorphism of $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}[\frac{1}{p}]$ -modules

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M[\frac{1}{p}] \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V),$$

compatible with Frobenius, filtration, connection and the action of Γ_R on each side. This proves the second claim and we are left to show that V is crystalline and $M[\frac{1}{p}] \xrightarrow{\sim} \mathcal{O}\mathbf{D}_{\text{cris}}(V)$ compatible with supplementary structures. Also note from Proposition 4.28 that we already have the inclusion of projective $R[\frac{1}{p}]$ -modules of rank $h = \dim_{\mathbb{Q}_p} V$, $M[\frac{1}{p}] \subset \mathcal{O}\mathbf{D}_{\text{cris}}(V)$. So we are left to show that this inclusion is bijective and compatible with supplementary structures.

First, we will show that V is crystalline and the inclusion described above is in fact bijective. Extending scalars along $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}[\frac{1}{p}] \hookrightarrow \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ for the isomorphism $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M[\frac{1}{p}] \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)$, we obtain an isomorphism of $\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ -modules

$$\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{R[\frac{1}{p}]} M[\frac{1}{p}] \xrightarrow{\sim} \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbf{B}_R^+} \mathbf{N}(V),$$

compatible with Frobenius, filtration, connection and G_R -action. Now, recall that from the definitions we have a natural inclusion of free \mathbf{A}^+ -modules $\mathbf{A}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(V) \hookrightarrow \mathbf{A}^+ \otimes_{\mathbf{A}_R^+} V$ compatible with supplementary structures and the cokernel of this inclusion is killed by π^s (see Proposition 4.11). Since π is invertible in $\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$, extending scalars along $\mathbf{A}^+ \hookrightarrow \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$, we obtain an isomorphism of $\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ -modules

$$\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbf{B}_R^+} \mathbf{N}(V) \xrightarrow{\sim} \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} V,$$

compatible with Frobenius, connection and G_R -action. Finally, since $R[\frac{1}{p}] \rightarrow \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ is faithfully flat (see [Bri08, Théorème 6.3.8]), we obtain an inclusion of $\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ -modules

$$\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{R[\frac{1}{p}]} M[\frac{1}{p}] \subset \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{R[\frac{1}{p}]} \mathcal{O}\mathbf{D}_{\text{cris}}(V),$$

compatible with Frobenius, connection and the action of G_R . In particular, we have a commutative diagram

$$\begin{array}{ccc}
\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{R[\frac{1}{p}]} M[\frac{1}{p}] & \xrightarrow{\sim} & \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbf{B}_R^+} \mathbf{N}(V) \\
\downarrow & & \downarrow \sim \\
\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{R[\frac{1}{p}]} \mathcal{O}\mathbf{D}_{\text{cris}}(V) & \xrightarrow{\sim} & \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} V,
\end{array}$$

compatible with Frobenius, connection and G_R -action. As the top horizontal arrow and right vertical arrow are bijections, it is immediately clear from the diagram that the left vertical arrow and bottom horizontal arrow must be bijective as well. The bijection of bottom horizontal arrow implies that V is a crystalline representation of G_R . Moreover, since $R[\frac{1}{p}] \rightarrow \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ is faithfully flat (see [Bri08, Théorème 6.3.8]), we obtain an isomorphism of $R[\frac{1}{p}]$ -modules $M[\frac{1}{p}] \xrightarrow{\sim} \mathcal{O}\mathbf{D}_{\text{cris}}(V)$.

Finally, we note that the isomorphism $M[\frac{1}{p}] \xrightarrow{\sim} \mathcal{O}\mathbf{D}_{\text{cris}}(V)$ is compatible with supplementary structures. From Proposition 4.28 it is clear that this isomorphism is compatible with Frobenius and connection. Combining Proposition 4.49 with observations made before, we obtain that the isomorphism of $R[\frac{1}{p}]$ -modules $M[\frac{1}{p}] \xrightarrow{\sim} \mathcal{O}\mathbf{D}_{\text{cris}}(V)$ is compatible with Frobenius, filtration and connection on each side.

Finally, we can compose these natural maps as

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R \mathcal{O}\mathbf{D}_{\text{cris}}(V) \xleftarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V))^{\Gamma_R} \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V),$$

where the second map is compatible with the Frobenius, filtration, connection and the action of Γ_R on each side (see Proposition 4.28). This proves the theorem. \blacksquare

Remark 4.47. In the case when $\mathbf{N}(T)$ is a free \mathbf{A}_R^+ -module of rank h , from Proposition 4.28 we obtain that $M[\frac{1}{p}] \xrightarrow{\sim} \mathcal{O}\mathbf{D}_{\text{cris}}(V)$ is a free $R[\frac{1}{p}]$ -module of rank h . In particular, for finite q -height representations there exists a finite étale extension R' over R such that $R'[\frac{1}{p}] \otimes_{R[\frac{1}{p}]} \mathcal{O}\mathbf{D}_{\text{cris}}(V)$ is free of rank h .

Remark 4.48. For $0 \leq i \leq d$, one can define $[\varepsilon]$ -derivatives by the formula $\frac{\gamma_i - 1}{\pi} : \mathbf{N}(T) \rightarrow \mathbf{N}(T)$. Considering the reduction modulo π of Frobenius, filtration and $[\varepsilon]$ -connection on $\mathbf{N}(T)$ defined above, we conjecture that we have $(\mathbf{N}(T)/\pi\mathbf{N}(T))[\frac{1}{p}] \xrightarrow{\sim} \mathcal{O}\mathbf{D}_{\text{cris}}(V)$ as filtered (φ, ∂) -modules over $R[\frac{1}{p}]$. Details on this line of thought and its connection with [BS19] and [GLQ20] will appear elsewhere.

4.5.1. Compatibility between filtrations. Recall that using Definition 4.16 and Remark 4.21 (ii), the filtration on $M[\frac{1}{p}]$ is given as

$$\text{Fil}^k M[\frac{1}{p}] = \left(\sum_{i \in \mathbb{N}} \text{Fil}^i \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \widehat{\otimes}_{\mathbf{A}_R^+} \text{Fil}^{k-i} \mathbf{N}(V) \right)^{\Gamma_R}.$$

Proposition 4.49. *We have $\text{Fil}^k M[\frac{1}{p}] = \text{Fil}^k \mathcal{O}\mathbf{D}_{\text{cris}}(V)$ for $k \in \mathbb{Z}$*

Proof. We only need to show the claim for $k \geq 1$. Note that from (4.1) we have

$$\text{Fil}^k M[\frac{1}{p}] = (\text{Fil}^k (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)))^{\Gamma_R} \subset (\text{Fil}^k (\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} V))^{G_R} = \text{Fil}^k \mathcal{O}\mathbf{D}_{\text{cris}}(V).$$

Conversely, let $\{e_1, \dots, e_h\}$ denote a \mathbb{Q}_p -basis of V and let $x \in \text{Fil}^k \mathcal{O}\mathbf{D}_{\text{cris}}(V) \setminus \text{Fil}^{k+1} \mathcal{O}\mathbf{D}_{\text{cris}}(V)$. Since $x \neq 0$, we can write $x = \sum_{i=1}^h b_i e_i$ where either $b_i = 0$ or $b_i \in \text{Fil}^k \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \setminus \text{Fil}^{k+1} \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ for each $1 \leq i \leq h$ and at least one $b_i \neq 0$. Moreover, we have $M[\frac{1}{p}] \xrightarrow{\sim} \mathcal{O}\mathbf{D}_{\text{cris}}(V)$ as $R[\frac{1}{p}]$ -modules, so we take $r \leq k$ to be the largest integer such that $x \in \text{Fil}^r M[\frac{1}{p}]$, in particular $x \notin \text{Fil}^{r+1} M[\frac{1}{p}]$. Let us write $x = \sum_{j \in \mathbb{N}} c_j \otimes f_{r-j}$

with $c_j \in \text{Fil}^j \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ and $f_{r-j} \in \text{Fil}^{r-j} \mathbf{N}(V)$ for all $j \in \mathbb{N}$. By assumption on x there exists $\emptyset \neq I \subset \mathbb{N}$ such that for each $j \in I$ we have $c_j \in \text{Fil}^j \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \setminus \text{Fil}^{j+1} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$, $f_{r-j} \in \text{Fil}^{r-j} \mathbf{N}(V) \setminus \text{Fil}^{r-j+1} \mathbf{N}(V)$ with

$$\begin{aligned} \sum_{j \in I} c_j \otimes f_{r-j} &\in \text{Fil}^r(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)) \setminus \text{Fil}^{r+1}(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)) \text{ and} \\ \sum_{j \in \mathbb{N} \setminus I} c_j \otimes f_{r-j} &\in \text{Fil}^{r+1}(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)). \end{aligned}$$

Equip \mathbf{B}^+ with the induced filtration $\text{Fil}^n \mathbf{B}^+ := \mathbf{B}^+ \cap \text{Fil}^n \mathbf{B}_{\text{cris}}(\overline{R}) := \mathbf{B}^+ \cap \text{Fil}^n(\mathbf{A}_{\text{inf}}(\overline{R})[\frac{1}{p}])$ for $n \in \mathbb{N}$. Using the definition of filtration on $\mathbf{N}(V)$ (see Definition 4.16) and Lemma 4.53, we have that $\text{Fil}^{r-j} \mathbf{N}(V) = (\text{Fil}^{r-j} \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V) \cap \mathbf{N}(V)$ for all $j \in \mathbb{N}$. Therefore, in the expression $\sum_{j \in I} c_j \otimes f_{r-j}$ we must have $f_{r-j} \in (\text{Fil}^{r-j} \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V) \setminus (\text{Fil}^{r-j+1} \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V)$ for all $j \in I$. This implies that in the basis of V we can write $f_{r-j} = \sum_{i=1}^h f_{r-j}^{(i)} e_i$ with $f_{r-j}^{(i)} \in \text{Fil}^{r-j} \mathbf{B}^+ \setminus \text{Fil}^{r-j+1} \mathbf{B}^+$ for all $j \in I$ and all $1 \leq i \leq h$. In conclusion, we obtain

$$x - \sum_{j \in \mathbb{N} \setminus I} c_j \otimes f_{r-j} = \sum_{j \in I} c_j \otimes \left(\sum_{i=1}^h f_{r-j}^{(i)} e_i \right) = \sum_{i=1}^h \left(\sum_{j \in I} c_j \otimes f_{r-j}^{(i)} \right) e_i, \quad (4.5)$$

with $c_j \in \text{Fil}^j \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \setminus \text{Fil}^{j+1} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ and $f_{r-j}^{(i)} \in \text{Fil}^{r-j} \mathbf{B}^+ \setminus \text{Fil}^{r-j+1} \mathbf{B}^+$ for all $1 \leq i \leq h$ and $j \in I$.

Let us set $g_i = \sum_{j \in I} c_j \otimes f_{r-j}^{(i)}$ for $1 \leq i \leq h$. Then by the discussion above we have that $g_i \in \text{Fil}^r(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{B}^+)$ for $1 \leq i \leq h$, where $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{B}^+$ is equipped with the tensor product filtration. Note that $x \in \text{Fil}^r M[\frac{1}{p}] \setminus \text{Fil}^{r+1} M[\frac{1}{p}]$ and $\sum_{j \in \mathbb{N} \setminus I} c_j \otimes f_{r-j} \in \text{Fil}^{r+1}(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V))$. Moreover, from Lemma 4.50 and Remark 4.51 we deduce that for $n \in \mathbb{N}$ and inside $\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} V$ we have

$$\text{Fil}^n(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)) = (\text{Fil}^n(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{B}^+) \otimes_{\mathbb{Q}_p} V) \cap (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)).$$

Therefore, we conclude that we must have at least one $i = i_0$ such that $g_{i_0} \in \text{Fil}^r(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{B}^+) \setminus \text{Fil}^{r+1}(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{B}^+)$. Now using Lemma 4.52 and Remark 4.51 we further note that for $n \in \mathbb{N}$

$$\text{Fil}^n(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{B}^+) = (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{B}^+) \cap \text{Fil}^n \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \subset \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}).$$

Therefore, we get that $g_i \in \text{Fil}^r \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ for all $1 \leq i \leq h$ and $g_{i_0} \in \text{Fil}^r \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \setminus \text{Fil}^{r+1} \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$. For convenience, let us write $\sum_{j \in \mathbb{N} \setminus I} c_j \otimes f_{r-j} = \sum_{i=1}^h d_i e_i$ with $d_i \in \text{Fil}^{r+1} \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ for all $1 \leq i \leq h$. In particular, comparing (4.5) with the expression $x = \sum_{i=1}^h b_i e_i$ at the start of the proof, we get $b_{i_0} = g_{i_0} + d_{i_0}$.

Finally, since $r \leq k$, consider the following commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Fil}^{k+1} \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) & \longrightarrow & \text{Fil}^k \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) & \longrightarrow & \text{gr}^k \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \text{Fil}^{r+1} \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) & \longrightarrow & \text{Fil}^r \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) & \longrightarrow & \text{gr}^r \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \longrightarrow 0, \end{array}$$

where the left and middle vertical arrows are injective and the right vertical arrow is non-trivial if and only if $r = k$. From the fact that $g_{i_0} \in \text{Fil}^r \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \setminus \text{Fil}^{r+1} \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$, we see that the image of b_{i_0} is non-zero in $\text{gr}^r \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$. But we already have that image of b_{i_0} is non-zero in $\text{gr}^k \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$. Therefore, the left vertical arrow must be non-trivial, i.e. $r = k$. Hence $x \in \text{Fil}^k M[\frac{1}{p}]$. This proves the claim. \blacksquare

Lemma 4.50. *For $k \in \mathbb{N}$ we have*

$$\mathrm{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^{\mathrm{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)) = (\mathrm{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^{\mathrm{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{A}^+) \otimes_{\mathbb{Z}_p} T) \cap (\mathcal{O}\mathbf{A}_{R,\varpi}^{\mathrm{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)).$$

Proof. From §3.1 we have rings $\mathbf{A}^+ \subset \mathbf{A}_{\varpi}^+ \subset \mathbf{A}_{\mathrm{inf}}(\overline{R})$ equipped with an induced filtration from $\mathbf{A}_{\mathrm{cris}}(\overline{R})$ and from Remark 3.18 we have an isomorphism $\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{A}^+ \xrightarrow{\sim} \mathbf{A}_{\varpi}^+$ compatible with Frobenius, filtration and G_R -action. So using Lemma 4.53, the fact that $\mathbf{A}_R^+ \rightarrow \mathbf{A}_{R,\varpi}^+$ is flat and $\mathrm{Fil}^i \mathbf{A}_{R,\varpi}^+ = \xi^i \mathbf{A}_{R,\varpi}^+$ we note that

$$\begin{aligned} \mathrm{Fil}^k(\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)) &= \sum_{i+j=k} \mathrm{Fil}^i \mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} ((\mathrm{Fil}^j \mathbf{A}^+ \otimes_{\mathbb{Z}_p} T) \cap \mathbf{N}(T)) \\ &= \left(\sum_{i+j=k} \mathrm{Fil}^i \mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathrm{Fil}^j \mathbf{A}^+ \otimes_{\mathbb{Z}_p} T \right) \cap (\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)) \quad (4.6) \\ &= (\mathrm{Fil}^k \mathbf{A}_{\varpi}^+ \otimes_{\mathbb{Z}_p} T) \cap (\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)). \end{aligned}$$

Next, from Definition 4.19 we have the ring $\mathcal{O}\mathbf{A}_{R,\varpi}^+$ flat over $\mathbf{A}_{R,\varpi}^+$ since

$$\mathcal{O}\mathbf{A}_{R,\varpi}^+ = \oplus_{\mathbf{k} \in \mathbb{N}^d} \mathbf{A}_{R,\varpi}^+ (X_1 - [X_1^b])^{k_1} \cdots (X_d - [X_d^b])^{k_d},$$

with the structure map $\mathbf{A}_{R,\varpi}^+ \rightarrow \mathcal{O}\mathbf{A}_{R,\varpi}^+$ being injective and its image is identified with term at index $\mathbf{k} = (0, \dots, 0)$. Let us set $\mathcal{O}\mathbf{A}_{\varpi}^+ := \mathcal{O}\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{A}^+ \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_{R,\varpi}^+} \mathbf{A}_{\varpi}^+$ equipped with natural filtration, Frobenius and G_R -action. Let $J = (X_1 - [X_1^b], \dots, X_d - [X_d^b]) \mathcal{O}\mathbf{A}_{R,\varpi}^+$ then the filtration on $\mathcal{O}\mathbf{A}_{\varpi}^+$ can also be given as $\mathrm{Fil}^k \mathcal{O}\mathbf{A}_{\varpi}^+ = \sum_{i+j=k} J^i \mathcal{O}\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_{R,\varpi}^+} \xi^j \mathbf{A}_{\varpi}^+$. Let us set $N_{R,\varpi}^+ = \mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ equipped with tensor product filtration. Then since J is flat as an $\mathbf{A}_{R,\varpi}^+$ -module an argument similar to (4.6) gives us that

$$\begin{aligned} \mathrm{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)) &= \mathrm{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_{R,\varpi}^+} N_{R,\varpi}^+) \\ &= \sum_{i+j=k} J^i \mathcal{O}\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_{R,\varpi}^+} ((\mathrm{Fil}^j \mathbf{A}_{\varpi}^+ \otimes_{\mathbb{Z}_p} T) \cap N_{R,\varpi}^+) \\ &= \left(\sum_{i+j=k} J^i \mathcal{O}\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_{R,\varpi}^+} \xi^j \mathbf{A}_{\varpi}^+ \otimes_{\mathbb{Z}_p} T \right) \cap (\mathcal{O}\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_{R,\varpi}^+} N_{R,\varpi}^+) \\ &= (\mathrm{Fil}^k \mathcal{O}\mathbf{A}_{\varpi}^+ \otimes_{\mathbb{Z}_p} T) \cap (\mathcal{O}\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)). \end{aligned} \quad (4.7)$$

Furthermore, let us set $\mathcal{O}\mathbf{A}_{\varpi}^{\mathrm{PD}} := \mathcal{O}\mathbf{A}_{R,\varpi}^{\mathrm{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{A}_{\varpi}^+ \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\mathrm{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{A}^+$ where the isomorphism is compatible with Frobenius, filtration, connection and G_R -action.

Now we will show our claim

$$\mathrm{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^{\mathrm{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)) = (\mathrm{Fil}^k \mathcal{O}\mathbf{A}_{\varpi}^{\mathrm{PD}} \otimes_{\mathbb{Z}_p} T) \cap (\mathcal{O}\mathbf{A}_{R,\varpi}^{\mathrm{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)).$$

Let $f \in \{\xi, X_1 - [X_1^b], \dots, X_d - [X_d^b]\}$ be one of the generators of the ideal $(\xi, X_1 - [X_1^b], \dots, X_d - [X_d^b]) \mathcal{O}\mathbf{A}_{\varpi}^{\mathrm{PD}}$. Then to obtain our claim, it is enough to show that if $f^{[k]}x \in (f^{[k]} \mathcal{O}\mathbf{A}_{\varpi}^{\mathrm{PD}} \otimes_{\mathbb{Z}_p} T) \setminus (f^{[k+1]} \mathcal{O}\mathbf{A}_{\varpi}^{\mathrm{PD}} \otimes_{\mathbb{Z}_p} T)$ such that $f^{[k]}x \in (\mathcal{O}\mathbf{A}_{R,\varpi}^{\mathrm{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))$ then $f^{[k]}x \in \mathrm{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^{\mathrm{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))$.

Note that the claim is true for $k = 0$. So let $k \geq 1$ and f as above. Let $f^{[k]}x \in (f^{[k]} \mathcal{O}\mathbf{A}_{\varpi}^{\mathrm{PD}} \otimes_{\mathbb{Z}_p} T) \setminus (f^{[k+1]} \mathcal{O}\mathbf{A}_{\varpi}^{\mathrm{PD}} \otimes_{\mathbb{Z}_p} T)$ such that $f^{[k]}x \in (\mathcal{O}\mathbf{A}_{R,\varpi}^{\mathrm{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))$. Since $x \neq 0$, by induction on k we may assume that $x = \sum_{i=1}^h x_i e_i \in \mathcal{O}\mathbf{A}_{\varpi}^+ \otimes_{\mathbb{Z}_p} T$ with either $x_i = 0$ or $x_i \in f^k \mathcal{O}\mathbf{A}_{\varpi}^+ \setminus f^{k+1} \mathcal{O}\mathbf{A}_{\varpi}^+$

for each $1 \leq i \leq h$ and at least one $x_i \neq 0$. Recall that we have $\pi^s \mathbf{A}^+ \otimes_{\mathbb{Z}_p} T \subset \mathbf{A}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$, therefore $\pi^s x \in \mathcal{O}\mathbf{A}_{\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$. But then inside $\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ we must have

$$f^k x = k! f^{[k]} x \in (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)) \cap \frac{1}{\pi^s} (\mathcal{O}\mathbf{A}_{\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)) = \mathcal{O}\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T).$$

Therefore, $f^k x \in (f^k \mathcal{O}\mathbf{A}_{\varpi}^+ \otimes_{\mathbb{Z}_p} T) \cap (\mathcal{O}\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)) = \text{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))$ where the last equality follows from (4.7). Hence, inside $\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)$ we have $f^{[k]} x \in \frac{1}{k!} \text{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)) \cap (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T)) = \text{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))$. ■

Remark 4.51. From the proof of Lemma 4.50 it easily follows that the corresponding rational version of the statement is also true, i.e. inside $\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} V$ we have

$$\begin{aligned} \text{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)) \\ = (\text{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{B}^+) \otimes_{\mathbb{Q}_p} V) \cap (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V)). \end{aligned}$$

Similarly, from Lemma 4.52 it easily follows that inside $\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ we have

$$\text{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{B}^+) = (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{B}^+) \cap \text{Fil}^k \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}).$$

Lemma 4.52. *For $k \in \mathbb{N}$ we have*

$$\text{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{A}^+) = (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{A}^+) \cap \text{Fil}^k \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R}) \subset \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R}).$$

Proof. Recall that filtrations on $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ and $\mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})$ are compatible (see Remark 4.23). Moreover, from §3.1 the inclusion of rings $\mathbf{A}^+ \subset \mathbf{A}_{\varpi}^+ \subset \mathbf{A}_{\text{inf}}(\overline{R})$ is compatible with induced filtration from $\mathbf{A}_{\text{cris}}(\overline{R})$. From the discussion in Lemma 4.50 we have an isomorphism of rings $\mathcal{O}\mathbf{A}_{\varpi}^{\text{PD}} = \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{A}^+ \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{A}_{\varpi}^+$ compatible with tensor product filtrations. Now by the description of filtration on the rightmost term we get that $\mathcal{O}\mathbf{A}_{\varpi}^{\text{PD}}$ is equipped with filtration by divided powers of the ideal $(\xi, X_1 - [X_1^b], \dots, X_d - [X_d^b])\mathcal{O}\mathbf{A}_{\varpi}^{\text{PD}}$. Finally, the natural multiplication map $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{A}_{\varpi}^+ \rightarrow \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})$ is injective. Hence, it follows that for $k \in \mathbb{N}$

$$\text{Fil}^k(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{A}^+) = (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{A}^+) \cap \text{Fil}^k \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R}) \subset \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R}).$$
■

Lemma 4.53. *For $k \in \mathbb{N}$ we have $(\text{Fil}^k \mathbf{A}^+ \otimes_{\mathbb{Z}_p} T) \cap \mathbf{N}(T) = \text{Fil}^k \mathbf{N}(T)$.*

Proof. It is enough to show that $(\text{Fil}^k \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V) \cap \mathbf{N}(V) = \text{Fil}^k \mathbf{N}(V)$. Indeed, from Definition 4.16 we have $\text{Fil}^k \mathbf{N}(T) = \text{Fil}^k \mathbf{N}(V) \cap \mathbf{N}(T) = (\text{Fil}^k \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V) \cap \mathbf{N}(V) \cap \mathbf{N}(T) = (\text{Fil}^k \mathbf{A}^+ \otimes_{\mathbb{Q}_p} T) \cap \mathbf{N}(T)$ since $\text{Fil}^k \mathbf{B}^+ \cap \mathbf{A}^+ = \text{Fil}^k \mathbf{A}^+$.

Now let us show the modified claim. The inclusion $\text{Fil}^k \mathbf{N}(V) \subset (\text{Fil}^k \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V)$ is obvious. For the converse, we claim that it is enough to show that $(q^k \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V) \cap \mathbf{N}(V) = q^k \mathbf{N}(V)$. Indeed, if we have $x \in (\text{Fil}^k \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V) \cap \mathbf{N}(V)$ then $\varphi(x) \in (q^k \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V) \cap \mathbf{N}(V) = q^k \mathbf{N}(V)$, i.e. $x \in \text{Fil}^k \mathbf{N}(V)$.

The inclusion $q^k \mathbf{N}(V) \subset (q^k \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V) \cap \mathbf{N}(V)$ is obvious. To show the converse, first let us assume that $\mathbf{N}(V)$ is free with $\{f_1, f_2, \dots, f_h\}$ as a \mathbf{B}_R^+ -basis, and let $\{e_1, \dots, e_h\}$ be a \mathbb{Q}_p -basis of V . Now let $q^k x \in (q^k \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V) \cap \mathbf{N}(V)$ for $x = \sum_{i=1}^h x_i e_i \in \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V$. We

can also write $q^k x = \sum_{i=1}^h y_i f_i \in \mathbf{N}(V)$ with $y_i \in \mathbf{B}_R^+$. Next, from Proposition 4.11 we have $\pi^s \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V \subset \mathbf{B}^+ \otimes_{\mathbf{B}_R^+} \mathbf{N}(V)$, so we can write

$$q^k x = \pi^{-s} q^k \sum_{i=1}^h x_i \pi^s e_i = \pi^{-s} q^k \sum_{i=1}^h x_i \sum_{j=1}^h z_{ij} f_j = \pi^{-s} q^k \sum_{i=1}^h \left(\sum_{j=1}^h x_j z_{ji} \right) f_i,$$

with $z_{ij} \in \mathbf{B}^+$. But then we must have $\pi^{-s} q^k \sum_{j=1}^h x_j z_{ji} = y_i$ for all $1 \leq i \leq h$. Since H_R acts trivially on π , q and y_i , we get that $w_i := \sum_{j=1}^h x_j z_{ji} \in \mathbf{B}_R^+$. But $y_i \in \mathbf{B}_R^+$ and π and q are coprime in \mathbf{B}_R^+ (since $q \equiv p \pmod{\pi \mathbf{B}_R^+}$), therefore we obtain that $w_i \in \pi^s \mathbf{B}_R^+$. In particular, $y_i \in q^k \mathbf{B}_R^+$, therefore $q^k x = \sum_{i=1}^h y_i f_i \in q^k \mathbf{N}(V)$. Hence, $(q^k \mathbf{B}^+ \otimes_{\mathbb{Q}_p} V) \cap \mathbf{N}(V) = q^k \mathbf{N}(V)$.

Next, if $\mathbf{N}(V)$ is projective (and not free) over \mathbf{B}_R^+ , let R' be the p -adic completion of a finite étale algebra over R such that the scalar extension $\mathbf{B}_{R'}^+ \otimes_{\mathbf{B}_R^+} \mathbf{N}(V)$ is a free module over $\mathbf{B}_{R'}^+$ and $R'[\frac{1}{p}]/R[\frac{1}{p}]$ is Galois (see Definition 4.9). Then we can argue as above and conclude by taking $\text{Gal}(R'[\frac{1}{p}]/R[\frac{1}{p}])$ -invariants of $q^k \mathbf{B}_{R'}^+ \otimes_{\mathbf{B}_R^+} \mathbf{N}(V)$. ■

4.6. One-dimensional representations. In this section we will show that all one-dimensional crystalline representations are of finite q -height by writing down the corresponding Wach modules precisely.

Proposition 4.54. *All one-dimensional crystalline representations of G_R are of finite q -height. Furthermore, for a one-dimensional crystalline representation V we have an isomorphism of $R[\frac{1}{p}]$ -modules*

$$(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V))^{\Gamma_R} \xrightarrow{\sim} \mathcal{O}\mathbf{D}_{\text{cris}}(V).$$

Therefore, there exists natural isomorphisms

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R \mathcal{O}\mathbf{D}_{\text{cris}}(V) \xleftarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V))^{\Gamma_R} \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V),$$

compatible with Frobenius, filtration and the action of Γ_R .

Proof. The structure of one-dimensional crystalline representations of G_R is well-known (see [Bri08, §8.6]). From Proposition 2.3 we have that for $\eta : G_R \rightarrow \mathbb{Z}_p^\times$, a continuous character, $V = \mathbb{Q}_p(\eta)$ is crystalline if and only if we can write $\eta = \eta_f \eta_{\text{ur}} \chi^n$ with $n \in \mathbb{Z}$, and where η_f is a finite unramified character, η_{ur} is an unramified character taking values in $1 + p\mathbb{Z}_p$ and trivialized by an element $\alpha \in 1 + p\widehat{R}^{\text{ur}}$, and χ is the p -adic cyclotomic character. Recall that a p -adic representation of G_R is unramified if the action of G_R factorizes through the quotient G_R^{ur} (see §2.3). Moreover, if η_f is trivial then $\mathcal{O}\mathbf{D}_{\text{cris}}(V)$ is a free $R[\frac{1}{p}]$ -module of rank 1.

In Lemma 4.55 below, we show that crystalline representations $V_1 := \mathbb{Q}_p(\eta_f \eta_{\text{ur}})$ and $V_2 := \mathbb{Q}_p(\chi^n)$ are of finite q -height. For a one-dimensional crystalline representation $V := \mathbb{Q}_p(\eta) = \mathbb{Q}_p(\eta_f \eta_{\text{ur}}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(\chi^n) = V_1 \otimes_{\mathbb{Q}_p} V_2$ as above, by compatibility of tensor products in Propositions 4.14 we get that V is a finite q -height representation as well with $\mathbf{N}(V) = \mathbf{N}(V_1) \otimes_{\mathbf{B}_R^+} \mathbf{N}(V_2)$.

From the isomorphisms of $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ -modules in Lemma 4.55 and compatibility of Wach modules with tensor product in Proposition 4.14 and compatibility of the functor $\mathcal{O}\mathbf{D}_{\text{cris}}$ with tensor products in §2.3 (see also [Bri08, Théorème 8.4.2]), we get a string of isomorphisms of $\mathcal{O}\mathbf{B}_{R,\varpi}^{\text{PD}} := \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}[\frac{1}{p}]$ -modules compatible with Frobenius, filtration and the action of Γ_R ,

$$\begin{aligned} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R \mathcal{O}\mathbf{D}_{\text{cris}}(V) &\xrightarrow{\sim} (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R \mathcal{O}\mathbf{D}_{\text{cris}}(V_1)) \otimes_{\mathcal{O}\mathbf{B}_{R,\varpi}^{\text{PD}}} (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R \mathcal{O}\mathbf{D}_{\text{cris}}(V_2)) \\ &\xleftarrow{\sim} (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V_1)) \otimes_{\mathcal{O}\mathbf{B}_{R,\varpi}^{\text{PD}}} (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V_2)) \\ &\xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V_1) \otimes_{\mathbf{B}_R^+} \mathbf{N}(V_2) \\ &\xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V_1 \otimes_{\mathbb{Q}_p} V_2) \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V). \end{aligned}$$

Taking Γ_R -invariants of the first and the last term gives us that $\mathcal{O}\mathbf{D}_{\text{cris}}(V) \xrightarrow{\sim} (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V))^{\Gamma_R}$, compatible with Frobenius and filtration. \blacksquare

Following claim was used above:

Lemma 4.55. (i) *Let $\eta : G_R \rightarrow \mathbb{Z}_p^\times$ be a continuous unramified character. Then the p -adic representation $\mathbb{Q}_p(\eta)$ is a finite q -height representation.*

(ii) *Let χ be the p -adic cyclotomic character then for $n \in \mathbb{Z}$, the p -adic representation $\mathbb{Q}_p(n)$ is a finite q -height representation.*

Further, for $V = \mathbb{Q}_p(\eta), \mathbb{Q}_p(n)$ we have an isomorphism of $R[\frac{1}{p}]$ -modules

$$(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V))^{\Gamma_R} \xrightarrow{\sim} \mathcal{O}\mathbf{D}_{\text{cris}}(V).$$

Therefore, there exists natural isomorphisms

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R \mathcal{O}\mathbf{D}_{\text{cris}}(V) \xleftarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V))^{\Gamma_R} \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(V),$$

compatible with Frobenius, filtration and the action of Γ_R .

Proof. Let $\eta = \eta_f \eta_{\text{ur}}$, where η_f is an unramified character of finite order and η_{ur} is an unramified character taking values in $1 + p\mathbb{Z}_p$ and trivialised by an element $\alpha \in 1 + p\widehat{R}^{\text{ur}}$ (see Proposition 2.3).

First, let us consider the finite unramified character η_f . Set $T = \mathbb{Z}_p(\eta_f) = \mathbb{Z}_p e$, such that $g(e) = \eta_f(g)e$. We have

$$\mathbf{D}^+(\mathbb{Z}_p(\eta_f)) = (\mathbf{A}^+ \otimes_{\mathbb{Z}_p} \mathbb{Z}_p(\eta_f))^{H_R} \xrightarrow{\sim} \{a \otimes e, \text{ with } a \in \mathbf{A}^+ \text{ such that } g(a) = \eta_f^{-1}(g)a, \text{ for } g \in H_R\}.$$

Since η_f is a finite unramified character, it trivializes over a finite Galois extension S over R (see [Bri08, Proposition 8.6.1]), and we have that $\text{Gal}(S[\frac{1}{p}]/R[\frac{1}{p}]) = G_R/G_S = H_R/H_S = \Gamma_R/\Gamma_S$. As S is finite étale over R the construction of previous chapters apply and we obtain that the \mathbf{A}_S^+ -module $\mathbf{D}_S^+(\mathbb{Z}_p(\eta_f)) = (\mathbf{A}^+ \otimes_{\mathbb{Z}_p} \mathbb{Z}_p(\eta_f))^{H_S} = \mathbf{A}_S^+(\eta_f) = \mathbf{A}_S^+ e$ is free of rank 1. Further, we know that $\mathbf{D}^+(\mathbb{Z}_p(\eta_f)) = \mathbf{D}_S^+(\mathbb{Z}_p(\eta_f))^{H_R/H_S}$, which implies that the natural inclusion

$$\mathbf{A}_S^+ \otimes_{\mathbf{A}_R^+} \mathbf{D}^+(\mathbb{Z}_p(\eta_f)) \longrightarrow \mathbf{D}_S^+(\mathbb{Z}_p(\eta_f)),$$

is bijective. Since $\mathbf{A}_R^+ \rightarrow \mathbf{A}_S^+$ is faithfully flat, we obtain that $\mathbf{D}^+(\mathbb{Z}_p(\eta_f))$ is projective of rank 1. Moreover, $\mathbf{D}^+(\mathbb{Z}_p(\eta_f))$ admits a Frobenius-semilinear endomorphism φ such that $\mathbf{D}^+(\mathbb{Z}_p(\eta_f)) \xrightarrow{\sim} \varphi^*(\mathbf{D}^+(\mathbb{Z}_p(\eta_f)))$ (one can obtain this after faithfully flat scalar extension $\mathbf{A}_R^+ \rightarrow \mathbf{A}_S^+$ and applying descent as above, since φ commutes with G_R -action). The action of Γ_R is trivial on $\mathbf{D}^+(\mathbb{Z}_p(\eta_f))$. Now, we can take $\mathbf{N}(\mathbb{Z}_p(\eta_f)) = \mathbf{D}^+(\mathbb{Z}_p(\eta_f))$. From the discussion above, $\mathbf{N}(\mathbb{Z}_p(\eta_f))$ clearly satisfies the conditions of Definition 4.9. Also, we have that $\mathbf{N}(\mathbb{Q}_p(\eta_f)) = \mathbf{D}^+(\mathbb{Q}_p(\eta_f))$. On the other hand, we have

$$\mathcal{O}\mathbf{D}_{\text{cris}}(\mathbb{Q}_p(\eta_f)) = (\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(\eta_f))^{G_R} = \{b \otimes e, \text{ with } b \in \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \text{ such that } g(b) = \eta_f(g)b\}.$$

Since η_f trivializes over the finite Galois extension S over R , we have

$$(\mathcal{O}\mathbf{A}_{S,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(\mathbb{Q}_p(\eta_f)))^{\Gamma_S} = S_0[\frac{1}{p}]e = (\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{S}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(\eta_f))^{G_S},$$

where the rings $\mathcal{O}\mathbf{A}_{S,\varpi}^{\text{PD}}$ and $\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{S})$ are defined for S over which all the construction of previous sections apply (since S is finite étale over R). Now taking invariants under the finite Galois group $\text{Gal}(S[\frac{1}{p}]/R[\frac{1}{p}]) = G_R/G_S$, gives us

$$(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(\mathbb{Q}_p(\eta_f)))^{\Gamma_R} = \mathcal{O}\mathbf{D}_{\text{cris}}(\mathbb{Q}_p(\eta_f)).$$

Clearly, the natural maps

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R \mathcal{O}\mathbf{D}_{\text{cris}}(\mathbb{Q}_p(\eta_f)) \xleftarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(\mathbb{Q}_p(\eta_f)))^{\Gamma_R} \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(\mathbb{Q}_p(\eta_f)),$$

are isomorphisms compatible with Frobenius, filtration and the action of Γ_R .

Next, let us consider the unramified character η_{ur} which takes values in $1+p\mathbb{Z}_p$ and trivialised by an element $\alpha \in 1+p\widehat{R^{\text{ur}}}$ (see Proposition 2.3). Set $T = \mathbb{Z}_p(\eta_{\text{ur}}) = \mathbb{Z}_p e$, such that $g(e) = \eta_{\text{ur}}(g)e$. We have

$$\mathbf{D}^+(\mathbb{Z}_p(\eta_{\text{ur}})) = (\mathbf{A}^+ \otimes_{\mathbb{Z}_p} \mathbb{Z}_p(\eta_{\text{ur}}))^{H_R} = \mathbf{A}_R^+ \alpha e.$$

So we take $\mathbf{N}(\mathbb{Z}_p(\eta_{\text{ur}})) = \mathbf{D}^+(\mathbb{Z}_p(\eta_{\text{ur}})) = \mathbf{A}_R^+ \alpha e$. This clearly satisfies the conditions of Definition 4.9. Also, we have that $\mathbf{N}(\mathbb{Q}_p(\eta_{\text{ur}})) = \mathbf{D}^+(\mathbb{Q}_p(\eta_{\text{ur}}))$. On the other hand, we have

$$\begin{aligned} \mathcal{O}\mathbf{D}_{\text{cris}}(\mathbb{Q}_p(\eta_{\text{ur}})) &= (\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(\eta_{\text{ur}}))^{G_R} \\ &= \{b \otimes e, \text{ with } b \in \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \text{ such that } g(b) = \eta_{\text{ur}}(g)b\} = R[\tfrac{1}{p}] \alpha e. \end{aligned}$$

Therefore, we obtain

$$(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(\mathbb{Q}_p(\eta_{\text{ur}})))^{\Gamma_R} = R[\tfrac{1}{p}] \alpha e = (\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(\eta_{\text{ur}}))^{G_R}.$$

Clearly, the natural maps

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R \mathcal{O}\mathbf{D}_{\text{cris}}(\mathbb{Q}_p(\eta_{\text{ur}})) \xleftarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(\mathbb{Q}_p(\eta_{\text{ur}})))^{\Gamma_R} \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(\mathbb{Q}_p(\eta_{\text{ur}})),$$

are isomorphisms compatible with Frobenius, filtration and the action of Γ_R .

Finally, let $T = \mathbb{Z}_p(n) = \mathbb{Z}_p e_n$ such that $g(e_n) = \chi(g)^n e_n$, then $V = \mathbb{Q}_p \otimes_{\mathbb{Z}_p} T$ is a crystalline representation and we can take $\mathbf{N}(\mathbb{Z}_p(n)) = \mathbf{A}_R^+ \pi^{-n} e_n$. Note that for $n \leq 0$, we have that $\mathbf{N}(\mathbb{Z}_p(n))/\varphi^*(\mathbf{N}(\mathbb{Z}_p(n)))$ is killed by q^{-n} , where $q = \frac{\varphi(\pi)}{\pi}$. It can easily be verified that Γ_R acts trivially modulo π on $\mathbf{N}(T)$. So, we set $\mathbf{N}(\mathbb{Q}_p(n)) = \mathbf{B}_R^+ \pi^{-n} e_n$. Similarly,

$$\mathcal{O}\mathbf{D}_{\text{cris}}(\mathbb{Q}_p(n)) = (\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(n))^{G_R} = R[\tfrac{1}{p}] t^{-n} e_n,$$

and $(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(\mathbb{Q}_p(n)))^{\Gamma_R} = R[\tfrac{1}{p}] t^{-n} e_n = \mathcal{O}\mathbf{D}_{\text{cris}}(\mathbb{Q}_p(n))$ compatible with Frobenius, filtration and connection on each side. Finally, the map

$$\begin{aligned} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R \mathcal{O}\mathbf{D}_{\text{cris}}(\mathbb{Q}_p(n)) &\longrightarrow \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(\mathbb{Q}_p(n)) \\ t^{-n} e_n &\longmapsto \frac{\pi^n}{t^n} \pi^{-n} e_n. \end{aligned}$$

is trivially an isomorphism compatible with Frobenius, filtration and the action of Γ_R , since $\frac{\pi^n}{t^n} \in \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ are units for $n \in \mathbb{Z}$ (see Lemma 3.14). This proves the lemma. \blacksquare

Remark 4.56. Note that for $T = \mathbb{Z}_p(\eta_f \eta_{\text{ur}})$ or $\mathbb{Z}_p(n)$, we even have an isomorphism on the integral level

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T))^{\Gamma_R} \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_R^+} \mathbf{N}(T).$$

5. Relative Fontaine-Laffaille modules

In this section we will consider relative Fontaine-Laffaille data and construct Wach modules given such data. Carrying out such a process would involve starting with a module over R and constructing modules over the ring $\mathbf{A}_{R,\varpi}^{\text{PD}}$ and $\mathbf{A}_{R,\varpi}^+$, and finally descending over to the ring \mathbf{A}_R^+ .

Explicitly, we will work with objects in the category $\text{MF}_{[0,p-2],\text{free}}(R, \Phi, \partial)$, defined by [Tsu20, §4] as a full subcategory of the abelian category $\mathfrak{MF}_{[0,p-2],\text{free}}^\nabla(R)$ introduced by Faltings in [Fal89, §II]. In particular,

Definition 5.1. Define the category of *free relative Fontaine-Laffaille* modules of level $[0, p-2]$, denoted by $\text{MF}_{[0,p-2],\text{free}}(R, \Phi, \partial)$, as follows:

An object with weights in the interval $[0, p-2]$ is a quadruple $(M, \text{Fil}^\bullet M, \partial, \Phi)$ such that,

- (i) M is a free R -module of finite rank.
- (ii) M is equipped with a decreasing filtration $\{\text{Fil}^k M\}_{k \in \mathbb{Z}}$ by finite R -submodules with $\text{Fil}^0 M = M$ and $\text{Fil}^{s+1} M = 0$ such that $\text{gr}_{\text{Fil}}^k M$ is a finite free R -module for every $k \in \mathbb{Z}$.
- (iii) The connection $\partial : M \rightarrow M \otimes_R \Omega_R^1$ is p -adically quasi-nilpotent and integrable, and satisfies Griffiths transversality with respect to the filtration, i.e. $\partial(\text{Fil}^k M) \subset \text{Fil}^{k-1} M \otimes_R \Omega_R^1$ for $k \in \mathbb{Z}$.
- (iv) Let $(\varphi^*(M), \varphi^*(\partial))$ denote the pullback of (M, ∂) by $\varphi : R \rightarrow R$, and equip it with a decreasing filtration defined as $\text{Fil}_p^k(\varphi^*(M)) = \sum_{i \in \mathbb{N}} p^{[i]} \varphi^*(\text{Fil}^{k-i} M)$ for $k \in \mathbb{Z}$. We suppose that there is an R -linear morphism $\Phi : \varphi^*(M) \rightarrow M$ such that Φ is compatible with connections, $\Phi(\text{Fil}_p^k(\varphi^*(M))) \subset p^k M$ for $0 \leq k \leq s$, and we have $\sum_{k=0}^s p^{-k} \Phi(\text{Fil}_p^k(\varphi^*(M))) = M$. We denote the composition $M \rightarrow \varphi^*(M) \xrightarrow{\Phi} M$ by φ .

A morphism between two objects of the category $\text{MF}_{[0,p-2],\text{free}}(R, \Phi, \partial)$ is a continuous R -linear map compatible with the homomorphism Φ , the connection ∂ and filtration on each side.

Notation 5.2. Abusing notations, we will denote $(M, \text{Fil}^\bullet M, \partial, \Phi) \in \text{MF}_{[0,p-2],\text{free}}(R, \Phi, \partial)$ by M and say that it is of level $[0, p-2]$.

To an object $M \in \text{MF}_{[0,p-2],\text{free}}(R, \Phi, \partial)$, we associate a \mathbb{Z}_p -module as

$$T_{\text{cris}}^*(M) := \text{Hom}_{R, \text{Fil}, \varphi, \partial}(M, \mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})), \quad (5.1)$$

i.e. R -linear maps from M to $\mathcal{O}\mathbf{A}_{\text{cris}}(\overline{R})$ compatible with Frobenius, filtration and connection, where we have $\varphi : M \rightarrow \varphi^*(M) \xrightarrow{\Phi} M$.

Proposition 5.3. (i) *For a free Fontaine-Laffaille module M of level $[0, p-2]$, the \mathbb{Z}_p -module $T_{\text{cris}}^*(M)$ is a free module of rank $= \text{rk}_R M$ equipped with a continuous action of G_R . Further, the p -adic representation $V_{\text{cris}}^*(M) := \mathbb{Q}_p \otimes_{\mathbb{Z}_p} T_{\text{cris}}^*(M)$ is a crystalline representation of G_R with Hodge-Tate weights in the interval $[0, p-2]$.*

(ii) *The contravariant \mathbb{Z}_p -linear functor*

$$T_{\text{cris}}^* : \text{MF}_{[0,p-2],\text{free}}(R, \Phi, \partial) \longrightarrow \text{Rep}_{\mathbb{Z}_p, \text{free}}(G_R),$$

is fully faithful. Here $\text{Rep}_{\mathbb{Z}_p, \text{free}}(G_R)$ denotes the category of finite free \mathbb{Z}_p -modules equipped with a continuous action of G_R .

Proof. The claim in (i) follows from [Fal89, Theorem 2.4] and [Tsu20, Proposition 66]. Further, the claim in (ii) follows from [Fal89, Theorem 2.4] and [Tsu20, Theorem 77]. ■

Definition 5.4. Let M be a free relative Fontaine-Laffaille module of level $[0, p-2]$, and set

$$T_{\text{cris}}(M) := \text{Hom}_{\mathbb{Z}_p}(T_{\text{cris}}^*(M), \mathbb{Z}_p),$$

which is a free \mathbb{Z}_p -module of rank $= \text{rk}_R M$, admitting a continuous action of G_R .

The main result of this section is as follows:

Theorem 5.5. *For a free relative Fontaine-Laffaille module M over R of level $[0, p-2]$, the associated representation $V_{\text{cris}}(M) := \mathbb{Q}_p \otimes_{\mathbb{Z}_p} T_{\text{cris}}(M)$ is a positive finite q -height representation (in the sense of Definition 4.9).*

The proof crucially exploits the computation of Fontaine [Fon94], Wach [Wac97] and Tsuji [Tsu20]. It follows in three steps: First, starting with a Fontaine-Laffaille module, we obtain an $\mathbf{A}_{R,\varpi}^{\text{PD}}$ -module using formal consequences of crystalline site for maps $\theta : \mathbf{A}_{R,\varpi}^{\text{PD}} \rightarrow R[\varpi]$, and $\theta_R : \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \rightarrow R[\varpi]$ (see Proposition 5.25, we also give an alternate proof of the proposition). Next, we exploit equivalence of categories obtained in Theorem 5.21 by extending scalars along $\mathbf{A}_{R,\varpi}^{\text{PD}} \rightarrow \mathbf{A}_{R,\varpi}^{\text{PD}}/I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}} \xleftarrow{\sim} \mathbf{A}_{R,\varpi}^+ / I^{(p-1)}\mathbf{A}_{R,\varpi}^+ \leftarrow \mathbf{A}_{R,\varpi}^+$. This gives us an $\mathbf{A}_{R,\varpi}^+$ -module with precise description of the Frobenius and the action of Γ_R (see Proposition 5.30). Finally, we descend over to the ring \mathbf{A}_R^+ by exploiting the Frobenius and Γ_R -action, thus obtaining a Wach module over \mathbf{A}_R^+ and proving the theorem (see §5.3.2).

For clarity of exposition and notational convenience in explaining the result of the first step, we start with preliminaries on some ideals of $\mathbf{A}_{R,\varpi}^+$ and $\mathbf{A}_{R,\varpi}^{\text{PD}}$ (appearing in the second step in the paragraph above) which will help us in proving categorical equivalence between certain modules over the concerned rings.

5.1. Some ideals of $\mathbf{A}_{R,\varpi}^+$ and $\mathbf{A}_{R,\varpi}^{\text{PD}}$. In this section, we will collect some technical results about the rings $\mathbf{A}_{R,\varpi}^+$ and $\mathbf{A}_{R,\varpi}^{\text{PD}}$ and some of their ideals. The results are motivated by the corresponding results over $\mathbf{A}_{\text{inf}}(\overline{R})$ and $\mathbf{A}_{\text{cris}}(\overline{R})$ and their respective ideals, studied in [Fon94, §5].

Lemma 5.6. *Let $a \in \mathbf{A}_{R,\varpi}^+$ such that $\mathbf{A}_{R,\varpi}^+ / p\mathbf{A}_{R,\varpi}^+$ is a -torsion free and a -adically complete. Then,*

- (i) $\mathbf{A}_{R,\varpi}^+$ is (p, a) -adically complete.
- (ii) For $n \in \mathbb{N}$, the rings $\mathbf{A}_{R,\varpi}^+ / a^n \mathbf{A}_{R,\varpi}^+$ are p -torsion free and p -adically complete.
- (iii) For $n \in \mathbb{N}$, $\mathbf{A}_{R,\varpi}^+$ and $\mathbf{A}_{R,\varpi}^+ / p^n \mathbf{A}_{R,\varpi}^+$ are a -torsion free and a -adically complete.
- (iv) The (p, a) -adic topology coincides with (p, π_m) -adic topology.

Proof. As $\mathbf{A}_{R,\varpi}^+$ is a flat \mathbb{Z}_p -algebra, claims (i), (ii) and (iii) follow from [Tsu20, Lemma 2]. The last claim follows from [Tsu20, Lemma 1] and the fact that $\mathbf{A}_{R,\varpi}^+ \subset \mathbf{A}_{\text{inf}}(\overline{R})$, where the former ring is equipped with the induced topology. ■

For $n \in \mathbb{N}$, let us write $n = (p-1)f(n) + r(n)$, with $r(n), f(n) \in \mathbb{N}$ and $0 \leq r(n) < p-1$. Let $t^{\{n\}} := \frac{t^n}{p^{f(n)}f(n)!}$ (resp. $t^{\{n\}} := \frac{t^n}{p^n n!}$ if $p=2$).

Lemma 5.7. *We have $t^{p-1} \in p\mathbf{A}_{R,\varpi}^{\text{PD}}$, therefore $t^{\{n\}} \in \mathbf{A}_{R,\varpi}^{\text{PD}}$.*

Proof. Note that we have $q = \frac{\varphi(\pi)}{\pi} = p\varphi\left(\frac{\pi}{t}\right)\frac{t}{\pi}$. Since $\frac{t}{\pi}$ is a unit in $\mathbf{A}_{R,\varpi}^{\text{PD}}$ (see Lemma 3.14), we get that q and p are associates in $\mathbf{A}_{R,\varpi}^{\text{PD}}$. But also, $q = \frac{\varphi(\pi)}{\pi} = \pi^{p-1} + p(\pi^{p-2} + \cdots + 1)$, i.e. $\pi^{p-1} \in p\mathbf{A}_{R,\varpi}^{\text{PD}}$. Again, using Lemma 3.14, we get that $t^{p-1} \in p\mathbf{A}_{R,\varpi}^{\text{PD}}$. ■

Note that we also have $\pi = \exp(t) - 1 = \sum_{n \geq 1} \frac{t^n}{n!} = \sum_{n \geq 1} c_n t^{\{n\}}$, where $c_n = \frac{p^{f(n)} f(n)!}{n!}$ (resp. $c_n = 2^n$ if $p = 2$) such that $c_n \rightarrow 0$ as $n \rightarrow +\infty$ (see [Fon94, §5.2.4]). Let

$$\Lambda := \left\{ \sum_{n \in \mathbb{N}} a_n t^{\{n\}} \text{ with } a_n \in O_F \text{ such that } a_n = 0 \text{ if } (p-1) \nmid n \text{ (resp. } 2 \nmid n \text{ if } p = 2) \right\}$$

be a ring and let $z = \sum_{a \in \mathbb{F}_p} [\varepsilon]^{[a]}$ (resp. $z = [\varepsilon] + [\varepsilon]^{-1}$ if $p = 2$) and $\pi_0 = z - p$, then we have $\pi_0 = (p-1) \sum_{n \geq 1, p-1 \mid n} \frac{t^n}{n!} \in \Lambda$ (resp. $\pi_0 = 2 \sum_{n \geq 1, 2 \mid n} \frac{t^n}{n!} \in \Lambda$ if $p = 2$). Further, we have that $\pi_0 \in p\Lambda$ (resp. $\pi_0 \in 8\Lambda$ if $p = 2$) and there exists $v \in \Lambda^\times$ such that $\frac{\pi_0}{p} = v \frac{t^{p-1}}{p}$ (resp. $\frac{\pi_0}{8} = v \frac{t^2}{8}$ if $p = 2$), see [Fon94, §5.2.5].

Next, recall that the filtration on $\mathbf{A}_{\text{cris}}(\overline{R})$ is given as $\text{Fil}^k \mathbf{A}_{\text{cris}}(\overline{R}) = \langle \xi^{[n]}, n \geq k \rangle \subset \mathbf{A}_{\text{cris}}(\overline{R})$, for $k \in \mathbb{N}$ (see §2.2). The filtration on $\mathbf{A}_{\text{inf}}(\overline{R})$ is defined as the induced filtration, i.e. $\text{Fil}^k \mathbf{A}_{\text{inf}}(\overline{R}) = \text{Fil}^k \mathbf{A}_{\text{cris}}(\overline{R}) \cap \mathbf{A}_{\text{inf}}(\overline{R}) = \xi^k \mathbf{A}_{\text{inf}}(\overline{R})$. Similarly, the filtration on $\mathbf{A}_{R,\varpi}^{\text{PD}}$ is again given by divided powers of ξ , i.e. $\text{Fil}^k \mathbf{A}_{R,\varpi}^{\text{PD}} = \langle \xi^{[n]}, n \geq k \rangle \subset \mathbf{A}_{R,\varpi}^{\text{PD}}$, for $k \in \mathbb{N}$ (see Definition 3.11). The filtration on $\mathbf{A}_{R,\varpi}^+$ is defined as the induced filtration, i.e. $\text{Fil}^k \mathbf{A}_{R,\varpi}^+ = \text{Fil}^k \mathbf{A}_{R,\varpi}^{\text{PD}} \cap \mathbf{A}_{R,\varpi}^+ = \xi^k \mathbf{A}_{\text{inf}}(\overline{R})$.

Now, for $k \in \mathbb{N}$ let us define an ideal of $\mathbf{A}_{\text{inf}}(\overline{R})$ as

$$I^{(k)} \mathbf{A}_{\text{inf}}(\overline{R}) = \{x \in \mathbf{A}_{\text{inf}}(\overline{R}) \text{ such that } \varphi^n(x) \in \text{Fil}^k \mathbf{A}_{\text{inf}}(\overline{R}) \text{ for } n \in \mathbb{N}\}.$$

Similarly, we can define respective ideals $I^{(k)} \mathbf{A}_{\text{cris}}(\overline{R}) \subset \mathbf{A}_{\text{cris}}(\overline{R})$, $I^{(k)} \mathbf{A}_{R,\varpi}^+ \subset \mathbf{A}_{R,\varpi}^+$ and $I^{(k)} \mathbf{A}_{R,\varpi}^{\text{PD}} \subset \mathbf{A}_{R,\varpi}^{\text{PD}}$. Since the natural map $\mathbf{A}_{R,\varpi}^+ \rightarrow \mathbf{A}_{\text{inf}}(\overline{R})$ is flat and we have $\mathbf{A}_{R,\varpi}^+ = \mathbf{A}_{\text{inf}}(\overline{R}) \cap \mathbf{A}_{R,\varpi} \subset W(\mathbb{C}(\overline{R})^\flat)$, we obtain that

Lemma 5.8. (i) *The ideal $I^{(k)} \mathbf{A}_{R,\varpi}^+$ is a principal ideal generated by π^k .*

(ii) *The element π_0 is a generator of $I^{(p-1)} \mathbf{A}_{R,\varpi}^+$ (resp. $I^{(2)} \mathbf{A}_{R,\varpi}^+$ if $p = 2$).*

(iii) *Let $S_0 = W[[\pi_0]]$ then there exists a unit $u \in S_0$ such that $\varphi(\pi_0) = u\pi_0 z^{p-1}$ (resp. $\varphi(\pi_0) = u\pi_0 z^2$ if $p = 2$).*

Proof. We only show the case $p \neq 2$, the claims for $p = 2$ follow analogously.

- (i) From the definitions it is clear that $I^{(k)} \mathbf{A}_{\text{inf}}(\overline{R}) \cap \mathbf{A}_{R,\varpi}^+ = I^{(k)} \mathbf{A}_{R,\varpi}^+$, where we take the intersection inside $\mathbf{A}_{\text{inf}}(\overline{R})$. Now, from [Fon94, §5.1.3, Proposition] we have that $I^{(k)} \mathbf{A}_{\text{inf}}(\overline{R}) = \pi^k \mathbf{A}_{\text{inf}}(\overline{R})$. Since the map $\mathbf{A}_{R,\varpi}^+ \rightarrow \mathbf{A}_{\text{inf}}(\overline{R})$ is flat and $\mathbf{A}_{R,\varpi}^+ = \mathbf{A}_{\text{inf}}(\overline{R}) \cap \mathbf{A}_{R,\varpi}$, we obtain that $I^{(k)} \mathbf{A}_{R,\varpi}^+ = I^{(k)} \mathbf{A}_{\text{inf}}(\overline{R}) \cap \mathbf{A}_{R,\varpi}^+ = \pi^k \mathbf{A}_{\text{inf}}(\overline{R}) \cap \mathbf{A}_{R,\varpi}^+ = \pi^k \mathbf{A}_{R,\varpi}^+$.
- (ii) Since $\pi_0 \in \mathbf{A}_{R,\varpi}^+$, the map $\mathbf{A}_{R,\varpi}^+ \rightarrow \mathbf{A}_{\text{inf}}(\overline{R})$ is flat and $\mathbf{A}_{R,\varpi}^+ = \mathbf{A}_{\text{inf}}(\overline{R}) \cap \mathbf{A}_{R,\varpi}$, we have $\pi_0 \mathbf{A}_{\text{inf}}(\overline{R}) \cap \mathbf{A}_{R,\varpi}^+ = \pi_0 \mathbf{A}_{R,\varpi}^+$. Now, from [Fon94, §5.2.6, Proposition (i)] we have that $I^{(p-1)} \mathbf{A}_{\text{inf}}(\overline{R}) = \pi_0 \mathbf{A}_{\text{inf}}(\overline{R})$. So we obtain that $I^{(p-1)} \mathbf{A}_{R,\varpi}^+ = I^{(p-1)} \mathbf{A}_{\text{inf}}(\overline{R}) \cap \mathbf{A}_{R,\varpi}^+ = \pi_0 \mathbf{A}_{\text{inf}}(\overline{R}) \cap \mathbf{A}_{R,\varpi}^+ = \pi_0 \mathbf{A}_{R,\varpi}^+$.
- (iii) This follows from [Fon94, §5.2.6, Proposition (ii)].

■

Proposition 5.9. *The continuous morphism of $\mathbf{A}_{R,\varpi}^+$ -algebras*

$$\begin{aligned} \alpha : \mathbf{A}_{R,\varpi}^+ \widehat{\otimes}_{S_0} \Lambda &\longrightarrow \mathbf{A}_{R,\varpi}^{\text{PD}} \\ \sum_{n \in \mathbb{N}} a_n \otimes \left(\frac{\pi_0}{p}\right)^{[n]} &\longmapsto \sum_{n \in \mathbb{N}} a_n \left(\frac{\pi_0}{p}\right)^{[n]}, \end{aligned}$$

is an isomorphism.

Proof. The proof follows in a manner similar to the proof of [Fon94, §5.2.7, Théorème]. We will only show the case $p \neq 2$, the claim for $p = 2$ follows analogously.

The homomorphism α in the claim is well defined and continuous since $\frac{\pi_0}{p} \in \text{Fil}^1 \mathbf{A}_{R,\varpi}^{\text{PD}}$. So we are left to show that α is an isomorphism. Since the source and targets are p -adically complete p -torsion-free rings, it is enough to show that α is an isomorphism modulo p .

Let $z_1 = \varphi^{-1}(z) \in \mathbf{A}_{R,\varpi}^+$. Note that $\mathbf{A}_{R,\varpi}^{\text{PD}}$ modulo p is the divided power envelope of $\mathbf{E}_{R,\varpi}^+$ with respect to the ideal generated by $\bar{z}_1 \equiv \bar{\xi} \pmod{p}$. Therefore, it is a free module over $\mathbf{E}_{R,\varpi}^+/\bar{z}_1^p$ with basis the images of $z_1^{[pn]}$, or equivalently $(\frac{z_1^p}{p})^{[n]}$. From Lemma 5.8 (iii), we have that $\varphi(\pi_0) = u\pi_0 z^{p-1}$, with $u \in S_0^\times$. Therefore, $\pi_0 = \varphi^{-1}(u)\varphi^{-1}(\pi_0)z_1^{p-1} = \varphi^{-1}(u)(z_1 - p)z_1^{p-1}$, which implies that $\mathbf{E}_{R,\varpi}^+/\bar{z}_1^p = \mathbf{E}_{R,\varpi}^+/\bar{\pi}_0$ and $\mathbf{A}_{R,\varpi}^{\text{PD}}$ modulo p is a free module over $\mathbf{E}_{R,\varpi}^+/\bar{\pi}_0$ with basis the images of $(\frac{\pi_0}{p})^{[n]}$. Since it is immediate that the same is true for $\mathbf{A}_{R,\varpi}^+ \otimes_{S_0} \Lambda$ modulo p , we get the claim. ■

Lemma 5.10. *For $k \in \mathbb{N}$ the ideal $I^{(k)} \mathbf{A}_{R,\varpi}^{\text{PD}}$ is a divided power ideal which is the associated $\mathbf{A}_{R,\varpi}^+$ -submodule of $\mathbf{A}_{R,\varpi}^{\text{PD}}$ generated by $t^{\{n\}}$ for $n \geq k$.*

Proof. The proof follows in a manner similar to the proof of [Fon94, §5.3.5, Proposition]. Let $J^{(k)}$ be the $\mathbf{A}_{R,\varpi}^+$ -submodule of $\mathbf{A}_{R,\varpi}^{\text{PD}}$ generated by $t^{\{n\}}$ for $n \geq k$. It is straightforward to check that $J^{(k)} \subset I^{(k)}$, and $J^{(k)}$ is a divided power ideal. Thus it remains to show that $I^{(k)} \subset J^{(k)}$. We will show this by induction on k . The case $k = 0$ is trivial.

Now suppose $k \geq 1$ and $x \in I^{(k)}$. The induction hypothesis allows us to write $x = \sum_{n \geq k-1} a_n t^{\{n\}}$ where $a_n \in \mathbf{A}_{R,\varpi}^+$ goes to 0 as $n \rightarrow +\infty$. If $b = a_{n-1}$, we have $a = bt^{\{n-1\}} + a'$ where $a' \in J^{(k)} \subset I^{(k)}$, thus $bt^{\{k-1\}} \in I^{(k)}$. But $\varphi^s(bt^{\{k-1\}}) = p^{(k-1)s} \varphi^s(b) t^{\{k-1\}} = c_{k,s} \varphi^s(b) t^{\{k-1\}}$, where $c_{k,s}$ is a nonzero rational number. Since $t^{k-1} \in \text{Fil}^{k-1} \mathbf{A}_{R,\varpi}^{\text{PD}} \setminus \text{Fil}^k \mathbf{A}_{R,\varpi}^{\text{PD}}$, one has $b \in I^{(1)} \mathbf{A}_{R,\varpi}^{\text{PD}} \cap \mathbf{A}_{R,\varpi}^+$, which is the principal ideal generated by π . Thus $bt^{\{k-1\}}$ belongs to an ideal of $\mathbf{A}_{R,\varpi}^{\text{PD}}$ generated by $\pi t^{\{k-1\}}$. But $\frac{t}{\pi} \in \mathbf{A}_{R,\varpi}^{\text{PD}}$ is a unit (see Lemma 3.14). Hence, $bt^{\{k-1\}}$ belongs to an ideal generated by $t \cdot t^{\{k-1\}}$, which is contained in $J^{(k)}$. ■

Following is an immediate consequence of Lemma 5.10:

Corollary 5.11. *For $k \in \mathbb{N}$, consider the homomorphism $\mathbf{A}_{R,\varpi}^+ \rightarrow I^{(k)} \mathbf{A}_{R,\varpi}^{\text{PD}}$ sending $x \mapsto x \cdot t^{\{k\}}$. Then, the induced map $\mathbf{A}_{R,\varpi}^+ / I^{(1)} \mathbf{A}_{R,\varpi}^+ \rightarrow I^{(k)} \mathbf{A}_{R,\varpi}^{\text{PD}} / I^{(k+1)} \mathbf{A}_{R,\varpi}^{\text{PD}}$ is bijective.*

Now, from [Fon94, §5.3.5, Proposition], we have a natural isomorphism $\mathbf{A}_{\text{inf}}(\bar{R}) / I^{(k)} \mathbf{A}_{\text{inf}}(\bar{R}) \xrightarrow{\sim} \mathbf{A}_{\text{cris}}(\bar{R}) / I^{(k)} \mathbf{A}_{\text{cris}}(\bar{R})$, for $0 \leq k \leq p-1$. A similar statement is true in our setting:

Proposition 5.12. *For $k \in \mathbb{N}$, the rings $\mathbf{A}_{R,\varpi}^+ / I^{(k)} \mathbf{A}_{R,\varpi}^+$ and $\mathbf{A}_{R,\varpi}^{\text{PD}} / I^{(k)} \mathbf{A}_{R,\varpi}^{\text{PD}}$ are p -torsion free. Moreover, if $0 \leq k \leq p-1$, then the natural map $\mathbf{A}_{R,\varpi}^+ / I^{(k)} \mathbf{A}_{R,\varpi}^+ \rightarrow \mathbf{A}_{R,\varpi}^{\text{PD}} / I^{(k)} \mathbf{A}_{R,\varpi}^{\text{PD}}$ is an isomorphism.*

Proof. The proof follows from arguments similar to the proof of [Fon94, §5.3.5, Proposition]. First, note that for every $k \in \mathbb{N}$, $\mathbf{A}_{R,\varpi}^{\text{PD}} / \text{Fil}^k \mathbf{A}_{R,\varpi}^{\text{PD}}$ is torsion free. Further, the kernel of the map

$$\begin{aligned} \mathbf{A}_{R,\varpi}^{\text{PD}} &\longrightarrow (\mathbf{A}_{R,\varpi}^{\text{PD}} / \text{Fil}^k \mathbf{A}_{R,\varpi}^{\text{PD}})^{\mathbb{N}} \\ x &\longmapsto (\varphi^n(x) \pmod{\text{Fil}^k \mathbf{A}_{R,\varpi}^{\text{PD}}})_{k \in \mathbb{N}}, \end{aligned}$$

is $I^{(k)} \mathbf{A}_{R,\varpi}^{\text{PD}}$. Therefore, $\mathbf{A}_{R,\varpi}^+ / I^{(k)} \mathbf{A}_{R,\varpi}^+ \hookrightarrow \mathbf{A}_{R,\varpi}^{\text{PD}} / I^{(k)} \mathbf{A}_{R,\varpi}^{\text{PD}} \hookrightarrow (\mathbf{A}_{R,\varpi}^{\text{PD}} / \text{Fil}^k \mathbf{A}_{R,\varpi}^{\text{PD}})^{\mathbb{N}}$, which implies that the former two rings are torsion free.

Now from Proposition 5.9 and Lemma 5.10, it follows that as $\mathbf{A}_{R,\varpi}^+$ -module, $\mathbf{A}_{R,\varpi}^{\text{PD}} / I^{(k)} \mathbf{A}_{R,\varpi}^{\text{PD}}$ is generated by the images of $(\frac{\pi_0}{p})^{[n]}$ for $0 \leq (p-1)n < k$. For $0 \leq k \leq p-1$, we have that $(\frac{\pi_0}{p})^{[n]} \in \mathbf{A}_{R,\varpi}^+$, hence we get the claim. ■

Next, we mention a lemma useful for the proof of Proposition 5.20.

Lemma 5.13. (i) For $0 \leq k < j$, we have that $I^{(k)}\mathbf{A}_{R,\varpi}^{\text{PD}}/I^{(j)}\mathbf{A}_{R,\varpi}^{\text{PD}}$ is p -torsion free.

(ii) For $k \in \mathbb{N}$, we have that $I^{(k)}\mathbf{A}_{R,\varpi}^{\text{PD}}$ is p -adically complete.

Proof. (i) The proof is similar to the proof of [Tsu99, Lemma A3.19 (1)]. Let $x \in I^{(k)}\mathbf{A}_{R,\varpi}^{\text{PD}}$ and assume that $px \in I^{(j)}\mathbf{A}_{R,\varpi}^{\text{PD}}$. Then $p\varphi^i(x) \in \text{Fil}^j\mathbf{A}_{R,\varpi}^{\text{PD}}$ for all $i \in \mathbb{N}$. Since $\mathbf{A}_{R,\varpi}^{\text{PD}}/\text{Fil}^j\mathbf{A}_{R,\varpi}^{\text{PD}} \subset \mathbf{A}_{\text{cris}}(\overline{R})/\text{Fil}^j\mathbf{A}_{\text{cris}}(\overline{R})$ is p -torsion free (see [Tsu99, Lemma A2.11 (2)]), we get that $\varphi^i(x) \in \text{Fil}^j\mathbf{A}_{R,\varpi}^{\text{PD}}$ for all $i \in \mathbb{N}$, i.e. $x \in I^{(j)}\mathbf{A}_{R,\varpi}^{\text{PD}}$.

(ii) The proof is similar to the proof of [Tsu99, Lemma A3.27]. We will prove the statement by induction on k . For $k = 0$, the statement is trivial by the definition of $\mathbf{A}_{R,\varpi}^{\text{PD}}$. Next, from part (i) and Corollary 5.11, we have that $I^{(k)}\mathbf{A}_{R,\varpi}^{\text{PD}}/I^{(k+1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$ is p -torsion free and p -adically complete. Therefore, we obtain exact sequences

$$0 \longrightarrow \lim_n (I^{(k+1)}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes \mathbb{Z}/p^n\mathbb{Z}) \longrightarrow \lim_n (I^{(k)}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes \mathbb{Z}/p^n\mathbb{Z}) \longrightarrow I^{(k)}\mathbf{A}_{R,\varpi}^{\text{PD}}/I^{(k+1)}\mathbf{A}_{R,\varpi}^{\text{PD}} \longrightarrow 0.$$

The statement now follows by induction on k . ■

5.2. Equivalence of categories. In [Tsu20], Tsuji has established a relationship between free relative Fontaine-Laffaille modules (see Definition 5.1) and $\mathbf{A}_{\text{inf}}(\overline{R})$ -representations as well as $\mathbf{A}_{\text{cris}}(\overline{R})$ -representations of G_R (in a precise functorial manner). Tsuji's computations are motivated by computations of Wach in [Wac97] for the arithmetic case.

Recall from §5.1 that for $k \in \mathbb{N}$ we have the ideal

$$I^{(k)}\mathbf{A}_{\text{inf}}(\overline{R}) = \{x \in \mathbf{A}_{\text{inf}}(\overline{R}) \text{ such that } \varphi^n(x) \in \text{Fil}^k\mathbf{A}_{\text{inf}}(\overline{R}) \text{ for } n \in \mathbb{N}\}.$$

Similarly, we define respective ideals $I^{(k)}\mathbf{A}_{\text{cris}}(\overline{R}) \subset \mathbf{A}_{\text{cris}}(\overline{R})$, $I^{(k)}\mathbf{A}_{R,\varpi}^+ \subset \mathbf{A}_{R,\varpi}^+$ and $I^{(k)}\mathbf{A}_{R,\varpi}^{\text{PD}} \subset \mathbf{A}_{R,\varpi}^{\text{PD}}$. Given a free Fontaine-Laffaille module, in [Tsu20, §5] Tsuji functorially obtains an $\mathbf{A}_{\text{cris}}(\overline{R})$ -module (in a manner similar to Proposition 5.25). Further, he exploits the isomorphism $\mathbf{A}_{\text{inf}}(\overline{R})/I^{(p-1)}\mathbf{A}_{\text{inf}}(\overline{R}) \xrightarrow{\sim} \mathbf{A}_{\text{cris}}(\overline{R})/I^{(p-1)}\mathbf{A}_{\text{cris}}(\overline{R})$, to construct an $\mathbf{A}_{\text{inf}}(\overline{R})$ -representation of G_R . The last step is carried out by establishing certain equivalence of categories. Tsuji's computations are general and follows from certain assumptions on the structure of the rings and modules, one is studying. In this section, we will recall and verify those assumptions in our case, which would help us in establishing equivalence between several categories (see Theorem 5.21).

Let $A = \mathbf{A}_{R,\varpi}^+$, $\mathbf{A}_{R,\varpi}^+/I^{(p-1)}\mathbf{A}_{R,\varpi}^+$, $\mathbf{A}_{R,\varpi}^{\text{PD}}$, or $\mathbf{A}_{R,\varpi}^{\text{PD}}/I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$.

Lemma 5.14. Let $q = \frac{\varphi(\pi)}{\pi} \in A$, then q is a non-zero-divisor in A .

Proof. For $A = \mathbf{A}_{R,\varpi}^+$ and $\mathbf{A}_{R,\varpi}^{\text{PD}}$, the claim follows from the definitions. Next, note that we have $q = \frac{\varphi(\pi)}{\pi} = \pi^{p-1} + pu \in \mathbf{A}_{R,\varpi}^+$ for some unit $u \in \mathbf{A}_{R,\varpi}^+$, in particular, $q \equiv pu \pmod{\pi^{p-1}}$. Now since $I^{(p-1)}\mathbf{A}_{R,\varpi}^+ = \pi^{p-1}\mathbf{A}_{R,\varpi}^+$ by Lemma 5.8 (ii), we obtain that q and p are associates in $\mathbf{A}_{R,\varpi}^+/I^{(p-1)}\mathbf{A}_{R,\varpi}^+ \xrightarrow{\sim} \mathbf{A}_{R,\varpi}^{\text{PD}}/I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$. As the rings $\mathbf{A}_{R,\varpi}^+/I^{(p-1)}\mathbf{A}_{R,\varpi}^+$ and $\mathbf{A}_{R,\varpi}^{\text{PD}}/I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$ are p -torsion free by Proposition 5.12, we get the claim. ■

Next, note that we have $\text{Fil}^0 A = A$ and $\text{Fil}^i A \cdot \text{Fil}^j A \subset \text{Fil}^{i+j} A$ for $i, j \in \mathbb{Z}$, and $\varphi(\text{Fil}^k A) \subset q^k A$ for $k \in \mathbb{N}$. In particular, we see that our choice of A and q satisfies [Tsu20, Condition 39].

Definition 5.15. Define the category $\text{MF}_{[0,p-2],\text{free}}^q(A, \varphi, \Gamma_R)$ as follows: An object is a triplet $(N, \text{Fil}^k N, \varphi)$ such that,

- (i) N is a free A -module of rank h .
- (ii) The filtration $\text{Fil}^k N$ is decreasing and there exists an A -basis of N as $\{e_1, \dots, e_h\}$ and integers $k_1, \dots, k_h \in \mathbb{N}_{\leq p-2}$ such that we have $\text{Fil}^k N = \sum_{i=1}^h \text{Fil}^{k-k_i} A e_i$ for $0 \leq k \leq p-2$.
- (iii) A Frobenius-semilinear endomorphism $\varphi : N \rightarrow N$ such that we have $\varphi(\text{Fil}^k N) \subset q^k N$ for $0 \leq k \leq p-2$, and $\sum_{k=0}^{p-2} A \cdot q^{-k} \varphi(\text{Fil}^k N) = N$.
- (iv) N is equipped with a continuous action of Γ_R such that $\text{Fil}^k N$ is stable under this action, and the endomorphism φ commutes with the action of Γ_R .

A morphism between two objects of the category $\text{MF}_{[0, p-2], \text{free}}^q(A, \varphi, \Gamma_R)$ is a continuous A -linear morphism commuting with the endomorphism φ and the action of Γ_R on each side.

Notation 5.16. Abusing notations, we will denote $(N, \text{Fil}^k N, \partial, \Phi) \in \text{MF}_{[0, p-2], \text{free}}^q(A, \varphi, \Gamma_R)$ by N and say that it has filtration of level $[0, p-2]$.

Remark 5.17. In $\mathbf{A}_{R, \varpi}^{\text{PD}}$, note that we can write $q = p \frac{t}{\pi} \varphi(\frac{\pi}{t})$, and since $\frac{t}{\pi}$ is a unit in $\mathbf{A}_{R, \varpi}^{\text{PD}}$ (see Lemma 3.14), we obtain that q and p are associates in $\mathbf{A}_{R, \varpi}^{\text{PD}}$. Therefore, for $A = \mathbf{A}_{R, \varpi}^{\text{PD}}$ in Definition 5.15, we can replace q by p . Further, since $q = \pi^{p-1} + pu$ for $u \in (\mathbf{A}_{R, \varpi}^+)^{\times}$ and π^{p-1} generates $I^{(p-1)} \mathbf{A}_{R, \varpi}^+$ (see Lemma 5.8 (ii)), we obtain that $q \equiv pu \pmod{I^{(p-1)} \mathbf{A}_{R, \varpi}^+}$, i.e. q and p are associates in $\mathbf{A}_{R, \varpi}^+ / I^{(p-1)} \mathbf{A}_{R, \varpi}^+ \xrightarrow{\sim} \mathbf{A}_{R, \varpi}^{\text{PD}} / I^{(p-1)} \mathbf{A}_{R, \varpi}^{\text{PD}}$. Therefore, for $A = \mathbf{A}_{R, \varpi}^+ / I^{(p-1)} \mathbf{A}_{R, \varpi}^+$ in Definition 5.15, we can replace q by p , and similarly for $\mathbf{A}_{R, \varpi}^{\text{PD}} / I^{(p-1)} \mathbf{A}_{R, \varpi}^{\text{PD}}$.

Lemma 5.18 ([Tsu20, Lemma 41]). *Let $(N, \text{Fil}^k N)$ be as in Definition 5.15 (i), (ii). Then a Frobenius-semilinear endomorphism $\varphi : N \rightarrow N$ satisfies the conditions in Definition 5.15 (iii) if and only if $\varphi(e_i) \in q^{k_i} N$ for $1 \leq i \leq h$ and $\{q^{-k_1} \varphi(e_1), \dots, q^{-k_h} \varphi(e_h)\}$ is an A -basis of N .*

Proof. Let us assume that $(N, \text{Fil}^k N)$ satisfies the condition in Definition 5.15 (iii). Then, since $e_i \in \text{Fil}^{k_i} N$, we have $\varphi(e_i) \in q^{k_i} N$ for $1 \leq i \leq h$. Now for $0 \leq k \leq p-2$, we have

$$\varphi(\text{Fil}^{k-k_i} A e_i) = \varphi(\text{Fil}^{k-k_i} A) \varphi(e_i) \subset q^k A \cdot q^{-k_i} \varphi(e_i) \subset q^k N.$$

Therefore, from the identity $\sum_{k=0}^{p-2} A \cdot q^{-k} \varphi(\text{Fil}^k N) = N$, we obtain that $\{q^{-k_1} \varphi(e_1), \dots, q^{-k_h} \varphi(e_h)\}$ generate N as an A -module. Since N is free of rank h over A , we get that $\{q^{-k_1} \varphi(e_1), \dots, q^{-k_h} \varphi(e_h)\}$ is indeed a basis.

Conversely, assume that $\varphi(e_i) \in q^{k_i} N$ for $1 \leq i \leq h$ such that elements $\{q^{-k_1} \varphi(e_1), \dots, q^{-k_h} \varphi(e_h)\}$ form an A -basis of N . Then, from Definition 5.15 (ii), we have

$$\varphi(\text{Fil}^k N) = \varphi\left(\sum_{i=1}^h \text{Fil}^{k-k_i} A e_i\right) \subset \sum_{i=1}^h q^{k-k_i} A \varphi(e_i) = q^k \sum_{i=1}^h A \cdot q^{-k_i} \varphi(e_i) = q^k N.$$

Further, since $\{q^{-k_1} \varphi(e_1), \dots, q^{-k_h} \varphi(e_h)\} \in \sum_{k=0}^{p-2} A \cdot q^{-k} \varphi(\text{Fil}^k N)$, we obtain the last equality in Definition 5.15 (iii). \blacksquare

Now we introduce some necessary conditions in order to adapt Tsuji's results from [Tsu20, §4 - §8].

Condition 5.19. Let $A = \mathbf{A}_{R, \varpi}^+$, $\mathbf{A}_{R, \varpi}^{\text{PD}}$ and $q = \frac{\varphi(\pi)}{\pi} \in A$. Consider the projection map $A \twoheadrightarrow A/J$ for some ideal $J \subset A$ and assume that

- (i) The ideal J is contained in the Jacobson radical of A , and $J \subset \text{Fil}^{p-2} A$. Moreover, $\varphi(J) \subset J$ and $\varphi(J) \subset q^{p-1} A$. Further, the ideal J is preserved under the action of Γ_R .
- (ii) The ideal J is closed as a submodule of A .

- (iii) There exists a decreasing sequence of ideals $\cdots \subset H_{n+1} \subset H_n \subset \cdots \subset H_0 \subset A$ for $n \in \mathbb{N}$, such that H_n form a fundamental system of neighborhoods of 0 in A , the homomorphism $A \rightarrow \lim_n A/H_n$ is an isomorphism, and $q^{-(p-1)}\varphi(H_n \cap J) \subset H_n \cap J$ for every $n \in \mathbb{N}$.
- (iv) The image of q in A/J is a non-zero-divisor. Moreover, the sequence $\prod_{k=0}^n \varphi^k(q) \in A$ converges to 0 as $n \rightarrow +\infty$.
- (v) The homomorphism $\varphi : A \rightarrow A$ is continuous and multiplication by q induces a homeomorphism $A \rightarrow qA$, where the latter is equipped with the induced topology.

Proposition 5.20. (i) Let $A = \mathbf{A}_{R,\varpi}^+$ with $J = I^{(p-1)}\mathbf{A}_{R,\varpi}^+$, and $H_n = p^n\mathbf{A}_{R,\varpi}^+ + \pi^{n+p-1}\mathbf{A}_{R,\varpi}^+$. Then $\mathbf{A}_{R,\varpi}^+$ satisfies Condition 5.19.

- (ii) Let $A = \mathbf{A}_{R,\varpi}^{\text{PD}}$ with $J = I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$, and $H_n = p^n\mathbf{A}_{R,\varpi}^{\text{PD}}$. Then $\mathbf{A}_{R,\varpi}^{\text{PD}}$ satisfies Condition 5.19.

Proof. The proof follows in a manner similar to the proof of [Tsu20, Proposition 59].

- (i) The ring $\mathbf{A}_{R,\varpi}^+$ is π -adically complete, and since $I^{(p-1)}\mathbf{A}_{R,\varpi}^+ = \pi^{p-1}\mathbf{A}_{R,\varpi}^+ \subset \pi\mathbf{A}_{R,\varpi}^+$, we see that $I^{(p-1)}\mathbf{A}_{R,\varpi}^+$ is contained in the Jacobson radical of $\mathbf{A}_{R,\varpi}^+$. Moreover, note that we have inclusions $\varphi(\pi^{p-1}\mathbf{A}_{R,\varpi}^+) \subset q^{p-1}\pi^{p-1}\mathbf{A}_{R,\varpi}^+ \subset \pi^{p-1}\mathbf{A}_{R,\varpi}^+$, therefore we get $I^{(p-1)}\mathbf{A}_{R,\varpi}^+ \subset \text{Fil}^{p-2}\mathbf{A}_{R,\varpi}^+$. It is clear that $I^{(p-1)}\mathbf{A}_{R,\varpi}^+$ is stable under the action of Γ_R . Therefore, Condition 5.19 (i) is satisfied.

Now we have $H_n = p^n\mathbf{A}_{R,\varpi}^+ + \pi^{n+p-1}\mathbf{A}_{R,\varpi}^+$ for $n \in \mathbb{N}$, which is a fundamental system of neighborhoods of $0 \in \mathbf{A}_{R,\varpi}^+$ and $\mathbf{A}_{R,\varpi}^+ = \lim_n \mathbf{A}_{R,\varpi}^+/H_n$ (see Lemma 5.6). Further, since $\mathbf{A}_{R,\varpi}^+/I^{(p-1)}\mathbf{A}_{R,\varpi}^+$ is p -torsion free, we obtain that $H_n \cap I^{(p-1)}\mathbf{A}_{R,\varpi}^+ = (p^n\mathbf{A}_{R,\varpi}^+ + \pi^{n+p-1}\mathbf{A}_{R,\varpi}^+) \cap I^{(p-1)}\mathbf{A}_{R,\varpi}^+ = p^n I^{(p-1)}\mathbf{A}_{R,\varpi}^+ + \pi^n I^{(p-1)}\mathbf{A}_{R,\varpi}^+$. The Condition 5.19 (iii) now follows from this. Moreover, $I^{(p-1)}\mathbf{A}_{R,\varpi}^+$ is a free $\mathbf{A}_{R,\varpi}^+$ -module of rank 1, so it follows that J is a closed submodule of $\mathbf{A}_{R,\varpi}^+$ by Lemma 5.6 (i) & (iv), verifying Condition 5.19 (ii).

Next, from Lemma 5.14, it follows that q is a non-zero-divisor in $\mathbf{A}_{R,\varpi}^+/I^{(p-1)}\mathbf{A}_{R,\varpi}^+$. Further, for $k \in \mathbb{N}$, we have $\varphi^k(q) = \varphi^{k+1}(\xi) \in \varphi^{k+1}(\text{Fil}^1\mathbf{A}_{R,\varpi}^+) \subset \varphi^{k+1}(p\mathbf{A}_{R,\varpi}^+ + \pi\mathbf{A}_{R,\varpi}^+) \subset p\mathbf{A}_{R,\varpi}^+ + \pi\mathbf{A}_{R,\varpi}^+$. Therefore, $\prod_{k=0}^n \varphi^k(q)$ converges to 0 as $n \rightarrow +\infty$, and Condition 5.19 (iv) has been verified.

By the definition of φ in §3.3, we see that it is continuous. Further, from Lemma 5.6 (iii), it follows that $\mathbf{A}_{R,\varpi}^+/q\mathbf{A}_{R,\varpi}^+$ is p -torsion free. Therefore, we have $(p^n\mathbf{A}_{R,\varpi}^+ + q^{n+1}\mathbf{A}_{R,\varpi}^+) \cap q\mathbf{A}_{R,\varpi}^+ = p^n(q\mathbf{A}_{R,\varpi}^+) + q^{n+1}\mathbf{A}_{R,\varpi}^+$. By Lemma 5.6 (i), it follows that $\mathbf{A}_{R,\varpi}^+ \xrightarrow{\times q} q\mathbf{A}_{R,\varpi}^+$ is a homeomorphism, which verifies Condition 5.19 (v).

- (ii) Note that we have $q = p\varphi(\frac{\pi}{t})\frac{t}{\pi}$, which implies that q and p are associates in $\mathbf{A}_{R,\varpi}^{\text{PD}}$ (see Lemma 3.14). Therefore, it is enough to verify Condition 5.19, with q replaced by p everywhere.

We have $I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}} \subset \text{Fil}^1\mathbf{A}_{R,\varpi}^{\text{PD}} + p\mathbf{A}_{R,\varpi}^{\text{PD}}$, $\mathbf{A}_{R,\varpi}^{\text{PD}}$ is p -adically complete and $\text{Fil}^1\mathbf{A}_{R,\varpi}^{\text{PD}}/p\text{Fil}^1\mathbf{A}_{R,\varpi}^{\text{PD}}$ is a nil ideal of $\mathbf{A}_{R,\varpi}^{\text{PD}}/p\mathbf{A}_{R,\varpi}^{\text{PD}}$. Therefore, $I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$ is contained in the Jacobson radical of $\mathbf{A}_{R,\varpi}^{\text{PD}}$. Further, by definitions we have $I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}} \subset \text{Fil}^{p-2}\mathbf{A}_{R,\varpi}^{\text{PD}}$ and $\varphi(I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}) \subset I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$. Also, $\varphi(I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}) \subset q^{p-1}\mathbf{A}_{R,\varpi}^{\text{PD}} = p^{p-1}\mathbf{A}_{R,\varpi}^{\text{PD}}$. It is clear that $I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$ is stable under the action of Γ_R . Therefore, Condition 5.19 (i) is satisfied.

Next, we know that $\mathbf{A}_{R,\varpi}^{\text{PD}}$ is p -adically complete and the ring $\mathbf{A}_{R,\varpi}^{\text{PD}}/I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$ is p -torsion free by Proposition 5.12, therefore $p^n\mathbf{A}_{R,\varpi}^{\text{PD}} \cap I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}} = p^n I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$. This

gives us Condition 5.19 (iii). Further, $I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$ is p -adically complete by Lemma 5.13 (ii), so we get Condition 5.19 (ii).

Condition 5.19 (iv) & (v) follow trivially from the fact that $\mathbf{A}_{R,\varpi}^{\text{PD}} = \lim_n \mathbf{A}_{R,\varpi}^{\text{PD}}/p^n \mathbf{A}_{R,\varpi}^{\text{PD}}$. ■

Finally, we come to the main result of this section. Note that the categories MF^p below are defined by combining Definition 5.15 and Remark 5.17.

Theorem 5.21. *The natural maps $\mathbf{A}_{R,\varpi}^+ \rightarrow \mathbf{A}_{R,\varpi}^+/I^{(p-1)}\mathbf{A}_{R,\varpi}^+ \xrightarrow{\sim} \mathbf{A}_{R,\varpi}^{\text{PD}}/I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}} \leftarrow \mathbf{A}_{R,\varpi}^{\text{PD}}$, induce equivalence of categories:*

$$\begin{aligned} \text{MF}_{[0,p-2],\text{free}}^p(\mathbf{A}_{R,\varpi}^{\text{PD}}, \varphi, \Gamma_R) &\xrightarrow[\sim]{(1)} \text{MF}_{[0,p-2],\text{free}}^p(\mathbf{A}_{R,\varpi}^{\text{PD}}/I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}, \varphi, \Gamma_R) \\ &\xleftarrow[\sim]{(2)} \text{MF}_{[0,p-2],\text{free}}^p(\mathbf{A}_{R,\varpi}^+/I^{(p-1)}\mathbf{A}_{R,\varpi}^+, \varphi, \Gamma_R) \\ &\xleftarrow[\sim]{(3)} \text{MF}_{[0,p-2],\text{free}}^q(\mathbf{A}_{R,\varpi}^+, \varphi, \Gamma_R). \end{aligned}$$

Proof. The natural projection map $\mathbf{A}_{R,\varpi}^+ \rightarrow \mathbf{A}_{R,\varpi}^+/I^{(p-1)}\mathbf{A}_{R,\varpi}^+$ is compatible with Frobenius and the action of Γ_R and we have $q \equiv pu \pmod{I^{(p-1)}\mathbf{A}_{R,\varpi}^+}$ for $u \in (\mathbf{A}_{R,\varpi}^+)^\times$ (see also Remark 5.17), i.e. q and p are associates in $\mathbf{A}_{R,\varpi}^+/I^{(p-1)}\mathbf{A}_{R,\varpi}^+$. Further, $\mathbf{A}_{R,\varpi}^+$ satisfies Condition 5.19. Therefore, from [Tsu20, Proposition 56], we obtain that the functor in (3) is an equivalence of categories.

Next, from Proposition 5.12, we have an isomorphism of rings $\mathbf{A}_{R,\varpi}^+/I^{(p-1)}\mathbf{A}_{R,\varpi}^+ \sim \mathbf{A}_{R,\varpi}^{\text{PD}}/I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$, compatible with Frobenius and the action of Γ_R . Therefore, we obtain that the functor in (2) is an equivalence of categories.

Finally, the natural projection map $\mathbf{A}_{R,\varpi}^{\text{PD}} \rightarrow \mathbf{A}_{R,\varpi}^{\text{PD}}/I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$ is compatible with Frobenius and the action of Γ_R and $q \equiv pu \pmod{I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}}$ for some $u \in (\mathbf{A}_{R,\varpi}^{\text{PD}})^\times$ (see also Remark 5.17), i.e. q and p are associates in $\mathbf{A}_{R,\varpi}^{\text{PD}}/I^{(p-1)}\mathbf{A}_{R,\varpi}^{\text{PD}}$. Further, $\mathbf{A}_{R,\varpi}^{\text{PD}}$ satisfies Condition 5.19. Therefore, from [Tsu20, Proposition 56], we obtain that the functor in (1) is an equivalence of categories. ■

5.3. Wach modules from Fontaine-Laffaille data. In this section, we will work with objects of $\text{MF}_{[0,p-2],\text{free}}(R, \Phi, \partial)$ (see Definition 5.1) and using these objects we will obtain Wach modules over \mathbf{A}_R^+ (see Definition 4.8). In §5.3.1, starting with a Fontaine-Laffaille module, we will first obtain a free module over $\mathbf{A}_{R,\varpi}^+$ with desired properties and in §5.3.2 we will descend over to \mathbf{A}_R^+ . Note that in §5.3.1, we will first establish a mod p^n statement (see (5.2)) and as a consequence deduce a p -adic statement (see Proposition 5.25). However, it is possible to prove the p -adic statement directly (see another proof of Proposition 5.25). Readers interested only in the p -adic statement can directly skip to Proposition 5.25.

5.3.1. From Fontaine-Laffaille modules to $\mathbf{A}_{R,\varpi}^+$ -modules. Following [Tsu20, §4], for $n \in \mathbb{N}_{>0}$ we set $X_n = \text{Spec}(R/p^n)$ and $\Sigma_n = \text{Spec}(\mathcal{O}_F/p^n)$ and consider big crystalline sites $\text{CRIS}(X_n, \Sigma_n)$ and $\text{CRIS}(X_1, \Sigma_n)$, and respective topos $(X_n/\Sigma_n)_{\text{CRIS}}$ and $(X_1/\Sigma_n)_{\text{CRIS}}$, with the PD-ideal $(p(\mathcal{O}_F/p^n), [])$. Let $F_{\Sigma_n} : \Sigma_n \rightarrow \Sigma_n$ denote a lifting of the absolute Frobenius of Σ_1 , such that it is a PD-morphism with respect to the PD-structure. The absolute Frobenius F_{X_1} of X_1 and F_{Σ_n} define a morphism of PD-ringtopos $F_{X_1/\Sigma_n, \text{CRIS}} : (X_1/\Sigma_n)_{\text{CRIS}} \rightarrow (X_1/\Sigma_n)_{\text{CRIS}}$.

Let $(M, \text{Fil}^\bullet M, \partial, \Phi) \in \text{MF}_{[0,p-2],\text{free}}(R, \Phi, \partial)$ be a free relative Fontaine-Laffaille module (see Definition 5.1), and let $(M_n, \text{Fil}^\bullet M_n, \partial, \Phi)$ denote its modulo p^n reduction. Then, by [Tsu20, Definition 26, Theorems 17 & 29] this data corresponds to a quasi-coherent filtered crystal $(\mathcal{F}_n, \text{Fil}^\bullet \mathcal{F}_n)$ on $\text{CRIS}(X_n/\Sigma_n)$. Similarly, by [Tsu20, Definition 26, Theorems 22 & 29] this

data also corresponds to a quasi-coherent crystal \mathcal{G}_n on $\text{CRIS}(X_1/\Sigma_n)$. The reduction modulo p^n of $\Phi : \varphi^*M \rightarrow M$ equip \mathcal{G}_n with a morphism $\Phi_{\mathcal{G}_n} : F_{X_1/\Sigma_n, \text{CRIS}}^*(\mathcal{G}_n) \rightarrow \mathcal{G}_n$. Further, for the morphism of ringed topoi $i_{n, \text{CRIS}} : (X_1/\Sigma_n)_{\text{CRIS}} \rightarrow (X_n/\Sigma_n)_{\text{CRIS}}$ induced by the closed immersion $i_n : X_1 \rightarrow X_n$ over id_{Σ_n} , we have $i_{n, \text{CRIS}}^*(\mathcal{F}_n) = \mathcal{G}_n$ (see [Tsu20, Propositions 25 & 32]). Moreover, we have similar statements for the morphism of ringed topoi induced by $X_n \rightarrow X_{n+1}$ and $\Sigma_n \rightarrow \Sigma_{n+1}$.

Now, for $n \in \mathbb{N}_{>0}$ let $X'_n := \text{Spec}(R[\varpi]/p^n)$, $D_n := \text{Spec}(\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n)$ and $F_{D_n} : D_n \rightarrow D_n$ be the lifting of the absolute Frobenius on D_1 defined by φ of $\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n$. We have the surjective map $\theta : \mathbf{A}_{R, \varpi}^+ \twoheadrightarrow R[\varpi]$. So taking mod p^n reduction, we obtain an embedding $X'_n \hookrightarrow \text{Spec}(\mathbf{A}_{R, \varpi}^+/p^n)$ and taking divided power envelope, we obtain a closed immersion $X'_n \hookrightarrow D_n$ (resp. $X'_1 \hookrightarrow D_n$) which can naturally be regarded as an object of the site $\text{CRIS}(X_n/\Sigma_n)$ (resp. $\text{CRIS}(X_1/\Sigma_n)$), endowed with a right action of Γ_R .

Definition 5.22. Define an $\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n$ -module by setting $N_n^{\text{PD}} := \Gamma(X'_n \hookrightarrow D_n, \mathcal{F}_n) \xrightarrow{\sim} \Gamma(X'_1 \hookrightarrow D_n, \mathcal{G}_n)$.

The right action of Γ_R on D_n induces a left action on N_n^{PD} . The filtration on \mathcal{F}_n induces a filtration by $\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n$ -submodules on N_n^{PD} , which is stable under the Γ_R -action. Then N_n^{PD} is a finite free filtered $\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n$ -module of level $[0, p-2]$ (see [Tsu20, Lemma 20]). The Frobenius $\Phi_{\mathcal{G}_n}$ of \mathcal{G}_n and the lifting of Frobenius F_{D_n} on D_n define a semilinear Γ_R -equivariant endomorphism of $\Gamma(X'_1 \hookrightarrow D_n, \mathcal{G}_n)$ and hence that of N_n^{PD} as $\Gamma(X'_1 \hookrightarrow D_n, \mathcal{G}_n) \rightarrow \Gamma(X'_1 \hookrightarrow D_n, F_{X_1, \text{CRIS}}^*\mathcal{G}_n) \xrightarrow{\Phi_{\mathcal{G}_n}} \Gamma(X'_1 \hookrightarrow D_n, \mathcal{G}_n)$, where the first homomorphism is induced by $F_{X'_1}$ and F_{D_n} .

Let $[\]$ denote the PD-structure on the ideal $p(\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n) + \text{Fil}^1 \mathbf{A}_{R, \varpi}^{\text{PD}}/p^n$ of $\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n$. Then we have the big crystalline sites $\text{CRIS}(X'_n/D_n)$ and $\text{CRIS}(X'_1/D_n)$, and the respective topoi $(X'_n/D_n)_{\text{CRIS}}$ and $(X'_1/D_n)_{\text{CRIS}}$ with the PD-ideal $(p(\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n) + \text{Fil}^1 \mathbf{A}_{R, \varpi}^{\text{PD}}/p^n, [\])$ of $\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n$. By taking the pullback of $(\mathcal{F}_n, \text{Fil}^\bullet \mathcal{F}_n)$ (resp. \mathcal{G}_n) under the morphism of ringed topoi $(X'_n/D_n)_{\text{CRIS}} \rightarrow (X_n/\Sigma_n)_{\text{CRIS}}$ (resp. $(X'_1/D_n)_{\text{CRIS}} \rightarrow (X_1/\Sigma_n)_{\text{CRIS}}$), we obtain a quasi-coherent filtered crystal $(\mathcal{F}'_n, \text{Fil}^\bullet \mathcal{F}'_n)$ (resp. a quasi-coherent crystal \mathcal{G}'_n with a morphism $\Phi_{\mathcal{G}'_n} : F_{X'_1/D_n, \text{CRIS}}^*(\mathcal{G}'_n) \rightarrow \mathcal{G}'_n$), endowed with compatible Γ_R -action. Since $X'_n \hookrightarrow D_n$ (resp. $X'_1 \hookrightarrow D_n$) is a final object of $\text{CRIS}(X'_n/D_n)$ (resp. $\text{CRIS}(X'_1/D_n)$), we have canonical $\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n$ -linear isomorphisms $N_n^{\text{PD}} \xrightarrow{\sim} \Gamma((X'_n/D_n)_{\text{CRIS}}, \mathcal{F}'_n) \xrightarrow{\sim} \Gamma((X'_1/D_n)_{\text{CRIS}}, \mathcal{G}'_n)$ compatible with supplementary structures (see [Tsu20, p. 188-189]).

Next, for $n \in \mathbb{N}_{>0}$, similar to above let $E_n := \text{Spec}(\mathcal{O}\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n)$ and $F_{E_n} : E_n \rightarrow E_n$ be the lifting of the absolute Frobenius on E_1 defined by φ of $\mathcal{O}\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n$. We have the surjective map $\theta_R : R \otimes_{\mathbb{Z}} \mathbf{A}_{R, \varpi}^+ \twoheadrightarrow R[\varpi]$. So taking mod p^n reduction, we have an embedding $X'_n \hookrightarrow \text{Spec}(R \otimes_{\mathbb{Z}} \mathbf{A}_{R, \varpi}^+/p^n)$ and taking divided power envelope, we obtain a closed immersion $X'_n \hookrightarrow E_n$ (resp. $X'_1 \hookrightarrow E_n$) which can naturally be regarded as an object of the site $\text{CRIS}(X'_n/D_n)$ (resp. $\text{CRIS}(X'_1/D_n)$), endowed with a right action of Γ_R .

Definition 5.23. Define an $\mathcal{O}\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n$ -module by setting $\mathcal{O}N_n^{\text{PD}} := \Gamma(X'_n \hookrightarrow E_n, \mathcal{F}'_n) \xrightarrow{\sim} \Gamma(X'_1 \hookrightarrow E_n, \mathcal{G}'_n)$.

The right action of Γ_R on E_n induces a left action on $\mathcal{O}N_n^{\text{PD}}$. The filtration on \mathcal{F}'_n induces a filtration by $\mathcal{O}\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n$ -submodules on $\mathcal{O}N_n^{\text{PD}}$, which is stable under the Γ_R -action. Then $\mathcal{O}N_n^{\text{PD}}$ is a finite free filtered $\mathcal{O}\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n$ -module of level $[0, p-2]$ (see [Tsu20, Lemma 20]). Further, by [Tsu20, Theorem 29, Proposition 32] $\mathcal{O}N_n^{\text{PD}}$ is equipped with a Γ_R -equivariant integrable connection compatible with the connection on $\mathcal{O}\mathbf{A}_{R, \varpi}^{\text{PD}}/p^n$ and satisfying Griffiths transversality with the respect to the filtration. Moreover, this the Γ_R -action and connection are compatible with the respective structures on $\Gamma(X'_1 \hookrightarrow E_n, \mathcal{G}'_n)$ (see [Tsu20, Propositions 25 & 32]). The Frobenius $\Phi_{\mathcal{G}'_n}$ of \mathcal{G}'_n and the lifting of Frobenius F_{E_n} on E_n define a semilinear Γ_R -equivariant endomorphism φ of $\Gamma(X'_1 \hookrightarrow E_n, \mathcal{G}'_n)$ and hence that of $\mathcal{O}N_n^{\text{PD}}$. Further, the Frobenius-semilinear endomorphism φ commutes with the connection on $\mathcal{O}N_n^{\text{PD}}$.

From [Ber74, Proposition 4.1.4] and [BO78, Theorem 7.1], we have a description of the global sections of a crystal in terms of horizontal sections of the corresponding module with an integrable connection on the PD-envelope of an embedding into a smooth scheme. In other words, we have an $\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n$ -linear isomorphism

$$N_n^{\text{PD}} \xrightarrow{\sim} (\mathcal{O}N_n^{\text{PD}})^{\partial=0},$$

compatible with filtration, Frobenius and the action of Γ_R on each side (see [Tsu20, p. 190]). Since $X'_n \rightarrow D_n$ (resp. $X'_1 \rightarrow D_n$) is a final object of $\text{CRIS}(X'_n/D_n)$ (resp. $\text{CRIS}(X'_1/D_n)$), we obtain a canonical $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n$ -linear isomorphism

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n \otimes_{\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n} N_n^{\text{PD}} \xrightarrow{\sim} \mathcal{O}N_n^{\text{PD}},$$

compatible with Frobenius, filtration, connection and the action of Γ_R on each side. Here the connection on the tensor product on the left is given as $\partial_{\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}} \otimes 1$. Moreover, from [Tsu20, Propositions 24, 25 & 32], we obtain an $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n$ -linear isomorphism

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n \otimes_{R/p^n} M/p^n \xrightarrow{\sim} \mathcal{O}N_n^{\text{PD}},$$

compatible with Frobenius, filtration, connection and the action of Γ_R on each side (see [Tsu20, p. 191]). Here the connection on the tensor product on the left is given as $\partial_{\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}} \otimes 1 + 1 \otimes \partial_M$. Combining the two isomorphisms above, we obtain an $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n$ -linear isomorphism

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n \otimes_{\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n} N_n^{\text{PD}} \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n \otimes_{R/p^n} M/p^n, \quad (5.2)$$

compatible with Frobenius, filtration, connection and the action of Γ_R on each side. Therefore, we also have an $\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n$ -linear isomorphism

$$N_n^{\text{PD}} \xrightarrow{\sim} (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}/p^n \otimes_{R/p^n} M/p^n)^{\partial=0}$$

compatible with Frobenius, filtration and the action of Γ_R on each side.

Definition 5.24. Define an $\mathbf{A}_{R,\varpi}^{\text{PD}}$ -module as $N^{\text{PD}}(M) := \lim_n N_n^{\text{PD}}$, equipped with a semilinear and continuous action of Γ_R , a filtration given as $\text{Fil}^k N^{\text{PD}}(M) := \lim_n \text{Fil}^k N_n^{\text{PD}}$, which is stable under the action of Γ_R , and a Frobenius-semilinear Γ_R -equivariant endomorphism φ .

Passing to the limit in (5.2) we obtain an $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ -linear isomorphism

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_{R,\varpi}^{\text{PD}}} N^{\text{PD}}(M) \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M,$$

compatible with Frobenius, filtration, connection and the action of Γ_R on each side. Therefore, we have the following conclusion:

Proposition 5.25. *Let M be a free relative Fontaine-Laffaille module. Then*

$$N^{\text{PD}}(M) := (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M)^{\partial=0},$$

is a finite free $\mathbf{A}_{R,\varpi}^{\text{PD}}$ -module equipped with a decreasing filtration of level $[0, p-2]$, a Frobenius-semilinear endomorphism $\varphi : N^{\text{PD}}(M) \rightarrow N^{\text{PD}}(M)$ and a continuous action of Γ_R on each side. In particular, $N^{\text{PD}}(M) \in \text{MF}_{[0,p-2], \text{free}}^p(\mathbf{A}_{R,\varpi}^{\text{PD}}, \varphi, \Gamma_R)$. Further, we have a natural isomorphism

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_{R,\varpi}^{\text{PD}}} N^{\text{PD}}(M) \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M, \quad (5.3)$$

compatible with the Frobenius, filtration, connection and the action of Γ_R on each side.

Another proof of Proposition 5.25. Let us consider the injective map $R \rightarrow \mathbf{A}_{R,\varpi}^{\text{PD}}$ sending $X_i \rightarrow [X_i^b]$.

Lemma 5.26. *We have an $\mathbf{A}_{R,\varpi}^{\text{PD}}$ -linear isomorphism $\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M \xrightarrow{\sim} (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M)^{\partial=0}$.*

Proof. Let $J = ([X_1^b] - X_1, \dots, [X_d^b] - X_d)\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ and let $J^{[n]}$ denote its n -th divided power for $n \geq 1$. We have the projection map,

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M \longrightarrow \mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M,$$

via the map $X_i \mapsto [X_i^b]$ and the kernel is given as $J^{[1]}\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M$. Moreover, we have an $\mathbf{A}_{R,\varpi}^{\text{PD}}$ -linear section of the projection above given as

$$\begin{aligned} \mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M &\longrightarrow \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M \\ 1 \otimes d &\longmapsto \sum_{\mathbf{k} \in \mathbb{N}^d} \prod_{i=1}^d \partial_i^{k_i}(d) \prod_{i=1}^d ([X_i^b] - X_i)^{[k_i]}. \end{aligned} \quad (5.4)$$

Note that the image of the section lies in $(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M)^{\partial=0}$. Now let $Q = J^{[1]}\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M$ and $Q' = (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M)/(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M)^{\partial=0}$ and we consider the following diagram with exact rows (the top row is split exact)

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M & \longrightarrow & \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M & \longrightarrow & Q \longrightarrow 0 \\ & & \downarrow & & \parallel & & \downarrow \\ 0 & \longrightarrow & (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M)^{\partial=0} & \longrightarrow & \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M & \longrightarrow & Q' \longrightarrow 0. \end{array}$$

Note that the left vertical arrow is an injection and the right vertical arrow is a surjection. To get that the left vertical arrow is a bijection we need to show that the right vertical arrow is an injection. We have

$$(J^{[1]}\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M)^{\partial=0} \subset (J^{[1]}\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{R[\frac{1}{p}]} \mathcal{O}\mathbf{D}_{\text{cris}}(V))^{\partial=0} = (J^{[1]}\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} V)^{\partial=0},$$

where $V = V_{\text{cris}}(M)$ is crystalline (see Proposition 5.3 (i)) and the divided power ideal $J^{[1]}\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \subset \mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R})$ is generated by $([X_1^b] - X_1, \dots, [X_d^b] - X_d)$. Then it easily follows that $(J^{[1]}\mathcal{O}\mathbf{B}_{\text{cris}}(\overline{R}) \otimes_{\mathbb{Q}_p} V)^{\partial=0} = 0$ and we conclude that $\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M \xrightarrow{\sim} (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M)^{\partial=0}$. ■

From the identification $N^{\text{PD}}(M) = (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M)^{\partial=0} \xrightarrow{\sim} \mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M$ (where the rightmost term is equipped with a Γ_R -action as in Remark 5.27), it easily follows that $N^{\text{PD}}(M) \in \text{MF}_{[0,p-2],\text{free}}^p(\mathbf{A}_{R,\varpi}^{\text{PD}}, \varphi, \Gamma_R)$. Next, we can $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ -linearly extend the map in (5.4) to obtain

$$\begin{aligned} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_{R,\varpi}^{\text{PD}}} (\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M) &\longrightarrow \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M \\ 1 \otimes d &\longmapsto \sum_{\mathbf{k} \in \mathbb{N}^d} \prod_{i=1}^d \partial_i^{k_i}(d) \prod_{i=1}^d ([X_i^b] - X_i)^{[k_i]}. \end{aligned} \quad (5.5)$$

We equip the left term with a Γ_R -action as in Remark 5.27. Choosing a basis of M it is easy to see that the determinant of the map in (5.5) is invertible in $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$, i.e. the map (5.5) is bijective. Moreover, it is compatible with Frobenius, filtration, connection and the action of Γ_R . Now we have a natural injective map

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_{R,\varpi}^{\text{PD}}} N^{\text{PD}}(M) \longrightarrow \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M,$$

compatible with the Frobenius, filtration, connection and the action of Γ_R on each side. The map above is bijective because of the following commutative diagram

$$\begin{array}{ccc}
\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_{R,\varpi}^{\text{PD}}} N^{\text{PD}}(M) & \xrightarrow{\quad} & \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M \\
\downarrow \wr & & \parallel \\
\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_{R,\varpi}^{\text{PD}}} (\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M) & \xrightarrow{\sim} & \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M,
\end{array}$$

where the bottom horizontal arrow is the isomorphism in (5.5). This concludes the proof. \blacksquare

Remark 5.27. Using the $\mathbf{A}_{R,\varpi}^{\text{PD}}$ -linear isomorphism in Lemma 5.26, $\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M \xrightarrow{\sim} (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M)^{\partial=0}$, we can describe the action of Γ_R on the left term explicitly. The action can be given by the formula $g(a \otimes d) = g(a) \otimes \sum_{\mathbf{k} \in \mathbb{N}^d} \prod_{i=1}^d \partial_i^{k_i}(d) \prod_{i=1}^d (g([X_i^b]) - [X_i^b])^{[k_i]}$, for $g \in \Gamma_R$.

Lemma 5.28. *Let $N^{\text{PD}}(M)$ as in Proposition 5.25. Then, the action of $\Gamma_{R,\varpi}$ is trivial on $N^{\text{PD}}(M)/\pi N^{\text{PD}}(M)$, whereas $\Gamma_R/\Gamma_{R,\varpi}$ acts trivially over $N^{\text{PD}}(M)/\pi_m N^{\text{PD}}(M)$.*

Proof. This follows from the Γ_R -equivariant isomorphism in (5.3) (or from Lemma 5.26 and Remark 5.27) and the action of Γ_R on $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ (see Lemma 4.24 (i)). \blacksquare

Proposition 5.29. *The functor*

$$\begin{aligned}
N^{\text{PD}} : \text{MF}_{[0,p-2],\text{free}}(R, \Phi, \partial) &\longrightarrow \text{MF}_{[0,p-2],\text{free}}^p(\mathbf{A}_{R,\varpi}^{\text{PD}}, \varphi, \Gamma_R) \\
M &\longmapsto N^{\text{PD}}(M) = (\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M)^{\partial=0},
\end{aligned}$$

is fully faithful.

Proof. By taking Γ_R -invariants in (5.3), we obtain an R -linear isomorphism $(\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_{R,\varpi}^{\text{PD}}} N^{\text{PD}}(M))^{\Gamma_R} \xrightarrow{\sim} M$ compatible with Frobenius, filtration, connection on each side, and functorial in M . \blacksquare

Having obtained a finite free module with desired structures over the ring $\mathbf{A}_{R,\varpi}^{\text{PD}}$, we will now pass to the ring $\mathbf{A}_{R,\varpi}^+$. Let $M \in \text{MF}_{[0,p-2],\text{free}}(R, \Phi, \partial)$ and $N^{\text{PD}}(M) \in \text{MF}_{[0,p-2],\text{free}}^p(\mathbf{A}_{R,\varpi}^{\text{PD}}, \varphi, \Gamma_R)$ the $\mathbf{A}_{R,\varpi}^{\text{PD}}$ -module obtained under the functor of Proposition 5.29.

Next, from Theorem 5.21, we have an equivalence of categories $\text{MF}_{[0,p-2],\text{free}}^p(\mathbf{A}_{R,\varpi}^{\text{PD}}, \varphi, \Gamma_R) \xrightarrow{\sim} \text{MF}_{[0,p-2],\text{free}}^q(\mathbf{A}_{R,\varpi}^+, \varphi, \Gamma_R)$ sending $N^{\text{PD}}(M) \mapsto N^+(M)$ for M as above. Combining this with Propositions 5.25 & 5.29, we obtain:

Proposition 5.30. *The functor*

$$\begin{aligned}
N^+ : \text{MF}_{[0,p-2],\text{free}}(R, \Phi, \partial) &\longrightarrow \text{MF}_{[0,p-2],\text{free}}^q(\mathbf{A}_{R,\varpi}^+, \varphi, \Gamma_R) \\
M &\longmapsto N^+(M),
\end{aligned}$$

is fully faithful. Further, for M and $N^+(M)$ as above, we have a natural isomorphism

$$\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_{\mathbf{A}_{R,\varpi}^+} N^+(M) \xrightarrow{\sim} \mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}} \otimes_R M, \quad (5.6)$$

compatible with the Frobenius, filtration, connection and the action of Γ_R on each side.

Lemma 5.31. *Let $N^+(M)$ as in Proposition 5.30. Then, the action of $\Gamma_{R,\varpi}$ is trivial on $N^+(M)/\pi N^+(M)$, whereas $\Gamma_R/\Gamma_{R,\varpi}$ acts trivially over $N^+(M)/\pi_m N^+(M)$.*

Proof. This follows from the Γ_R -equivariant isomorphism in (5.6) and the action of Γ_R on $\mathcal{O}\mathbf{A}_{R,\varpi}^{\text{PD}}$ (see Lemma 4.24 (i)). \blacksquare

5.3.2. Obtaining Wach modules. For the rest of this section we will fix $m = 1$ (fix $m = 2$ if $p = 2$), i.e. we take $K = F(\zeta_p)$ (take $K = F(\zeta_{p^2})$ if $p = 2$). Consider the localization $S = \mathbf{A}_{R,\varpi}^+[\frac{1}{\pi_1}]$. Let M and M' be free relative Fontaine-Laffaille modules and $N^+(M)$ and $N^+(M')$ the respective $\mathbf{A}_{R,\varpi}^+$ -modules obtained by the functor in Proposition 5.30.

Lemma 5.32. *We have a natural bijection*

$$\mathrm{Hom}_{\mathbf{A}_{R,\varpi}^+, \Gamma_R}(N^+(M), N^+(M')) \xrightarrow{\sim} \mathrm{Hom}_{S, \Gamma_R}(N^+(M)[\frac{1}{\pi_1}], N^+(M')[\frac{1}{\pi_1}]). \quad (5.7)$$

Proof. As we are working with free modules and the morphism of rings $\mathbf{A}_{R,\varpi}^+ \rightarrow \mathbf{A}_{R,\varpi}^+[\frac{1}{\pi_1}] = S$ is flat, we obtain that (5.7) is injective. To check surjectivity, let $f : N^+(M)[\frac{1}{\pi_1}] \rightarrow N^+(M')[\frac{1}{\pi_1}]$ be an S -linear and Γ_R -equivariant morphism. We need to show that $f(N^+(M)) \subset N^+(M')$. Assume $f(N^+(M)) \subset \pi_1^{-k} N^+(M')$ for $k \in \mathbb{N}$, and consider the reduction of f modulo π , which is again Γ_R -equivariant. Now from Lemma 5.31, we have that Γ_R acts trivially over $N^+(M)/\pi_1 N^+(M)$, whereas the action of Γ_R is non-trivial over $\pi_1^{-k} N^+(M')/\pi_1^{-k+1} N^+(M')$ for $k \neq 0$ (the action of $\gamma_0 \in \Gamma_K$ is non-trivial for $k \neq 0$). Hence, we must have $k = 0$, i.e. $f(N^+(M)) \subset N^+(M')$, which shows the claim. \blacksquare

Note that we have a morphism $\varphi : S = \mathbf{A}_{R,\varpi}^+[\frac{1}{\pi_1}] \rightarrow \mathbf{A}_{R,\varpi}^+[\frac{1}{\pi}]$. The respective Frobenius-semilinear endomorphisms φ on $N^+(M)$ and $N^+(M')$ induce semilinear morphisms $\varphi : N^+(M)[\frac{1}{\pi_1}] \rightarrow N^+(M)[\frac{1}{\pi}]$ and $\varphi : N^+(M')[\frac{1}{\pi_1}] \rightarrow N^+(M')[\frac{1}{\pi}]$. Let $f \in \mathrm{Hom}_{S, \Gamma_R}(N^+(M)[\frac{1}{\pi_1}], N^+(M')[\frac{1}{\pi_1}])$ be a morphism, such that the following diagram commutes

$$\begin{array}{ccc} N^+(M)[\frac{1}{\pi_1}] & \xrightarrow{f} & N^+(M')[\frac{1}{\pi_1}] \\ \downarrow \varphi & & \downarrow \varphi \\ N^+(M)[\frac{1}{\pi}] & \xrightarrow{f} & N^+(M')[\frac{1}{\pi}], \end{array}$$

where the bottom horizontal arrow is well-defined due to Lemma 5.32. We will call such a morphism f to be (φ, Γ_R) -equivariant.

Lemma 5.33. *We have a natural bijection*

$$\mathrm{Hom}_{\mathbf{A}_{R,\varpi}^+, \varphi, \Gamma_R}(N^+(M), N^+(M')) \xrightarrow{\sim} \mathrm{Hom}_{S, \varphi, \Gamma_R}(N^+(M)[\frac{1}{\pi_1}], N^+(M')[\frac{1}{\pi_1}]).$$

Proof of Theorem 5.5. Let $M \in \mathrm{MF}_{[0,p-2], \mathrm{free}}(R, \Phi, \partial)$ and let $N^+(M)$ denote the $\mathbf{A}_{R,\varpi}^+$ -module obtained from M from the functor of Proposition 5.30. We will show that a basis of $N^+(M)$ descends over to \mathbf{A}_R^+ .

In the notation of Definition 5.15, let $\{e_1, \dots, e_h\}$ denote an $\mathbf{A}_{R,\varpi}^+$ -basis of $N^+(M)$. Then from Lemma 5.18, we get that $\{q^{-k_1}\varphi(e_1), \dots, q^{-k_h}\varphi(e_h)\}$ is also an $\mathbf{A}_{R,\varpi}^+$ -basis of $N^+(M)$. Without loss of generality, we may further assume that $k_h \leq k_{h-1} \leq \dots \leq k_1$. Let us set $s := k_1$, so we get that $N^+(M)/\varphi^*(N^+(M))$ is killed by q^s and $s \in \mathbb{N}$ is the smallest such number.

Let $D(M) := N^+(M)[\frac{1}{\pi_1}]^\wedge$, where $^\wedge$ denotes the p -adic completion. Then $D(M)$ is an étale $(\varphi, \Gamma_{R,\varpi})$ -module over $\mathbf{A}_{R,\varpi} = \mathbf{A}_{R,\varpi}^+[\frac{1}{\pi_1}]^\wedge$, free of rank h . Since $N^+(M)$ is free, it follows that $N^+(M)[\frac{1}{\pi_1}]/p^n \xrightarrow{\sim} D(M)/p^n$. Similar to the proof of Lemma 5.32 and using dévissage we obtain that the functor $N^+(M) \mapsto D(M)$ is fully faithful. Therefore, using Proposition 5.30 we conclude that the functor

$$\begin{aligned} \mathrm{MF}_{[0,p-2], \mathrm{free}}(R, \Phi, \partial) &\longrightarrow (\varphi, \Gamma_R)\text{-Mod}_{\mathbf{A}_{R,\varpi}}^{\mathrm{ét}} \\ M &\longmapsto N^+(M)[\frac{1}{\pi_1}]^\wedge, \end{aligned}$$

is fully faithful.

Now, from Proposition 5.3 and Definition 5.4 we have that $T := T_{\text{cris}}(M)$ is a free \mathbb{Z}_p -representation of G_R . Considering T as a representation of $G_{R,\varpi}$, we have the associated $(\varphi, \Gamma_{R,\varpi})$ -module $\mathbf{D}_{R,\varpi}(T)$ over $\mathbf{A}_{R,\varpi}$. By the full faithfulness of the functor above and equivalence of categories in (3.2), we conclude that $D(M) \xrightarrow{\sim} \mathbf{D}_{R,\varpi}(T)$ as étale $(\varphi, \Gamma_{R,\varpi})$ -module over $\mathbf{A}_{R,\varpi}$. Also, we have $\varphi(\mathbf{D}_{R,\varpi}(T)) \subset \mathbf{D}(T)$, where the latter module is an étale (φ, Γ_R) -module over \mathbf{A}_R , free of rank h .

Next, let $N := N^+(M) \cap \mathbf{D}(T)$ where we take the intersection inside $\mathbf{D}_{R,\varpi}(T)$. Note that N is equipped with a Frobenius-semilinear endomorphism φ and it is stable under the action of Γ_R . We claim that

Lemma 5.34. *The elements $\{q^{-k_1}\varphi(e_1), \dots, q^{-k_h}\varphi(e_h)\}$ form a basis of N .*

Proof. Let us set $N' := \sum_{i=1}^h \mathbf{A}_R^+ q^{-k_i} \varphi(e_i)$. Since we have $q^{-k_i} \varphi(e_i) \in N^+(M) \cap \mathbf{D}(T) = N$, therefore $N' \subset N$. This also implies that $\varphi(e_i) \in q^{k_i} N$. Extending scalars along the faithfully flat morphism of rings $\mathbf{A}_R^+ \rightarrow \mathbf{A}_{R,\varpi}^+$, we get that $N^+(M) = \mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} N' \subset \mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} N \subset N^+(M)$. Therefore, $\mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} N' \xrightarrow{\sim} \mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} N$. But since the map $\mathbf{A}_R^+ \rightarrow \mathbf{A}_{R,\varpi}^+$ is faithfully flat, we obtain that $N' \xrightarrow{\sim} N$. \blacksquare

We will now verify the conditions of Definition 4.9 for $V = \mathbb{Q}_p \otimes_{\mathbb{Z}_p} T$. Since V arises from a Fontaine-Laffaille module of level $[0, p-2]$, we have that V is crystalline with non-positive Hodge-Tate weights. We have that N is a free \mathbf{A}_R^+ -module of rank h stable under φ and Γ_R , and such that $N \subset \mathbf{D}^+(T)$ as well as $\mathbf{A}_R \otimes_{\mathbf{A}_R^+} N \xrightarrow{\sim} \mathbf{D}(T)$. Next, we want to show that $q^s N \subset \varphi^*(N)$ as \mathbf{A}_R^+ -modules, where $s = k_1$. Since $\mathbf{A}_R^+ \rightarrow \mathbf{A}_{R,\varpi}^+$ is faithfully flat, it is equivalent to showing that $q^s \mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} N \subset \mathbf{A}_{R,\varpi}^+ \otimes_{\mathbf{A}_R^+} \varphi^*(N)$. But the latter inclusion can be re-expressed as $q^s N^+(M) \subset \varphi^*(N^+(M))$ as $\mathbf{A}_{R,\varpi}^+$ -modules, which was established above by showing that $N^+(M)/\varphi^*(N^+(M))$ is killed by q^s . Therefore, we conclude that $N/\varphi^*(N)$ is killed by q^s and $s \in \mathbb{N}$ is the smallest such number.

Next, we look at the action of Γ_R over N . Recall from §3.1 that we have $\{\gamma_0, \gamma_1, \dots, \gamma_d\}$ as topological generators of $\Gamma_{R,\varpi}$, where γ_0 is a lift of a topological generator of Γ_K . The action of γ_j on the basis elements of $N^+(M)$ can be given as

$$\gamma_j(e_i) = e_i + \pi x_{i,j} \text{ for } 1 \leq i \leq h, 0 \leq j \leq d \text{ and } x_{i,j} \in \mathbf{A}_{R,\varpi}^+.$$

Since φ is Γ_R -equivariant, we get that $\gamma_j(\varphi(e_i)) = \varphi(e_i) + q\pi\varphi(x_{i,j})$, where $\varphi(x_{i,j}) \in \varphi(N^+(M)) \subset N^+(M) \cap \mathbf{D}(T) = N$. Now $\varphi(e_i) \in q^{k_i} N$, so we must have that $q\pi\varphi(x_{i,j}) \in q^{k_i} N \cap q\pi N = q^{k_i} \pi N \subset N$, for $1 \leq i \leq h$ and $0 \leq j \leq d$. Therefore, we get that

$$\gamma_j(q^{-k_i} \varphi(e_i)) \equiv q^{-k_i} \varphi(e_i) \pmod{\pi N} \text{ for } 1 \leq j \leq d.$$

For $j = 0$, recall that $\gamma_0(\pi) = \chi(\gamma_0)\pi u$ for some unit $u \in 1 + \pi\mathbf{A}_R^+$. Therefore, we have $\gamma_0(q) = q\varphi(u)u^{-1}$ and $\gamma_0(q^{-1}) = q^{-1}\varphi(u^{-1})u$. So we obtain

$$\gamma_0(q^{-k_i} \varphi(e_i)) = \gamma_0(q^{-k_i})\gamma_0(\varphi(e_i)) = q^{-k_i} \varphi(u^{-k_i})u^{k_i}(\varphi(e_i) + q\pi\varphi(x_{i,j})) \equiv q^{-k_i} \varphi(e_i) \pmod{\pi N}.$$

Finally, let $g \in \Gamma_R$ be a lift of a generator $\bar{g} \in \Gamma_R/\Gamma_{R,\varpi}$, a finite group of order $p-1$. Then we have $g(e_i) = e_i + \pi_1 y_i$ for $1 \leq i \leq h$ and $y_i \in N^+(M)$. Since φ is Γ_R -equivariant, we get that $g(\varphi(e_i)) = \varphi(e_i) + \pi\varphi(y_i)$, where $\varphi(y_i) \in \varphi(N^+(M)) \subset N^+(M) \cap \mathbf{D}(T) = N$. Now $\varphi(e_i) \in q^{k_i} N$, so we must have that $\pi\varphi(y_i) \in q^{k_i} N \cap \pi N = q^{k_i} \pi N \subset N$, for $1 \leq i \leq h$. Further, we know that $g(\pi) = \chi(g)\pi v$ for some unit $v \in 1 + \pi\mathbf{A}_R^+$, which gives us that $g(q) = q\varphi(v)v^{-1}$. Therefore, $g(q^{-k_i} \varphi(e_i)) = q^{-k_i} \varphi(u^{-k_i})u^{k_i}(\varphi(e_i) + \pi\varphi(y_i)) \equiv q^{-k_i} \varphi(e_i) \pmod{\pi N}$, for $1 \leq i \leq h$. Hence, Γ_R acts trivially over $N/\pi N$.

Setting $\mathbf{N}(T) := N$, we see that conditions of Definition 4.9 have been satisfied. In particular, V is a positive finite q -height representation. \blacksquare

References

- [Abh21] Abhinandan. “Représentations de hauteur finie et complexe syntomique”. Theses. Université de Bordeaux, Nov. 2021.
- [Abh22] Abhinandan. “Syntomic complex and p -adic nearby cycles”. In: *preprint* (2022). Available at: https://abhinandan.perso.math.cnrs.fr/syntomic_complex.pdf.
- [And06] Fabrizio Andreatta. “Generalized ring of norms and generalized (ϕ, Γ) -modules”. In: *Ann. Sci. École Norm. Sup. (4)* 39.4 (2006), pp. 599–647. ISSN: 0012-9593.
- [AB08] Fabrizio Andreatta and Olivier Brinon. “Surconvergence des représentations p -adiques: le cas relatif”. In: *Astérisque* 319 (2008). Représentations p -adiques de groupes p -adiques. I. Représentations galoisiennes et (ϕ, Γ) -modules, pp. 39–116. ISSN: 0303-1179.
- [AI08] Fabrizio Andreatta and Adrian Iovita. “Global applications of relative (φ, Γ) -modules. I”. In: *Astérisque* 319 (2008). Représentations p -adiques de groupes p -adiques. I. Représentations galoisiennes et (φ, Γ) -modules, pp. 339–420. ISSN: 0303-1179.
- [Ben00] Denis Benois. “On Iwasawa theory of crystalline representations”. In: *Duke Math. J.* 104.2 (2000), pp. 211–267. ISSN: 0012-7094.
- [BB08] Denis Benois and Laurent Berger. “Théorie d’Iwasawa des représentations cristallines. II”. In: *Comment. Math. Helv.* 83.3 (2008), pp. 603–677. ISSN: 0010-2571.
- [Ber02] Laurent Berger. “Représentations p -adiques et équations différentielles”. In: *Invent. Math.* 148.2 (2002), pp. 219–284. ISSN: 0020-9910.
- [Ber04] Laurent Berger. “Limites de représentations cristallines”. In: *Compos. Math.* 140.6 (2004), pp. 1473–1498. ISSN: 0010-437X.
- [BB10] Laurent Berger and Christophe Breuil. “Sur quelques représentations potentiellement cristallines de $\mathrm{GL}_2(\mathbb{Q}_p)$ ”. In: *Astérisque* 330 (2010), pp. 155–211. ISSN: 0303-1179.
- [Ber74] Pierre Berthelot. *Cohomologie cristalline des schémas de caractéristique $p > 0$* . Lecture Notes in Mathematics, Vol. 407. Springer-Verlag, Berlin-New York, 1974, p. 604.
- [BO78] Pierre Berthelot and Arthur Ogus. *Notes on crystalline cohomology*. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1978, pp. vi+243. ISBN: 0-691-08218-9.
- [BS19] Bhargav Bhatt and Peter Scholze. “Prisms and Prismatic Cohomology”. In: *arXiv e-prints* (May 2019). arXiv: 1905.08229 [math.AG].
- [BS21] Bhargav Bhatt and Peter Scholze. “Prismatic F -crystals and crystalline Galois representations”. In: *arXiv e-prints* (June 2021). arXiv: 2106.14735 [math.NT].
- [BGR84] S. Bosch, U. Güntzer, and R. Remmert. *Non-Archimedean analysis*. Vol. 261. Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. A systematic approach to rigid analytic geometry. Springer-Verlag, Berlin, 1984, pp. xii+436. ISBN: 3-540-12546-9.
- [Bri08] Olivier Brinon. “Représentations p -adiques cristallines et de de Rham dans le cas relatif”. In: *Mém. Soc. Math. Fr. (N.S.)* 112 (2008), pp. vi+159. ISSN: 0249-633X.
- [Col99] Pierre Colmez. “Représentations cristallines et représentations de hauteur finie”. In: *J. Reine Angew. Math.* 514 (1999), pp. 119–143. ISSN: 0075-4102.

- [CF00] Pierre Colmez and Jean-Marc Fontaine. “Construction des représentations p -adiques semi-stables”. In: *Invent. Math.* 140.1 (2000), pp. 1–43. ISSN: 0020-9910.
- [CN17] Pierre Colmez and Wiesława Nizioł. “Syntomic complexes and p -adic nearby cycles”. In: *Invent. Math.* 208.1 (2017), pp. 1–108. ISSN: 0020-9910.
- [DLMS22] Heng Du, Tong Liu, Yong Suk Moon, and Koji Shimizu. “Completed prismatic F -crystals and crystalline \mathbf{Z}_p -local systems”. In: *arXiv e-prints* (Mar. 2022). arXiv: 2203.03444 [math.NT].
- [Fal88] Gerd Faltings. “ p -adic Hodge theory”. In: *J. Amer. Math. Soc.* 1.1 (1988), pp. 255–299. ISSN: 0894-0347.
- [Fal89] Gerd Faltings. “Crystalline cohomology and p -adic Galois-representations”. In: *Algebraic analysis, geometry, and number theory (Baltimore, MD, 1988)*. Johns Hopkins Univ. Press, Baltimore, MD, 1989, pp. 25–80.
- [Fon82] Jean-Marc Fontaine. “Sur certains types de représentations p -adiques du groupe de Galois d’un corps local; construction d’un anneau de Barsotti-Tate”. In: *Ann. of Math. (2)* 115.3 (1982), pp. 529–577. ISSN: 0003-486X.
- [Fon90] Jean-Marc Fontaine. “Représentations p -adiques des corps locaux. I”. In: *The Grothendieck Festschrift, Vol. II*. Vol. 87. Progr. Math. Birkhäuser Boston, Boston, MA, 1990, pp. 249–309.
- [Fon94] Jean-Marc Fontaine. “Le corps des périodes p -adiques”. In: *Astérisque* 223 (1994). With an appendix by Pierre Colmez, Périodes p -adiques (Bures-sur-Yvette, 1988), pp. 59–111. ISSN: 0303-1179.
- [FL82] Jean-Marc Fontaine and Guy Laffaille. “Construction de représentations p -adiques”. In: *Ann. Sci. École Norm. Sup. (4)* 15.4 (1982), 547–608 (1983). ISSN: 0012-9593.
- [FW79a] Jean-Marc Fontaine and Jean-Pierre Wintenberger. “Extensions algébrique et corps des normes des extensions APF des corps locaux”. In: *C. R. Acad. Sci. Paris Sér. A-B* 288.8 (1979), A441–A444. ISSN: 0151-0509.
- [FW79b] Jean-Marc Fontaine and Jean-Pierre Wintenberger. “Le “corps des normes” de certaines extensions algébriques de corps locaux”. In: *C. R. Acad. Sci. Paris Sér. A-B* 288.6 (1979), A367–A370. ISSN: 0151-0509.
- [GLQ20] Michel Gros, Bernard Le Stum, and Adolfo Quirós. “Twisted differential operators and q -crystals”. In: *arXiv e-prints* (Apr. 2020). arXiv: 2004.14320 [math.AG].
- [Gro63] Alexander Grothendieck. *Revêtements étales et groupe fondamental. Fasc. I: Exposés 1 à 5*. Troisième édition, corrigée, Séminaire de Géométrie Algébrique, 1960/61. Institut des Hautes Études Scientifiques, Paris, 1963, iv+143 pp. (not consecutively paged) (loose errata).
- [GR22] Haoyang Guo and Emanuel Reinecke. “Prismatic F -crystals and crystalline local systems”. In: *arXiv e-prints* (Mar. 2022). arXiv: 2203.09490 [math.AG].
- [KL15] Kiran S. Kedlaya and Ruochuan Liu. “Relative p -adic Hodge theory: foundations”. In: *Astérisque* 371 (2015), p. 239. ISSN: 0303-1179.
- [Mat86] Hideyuki Matsumura. *Commutative ring theory*. Vol. 8. Cambridge Studies in Advanced Mathematics. Translated from the Japanese by M. Reid. Cambridge University Press, Cambridge, 1986, pp. xiv+320. ISBN: 0-521-25916-9.
- [Moo18] Yong Suk Moon. “Relative crystalline representations and weakly admissible modules”. In: *arXiv e-prints* (June 2018). arXiv: 1806.00867 [math.NT].

- [MT20] Matthew Morrow and Takeshi Tsuji. “Generalised representations as q -connections in integral p -adic Hodge theory”. In: *arXiv e-prints* (Oct. 2020). arXiv: 2010.04059 [math.NT].
- [Sch12] Peter Scholze. “Perfectoid spaces”. In: *Publ. Math. Inst. Hautes Études Sci.* 116 (2012), pp. 245–313. ISSN: 0073-8301.
- [Sch17] Peter Scholze. “Canonical q -deformations in arithmetic geometry”. In: *Ann. Fac. Sci. Toulouse Math. (6)* 26.5 (2017), pp. 1163–1192. ISSN: 0240-2963.
- [Tsu99] Takeshi Tsuji. “ p -adic étale cohomology and crystalline cohomology in the semi-stable reduction case”. In: *Invent. Math.* 137.2 (1999), pp. 233–411. ISSN: 0020-9910.
- [Tsu20] Takeshi Tsuji. “Crystalline \mathbb{Z}_p -representations and A_{inf} -Representations with Frobenius”. In: *Proceedings in Simons Symposium: p -adic Hodge theory*. Simons symposia (2020), pp. 161–319.
- [Wac96] Nathalie Wach. “Représentations p -adiques potentiellement cristallines”. In: *Bull. Soc. Math. France* 124.3 (1996), pp. 375–400. ISSN: 0037-9484.
- [Wac97] Nathalie Wach. “Représentations cristallines de torsion”. In: *Compositio Math.* 108.2 (1997), pp. 185–240. ISSN: 0010-437X.
- [Win83] Jean-Pierre Wintenberger. “Le corps des normes de certaines extensions infinies de corps locaux; applications”. In: *Ann. Sci. École Norm. Sup. (4)* 16.1 (1983), pp. 59–89. ISSN: 0012-9593.

ABHINANDAN

(1) UNIVERSITÉ DE BORDEAUX, FRANCE

(2) UNIVERSITÉ DE LILLE, FRANCE

(3) UNIVERSITY OF TOKYO, JAPAN

E-mail : abhinandan@math.cnrs.fr, abhi@ms.u-tokyo.ac.jp