

# ARMA GARCH Processes

*Abhinav Anand, IIMB*

*2018/07/30*

## The Autocorrelation Function (ACF)

The correlation between random variables  $X_1, X_2$  is a measure of their linear dependence and is defined as:

$$\rho_{12} := \frac{\text{cov}(X_1, X_2)}{\sqrt{\text{var}(X_1)\text{var}(X_2)}} = \frac{\sigma_{12}}{\sigma_1\sigma_2}$$

It lies between -1 and 1 and for normal random variables  $\rho_{12} = 0$  implies that the variables are independent.

If we have a sample  $\{x_{1,t}, x_{2,t}\}_{t=1}^T$  the correlation can be consistently estimated by computing sample correlation:

$$\hat{\rho}_{12} = \frac{\hat{\sigma}_{12}}{\hat{\sigma}_1\hat{\sigma}_2}$$

For a time series  $r_t$  which is weakly stationary, the lag- $l$  autocorrelation function is the correlation between  $r_t$  and  $r_{t-l}$ :

$$\rho_l = \frac{\sigma_{t,t-l}}{\sigma_t\sigma_{t-l}} = \frac{\sigma_{t,t-l}}{\sigma_t^2} = \frac{\gamma_l}{\gamma_0}$$

This follows from weak stationarity:  $\sigma_t^2 = \sigma_{t-l}^2 = \gamma_0$  and  $\text{cov}(r_t, r_{t-l}) = \gamma_l$ .

We claim that there is no autocorrelation if  $\rho_l = 0 \forall l > 0$ .

To estimate the autocorrelation function of lag (say) 1, we use its sample counterpart:

$$\hat{\rho}_1 = \frac{\sum_{t=2}^T (r_t - \bar{r})(r_{t-1} - \bar{r})}{\sum_{t=1}^T (r_t - \bar{r})^2}$$

In general for lag  $l$  we consistently estimate it as:

$$\hat{\rho}_l = \frac{\sum_{t=l+1}^T (r_t - \bar{r})(r_{t-l} - \bar{r})}{\sum_{t=1}^T (r_t - \bar{r})^2}$$

The statistic  $\hat{\rho}_1, \hat{\rho}_2, \dots$  is the *sample autocorrelation function* of  $r_t$  and is key to capturing the linear dependence nature of the time series in question.

## Autoregressive (AR) Processes

Perhaps last period's returns may have some significant impact on the value of the returns this period. If so, its lag-1 autocorrelation may be useful for predicting the current period's value:

$$r_t = \phi_0 + \phi_1 r_{t-1} + u_t$$

where  $u_t$  is weakly stationary with mean 0 and variance  $\sigma_u^2$ . This is simply equivalent to a regression where  $r_{t-1}$  is the explanatory or independent variable.

It's straightforward to check the conditional mean and variance of such a process:

$$\begin{aligned}\mathbb{E}(r_t | r_{t-1}) &= \phi_0 + \phi_1 r_{t-1} \\ \text{var}(r_t | r_{t-1}) &= \sigma_u^2\end{aligned}$$

And more generally there could be defined autoregressive processes of order  $p$  ( $AR(p)$ ):

$$r_t = \phi_0 + \phi_1 r_{t-1} + \dots + \phi_p r_{t-p} + u_t$$

### AR(1) processes

Is the AR(1) process  $r_t = \phi_0 + \phi_1 r_{t-1} + u_t$  weakly stationary? This will imply that its unconditional mean and variance must be fixed in time and lag- $l$  covariance must depend only on the lag length  $l$ .

$$\begin{aligned}\mathbb{E}(r_t) &= \phi_0 + \phi_1 \mathbb{E}(r_{t-1}) + \mathbb{E}(u_t) \\ \mathbb{E}(r_t) &= \phi_0 + \phi_1 \mu\end{aligned}$$

$$\mu = \frac{\phi_0}{1 - \phi_1}$$

This clearly implies that for the mean of an AR(1) process to exist,  $\phi_1 \neq 1$  and  $\phi_0 = \mu \cdot (1 - \phi_1) = \mu - \mu\phi_1$ .

Hence a weakly stationary AR(1) process is:

$$\begin{aligned} r_t &= \mu - \mu\phi_1 + \phi_1 r_{t-1} + u_t \\ r_t - \mu &= (r_{t-1} - \mu)\phi_1 + u_t \\ r_t - \mu &= ((r_{t-2} - \mu)\phi_1 + u_{t-1})\phi_1 + u_t \\ &\vdots \\ r_t - \mu &= u_t + \phi_1 u_{t-1} + \phi_1^2 u_{t-2} + \dots \\ r_t &= \mu + \sum_{i=0}^{\infty} \phi_1^i \cdot u_{t-i} \end{aligned}$$

Additionally,

$$\text{var}(r_t) = \phi_1^2 \text{var}(r_{t-1}) + \sigma_u^2$$

Since for weakly stationary AR(1) processes  $\text{var}(r_t) = \text{var}(r_{t-1}) = \gamma_0$  we have

$$\gamma_0 = \frac{\sigma_u^2}{1 - \phi_1^2}$$

Weak stationarity immediately implies that  $\phi_1 \in (-1, 1)$ .

Hence taken together, for an AR(1) process to be weakly stationary it is necessary and sufficient that  $\phi_1 \in (-1, 1)$ ; <sup>1</sup> and the canonical AR(1) series can be written as:


$$r_t = (1 - \phi_1)\mu + \phi_1 r_{t-1} + u_t$$

We can plot some hypothetical autoregressive processes by simulation via the function `arma.sim()` include in the `stats` package that loads by default.


---

<sup>1</sup>For a general AR( $p$ ) process, the corresponding condition is:  $|\phi_1| + |\phi_2| + \dots + |\phi_p| < 1$ .


```
# AR(0)
plot(rnorm(500, 0, 0.8), type = "l", col = "blue")
abline(h = 0)
```




```
# AR(1)
ar_1 <- arima.sim(n = 500, list(ar = c(0.8)), sd = 0.8)
plot(ar_1, col = "blue")
abline(h = 0)
```



```
# AR(2)
ar_2 <- arima.sim(n = 500, list(ar = c(0.8, 0.15)), sd = 0.8)
plot(ar_2, col = "blue")
abline(h = 0)
```



```
# AR(3)
ar_3 <- arima.sim(n = 500, list(ar = c(0.5, 0.3, 0.15)), sd = 0.8)
plot(ar_3, col = "blue")
abline(h = 0)
```



### Autocorrelation Function for AR(1) processes

We can easily check that for positive lags  $l > 0$ , the lagged covariance follows:

$$\gamma_l = \phi_1 \gamma_{l-1}$$


Hence it follows that for the autocorrelation function  $\rho_l = \phi_1 \rho_{l-1}$ ; and because

$\rho_0 = 1$ ,  $\rho_l = \phi_1^l$ . This implies that the autocorrelation function of an AR(1) series decays exponentially with rate  $\phi_1$  and starting value 1. If  $\phi_1 < 0$  the series alternates between positive and negative terms.

### Illustration


For example let's compute the sample autocorrelation function (ACF) for the financial market indices.

```
acf(ret_BSE, na.action = na.pass)
```




```
acf(ret_sp500, na.action = na.pass)
```

### Series ret\_sp500



```
acf(ret_nikkei, na.action = na.pass)
```

### Series ret\_nikkei




What about log-returns?


```
logret_BSE <- func_pr_to_logret(index_bse$Close)
logret_SP <- func_pr_to_logret(ind_sp500$SP500)
logret_Nikkei <- func_pr_to_logret(ind_nikkei$NIKKEI225)
```

```
ACF_BSE <- acf(logret_BSE, na.action = na.pass)
```

### Series logret\_BSE




```
barplot(head(ACF_BSE$acf))
```




```
ACF_SP <- acf(logret_SP, na.action = na.pass)
```



### Series logret\_SP




```
barplot(head(ACF_SP$acf))
```




```
ACF_Nikkei <- acf(logret_Nikkei, na.action = na.pass)
```

### Series logret\_Nikkei



```
barplot(head(ACF_Nikkei$acf))
```



### Partial Autocorrelation Functions (PACF)

Is there a way to know how many lags to include for an autoregressive return series?  
This issue is solved via the usage of *partial autocorrelation functions* as shown below.

Consider the following sequences of AR processes:

$$r_t = \phi_{01} + \phi_{11}r_{t-1} + u_{1t}$$

$$r_t = \phi_{02} + \phi_{12}r_{t-1} + \phi_{22}r_{t-2} + u_{2t}$$

$$r_t = \phi_{03} + \phi_{13}r_{t-1} + \phi_{23}r_{t-2} + \phi_{33}r_{t-3} + u_{3t}$$

$$r_t = \phi_{04} + \phi_{14}r_{t-1} + \phi_{24}r_{t-2} + \phi_{34}r_{t-3} + \phi_{44}r_{t-4} + u_{4t}$$

$$\vdots$$


These models are merely multiple regressions and can be estimated via the standard least squares method.

In these models,  $\hat{\phi}_{11}$  is called the lag 1 sample PACF of  $r_t$ ,  $\hat{\phi}_{22}$  of the second equation is the lag 2 sample PACF of  $r_t$  and so on. By construction, the lag 2  $\hat{\phi}_{22}$  is the marginal contribution of  $r_{t-2}$  in explaining  $r_t$  over the AR(1) model and so on. Hence if the underlying model is say AR( $p$ ) then all sample PACFs  $\hat{\phi}_{11}, \dots, \hat{\phi}_{pp}$  must be different from 0 but all sample PACFs from then on:  $\hat{\phi}_{p+1,p+1}, \dots = 0$ . This property can be used to find the order  $p$ .

Armed with this knowledge, let's compute the PACFs for the three financial market indices:


```
pacf(ret_BSE, na.action = na.pass)
```

### Series ret\_BSE




```
pacf(ret_sp500, na.action = na.pass)
```

### Series ret\_sp500



```
pacf(ret_nikkei, na.action = na.pass)
```

### Series ret\_nikkei



## Information Criteria

Apart from PACF, another way to find the number of lags is the use of likelihood based information criteria. Here we look at the two most famous ones: the Akaike Information Criterion (AIC) and the Bayesian Information Criterion (BIC).

$$AIC(l) = -\frac{2}{T} \cdot \ln(\text{likelihood}) + \frac{2}{T} \cdot (\# \text{parameters})$$

For a Gaussian AR( $l$ ),  $AIC = \ln(\hat{\sigma}_{u,MLE}^2) + 2\frac{l}{T}$ . The first term measures the goodness of fit of the model while the second penalizes the usage of parameters.

The Bayesian Information Criterion (BIC) uses a different penalty function. For a Gaussian AR( $l$ ) it takes the following form:

$$BIC(l) = \ln(\hat{\sigma}_{u,MLE}^2) + \ln(T) \cdot \frac{l}{T}$$

## Moving Average (MA) Processes

Consider an infinitely long autoregressive process:

$$r_t = \phi_0 + \phi_1 r_{t-1} + \phi_2 r_{t-2} + \dots + u_t$$

If this series is to be weakly stationary, the coefficients  $\phi_j$  must decay sufficiently fast. One way to ensure this is to assume that  $\phi_j = -\theta^j$  for some  $\theta \in (0, 1)$ .

$$r_t = \phi_0 - \theta_1 r_{t-1} - \theta_1^2 r_{t-2} - \dots + u_t$$

$$r_t + \sum_{j=1}^{\infty} \theta_1^j r_{t-j} = \phi_0 + u_t$$

The same form can be written for  $r_{t-2}$ :

$$r_{t-1} + \sum_{j=2}^{\infty} \theta_1^j r_{t-j} = \phi_0 + u_{t-1}$$

Solving for the above two equations, we get:

$$r_t = \phi_0(1 - \theta_1) + u_t - \theta_1 u_{t-1}$$

This indicates the AR model is a weighted average of shocks  $u_t, u_{t-1}$  and a constant. This is a *moving average* form of order 1 or MA(1). It's straightforward to check that unlike AR processes, MA processes are *always* stationary. (Can you see why?)

The general form for the MA( $q$ ) process is:

$$r_t = c_0 + u_t - \theta_1 u_{t-1} - \theta_2 u_{t-2} - \dots - \theta_q u_{t-q}$$

## Autocorrelation Function

Consider the MA(1) model with the constant term 0:

$$r_t = u_t - \theta_1 u_{t-1}$$

$$r_{t-l} r_t = u_t r_{t-l} - \theta_1 u_{t-1} r_{t-l}$$

$$\mathbb{E}(r_{t-l} r_t) = 0 - \theta_1 \mathbb{E}(u_{t-1} r_{t-l})$$

From this we see that:

$$\gamma_1 = -\theta_1 \sigma_u^2$$

$$\gamma_{l>1} = 0$$

Also,  $\text{var}(r_t) = \gamma_0 = (1 + \theta_1^2) \sigma_u^2$  and this implies that for  $\rho_l = \frac{\gamma_l}{\gamma_0}$  becomes:

$$\rho_0 = 1$$

$$\rho_1 = -\frac{\theta_1}{1 + \theta_1^2}$$

$$\rho_{l>1} = 0$$

Hence for MA(1) processes, while the first lag autocorrelation is nonzero, all further lags produce zero autocorrelation. This property can be exploited to locate the order of the MA process. In general, for an MA( $q$ ) process, the ACF cuts off at lag  $q$ . Since the MA( $q$ ) process only relies on its past  $q - 1$  realization, it's often called a 'finite memory' process.


To check the order of the MA series in question, we plot the ACF of the series. If  $\rho_q \neq 0$  but  $\rho_{l>q} = 0$  we may conclude that the series is MA( $q$ ).

## Illustrations


We simulate some moving average processes below:

```
n <- 500

# MA(0)
plot(rnorm(n), type = "l", col = "blue")
abline(h = 0)
```




```
# MA(1)
ma_1 <- arima.sim(n = n, list(ma = c(0.8)), innov=rnorm(n))
plot(ma_1, col = "blue")
abline(h = 0)
```




```
# MA(2)
ma_2 <- arima.sim(n = n, list(ma = c(0.8, 0.15)), innov=rnorm(n))
```

```
plot(ma_2, col = "blue")  
abline(h = 0)
```



```
# MA(3)  
ma_3 <- arima.sim(n = n, list(ma = c(0.5, 0.3, 0.15)), innov=rnorm(n))  
plot(ma_3, col = "blue")  
abline(h = 0)
```





## Autoregressive Moving Average (ARMA) Processes

While in principle we could fit empirical time series to  $AR(p)$  or  $MA(q)$  models exclusively, often the concomitant order is very high. In order to circumvent this problem, we combine AR and MA models into a composite  $ARMA(p, q)$  model with fewer parameters to estimate.

For example the  $ARMA(1,1)$  series is as follows:

$$r_t - \phi_1 r_{t-1} = \phi_0 + u_t - \theta_1 u_{t-1}$$

The LHS is the AR component while the RHS is MA component.

### Properties

Taking expectations we get:

$$\mathbb{E}(r_t) - \phi_1 \mathbb{E}(r_{t-1}) = \phi_0 + \mathbb{E}(u_t) - \theta_1 \mathbb{E}(u_{t-1})$$

$$\mathbb{E}(r_t) = \mu = \frac{\phi_0}{1 - \phi_1}$$

This is the same exact mean as that of an  $AR(1)$  process. Solving for the stationarity of the variance we get the same condition for the parameter  $\phi_1 \in (0, 1)$  as that of the  $AR(1)$  model. It is not hard to derive the autocorrelation function of the  $ARMA(1,1)$  process which behaves the same way as that of an  $AR(1)$  process.

In general for an  $ARMA(p, q)$  process we have the following definition:

$$r_t = \phi_0 + \sum_{i=1}^p \phi_i r_{t-i} + u_t - \sum_{j=1}^q \theta_j u_{t-j}$$


## Conditional Heteroskedasticity Models

In empirical time series, we do not observe volatility directly but only estimate it with the help of some other directly observed characteristics. However as discussed in

Cont (2001) volatility displays some striking regularities such as volatility clustering and differential reaction to price changes.


To investigate the volatility of empirical time series, we plot the daily returns and ACF of the Bombay stock exchange sensx.

```
plot(logret_BSE, type = "l", col = "blue")  
abline(h = 0)
```



```
plot(acf(ret_BSE, na.action = na.pass))
```

**Series ret\_BSE**




While there seems to be no serial autocorrelation there is some dependence structure

embedded in the series. We can check this for various functions of the return process such as squared returns or absolute returns:


```
ret_BSE_sq <- ret_BSE^2  
acf(ret_BSE_sq, na.action = na.pass)
```

**Series ret\_BSE\_sq**




```
pacf(ret_BSE_sq, na.action = na.pass)
```

**Series ret\_BSE\_sq**




```
ret_BSE_abs <- abs(ret_BSE)
acf(ret_BSE_abs, na.action = na.pass)
```

**Series ret\_BSE\_abs**



```
pacf(ret_BSE_abs, na.action = na.pass)
```


**Series ret\_BSE\_abs**



This behavior is also displayed by the other financial markets.


```
acf(ret_sp5002, na.action = na.pass)
```

### Series ret\_sp500<sup>2</sup>




```
pacf(ret_sp5002, na.action = na.pass)
```

### Series ret\_sp500<sup>2</sup>




```
acf(ret_nikkei2, na.action = na.pass)
```

**Series ret\_nikkei^2**



```
pacf(ret_nikkei^2, na.action = na.pass)
```

**Series ret\_nikkei^2**



The conditional mean volatility of the series may be written as:

$$\mu_t = \mathbb{E}(r_t | \mathbb{F}_{t-1})$$

$$\sigma_t^2 = \text{var}(r_t | \mathbb{F}_{t-1}) = \mathbb{E}((r_t - \mu_t)^2 | \mathbb{F}_{t-1}) = \text{var}(u_t | \mathbb{F}_{t-1})$$

Here  $\mathbb{F}(\cdot)$  denotes the ‘filtration’ or the information available at the given time.

## ARCH Effect

Consider  $r_t = \mu_t + u_t$ . The square of the residual series  $u_t = r_t - \mu_t$  is of interest for estimating conditional heteroskedasticity, also known as the ARCH effect (autoregressive conditional heteroskedasticity effect). A popular method of testing for ARCH effects is via Engel’s Lagrange multiplier test which is equivalent to the standard  $F$  statistic for testing  $\beta_i = 0$  in the multiple linear regression:

$$u_t^2 = \beta_0 + \beta_1 u_{t-1}^2 + \dots + \beta_m u_{t-m}^2 + \epsilon_t$$

## References

- Cont, Rama. 2001. “Empirical Properties of Asset Returns: Stylized Facts and Statistical Issues.” *Quantitative Finance* 1 (2): 223–36.
- Jondeau, Eric, Ser-Huang Poon, and Michael Rockinger. 2007. *Financial Modeling Under Non-Gaussian Distributions*. Springer Finance.
- Tsay, Ruey S. 2010. *Analysis of Financial Time Series*. Third Edition. John Wiley; Sons.