
## Financial Time Series

Abhinav Anand, IIMB 2018/07/21

## Background

Financial prices, indices, returns etc. are sequences of real numbers indexed by time. The study of their mathematical and statistical properties is vital for those aspiring to write papers in empirical finance.

For example, consider the following plot of a hypothetical market index: what can we say about the market conditions from its ups and downs?



The above series however generated via the command: plot(cumsum(rnorm(10000, 0, 1)) which plots the cumulative sum of ten thousand standard normal realizations—a random walk!

The random walk hypothesis is a theory that claims that stock market prices obey a random walk process. This hypothesis is consistent with the "Efficient Market Hypothesis". Roughly speaking, the efficient market hypothesis claims in its weak form that for publicly traded stocks, all public information is captured in the price of the stock. The strong form of the hypothesis claims that all information—both

public and private—is reflected the stock price. Hence according to this theory, any attempt to consistently beat the market in terms of risk-adjusted returns is doomed to failure.<sup>1</sup>

Hence in particular, any attempt to discern patterns in the hypothetical market's ups and downs as shown above is worthless.

Do empirical financial markets behave in the same way too?

### **Empirical Financial Time Series**


As an illustration we produce the daily time series for the closing value of the Bombay Stock Exchange index ("Sensex").

```
file_bse <- "SENSEX.csv"</pre>
index_bse <- readr::read_csv(file_bse) %>%
  dplyr::select(-empty)
index_bse$Date <- as.Date(index_bse$Date,</pre>
                            format = \mbox{"}\%d-\mbox{"}B-\mbox{"}Y"
                             ) #date reformat
plot(index_bse$Date,
     index_bse$Close,
     type = "1",
     col = "blue",
     xlab = "Year",
     ylab = "BSE Index",
     main = "Indian stock market performance"
     )
fit_lm <- lm(Close ~ Date,
              data = index_bse) #fit linear model
abline(fit_lm, #plot linear model line
       lty = "dotdash",
```

<sup>&</sup>lt;sup>1</sup>Or, in other words, if investors beat the market it's not perhaps due to their so-called investing skills but due to their being lucky.


```
col = "red",
lwd = 2
)
```

# Indian stock market performance



```
# via ggplot
ggplot(data = index_bse,
       aes(Date, Close)
       ) +
  geom_line(lwd = 0.3,
            color = "blue"
            ) +
  geom_smooth(method = "lm",
              lty = "dotdash",
              lwd = 0.6,
              color = "red",
              se = F) +
  theme_minimal() +
  labs(x = "Years",
       y = "BSE Sensex",
       title = "Indian stock market performance"
```






It seems that the level of the series is rising and the fluctuations are sometimes high and sometimes low.

This index series is an example of a *non-stationary* time series. This roughly means that the mean and the variance of such a series are functions of time.


Here is the index level time series for the US index S&P500 from 2008:

## Warning: Removed 92 rows containing non-finite values (stat\_smooth).



And here it for the Japanese Nikkei225:

```
NIKKEI225 = col_double()
                                              ),
                            na = c("", "NA", ".")
                             )
ggplot(ind_nikkei,
      aes(DATE, NIKKEI225)
      ) +
 geom_line(lwd = 0.3,
           color = "blue") +
 geom_smooth(method = "lm",
             lty = "dotdash",
             lwd = 0.6,
             se = F,
             color = "red"
             ) +
 theme_minimal() +
 labs(x = "Years",
      y = "Nikkei225"
```



## Returns

We observe prices in the financial markets empirically. However, due to their non-stationary nature, they are hard to analyze. Hence they are converted to return series which are usually stationary. There are many ways to construct different notions of returns from the same underlying price sequence. We discuss some prominent ones below.

### One-Period Simple Return

The simple one period return for holding some asset whose price is given by the sequence  $\{p_t\}_{t=1}^n$  is:

$$r_t := \frac{p_t - p_{t-1}}{p_{t-1}} = \frac{p_t}{p_{t-1}} - 1$$

### Multi-period Simple Return

$$r_t[k] := \frac{p_t - p_{t-k}}{p_{t-k}} = \frac{p_t}{p_{t-k}} - 1$$

$$r_t[k] := \frac{p_t}{p_{t-k}} - 1 = \frac{p_t}{p_{t-1}} \frac{p_{t-1}}{p_{t-2}} \dots \frac{p_{t-k+1}}{p_{t-k}} - 1$$

$$r_t[k] := (1 + r_t)(1 + r_{t-1}) \dots (1 + r_{t-k+1}) - 1$$

Multi-period returns are used to convert high frequency returns to low frequency returns—i.e., daily to monthly; or monthly to yearly etc.

For example to convert monthly returns to annual returns:

$$r_t[12] = \left[\prod_{j=0}^{12-1} (1 + r_{t-j})\right]^{1/12} - 1$$

Often (when working with daily returns especially)  $1 + r_t \approx r_t$ , in which case:

$$r_t[k] \approx \sum_{j=0}^{k-1} r_{t-j}$$

Additionally, the simple returns for a portfolio with fractional weights  $w_1, \ldots, w_n$  are:

$$r_{p,t} = \sum_{i=1}^{n} w_i r_{i,t}$$

#### Log Returns

We know that to compute yearly returns from say, monthly returns we use the following formula:

$$r_t[12] = \left[\prod_{j=0}^{12-1} (1 + r_{t-j})\right]^{1/12} - 1$$

In general, if a bank pays an annual interest of  $r_t^m$  m times a year, the interest rate for unit investment is  $r_t^m/m$  and after one year the value of the deposit is  $(1+\frac{r_t^m}{m})^m$ . If there is continous compounding,  $m\to\infty$ , in which case the value of the investment becomes:

$$\lim_{m \to \infty} (1 + \frac{r_t^m}{m})^m = e^{r_t}$$

Hence it must be that:

$$R_t = \log p_t - \log p_{t-1} = \log \frac{p_t}{p_{t-1}}$$

A particular advantage of log-returns are that multiperiod log returns are merely the sum of one period log returns:

$$R_t[k] = \log p_t - \log p_{t-k} = \log p_t - \log p_{t-1} + \log p_{t-1} - \dots \log p_{t-k}$$
$$R_t[k] = R_t[1] + R_t[2] + \dots + R_t[k-1]$$

### Computational Examples

To construct return series from prices, we write a function that accepts as input a price (or index) sequence and returns a sequence of simple one period returns by using the formula  $r_t = \frac{p_t - p_{t-1}}{p_{t-1}}$ . Since there is no return for the first entry of the price sequence, we append an NA at the beginning of the returns.

```
func_pr_to_ret <- function(price_vec)
{
    # This function takes a vector of prices and
    # returns a vector of simple one period returns
    1 <- length(price_vec)
    ret_num <- diff(price_vec) #numerator = change in prices
    ret_den <- price_vec[-1] #denominator = price series

return(c(NA, ret_num/ret_den)) #first return is NA
}

price_vec_1 <- seq(from = 1, to = 10, by = 2)
func_pr_to_ret(price_vec_1)</pre>
```

## [1] NA 2.0000000 0.6666667 0.4000000 0.2857143

Similarly log-returns can be calculated using the formula  $R_t = \log(p_t) - \log(p_{t-1})$ .

```
func_pr_to_logret <- function(p_vec)
{
    # This function takes a vector of prices and
    # returns a vector of log returns
    log_price <- log(p_vec)
    log_ret <- diff(log_price) #log ret = delta log p

    return(c(NA, log_ret)) #first return is NA
}


p_vec <- seq(from = 1, to = 10, by = 2)
func_pr_to_logret(p_vec)</pre>
```

## [1] NA 1.0986123 0.5108256 0.3364722 0.2513144

### Return Series for Market Indices


We can compute returns and log-returns for the Bombay Stock Exchange Index, S&P500 and the Nikkei 225 as follows:

```
ret_BSE <- func_pr_to_ret(index_bse$Close)
plot(index_bse$Date, #x variable
    ret_BSE, #y variable
    type = "l",
    lwd = 1.2,
    col = "blue",
    xlab = "Years",
    ylab = "BSE Returns"
    )
abline(h = 0)</pre>
```




We can do the same for the S&P500 and the Nikkei 225 series:

```
ret_sp500 <- func_pr_to_ret(ind_sp500$SP500)
plot(ind_sp500$DATE, #x variable
    ret_sp500, #y variable
    type = "l",
    lwd = 1.2,
    col = "red",
    xlab = "Years",
    ylab = "S&P500 Returns"
    )
abline(h = 0)</pre>
```



```
ret_nikkei <- func_pr_to_ret(ind_nikkei$NIKKEI225)
plot(ind_nikkei$DATE, #x variable
    ret_nikkei, #y variable
    type = "l",
    lwd = 1.2,
    col = "green",
    xlab = "Years",
    ylab = "Nikkei225 Returns"
    )
abline(h = 0)</pre>
```



One clearly sees that the index sequences have no trend now. It seems that the returns are fluctuating around a mean level 0. The fluctuations (volatility) around the mean are sometimes and sometimes low.

## Stationarity of Time Series

### Distribution of Returns

### **Summary Statistics of Empirical Returns**

min, max, mean, median, mode, std, variance, iqr, skewness, kurtosis

```
func_summ_stat_ind <- function(vec)</pre>
{
  summ_stat_ind <- data.frame(min = min(vec, na.rm = T),</pre>
                        max = max(vec, na.rm = T),
                        mean = mean(vec, na.rm = T),
                        median = median(vec, na.rm =T),
                        std = sd(vec, na.rm =T),
                        iqr = IQR(vec, na.rm=T),
                        var = var(vec, na.rm =T)
                        ) %>%
    signif(., digits = 2)
  return(summ_stat_ind)
}
table_summ_stat <- rbind(c("BSE", func_summ_stat_ind(ret_BSE)),</pre>
                          c("S&P500", func_summ_stat_ind(ret_sp500)),
                          c("Nikkei225", func_summ_stat_ind(ret_nikkei))
                          )
knitr::kable(table_summ_stat)
```


	min	max	mean	median	std	iqr	var
BSE	-0.11	0.17	0.00061	0.00092	0.014	0.014	2e-04
S&P500	-0.09	0.12	0.00038	0.00061	0.013	0.0093	0.00016
Nikkei225	-0.11	0.1	0.00014	0.00059	0.016	0.015	0.00024

### Histograms for returns for each index

```
hist(ret_BSE,
    breaks = 80,
```


```
col = "grey",
xlab = "Daily Returns",
ylab = "Frequency",
main = "Histogram for BSE"
)
```

# **Histogram for BSE**




```
hist(ret_sp500,
    breaks = 80,
    col = "grey",
    xlab = "Daily Returns",
    ylab = "Frequency",
    main = "Histogram for S&P500"
    )
```

# Histogram for S&P500



```
hist(ret_nikkei,
    breaks = 80,
    col = "grey",
    xlab = "Daily Returns",
    ylab = "Frequency",
    main = "Histogram for Nikkei225"
)
```

# Histogram for Nikkei225



### Stylized Facts

Stock prices, commodity prices, exchange rates etc. in empirical financial markets display many striking regularities discussed in Cont (2001).

- 1. **Fat Tails:** Unconditional return distributions have tails fatter than those of normal distribution. Conditional return distributions are also non-normal.
- 2. Asymmetry: Unconditional return distributions are negatively skewed.
- 3. **Aggregated Normality:** Lower frequency returns resemble normal distributions more than higher frequency returns.
- 4. **No Autocorrelation:** Except at high frequencies, returns generally do not display autocorrelation.
- 5. Volatility Clustering: Return volatility is autocorrelated.
- 6. **Time-Varying Cross Correlation:** Correlation between assets returns tends to be higher during high volatility periods especially during market crashes.

#### Aggregated Time Series Summary Stats

### References

Cont, Rama. 2001. "Empirical Properties of Asset Returns: Stylized Facts and Statistical Issues." Quantitative Finance 1 (2): 223–36.

Jondeau, Eric, Ser-Huang Poon, and Michael Rockinger. 2007. Financial Modeling Under Non-Gaussian Distributions. Springer Finance.

Tsay, Ruey S. 2010. Analysis of Financial Time Series. Third Edition. John Wiley; Sons.