

# Partitioning Theorems for Sets of Semi-Pfaffian Sets, with Applications

Abhiram Natarajan

Collaborators: Prof. Martin Lotz (Univ. of Warwick), Prof.  
Nicolai Vorobjov (Univ. of Bath)

# Polynomial Method' in Combinatorics

- **Real Algebraic Set:** The set of real zeros of a polynomial  $P$   
 $Z(P) := \{x \in \mathbb{R}^n \mid P(x) = 0\}$

# Polynomial Method' in Combinatorics

- **Real Algebraic Set:** The set of real zeros of a polynomial  $P$

$$Z(P) := \{x \in \mathbb{R}^n \mid P(x) = 0\}$$

$$Z(x^2 + y^2 - 1)$$



$$Z(y - x^2)$$



# Polynomial Method' in Combinatorics

- **Real Algebraic Set:** The set of real zeros of a polynomial  $P$

$$Z(P) := \{x \in \mathbb{R}^n \mid P(x) = 0\}$$

$$Z(x^2 + y^2 - 1) \quad Z(y - x^2)$$



- Incidence combinatorics studies combinatorial aspects of the intersections of geometric objects



# Polynomial Method' in Combinatorics

- **Real Algebraic Set:** The set of real zeros of a polynomial  $P$

$$Z(P) := \{x \in \mathbb{R}^n \mid P(x) = 0\}$$

$$Z(x^2 + y^2 - 1) \quad Z(y - x^2)$$



- Incidence combinatorics studies combinatorial aspects of the intersections of geometric objects
- **Algebra-geometric** techniques have been very effective

# Polynomial Method' in Combinatorics

- **Real Algebraic Set:** The set of real zeros of a polynomial  $P$

$$Z(P) := \{x \in \mathbb{R}^n \mid P(x) = 0\}$$

$$Z(x^2 + y^2 - 1) \quad Z(y - x^2)$$



- Incidence combinatorics studies combinatorial aspects of the intersections of geometric objects
- **Algebra-geometric** techniques have been very effective
- Technique called **polynomial partitioning** has helped solve **several** open problems in incidence geometry and other areas

# Cells Induced by a Polynomial

Cells induced by poly.  $P$ : Connected components of  $\mathbb{R}^n \setminus Z(P)$

# Cells Induced by a Polynomial

Cells induced by poly.  $P$ : Connected components of  $\mathbb{R}^n \setminus Z(P)$





# Cells Induced by a Polynomial

Cells induced by poly.  $P$ : Connected components of  $\mathbb{R}^n \setminus Z(P)$




# Cells Induced by a Polynomial

Cells induced by poly.  $P$ : Connected components of  $\mathbb{R}^n \setminus Z(P)$




# Cells Induced by a Polynomial

Cells induced by poly.  $P$ : Connected components of  $\mathbb{R}^n \setminus Z(P)$



# Cells Induced by a Polynomial


Cells induced by poly.  $P$ : Connected components of  $\mathbb{R}^n \setminus Z(P)$






# Cells Induced by a Polynomial

Cells induced by poly.  $P$ : Connected components of  $\mathbb{R}^n \setminus Z(P)$



# Cells Induced by a Polynomial

Cells induced by poly.  $P$ : Connected components of  $\mathbb{R}^n \setminus Z(P)$



# Polynomial Partitioning


Theorem (Guth and Katz [2015], Guth [2015])

Let  $\Gamma$  be a set of  $k$ -dimensional real algebraic sets in  $\mathbb{R}^n$ . For any  $D \geq 1$ , there is a polynomial  $P$  of degree  $\leq D$ , such that each cell induced by  $P$  intersects at most  $\sim \frac{|\Gamma|}{D^{n-k}}$  elements of  $\Gamma$ .

# Polynomial Partitioning

Theorem (Guth and Katz [2015], Guth [2015])

Let  $\Gamma$  be a set of  $k$ -dimensional real algebraic sets in  $\mathbb{R}^n$ . For any  $D \geq 1$ , there is a polynomial  $P$  of degree  $\leq D$ , such that each cell induced by  $P$  intersects at most  $\sim \frac{|\Gamma|}{D^{n-k}}$  elements of  $\Gamma$ .




a set  $\Gamma$  of 10  
curves in  $\mathbb{R}^2$




# Polynomial Partitioning

Theorem (Guth and Katz [2015], Guth [2015])

Let  $\Gamma$  be a set of  $k$ -dimensional real algebraic sets in  $\mathbb{R}^n$ . For any  $D \geq 1$ , there is a polynomial  $P$  of degree  $\leq D$ , such that each cell induced by  $P$  intersects at most  $\sim \frac{|\Gamma|}{D^{n-k}}$  elements of  $\Gamma$ .



a set  $\Gamma$  of 10  
curves in  $\mathbb{R}^2$




partitioning  
polynomial induces 5  
cells


# Polynomial Partitioning

Theorem (Guth and Katz [2015], Guth [2015])

Let  $\Gamma$  be a set of  $k$ -dimensional real algebraic sets in  $\mathbb{R}^n$ . For any  $D \geq 1$ , there is a polynomial  $P$  of degree  $\leq D$ , such that each cell induced by  $P$  intersects at most  $\sim \frac{|\Gamma|}{D^{n-k}}$  elements of  $\Gamma$ .



a set  $\Gamma$  of 10  
curves in  $\mathbb{R}^2$



partitioning  
polynomial induces 5  
cells



each cell intersects  
only few curves from  
 $\Gamma$

Szemerédi-Trotter - Proof using Poly. Part.

# Szemerédi-Trotter - Proof using Poly. Part.

- $m$  points and  $n$  lines in  $\mathbb{R}^2$ ;  $\mathcal{O}(m^{2/3}n^{2/3} + m + n)$  incidences



# Szemerédi-Trotter - Proof using Poly. Part.

- ▶  $m$  points and  $n$  lines in  $\mathbb{R}^2$ ;  $O(m^{2/3}n^{2/3} + m + n)$  incidences

Proof:

- ▶ (weak bound) incidences at most  $\min\{m + n^2, n + m^2\}$

# Szemerédi-Trotter - Proof using Poly. Part.

- ▶  $m$  points and  $n$  lines in  $\mathbb{R}^2$ ;  $O(m^{2/3}n^{2/3} + m + n)$  incidences

Proof:

- ▶ (weak bound) incidences at most  $\min\{m + n^2, n + m^2\}$
- ▶ Partition using polynomial  $P$  of degree  $D$ 
  - ▶  $D^2$  cells each containing  $\frac{m}{D^2}$  points, weak bound:  $nD + \frac{m^2}{D^2}$

# Szemerédi-Trotter - Proof using Poly. Part.

- ▶  $m$  points and  $n$  lines in  $\mathbb{R}^2$ ;  $O(m^{2/3}n^{2/3} + m + n)$  incidences

Proof:

- ▶ (weak bound) incidences at most  $\min\{m + n^2, n + m^2\}$
- ▶ Partition using polynomial  $P$  of degree  $D$ 
  - ▶  $D^2$  cells each containing  $\frac{m}{D^2}$  points, weak bound:  $nD + \frac{m^2}{D^2}$
  - ▶ each line intersects  $Z(P)$  in  $< D$  points:  $nD$

# Szemerédi-Trotter - Proof using Poly. Part.

- ▶  $m$  points and  $n$  lines in  $\mathbb{R}^2$ ;  $O(m^{2/3}n^{2/3} + m + n)$  incidences

Proof:

- ▶ (weak bound) incidences at most  $\min\{m + n^2, n + m^2\}$
- ▶ Partition using polynomial  $P$  of degree  $D$ 
  - ▶  $D^2$  cells each containing  $\frac{m}{D^2}$  points, weak bound:  $nD + \frac{m^2}{D^2}$
  - ▶ each line intersects  $Z(P)$  in  $< D$  points:  $nD$
  - ▶  $Z(P)$  can contain only  $D$  lines, weak bound:  $m + D^2$



# Szemerédi-Trotter - Proof using Poly. Part.

- ▶  $m$  points and  $n$  lines in  $\mathbb{R}^2$ ;  $O(m^{2/3}n^{2/3} + m + n)$  incidences

Proof:

- ▶ (weak bound) incidences at most  $\min\{m + n^2, n + m^2\}$
- ▶ Partition using polynomial  $P$  of degree  $D$ 
  - ▶  $D^2$  cells each containing  $\frac{m}{D^2}$  points, weak bound:  $nD + \frac{m^2}{D^2}$
  - ▶ each line intersects  $Z(P)$  in  $< D$  points:  $nD$
  - ▶  $Z(P)$  can contain only  $D$  lines, weak bound:  $m + D^2$
- ▶ Set  $D = \frac{m^{2/3}}{n^{1/3}}$  and sum up ■

# Szemerédi-Trotter - Proof using Poly. Part.

- ▶  $m$  points and  $n$  lines in  $\mathbb{R}^2$ ;  $O(m^{2/3}n^{2/3} + m + n)$  incidences

Proof:

- ▶ (weak bound) incidences at most  $\min\{m + n^2, n + m^2\}$
- ▶ Partition using polynomial  $P$  of degree  $D$ 
  - ▶  $D^2$  cells each containing  $\frac{m}{D^2}$  points, weak bound:  $nD + \frac{m^2}{D^2}$
  - ▶ each line intersects  $Z(P)$  in  $< D$  points:  $nD$
  - ▶  $Z(P)$  can contain only  $D$  lines, weak bound:  $m + D^2$
- ▶ Set  $D = \frac{m^{2/3}}{n^{1/3}}$  and sum up ■

Takeaway

*Polynomial partitioning and basic arguments worked!*

# Generalize Real Algebraic Geometry

- Real algebraic sets possess **tameness** properties such as stratifiability, triangulability, etc.

# Generalize Real Algebraic Geometry

- ▶ Real algebraic sets possess **tameness** properties such as stratifiability, triangulability, etc.
- ▶ *...investigate classes of sets with the tame topological properties of real algebraic sets... - Grothendieck, Esquisse d'un Programme*



# Generalize Real Algebraic Geometry

- ▶ Real algebraic sets possess **tameness** properties such as stratifiability, triangulability, etc.
- ▶ *...investigate classes of sets with the tame topological properties of real algebraic sets... - Grothendieck, Esquisse d'un Programme*
- ▶ **O-minimal geometry** (geometry of **definable sets**) is an axiomatic generalization of real algebraic geometry

# Why O-Minimal Structures?

- Real algebraic sets in  $\mathbb{R}^n$  are included in an o-minimal structure

# Why O-Minimal Structures?

- ▶ Real algebraic sets in  $\mathbb{R}^n$  are included in an o-minimal structure
- ▶ Other examples -  $\mathbb{R}$  with exp function (e.g.  $x^3 + e^{x+2y} = 0$ ), Pfaffian functions (e.g.  $x^{2.31} - e^{e^y} = 0$ )

# Why O-Minimal Structures?

- ▶ Real algebraic sets in  $\mathbb{R}^n$  are included in an o-minimal structure
- ▶ Other examples -  $\mathbb{R}$  with exp function (e.g.  $x^3 + e^{x+2y} = 0$ ), Pfaffian functions (e.g.  $x^{2.31} - e^{e^y} = 0$ )
- ▶ O-minimal incidence combinatorics is lagging behind algebraic incidence combinatorics



# Why O-Minimal Structures?

- ▶ Real algebraic sets in  $\mathbb{R}^n$  are included in an o-minimal structure
- ▶ Other examples -  $\mathbb{R}$  with exp function (e.g.  $x^3 + e^{x+2y} = 0$ ), Pfaffian functions (e.g.  $x^{2.31} - e^{e^y} = 0$ )
- ▶ O-minimal incidence combinatorics is lagging behind algebraic incidence combinatorics

## Question

*Can we generalize polynomial partitioning to the o-minimal setting?*

# Why O-Minimal Structures?

- ▶ Real algebraic sets in  $\mathbb{R}^n$  are included in an o-minimal structure
- ▶ Other examples -  $\mathbb{R}$  with exp function (e.g.  $x^3 + e^{x+2y} = 0$ ), Pfaffian functions (e.g.  $x^{2.31} - e^{e^y} = 0$ )
- ▶ O-minimal incidence combinatorics is lagging behind algebraic incidence combinatorics

## Question

*Can we generalize polynomial partitioning to the o-minimal setting? ... we make progress...*

# Pfaffian Functions

- $\vec{f} = (f_1, \dots, f_r)$  is a **Pfaffian chain** if there exist polynomials  $P_{i,j}$  verifying

$$\frac{\partial f_i}{\partial x_j} = P_{i,j}(x, f_1(x), \dots, f_i(x))$$

# Pfaffian Functions

- $\vec{f} = (f_1, \dots, f_r)$  is a **Pfaffian chain** if there exist polynomials  $P_{i,j}$  verifying

$$\frac{\partial f_i}{\partial x_j} = P_{i,j}(x, f_1(x), \dots, f_r(x))$$

- $g : \mathbb{R}^n \rightarrow \mathbb{R}$  is a **Pfaffian function** w.r.t.  $\vec{f}$  if there exists polynomial  $Q_g$  such that

$$g(x) = Q_g(x, f_1(x), \dots, f_r(x))$$



# Pfaffian Functions

- ▶  $\vec{f} = (f_1, \dots, f_r)$  is a **Pfaffian chain** if there exist polynomials  $P_{i,j}$  verifying

$$\frac{\partial f_i}{\partial x_j} = P_{i,j}(x, f_1(x), \dots, f_r(x))$$

- ▶  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  is a **Pfaffian function** w.r.t.  $\vec{f}$  if there exists polynomial  $Q_g$  such that

$$g(x) = Q_g(x, f_1(x), \dots, f_r(x))$$

- ▶  $\deg(Q_g)$  is **degree** of  $g$ ,  $r$  is **order** of  $g$

# Pfaffian Functions

- ▶  $\vec{f} = (f_1, \dots, f_r)$  is a **Pfaffian chain** if there exist polynomials  $P_{i,j}$  verifying

$$\frac{\partial f_i}{\partial x_j} = P_{i,j}(x, f_1(x), \dots, f_r(x))$$

- ▶  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  is a **Pfaffian function** w.r.t.  $\vec{f}$  if there exists polynomial  $Q_g$  such that

$$g(x) = Q_g(x, f_1(x), \dots, f_r(x))$$

- ▶  $\deg(Q_g)$  is **degree** of  $g$ ,  $r$  is **order** of  $g$
- ▶ Zero of  $g$  is called a **Pfaffian set**; such sets are **included in an o-minimal structure**

# Pfaffian Functions

- ▶  $\vec{f} = (f_1, \dots, f_r)$  is a **Pfaffian chain** if there exist polynomials  $P_{i,j}$  verifying

$$\frac{\partial f_i}{\partial x_j} = P_{i,j}(x, f_1(x), \dots, f_r(x))$$

- ▶  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  is a **Pfaffian function** w.r.t.  $\vec{f}$  if there exists polynomial  $Q_g$  such that

$$g(x) = Q_g(x, f_1(x), \dots, f_r(x))$$

- ▶  $\deg(Q_g)$  is **degree** of  $g$ ,  $r$  is **order** of  $g$
- ▶ Zero of  $g$  is called a **Pfaffian set**; such sets are **included in an o-minimal structure**
- ▶ e.g.  $e^x, e^{e^x}, \dots, \tan x, \ln x, x^\pi$

# Main Theorem

Theorem (Partitioning Pfaffian sets [Lotz-N-Vorobjov, 2024])

Let  $\Gamma$  be a collection of Pfaffian sets in  $\mathbb{R}^n$  of dimension  $k$ , where each  $\gamma \in \Gamma$  has order  $r$ .

1. For any  $D \geq 1$ , there is  $P \in \mathbb{R}[X_1, \dots, X_n]$  of degree  $D$ , such that *each cell induced by  $P$  intersects at most  $\frac{|\Gamma|}{D^{n-k-r}}$  elements of  $\Gamma$ .*



# Main Theorem

**Theorem (Partitioning Pfaffian sets [Lotz-N-Vorobjov, 2024])**

Let  $\Gamma$  be a collection of Pfaffian sets in  $\mathbb{R}^n$  of dimension  $k$ , where each  $\gamma \in \Gamma$  has order  $r$ .

1. For any  $D \geq 1$ , there is  $P \in \mathbb{R}[X_1, \dots, X_n]$  of degree  $D$ , such that *each cell induced by  $P$*  intersects at most  $\frac{|\Gamma|}{D^{n-k-r}}$  *elements of  $\Gamma$* .
2. For any  $D \geq 1$ , there is a Pfaffian function  $P'$  of degree  $D$  such that *each cell induced by  $P$*  intersects at most  $\frac{|\Gamma|}{D^{n-k}}$  *elements of  $\Gamma$* .

# Main Theorem

## Theorem (Partitioning Pfaffian sets [Lotz-N-Vorobjov, 2024])


Let  $\Gamma$  be a collection of Pfaffian sets in  $\mathbb{R}^n$  of dimension  $k$ , where each  $\gamma \in \Gamma$  has order  $r$ .

1. For any  $D \geq 1$ , there is  $P \in \mathbb{R}[X_1, \dots, X_n]$  of degree  $D$ , such that *each cell induced by  $P$*  intersects at most  $\frac{|\Gamma|}{D^{n-k-r}}$  elements of  $\Gamma$ .
2. For any  $D \geq 1$ , there is a Pfaffian function  $P'$  of degree  $D$  such that *each cell induced by  $P$*  intersects at most  $\frac{|\Gamma|}{D^{n-k}}$  elements of  $\Gamma$ .


## Takeaway

1. *Generalization* of Polynomial Partitioning to Pfaffians
2. New technique of *Pfaffian Partitioning*

# Proof - Main Technical Step




# Proof - Main Technical Step



the line intersects three cells induced by  $P$




# Proof - Main Technical Step



the line intersects three cells induced by  $P$

- Poly.  $P$  of deg.  $D$  in  $n$  variables induces **at most  $D^n$**  cells

# Proof - Main Technical Step



the line intersects three cells induced by  $P$

- ▶ Poly.  $P$  of deg.  $D$  in  $n$  variables induces **at most  $D^n$**  cells
- ▶ We show for a  $k$ -dimensional Pfaffian set  $\gamma$  of order  $r$   
 **$\gamma$  intersects at most  $D^{k+r}$  cells induced by  $P$**

# Applications of Partitioning Pfaffians

- (Pfaffian Szemerédi-Trotter)  $m$  points and  $n$  Pfaffian curves in  $\mathbb{R}^2$ :  $\mathcal{O}(m^{\frac{2r+2}{2r+3}+\varepsilon} n^{\frac{(r+2)}{2r+3}} + m + n)$  incidences

# Applications of Partitioning Pfaffians

- ▶ (Pfaffian Szemerédi-Trotter)  $m$  points and  $n$  Pfaffian curves in  $\mathbb{R}^2$ :  $\mathcal{O}(m^{\frac{2r+2}{2r+3}+\varepsilon} n^{\frac{(r+2)}{2r+3}} + m + n)$  incidences
- ▶ We also count joints between Pfaffian curves



# Applications of Partitioning Pfaffians

- ▶ (Pfaffian Szemerédi-Trotter)  $m$  points and  $n$  Pfaffian curves in  $\mathbb{R}^2$ :  $\mathcal{O}(m^{\frac{2r+2}{2r+3}+\varepsilon} n^{\frac{(r+2)}{2r+3}} + m + n)$  incidences
- ▶ We also count joints between Pfaffian curves
- ▶ More applications possible

# Applications of Partitioning Pfaffians

- ▶ (Pfaffian Szemerédi-Trotter)  $m$  points and  $n$  Pfaffian curves in  $\mathbb{R}^2$ :  $\mathcal{O}(m^{\frac{2r+2}{2r+3}+\varepsilon} n^{\frac{(r+2)}{2r+3}} + m + n)$  incidences
- ▶ We also count joints between Pfaffian curves
- ▶ More applications possible
- ▶ Our technique lends itself to generalizing to other o-minimal structures (caveat [Basu-Lerario-N, 2019])

## References

- S. Basu, A. Lerario, and A. Natarajan. Zeroes of polynomials on definable hypersurfaces: pathologies exist, but they are rare. *The Quarterly Journal of Mathematics*, 70(4):1397–1409, 2019.
- L. Guth. Polynomial partitioning for a set of varieties. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 159, pages 459–469. Cambridge University Press, 2015.
- L. Guth and N. H. Katz. On the erdős distinct distances problem in the plane. *Annals of Mathematics*, pages 155–190, 2015.
- M. Lotz, A. Natarajan, and N. Vorobjov. Partitioning theorems for sets of semi-pfaffian sets, with applications, 2024. URL <https://arxiv.org/abs/2412.02961>.