

Assignment 1

Abhiroop Chintalapudi - AI20BTECH11005

Download all python and latex codes from

<https://github.com/abhiroopchintalapudi03/EE3900.git>

1 PROBLEM 2.5

Check whether

$$\mathbf{A} = \begin{pmatrix} 5 \\ -2 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 6 \\ 4 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} 7 \\ -2 \end{pmatrix} \quad (1.0.1)$$

are the vertices of an isosceles triangle.

2 SOLUTION

Let,

$$\mathbf{A} = \begin{pmatrix} 5 \\ -2 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 6 \\ 4 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} 7 \\ -2 \end{pmatrix} \quad (2.0.1)$$

For the triangle to be isosceles triangle, one of

$$\|\mathbf{A} - \mathbf{B}\| = \|\mathbf{B} - \mathbf{C}\| \text{ or}$$

$$\|\mathbf{B} - \mathbf{C}\| = \|\mathbf{C} - \mathbf{A}\| \text{ or}$$

$$\|\mathbf{C} - \mathbf{A}\| = \|\mathbf{A} - \mathbf{B}\|$$

Now,

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} 5 - 6 \\ (-2) - 4 \end{pmatrix} = \begin{pmatrix} -1 \\ -6 \end{pmatrix}$$

$$\Rightarrow \|\mathbf{A} - \mathbf{B}\| = \sqrt{37}$$

$$\mathbf{B} - \mathbf{C} = \begin{pmatrix} 6 - 7 \\ 4 - (-2) \end{pmatrix} = \begin{pmatrix} -1 \\ 6 \end{pmatrix}$$

$$\Rightarrow \|\mathbf{B} - \mathbf{C}\| = \sqrt{37}$$

$$\mathbf{C} - \mathbf{A} = \begin{pmatrix} 7 - 5 \\ (-2) - (-2) \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$$

$$\Rightarrow \|\mathbf{C} - \mathbf{A}\| = 2$$

$$\text{As } \|\mathbf{A} - \mathbf{B}\| = \|\mathbf{B} - \mathbf{C}\| = \sqrt{37}$$

\Rightarrow In $\triangle ABC$ sides AB, BC are equal.

$\Rightarrow \triangle ABC$ is an isosceles triangle.

You can also see from the below diagram that the triangle is an isosceles triangle with sides AB, BC equal.




Fig. 0: $\triangle ABC$