

Frustration shapes multi-channel Kondo physics: a star graph perspective

Siddhartha Patra¹, Abhirup Mukherjee¹, Anirban Mukherjee¹, N. S. Vidhyadhiraja⁴, A. Taraphder⁵ and Siddhartha Lal¹

E-mail: sp14ip022@iiserkol.ac.in, am18ip014@iiserkol.ac.in, mukherjee.anirban.anirban@gmail.com, raja@jncasr.ac.in, arghya@phy.iitkgp.ernet.in, slal@iiserkol.ac.in

¹Department of Physical Sciences, Indian Institute of Science Education and Research-Kolkata, W.B. 741246, India

⁴Theoretical Sciences Unit, Jawaharlal Nehru Center for Advanced Scientific Research, Jakkur, Bengaluru 560064, India

⁵Department of Physics, Indian Institute of Technology Kharagpur, Kharagpur 721302, India

Abstract. We study the overscreened multi-channel Kondo (MCK) model using the recently developed unitary renormalization group (URG) technique. Our results display the importance of ground state degeneracy in explaining various important properties like the breakdown of screening and the presence of local non-Fermi liquids. The impurity susceptibility of the intermediate coupling fixed point Hamiltonian in the zero-bandwidth (or star graph) limit shows a power-law divergence at low temperature. Despite the absence of inter-channel coupling in the MCK fixed point Hamiltonian, the study of mutual information between any two channels shows non-zero correlation between them. A spectral flow analysis of the star graph reveals that the degenerate ground state manifold possesses topological quantum numbers. Upon disentangling the impurity spin from its partners in the star graph, we find the presence of a local Mott liquid arising from inter-channel scattering processes. The low energy effective Hamiltonian obtained upon adding a finite non-zero conduction bath dispersion to the star graph Hamiltonian for both the two and three-channel cases displays the presence of local non-Fermi liquids arising from inter-channel quantum fluctuations. Specifically, we confirm the presence of a local marginal Fermi liquid in the two channel case, whose properties show logarithmic scaling at low temperature as expected. Discontinuous behaviour is observed in several measures of ground state entanglement, signalling the underlying orthogonality catastrophe associated with the degenerate ground state manifold. We extend our results to underscreened and perfectly screened MCK models through duality arguments. A study of channel anisotropy under renormalisation flow reveals a series of quantum phase transitions due to the change in ground state degeneracy. Our work thus presents a template for the study of how a degenerate ground state manifold arising from symmetry and duality properties in a multichannel quantum impurity model can lead to novel multicritical phases at intermediate coupling.

1. Introduction

A local antiferromagnetic exchange interaction between a spin- $\frac{1}{2}$ impurity and its host metal gives rise to the well-understood phenomenon of the Kondo effect ([1], see [2] for an extensive discussion). Here, the impurity local moment is screened by the conduction electrons at temperatures below a certain scale called the Kondo temperature [3, 4, 5, 6, 7, 8, 9, 10], leading to a spin singlet ground state and local Fermi liquid excitations above that ground state [11, 12]. The screening manifests as an initial increase at temperatures below the Kondo temperature, followed by a saturation of the resistivity of the metal [1, 13], and in the saturation of the impurity contribution to the magnetic susceptibility at low temperatures [6, 7, 8]. Generalisations of this model are obtained by taking impurities of higher spin [2, 14, 9], adding interactions between them [15, 16, 17, 18, 19, 20, 21], by promoting the model to a lattice of Kondo impurities [22, 23, 24, 25, 26] or by considering the case of a correlated host metal [27]. One can also construct multi-channel Kondo (MCK) models by allowing K conduction electron channels ($K > 1$) to interact with a spin- S_d impurity [14, 28, 29] via a common exchange coupling \mathcal{J} . Using conformal field theory [30, 31, 32, 9, 33, 34], bosonization [35, 36, 37, 38, 39], numerical renormalization group [40, 32, 41], Bethe ansatz [42, 43, 44, 45, 46] and other methods [47, 48, 49, 50], it has been shown that for the over-screened systems ($K > 2S_d$), the low energy physics is of the non-Fermi liquid type obtained at an intermediate coupling fixed point of the RG flows [14]. Further, the system displays anomalous behaviour in thermodynamic properties near $T = 0$ and a fractional zero temperature entropy in the thermodynamic limit (upon ensuring that temperature is taken to zero after the system size is taken to infinity [51, 38]). This anomalous divergent behaviour is a signature of the fact that the MCK system with a single channel-symmetric exchange coupling is quantum critical: the ground state is susceptible to perturbations that introduce channel anisotropy in the exchange couplings [14, 45, 32, 37, 52]. This makes the experimental realisation of such states quite challenging. Nevertheless, several features of the two-channel Kondo model have been reproduced using structural two-level systems [28, 53] interacting with a conduction bath [54, 55, 56, 57, 58]. More recently, it has become possible to tune a two-channel quantum dot system across the quantum phase transition [59, 60], revealing the fractionalisation of the low-energy degrees of freedom [61, 49, 62].

The importance of ground state degeneracy and quantum-mechanical *frustration* in the MCK problem has not received sufficient attention, even though they play crucial roles in determining the quantum

criticality and associated non-Fermi liquid physics of the system. This is, at least partially, due to the lack of an intermediate-coupling effective Hamiltonian that describes the low-energy physics of the MCK (i.e., reveals the nature of the excitations). Frustration in this context refers to the inability of the impurity spin to bind with the conduction channels, and be screened completely by forming a maximally-entangled singlet ground state. The absence of such frustration in the single-channel spin-1/2 Kondo model (and more generally, the exact screening variant of the MCK) means that imperfect screening ($K \neq 2S_d$) leads to dramatic differences in the multi-channel models. Indeed, we demonstrate that there exists an orthogonality catastrophe in the imperfectly screened MCK models that leads to the non-Fermi liquid behaviour of its low-energy excitations, and that this is directly related to the existence of a degenerate ground state manifold. The effects of these non-Fermi liquid low energy excitations are expected to appear in measures of many-particle entanglement; the lack of an intermediate coupling Hamiltonian has, however, prevented the study of the renormalisation group (RG) evolution of such measures [63, 64]. Importantly, our approach enables a comparison of, for instance, the inter-channel mutual information along the RG flow and as functions of the excitation energies, providing insight into the nature of the fixed point theory.

To obtain the RG flow of the MCK model, we have employed the unitary renormalization group (URG) technique developed recently by some of us [65, 66]. The method has been applied on several fermionic and spin problems like the single-channel Kondo model [67], the kagome antiferromagnet [68], the 1D [69] and 2D Hubbard models [70, 71, 72], the reduced-BCS model [73] as well as other generalised models of interacting electrons with and without translation invariance [66]. The URG proceeds by resolving the number fluctuations in the electronic Fock states at high energies by the iterative application of many-particle unitary transformations, leading to their successive decoupling from the states lying at lower energies. Applying this URG method to the channel-isotropic MCK model leads to a low-energy fixed point Hamiltonian at the intermediate (Kondo) coupling fixed point of Ref.[14]. In further analysing the effective Hamiltonian, we focus initially on the zero bandwidth limit of this Hamiltonian (which corresponds to a frustrated quantum spin model on a star graph). We show thereby that several important properties of the MCK problem, such as the ground state degeneracy and breakdown of screening, can be understood from the zero bandwidth problem. We then proceed to studying the low energy effective Hamiltonian for the non-Fermi liquid

excitations lying above the degenerate ground state manifold. This enables the computation of a plethora of quantities, such as various thermodynamic measures (e.g., specific heat and susceptibility), many-particle entanglement and the self-energy of the propagating degrees of freedom. We also explore various signatures of criticality, and various duality transformations of the MCK Hamiltonian. Finally, we investigate the evolution of ground state degeneracy under RG of the channel anisotropic MCK model.

Summary of Main Results

We start in section 2 by describing the URG flows for the MCK, the intermediate coupling URG fixed point Hamiltonian and the features of the zero bandwidth (star graph) limit of the effective Hamiltonian. In section 3, we explore some properties of the zero bandwidth model: namely, the degree of compensation, impurity magnetisation and impurity susceptibility. The degree of compensation is simply the average correlation between the impurity spin and the conduction channel local spins; it is found to be maximum at exact screening, and decreases for both over- and under-screening. The magnetisation and susceptibility of the star graph model show a discontinuity and a T^{-1} divergence respectively. We also explore the topological properties of the degenerate ground state manifold of the star graph model using non-local twist and translation operators. On decoupling the impurity spin from the conduction spin degrees of freedom within the star graph, we obtain an all-to-all Hamiltonian that represents a local Mott liquid phase. In section 4, we obtain the low energy effective Hamiltonian for the low-lying excitations by including a dispersion into the conduction bath. Notable features of this effective Hamiltonian include (i) the complete absence of local Fermi liquid terms, and (ii) the emergence of local non-Fermi liquid terms. Both are shown to result from the ground state degeneracy of the underlying star graph model. The excitations of the two-channel Kondo problem are found to pertain to a local marginal Fermi liquid, signalling an orthogonality catastrophe.

In section 5, we study various entanglement signatures (e.g., von-Neumann entanglement entropy, mutual information, multipartite information, Bures distance) of the low energy fixed-point ground state of the MCK problem, with and without dispersion into the bath. The discontinuous behaviour observed in these measures for the MCK problem reveal clear distinctions from the single-channel case. In 6, we discuss two duality transformations that exist for the MCK model - a strong-weak duality, and an over-screened-under-screened duality. It is found that the transformations are constrained by the fixed point

structure, and impose relations between the under-screened and over-screened models. In 7, we show the robustness of the ground state degeneracy of the star graph model against the impurity-channel coupling anisotropy, and correlate it with a URG treatment of the channel anisotropic MCK. The isotropic fixed point is found to be unstable to asymmetry, leading to impurity phase transitions that lead to dramatic changes in the ground state degeneracy. For the sake of brevity, some supporting evidence and details of certain lengthy calculations are presented in Supplementary Materials [74]. Finally, we conclude in section 8.

2. Fixed point theory of over-screened MCK model

2.1. RG flows towards intermediate coupling

We start with the K - conduction channel Kondo model Hamiltonian with isotropic couplings [14]:

$$H = \sum_{\substack{l,k \\ \alpha=\uparrow,\downarrow}} \epsilon_{k,l} \hat{n}_{k\alpha,l} + \frac{\mathcal{J}}{2} \sum_{\substack{l,k,k' \\ \alpha,\alpha'=\uparrow,\downarrow}} \vec{S}_d \cdot \vec{\sigma}_{\alpha\alpha'} c_{k\alpha,l}^\dagger c_{k'\alpha',l}, \quad (1)$$

where \mathcal{J} is the Kondo spin-exchange coupling, $c_{k\alpha,l}$ is the fermionic field operator at momentum k , spin α and channel l , $\epsilon_{k,l}$ represents the dispersion of the l^{th} conduction channel, $\vec{\sigma}$ is the vector of Pauli matrices and $\vec{S}_d = \frac{1}{2}\vec{\sigma}_d$ is the impurity spin operator. Here, l sums over the K channels of the conduction bath, k, k' sum over all the momentum states of the bath and α, α' sum over the two spin indices of a single electron.

We now perform a renormalisation group analysis of the MCK Hamiltonian using the recently developed URG method [70, 71, 65, 66, 73, 68, 69]. The RG proceeds by applying unitary transformations in order to block-diagonalize the Hamiltonian by removing number fluctuations of the high energy degrees of freedom. At the j^{th} RG step, we focus on decoupling the highest energy electronic state denoted by $|j\rangle$. The Hamiltonian will in general not conserve the number of particles in this state: $[H_{(j)}, \hat{n}_j] \neq 0$, and the unitary transformation $U_{(j)}$ causes this number fluctuation to vanish [65, 66]:

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger, \quad [H_{(j-1)}, \hat{n}_j] = 0. \quad (2)$$

The unitary transformations are given in terms of a fermionic generator $\eta_{(j)}$ [65, 66]:

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 + \eta_{(j)} - \eta_{(j)}^\dagger \right), \quad \left\{ \eta_{(j)}, \eta_{(j)}^\dagger \right\} = 1, \quad (3)$$

where $\{\cdot\}$ is the anticommutator. Further, $[\eta_{(j)}, \eta_{(j)}^\dagger] = 2\hat{n}_j - 1$. The unitary operator $U_{(j)}$ mentioned in Eq. (3) can be thought of as a special case of the most general form $U = e^{\mathcal{S}}$ of a unitary operator defined by an

anti-Hermitian generator \mathcal{S} . For our case, the generator turns out to be $\mathcal{S} = \frac{\pi}{4} (\eta_{(j)}^\dagger - \eta_{(j)})$, such that the unitary operator is $U_{(j)} = e^{\frac{\pi}{4} (\eta_{(j)}^\dagger - \eta_{(j)})}$ [65]. If one then uses the anti-commutation and commutation properties of $\eta_{(j)}, \eta_{(j)}^\dagger$ [65, 66], the exponential can be expanded in its Taylor series and then resummed into the form in Eq. (3).

The generator itself is given by [65, 66]

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - \text{Tr}(H_{(j)}\hat{n}_j)} c_j^\dagger \text{Tr}(H_{(j)}c_j) . \quad (4)$$

The operators $\eta_{(j)}, \eta_{(j)}^\dagger$ can be thought of as the many-particle analogue of the single-particle field operators c_j, c_j^\dagger that change the occupation number of the single-particle Fock space, $|n_j\rangle$. The presence of the operators in $\eta_{(j)}$ other than c_j are required to ensure that the decoupling of the specific Fock state under consideration is carried out through a unitary modification of the spectrum of the remnant degrees of freedom. Since the j^{th} degree of freedom is coupled with many others, a rotation of the basis of j also involves a rotation of the many-particle basis of the remaining degrees of freedom in the Hamiltonian $H_{(j)}$; this is encoded within the $\text{Tr}(H_{(j)}c_j)$ term. The operator $\hat{\omega}_{(j)}$ encodes the quantum fluctuation scales arising from the interplay of the kinetic energy terms and the interaction terms in the Hamiltonian:

$$\hat{\omega}_{(j)} = H_{(j-1)} - H_{(j)}^i . \quad (5)$$

$H_{(j)}^i$ is that part of $H_{(j)}$ that commutes with \hat{n}_j but does not commute with at least one \hat{n}_l for $l < j$. The RG flow continues till a fixed point is reached at an energy D^* . The Greens function in front of c_j (in Eq. 4) accounts for the dynamical cost of this rotation in terms of an operator for quantum fluctuations ($\hat{\omega}$), as well as the kinetic and self-energies $\text{Tr}(H_{(j)}\hat{n}_j)$.

There exist other renormalisation group schemes that depend on canonical transformations, such as Wegner's continuous unitary transformations (CUT) RG [75] and the similarity RG method by Glazek and Wilson [76]. There are at least two specific distinctions between the CUT RG and the URG method adopted in the present work.

- While most RG methods that apply successive unitary transformations rely on a gradual decrease in the off-diagonal content of the Hamiltonian, the URG proceeds by decoupling each discrete Fock space node at a time. In other words, the URG achieves a block-diagonalisation of the Hamiltonian at each RG step, while the CUT RG involves a gradual decay of the off-diagonal terms of the Hamiltonian in order to make it more band diagonal.

- The CUT RG involves choosing a truncation scheme that allows closing the RG equations. The URG, on the other hand, is able to resum the coupling expansion series through the denominator structure and the quantum fluctuation operator.

Detailed comparisons of the URG with other methods (e.g., the functional RG, spectrum bifurcation RG etc.) can be found in Refs. [70, 65].

The derivation of the RG equation for the over-screened regime ($2S < K$) of the spin- S -impurity K -channel Kondo problem is shown in detail in the Supplementary Materials [74]. On decoupling circular isoenergetic shells at energies $D_{(j)}$, the change in the Kondo coupling at the j^{th} RG step, $\Delta\mathcal{J}_{(j)}$, is given by

$$\Delta\mathcal{J}_{(j)} = -\frac{\mathcal{J}_{(j)}^2 \mathcal{N}_{(j)}}{\omega_{(j)} - \frac{D_{(j)}}{2} + \frac{\mathcal{J}_{(j)}}{4}} \left(1 - \frac{1}{2} \rho \mathcal{J}_{(j)} K\right) , \quad (6)$$

where $\mathcal{N}_{(j)}$ is the number of electronic states at the energy shell $D_{(j)}$. We work in the low quantum fluctuation regime $\omega_{(j)} < \frac{D_{(j)}}{2}$. There are three fixed points of the RG equation. One arises from the vanishing of the denominator, and was present in the single-channel Kondo RG equation as well [67]. As shown there, this fixed-point goes to $\mathcal{J}^* = \infty$ as the bare bandwidth of the conduction electrons is made large. The other fixed point is the trivial one at $\mathcal{J}^* = 0$. The third fixed point is reached when the numerator vanishes: $\mathcal{J}^* = \frac{2}{K\rho}$ [77, 78, 79, 14]. Only the intermediate fixed point is found to be stable. This is consistent with results from Bethe ansatz calculations [80, 42, 46, 45, 44, 43], CFT calculations [30, 9, 31], bosonization treatments [35, 38] and NRG analysis [41, 81].




Figure 1. The three fixed points of the over-screened RG equation. Only the intermediate one is stable.

The RG equation reduces to the perturbative form $\Delta\mathcal{J}_{(j)} \simeq \frac{\mathcal{J}_{(j)}^2 \mathcal{N}_{(j)}}{D_{(j)}} (1 - \frac{1}{2} \rho \mathcal{J}_{(j)} K)$ [78, 79, 14, 82] when one replaces $\omega_{(j)}$ with the ground state energy $-\frac{D_{(j)}}{2}$ and assumes $\mathcal{J} \ll D_{(j)}$.

2.2. The star graph as the zero-bandwidth limit of the fixed point Hamiltonian

The fixed point Hamiltonian takes the form

$$H^* = \sum_l \left[\sum_k^* \epsilon_{k,l} \hat{n}_{k\alpha,l} + \mathcal{J} \sum_{\substack{k,k' \\ \alpha,\alpha'}}^* \vec{S}_d \cdot \frac{1}{2} \vec{\sigma}_{\alpha\alpha'} c_{k\alpha,l}^\dagger c_{k'\alpha',l} \right] . \quad (7)$$

We have not explicitly written the decoupled degrees of freedom $D_{(j)} > D^*$ in the Hamiltonian. The $*$ over the summations indicate that only the momenta inside the window D^* enter the summation.

We will first study the zero bandwidth limit of the fixed point Hamiltonian, obtained by compressing the sum over the momentum states to a single state at the Fermi surface for each conduction channel (see Fig.2). Upon setting the chemical potential equal to the Fermi energy, the kinetic energy part vanishes and the zero bandwidth model becomes a Heisenberg spin-exchange Hamiltonian

$$H^* = \mathcal{J} \sum_l \sum_{\substack{k, k' \\ \alpha, \alpha'}}^* \vec{S}_d \cdot \frac{1}{2} \vec{\sigma}_{\alpha\alpha'} c_{k\alpha, l}^\dagger c_{k'\alpha', l} = \mathcal{J} \vec{S}_d \cdot \vec{S}. \quad (8)$$

At the last step, we defined the total bath local spin operator $S = \sum_l \vec{S}_l = \frac{1}{2} \sigma_l = \frac{1}{2} \sum_{k, k'}^* \sum_{\alpha, \alpha'} \vec{\sigma}_{\alpha\alpha'} c_{k\alpha, l}^\dagger c_{k'\alpha', l}$. The star graph com-




Figure 2. Zero bandwidth limit of the fixed point Hamiltonian. The central yellow node is the impurity; it is interacting with the K green outer nodes that represent the local spins of the channels.

mutates with several operators, including the total spin operator $J^z = S_d^z + S^z$ along z , the square of the total bath local spin operator (S^2) and the string operators

$$\pi^{x,y,z} = \sigma_d^{x,y,z} \otimes_{l=1}^K \sigma_l^{x,y,z}. \quad (9)$$

If we define the global spin operator $\vec{J} = \vec{S} + \vec{S}_d$, the star graph Hamiltonian can be written as $\frac{1}{2} \mathcal{J} [J^2 - S_d^2 - S^2]$.

- The ground state is achieved for the maximal value of S , $S = \frac{K}{2}$, and the corresponding minimal value of J , $J = |\frac{K}{2} - S_d|$. The ground state energy is therefore

$$E_g = \frac{1}{2} \mathcal{J} [J(J+1) - S_d(S_d+1) - S(S+1)] \quad (10)$$

$$= \begin{cases} -\mathcal{J} S_d (\frac{K}{2} + 1) & \text{when } K \geq 2S_d, \text{ and} \\ -\mathcal{J} \frac{K}{2} (S_d + 1) & \text{otherwise.} \end{cases} \quad (11)$$

- The value of J corresponds to a multiplicity of $2J+1 = |K - 2S_d| + 1$ in J^z , and since the Hamiltonian does not depend on J^z , these orthogonal states $|J^z\rangle$ constitute a degeneracy of $g_K^{S_d} = |K - 2S_d| + 1$.

The π^z acts on the eigenstates $|J^z\rangle$ and reveals the odd-even parity of the eigenvalue J^z , and is hence a parity operator. Interestingly, the string operator π^z is a Wilson loop operator [83] that wraps around all the nodes of the star graph:

$$\pi^z = \exp \left[i \frac{\pi}{2} \left(\sigma_d^z + \sum_{l=1}^K \sigma_l^z - K \right) \right] = e^{i\pi(J^z - \frac{1}{2}K)} \quad (12)$$

π^x and π^y are 't Hooft operators [83] and mix states of opposite parity. For example, it can be shown that $\pi^x |J^z\rangle = -|J^z\rangle$.

There are several reasons for working with the star graph in particular and zero mode Hamiltonians in general. In the single-channel Kondo model, the star graph is just the two spin Heisenberg problem, and it reveals the stabilization of the Kondo model ground state, as well as certain thermodynamic properties (e.g., the impurity contribution to the susceptibility) [84, 85, 86, 87, 84, 67]. Similarly, in the MCK model, the star graph captures accurately the nature of the RG flows. At weak coupling $\mathcal{J} \rightarrow 0^+$, the central spin is weakly coupled to the outer spins and prone to screening because of the S^\pm terms in the star graph; at strong coupling $\mathcal{J} \rightarrow \infty^-$, the outer spin-half objects tightly bind with the central spin-half object to form a single spin object that interacts with the remaining states through an exchange coupling which is RG relevant. This renders both the terminal fixed points unstable. The true stable fixed point must then lie somewhere in between, and we recover the schematic diagram of fig. 1.

Moreover, the RG flows of the MCK model have been shown to preserve the degeneracy of the ground state [41, 88, 89], and the star graph captures this degeneracy in its entirety. This is important, because it will be shown in a later section that the lowest excitations of the intermediate fixed point are described by a non-Fermi liquid phase, and it can be argued that this non-Fermi liquid physics arises solely from the ground state degeneracy of the underlying zero mode Hamiltonian. As mentioned previously, the ground state degeneracy of the more general star graph with a spin- S_d impurity and K channels is given by $g_K^{S_d} = |K - 2S_d| + 1$. The cases of $K = 2S_d$, $K < 2S_d$ and $K > 2S_d$ correspond to exactly screened, under-screened and over-screened regimes respectively. The latter two cases correspond to a multiply-degenerate manifold with $g_K^S > 1$, and simultaneously have non-Fermi liquid phases [14, 90, 35, 77, 80, 91, 33, 92, 93, 94, 30, 49, 9, 95, 96, 97, 98], while the first regime has a unique ground state with $g_K^S = 1$ and is described by a local Fermi liquid (LFL) phase [6, 11, 14, 7, 8], thereby substantiating the claim that a degeneracy greater than unity is closely tied to non-Fermi liquid physics.

We now comment on the validity of our study of the zero-bandwidth model at the intermediate coupling fixed point (ICFP) of the MCK problem. Noziers and Blandin indeed present an argument for why the zero-bandwidth model at the strong coupling fixed point ($J = \infty$) is unstable due to the existence of non-trivial RG relevant quantum fluctuations [14]. The resulting RG flow of the Kondo coupling J to the ICFP resolves these quantum fluctuations. The RG stability of the Kondo coupling J at the ICFP then allows for a renormalised perturbation theory approach to studying the fixed point Hamiltonian. Specifically, the zero-bandwidth model can be safely extracted (in the RG sense) as the zeroth approximation of a renormalised perturbation theory in the ratio of the conduction bath hopping amplitude (t) to J . A study of the zero-bandwidth model at the ICFP offers, in turn, new perspectives into the problem, e.g., the ground state degeneracy and its relation to the breakdown of exact screening etc., as described in the two preceding paragraphs. It also informs us on the ground state manifold about which to carry out a systematic renormalised perturbative expansion to obtain the low-lying excitations (see Sec.4 below, and Sec.III of the supplementary materials). Moreover, in the following sections, we will show how the inherent quantum-mechanical frustration of singlet order that is present in the Hamiltonian leads to the non-trivial physics of the fixed points in terms of non-Fermi liquid phase, diverging thermodynamic quantities, quantum criticality as well as emergent gauge theories.

3. Important properties of the star graph

3.1. Degree of compensation: a measure of the frustration

One can quantify the screening of the local moment at the impurity site by defining a degree of compensation Γ . Such a quantity also measures the inherent singlet frustration in the problem: the higher the degree of compensation, the better the spin can be screened into a singlet and lower is the frustration. It is given by the antiferromagnetic correlation existing between the impurity spin and conduction electron channels: $\Gamma \equiv -\langle \vec{S}_d \cdot \vec{S} \rangle$. The expectation value is calculated in the ground state. Since the inner product is simply the ground state energy of a spin- S_d impurity K -channel MCK model in units of the exchange coupling J , we have $\Gamma = \frac{1}{2} [l_{\text{imp}}^2 + l_c^2 - g_K^S (g_K^S - 1)]$, where $l_{\text{imp}}^2 = S_d(S_d + 1)$ is the length-squared of the impurity spin. Similarly, $l_c^2 = \frac{K}{2} (\frac{K}{2} + 1)$ is the length-squared of the total conduction bath spin. $g_K^S = |K - S_d| + 1$ is the ground state degeneracy. We will explore the three regimes of screening by defining $K = K_0 + \delta$, $S_d =$

$\frac{K_0}{2} - \delta$. $\delta = 0$ represents the exactly-screened case of $K = 2S_d = K_0$. Non-zero δ represents either over- or under-screening. In terms of K_0 and δ , the degree of compensation becomes

$$\Gamma = \frac{1}{4} [(K_0 + 1)^2 - (|\delta| + 1)^2]. \quad (13)$$

For a given K_0 , the degree of compensation Γ is maximised for exact screening $\delta = 0$, and is reduced for $\delta \neq 0$ (see Fig.3). This shows the inability of the system to form a unique singlet ground state and reveals the quantum-mechanical frustration inherent in the zero mode Hamiltonian and therefore in the entire problem. The degree of compensation is symmetric under the Hamiltonian transformation $\delta \rightarrow -\delta$, and this represents a duality transformation between over-screened and under-screened MCK models. This topic will be discussed in more detail later.




Figure 3. Variation of the degree of compensation Γ upon tuning from under-screening to over-screening by varying the deviation from exact screening $\delta = K_0/2 - S$. The maximum spin compensation occurs at exact-screening $\delta = 0$.

3.2. Impurity magnetization and susceptibility

The impurity magnetic susceptibility can be obtained by diagonalising the zero-mode Hamiltonian $H(h)$ in the presence of a local magnetic field h on the impurity:

$$H(h) = \mathcal{J}^* \vec{S}_d \cdot \vec{S} + h S_d^z. \quad (14)$$

The Hamiltonian commutes with \vec{S} , so it is already block-diagonal in terms of the eigenvalues M of S . M takes values in the range $[M_{\min}, M_{\max}]$, where $M_{\max} = K/2$ for a K -channel Kondo model, and $M_{\min} = 0$ if K is even, otherwise $\frac{1}{2}$. The diagonalisation of the Hamiltonian is shown in the Supplementary Materials [74]. Defining $\alpha = \frac{1}{2}(\mathcal{J}m + h) + \frac{\mathcal{J}}{4}$ and $x_m^M = M(M + 1) - m(m + 1)$, the partition function

can be written as

$$Z(h) = \sum_{M=M_{\min}}^{M_{\max}} r_M^K \left[\sum_{\substack{m=-M, \\ m \in \mathbb{Z}}}^{M-1} 2e^{\beta \frac{\mathcal{J}}{4}} \cosh \beta \sqrt{\mathcal{J}^2 x_m^M / 4 + \alpha^2} + 2e^{-\beta \mathcal{J} M/2} \cosh \beta h/2 \right]. \quad (15)$$

Here, $\beta = \frac{1}{k_B T}$, M sums over the eigenvalues of S while m sums over $\mathcal{J}^z - \frac{1}{2}$ and the additional degeneracy factor $r_M^K = K^{-1} C_{K/2-M}$ arises from the possibility that there are multiple subspaces defined by $S = M$. To calculate the impurity magnetic susceptibility, we will use the expression

$$\chi = \frac{1}{\beta} \lim_{h \rightarrow 0} \left[\frac{Z''(h)}{Z(h)} - \left(\frac{Z'(h)}{Z(h)} \right)^2 \right], \quad (16)$$

where the $'$ indicates derivative with respect to h . At low temperatures, the susceptibility takes the form

$$\chi \rightarrow \frac{\beta \Sigma_{\max}}{2M_{\max}(2M_{\max} + 1)^2} = \frac{\beta(K-1)}{12(K+1)} \sim \frac{1}{T}. \quad (17)$$




Figure 4. Variation of impurity susceptibility against temperature. The exactly screened case ($\delta = 0$) saturates to a constant value at low temperatures, indicating complete screening. The cases of inexact screening show a divergence of the susceptibility, which means there is a remnant local spin at the impurity. Since the axes are in log scale, the behaviour is $\log T \sim -\log \chi$ which translates to $\chi \sim 1/T$.

The χ is seen to diverge as T^{-1} at low temperatures. Such a non-analyticity in a response function is in contrast to the behaviour in the exactly-screened fixed point where the ground state is unique. There, the susceptibility becomes constant at low temperatures: $\chi(T \rightarrow 0) = \frac{W}{4T_K}$, T_K being the single-channel Kondo temperature and W the Wilson number [86, 11, 40, 67]. In Fig. 4, we have checked the case of general spin- S_d impurity by diagonalising the star graph Hamiltonian numerically, and the general conclusion is that all exactly-screened models show a constant impurity susceptibility at $T \rightarrow 0$, while the over-screened and under-screened cases show a

diverging impurity susceptibility in the same limit. A similar divergence is also seen in the susceptibility of the outer spins, calculated by inserting a magnetic field purely on the outer spins. Some additional results regarding susceptibility are presented in the Supplementary Materials [74].

Departure from single-channel behaviour is most apparent in the presence of the highly polarised states $|J^z = \pm \frac{K-1}{2}\rangle$ in the ground-state spectrum,

$$|J^z = \sigma \frac{K-1}{2}\rangle = \frac{1}{\sqrt{1+K}} |\sigma\rangle_d \otimes |S^z = \sigma(\frac{K}{2} - 1)\rangle - \frac{\sqrt{K}}{\sqrt{1+K}} |\bar{\sigma}\rangle_d \otimes |S^z = \sigma \frac{K}{2}\rangle, \quad (18)$$

where $\sigma = \pm 1$ and $|\sigma\rangle_d$ represents the up and down configurations of the impurity spin and $|S^z\rangle$ represents the configuration of the bath spins. These states lead to large unquenched magnetisation at zero temperatures. These states lead to singular behaviour of the impurity magnetisation in vanishing field, $m(h = 0^\pm)$. In the presence of a field, the ground-state becomes unique: $|\psi\rangle_{\text{gs}}(h) = \theta(h) |J^z = \frac{K-1}{2}\rangle + \theta(-h) |J^z = -\frac{K-1}{2}\rangle$, and the impurity magnetisation can be calculated from the state:

$$\begin{aligned} m(h = 0^\pm) &= \theta(h) \langle J^z = \frac{K-1}{2} | S_d^z | J^z = \frac{K-1}{2} \rangle \\ &\quad + \theta(-h) \langle J^z = -\frac{K-1}{2} | S_d^z | J^z = -\frac{K-1}{2} \rangle \\ &= \pm \frac{1}{2} \frac{1-K}{1+K}. \end{aligned} \quad (19)$$




Figure 5. Behaviour of the impurity magnetization for three values of $(K, 2S) = (2, 4), (3, 3), (4, 2)$. Only the case of $K = 2S = 3$ ($\delta = 0$) is analytic near zero. The non-analyticity of the other cases arises because of the frustration brought about by the degeneracy of the star graph ground state.

The magnetization is therefore discontinuous as $h \rightarrow 0$ for $K > 1$. The non-analyticity occurs because the magnetic field is able to flip the $J^z = \pm(K-1)/2$ state into the state of the opposite magnetisation. This available space for scattering is simply the frustration

that we discussed earlier. Indeed, we have checked (see Fig. 5) by diagonalising numerically the star graph Hamiltonian that the non-analyticity exists for all $\delta \neq 0$, where $\delta = \frac{K}{2} - S_d$ is the deviation from exact screening.

These polarised states can also be used to obtain the scattering phase shift suffered by the conduction channels at the Fermi surface. As shown in the Supplementary Materials [74], the impurity-bath hybridisation in the polarised states contribute an excess charge of $n_{\text{exc}}^{(l)} = 1/K$ to each conduction channel. Following Friedel's sum rule, this leads to a total scattering phase shift of π/K for each channel:

$$\delta^{(l)} = \pi n_{\text{exc}}^{(l)} = \frac{\pi}{K}, \quad l \in [1, K] \quad (20)$$

For $K > 1$, each channel therefore experiences an unrenormalised phase shift of value between 0 and π that, as we will show in a later section, is responsible for the non-Fermi liquid behaviour. For example, we will show that this phase shift are linked to the critical non-Fermi liquid exponents of thermodynamic quantities at the fixed point. Right now, some evidence of this role of the phase shift can be provided through its relation to the vanishing field impurity magnetisation m obtained in eq. 19:

$$m(h = 0^\pm) = \pm \frac{1}{2} \frac{\delta^{(l)} - \pi}{\delta^{(l)} + \pi}. \quad (21)$$

When the phase shift $\delta^{(l)}$ acquires a fractional value (in units of π), the impurity magnetisation also becomes fractional (in units of $\frac{1}{2}$). This shows that the unrenormalised phase shift value $0 < \delta^{(l)} < \pi$ is at the heart of the quantum frustration present in the star graph (and therefore the MCK problem) that neither allows the impurity to be screened, nor allows it to decouple from the conduction channels. The decoupling only happens in the limit of extreme screening ($K \rightarrow \infty$), and the impurity spin then becomes a local moment.

3.3. Topological properties of ground state manifold

We now present the non-local twist and translation operators which can be used to explore the degenerate ground state manifold of the star graph model. We begin by defining two operators \hat{T} and \hat{O} , which we will call the translation and twist operators respectively:

$$\hat{T} = e^{i\frac{2\pi}{K}\hat{\Sigma}}, \quad \hat{O} = e^{i\hat{\phi}}, \quad \hat{\Sigma} = [\hat{J}^z - (K-1)/2]. \quad (22)$$

One can see that the generators of these above operators commute with the Hamiltonian: $[H, J^x] = [H, J^z] = [H, J^z] = 0$. In the large K limit, we can perform a semi-classical approximation. Thus, we define

$$e^{i\hat{\phi}} = \frac{J_x}{\sqrt{J^2 - J_z^2}} + i \frac{J_y}{\sqrt{J^2 - J_z^2}}, \quad (23)$$

such $[e^{i\hat{\phi}}, \hat{H}] = 0$. We can label the ground states by the eigenvalues of the translation operator \hat{T} . We can also use the ground states labeled by the eigenvalues of J^z (say M). The ground states are now written as $|M_1\rangle, |M_2\rangle, |M_2\rangle, \dots, |M_K\rangle$.

$$(24)$$

The operations of the translation operators on these states are then given by

$$\hat{T}|M_i\rangle = e^{i\frac{2\pi}{K}[M_i - (K-1)/2]}|M_i\rangle = e^{i2\pi\frac{p_i}{K}}|M_i\rangle, \quad p_i \in [\mathbb{Z}] \quad (25)$$

The braiding rule between the twist and the translation operators is

$$\begin{aligned} \hat{T}\hat{O}\hat{T}^\dagger\hat{O}^\dagger &= e^{\frac{2\pi}{K}[i\hat{\Sigma}, i\hat{\phi}]} = e^{i\frac{2\pi}{K}}, \\ \hat{T}\hat{O}^m\hat{T}^\dagger\hat{O}^{\dagger m} &= e^{\frac{2\pi}{K}[i\hat{\Sigma}, im\hat{\phi}]} = e^{i2\pi\frac{m}{K}}. \end{aligned} \quad (26)$$

The states $\hat{O}^m|M_i\rangle$ (with distinct m) are labelled by different eigenvalues of the translation operations, and are thus orthogonal to each other

$$\hat{T}\hat{O}^m|M_i\rangle = \hat{O}^m\hat{T}e^{i2\pi\frac{m}{K}}|M_i\rangle = \hat{O}^me^{i\frac{2\pi(m+p_i)}{K}}|M_i\rangle \quad (27)$$

$$\hat{T}\left(\hat{O}^m|M_i\rangle\right) = e^{i\frac{2\pi(m+p_i)}{K}}\left(\hat{O}^m|M_i\rangle\right). \quad (28)$$

As the twist operator \hat{O} commutes with the Hamiltonian, we can write

$$\langle M_i|\hat{O}^m H \hat{O}^m|M_i\rangle = \langle M_i|H|M_i\rangle. \quad (29)$$

Thus, the energy eigenvalues of all the orthogonal states are equal, and they form the K fold degenerate ground state subspace. By the application of the twist operator \hat{O} , we can go from one degenerate ground state to another. We note that a similar exploration of the degenerate ground state manifold of spin-1/2 Heisenberg antiferromagnets on geometrically frustrated lattices via twist and translation operators has been conducted in Refs.[99, 100].

3.4. Local Mott liquid

In order to obtain the effective theory for the K zero modes, we will now use the URG method to decouple the impurity site from the star graph. The starting point is the star graph Hamiltonian

$$\begin{aligned} H &= \mathcal{J}\vec{S}_d \cdot \sum_i \vec{S}_i = \mathcal{J}S_d^z S^z + \frac{\mathcal{J}}{2}(S_d^+ S^- + S_d^- S^+), \\ H_D &= \mathcal{J}S_d^z S^z, \quad H_X = \frac{\mathcal{J}}{2}(S_d^+ S^- + S_d^- S^+). \end{aligned} \quad (30)$$

In order to remove the quantum fluctuations between the impurity spin and the rest, we perform one step of URG:

$$\Delta H = H_X(\hat{\omega} - H_D)^{-1}H_X \quad (31)$$

The total bath spin operator S^z is not a good quantum number for the zero mode ground state, and because there is no net S^z field, the ground state manifold has a vanishing expectation value of S^z , $\langle S^z \rangle = 0$. We use

this expectation value to replace the denominator of the above RG equation.

$$\beta_{\uparrow}(\mathcal{J}, \omega_{\uparrow}) = (\mathcal{J}^2 \Gamma_{\uparrow})/2, \quad \Gamma_{\uparrow} = (\omega_{\uparrow} - \mathcal{J}(S_d^z - 1))^{-1} \quad (32)$$

The effective Hamiltonian is therefore

$$H_{eff} = \frac{\beta_{\uparrow}(\mathcal{J}, \omega_{\uparrow})}{4} (S^+ S^- + S^- S^+) . \quad (33)$$

In terms of the electronics degree of freedom, this can be written as

$$\frac{\beta_{\uparrow}(\mathcal{J}, \omega_{\uparrow})}{4} \sum_{i \neq j} \sum_{\alpha_i, \beta_i} \sum_{\alpha_j, \beta_j} \left(\vec{\sigma}_{\alpha_i \beta_i} \vec{\sigma}_{\alpha_j \beta_j} c_{0\alpha_i}^{(i)\dagger} c_{0\beta_i}^{(i)} c_{0\alpha_j}^{(j)\dagger} c_{0\beta_j}^{(j)} + \text{h.c.} \right) . \quad (34)$$

The case $\frac{\beta_{\uparrow}(\mathcal{J}, \omega_{\uparrow})}{2} < 0$ leads to a ferromagnetic effective Hamiltonian. In this case, the ground state is realised for S being maximum and S^z being minimum. The effective Hamiltonian can be rewritten in this case as

$$H_{eff} = -|\beta_{\uparrow}(\mathcal{J}, \omega_{\uparrow})| \times (S^+ S^- + S^- S^+)/4 . \quad (35)$$

The complete set of commuting observables for this Hamiltonian contains H, S, S^z . In the ground state, S is maximum, and therefore $S = K/2$, and S^z can take $2S + 1 = K + 1$ values. Defining the dual operator $\hat{\phi}$ with the algebra $[\phi, \hat{S}^z] = i$, we get the twist operator $\hat{Q} = \exp(i\hat{\phi}\Phi/\Phi_0)$. Applying this operator, we get the twisted Hamiltonian

$$H(\Phi) = -\frac{|\beta_{\uparrow}(\mathcal{J}, \omega_{\uparrow})|}{2} \left[S^2 - \left(S^z - \frac{\Phi}{\Phi_0} \right)^2 \right] . \quad (36)$$

One can thus explore different S^z ground states via this spectral flow (flux insertion) argument.

For a K channel star graph problem the ground state is K fold degenerate associated with different J^z values $\{-(K-1)/2, -(K-3)/2, \dots, (K-1)/2\}$. After removing the quantum fluctuations between the impurity spin and the outer spins, we get an all-to-all model (eq.(36)) as the effective Hamiltonian. The eigenstates of this all-to-all Hamiltonian can be labeled by the eigenvalues of S^z . There are $K + 1$ such states made out of only the outer spins. The total state including the impurity spin can be written as $|J^z\rangle = |S_d^z\rangle \otimes |S^z\rangle$ labeled by $J^z = S_d^z + S^z$. As the all-to-all model has Z_2 symmetry in impurity sector, there are $2(K+1)$ total states. For $S_d^z = 1/2$, $\{J^z\} = \{-(K-1)/2, \dots, (K-1)/2, (K+1)/2\}$ and for $S_d^z = -1/2$, $\{J^z\} = \{-(K+1)/2, -(K-1)/2, \dots, (K-1)/2\}$. We can see that in both the cases, all K states of the star graph ground state manifold are present in the spectrum of the all-to-all model, one of them being the ground state. Using the twist operator, we can cycle between the various states. Some additional topological features of the Mott liquid are presented in the Supplementary Materials [74].

4. Local non-Fermi liquid excitations of the 2CK model

4.1. Effective Hamiltonian

We will now proceed to extract the nature of the low-energy excitations of the two-channel Kondo problem. This will be done by treating real-space hopping as a perturbation to the zero-bandwidth Hamiltonian, as it accounts for the lowest-energy electronic quantum fluctuations in the conduction bath [11]. The zero-bandwidth Hamiltonian obtained from the URG of the two-channel Kondo problem then acts as the zeroth-level Hamiltonian

$$H_{2CK}^{(0)} = \mathcal{J} \vec{S}_d \cdot (\vec{S}_1 + \vec{S}_2) , \quad \vec{S}_i = \frac{1}{2} c_{i,\alpha}^\dagger \vec{\sigma}_{\alpha,\beta} c_{i,\beta} . \quad (37)$$

Here, $\vec{S}_i = 1, 2$ represents the spin degree of freedom present at the origin of the i^{th} channel. This zeroth-level Hamiltonian has two degenerate ground states labeled by $|J^z = \pm 1/2\rangle$ with energy $-\mathcal{J}$ and 6 excited states [101]. The perturbation Hamiltonian is the real-space hopping:

$$H_X = -t \sum_{\langle 1, l_1 \rangle \langle 2, l_2 \rangle} (c_{1,\sigma}^\dagger c_{l_1,\sigma} + c_{2,\sigma}^\dagger c_{l_2,\sigma} + \text{h.c.}) , \quad (38)$$

and its effects will now be accounted for using degenerate perturbation theory. Here l_i represents the nearest site to the origin of the i^{th} channel.

Since the perturbation is a single-particle hopping that transfers electrons, we need to rewrite the spin operators in the zeroth Hamiltonian in terms of fermionic operators:

$$\mathcal{H} = \frac{\mathcal{J}\hbar}{2} \vec{S}_d \cdot \sum_{i=\{1,2\}} \sum_{\alpha,\beta \in \{\uparrow,\downarrow\}} c_{i\alpha}^\dagger \vec{\sigma}_{\alpha\beta} c_{i\beta} + H_X . \quad (39)$$

In this expanded basis, there are 32 degenerate ground states:

$$\{|\alpha_i\rangle\} = \{|J^z\rangle \otimes |n_{l_1,\uparrow}, n_{l_1,\downarrow}, n_{l_2,\uparrow}, n_{l_2,\downarrow}\rangle\} . \quad (40)$$

The 32-fold degeneracy arises from the two-fold degeneracy of J^z ($J^z = \pm 1/2$) multiplying the 16-fold degeneracy of the rest of the sites $n_{l_i,\sigma}$ (each of the four $n_{l_i,\sigma}$ can be 0 or 1, leading to a 2^4 -fold degeneracy).

The first and the second order corrections to the low energy effective Hamiltonian are

$$H^{(1)} = \sum_{ij} |\alpha_i\rangle \langle \alpha_i | V | \alpha_j \rangle \langle \alpha_j | .$$

$$H^{(2)} = \sum_{ij} \sum_l |\alpha_i\rangle \frac{\langle \alpha_i | V | \mu_l \rangle \langle \mu_l | V | \alpha_j \rangle}{E_0 - E_l} \langle \alpha_j | . \quad (41)$$

Here, $|\alpha_i\rangle$ represents the ground states with energy E_0 and is the set of states defined in eq. 40, and μ_l represents the excited states with energy E_l . It is easy to see that the diagonal contribution at any odd order is zero - the final state is never equal to the initial state. The off-diagonal part at first order is also zero. At second order we get both diagonal and off-diagonal contributions to the effective Hamiltonian.

4.1.1. Diagonal renormalisation We first calculate the diagonal renormalisation at second order coming from each of the states $|\tilde{\alpha}_0\rangle = |J^z = \frac{1}{2}\rangle \otimes |\psi_{\text{rest}}\rangle$, $|\tilde{\alpha}_1\rangle = |J^z = -\frac{1}{2}\rangle \otimes |\psi_{\text{rest}}\rangle$, where $|\psi_{\text{rest}}\rangle$ refers to the configuration of the remaining sites apart from the two that form the zeroth Hamiltonian. These second order corrections are given by

$$H_{\text{diag}}^{(2)}(\tilde{\alpha}_0) = \frac{2t^2}{E_0} \hat{I} - \Omega_l, \quad H_{\text{diag}}^{(2)}(\tilde{\alpha}_1) = \frac{2t^2}{E_0} \hat{I} + \Omega_l, \quad (42)$$

where $S_{l_i}^z = (n_{l_i\uparrow} - n_{l_i\downarrow})/2$ and $\Omega_l = \frac{2t^2}{3E_0}(S_{l_1}^z + S_{l_2}^z)$. The total diagonal second-order contribution to the effective Hamiltonian is obtained by adding these two contributions, and the result is a trivial shift: $H_{\text{diag}}^{(2)} = H_{\text{diag}}^{(2)}(\tilde{\alpha}_0) + H_{\text{diag}}^{(2)}(\tilde{\alpha}_1) = -4t^2 \hat{I}$. The fact there is no non-trivial diagonal renormalisation to the low-energy excitations shows the absence of (local) Fermi liquid terms.

4.1.2. Off-diagonal renormalisation The off-diagonal contribution to the second order effective Hamiltonian is obtained purely from $H_{\text{od}}^{(2)}$:

$$\begin{aligned} H_{\text{NFL}} &= \sum_{i \neq j} \sum_l |\alpha_i\rangle \frac{\langle \alpha_i | V | \mu_l \rangle \langle \mu_l | V | \alpha_j \rangle}{E_0 - E_l} \langle \alpha_j | \\ &= -\frac{8t^2}{3} [(S_1^z)^2 c_{2\uparrow}^\dagger c_{2\downarrow} (c_{l_1\uparrow}^\dagger c_{l_1\downarrow}^\dagger + c_{l_2\uparrow}^\dagger c_{l_2\downarrow}^\dagger) \\ &\quad + (S_2^z)^2 c_{1\uparrow}^\dagger c_{1\downarrow} (c_{l_1\uparrow}^\dagger c_{l_1\downarrow}^\dagger + c_{l_2\uparrow}^\dagger c_{l_2\downarrow}^\dagger)] + \text{h.c.} \quad (43) \end{aligned}$$

This effective Hamiltonian is purely of the non-Fermi liquid kind, arising due to the degeneracy of the ground state manifold. Using this low energy effective Hamiltonian, we have calculated different thermodynamic quantities like susceptibility, specific heat and the Wilson ratio. The first two measures show logarithmic behavior at low temperatures, in agreement with known results in the literature [14, 90, 35, 77, 80, 91, 33, 92, 93, 94, 30, 49, 9, 95, 96, 97, 98]. These results are shown in the Supplementary Material [74].

Following this method, we have also calculated the effective Hamiltonian for the three channel Kondo model. We again find the absence of all Fermi liquid terms up to 2^{nd} order, and the presence of non-Fermi liquid terms in the off-diagonal part of the effective Hamiltonian, and these are also shown in the Supplementary Material [74].

4.2. Local marginal Fermi liquid and orthogonality catastrophe

The real space local low energy Hamiltonian that takes into account the excitations above the ground state is

given by eq. 43 and can be written as

$$V_{\text{eff}} = \frac{2t^2}{\mathcal{J}^*} \left[(\sigma_{0,1}^z)^2 s_{0,2}^+ + (\sigma_{0,2}^z)^2 s_{0,1}^+ \right] (s_{1,1}^- + s_{1,2}^-) + \text{h.c.}; \quad (44)$$

here, $\sigma_{0,l}^z = \hat{n}_{0\uparrow,l} - \hat{n}_{0\downarrow,l}$, $s^+ = c_{0\uparrow,l}^\dagger c_{0\downarrow,l}$ and $s^- = (s^+)^\dagger$. The notation $0\sigma,l$ has the site index $i = 0, 1, 2, \dots$ as the first label, the spin index $\sigma = \uparrow, \downarrow$ as the second label and the channel index $l = 1, 2$ as the third label. Such non-Fermi liquid (NFL) terms in the effective Hamiltonian and the absence of any Fermi-liquid term at the same order should be contrasted with the local Fermi liquid excitations induced by the singlet ground state of the single-channel Kondo model [11, 86, 2]. We now take the MCK Hamiltonian to strong-coupling, and perform a perturbative treatment of the hopping. At $J \rightarrow \infty$, the perturbative coupling t^2/J is arbitrarily small and we again obtain Eq. 44. Such a change from the strong coupling model with parameter J to a weak coupling model with parameter t^2/J amounts to a duality transformation [88, 89]. It can be shown that the duality transformation leads to an identical MCK model [88] (self-duality), which implies we can have identical RG flows, and our transformation simply extracts the NFL piece from the dual model. The self-duality also ensures that the critical intermediate-coupling fixed point is unique and can be reached from either of the models.

The diagonal part of eq. 44 is

$$V_{\text{eff}} = \frac{2t^2}{J} \sum_{l=1,2} \left(\sum_{\sigma} \hat{n}_{0\sigma,l} \right) s_{0,\bar{l}}^+ s_{1,\bar{l}}^- + \text{h.c.}, \quad (45)$$

where $\bar{l} = 3 - l$ is the channel index complementary to l . We will Fourier transform this effective Hamiltonian into k -space. The NFL part becomes

$$\begin{aligned} &\frac{2t^2}{J} \sum_{\{k_i, k'_i\}} \sum_{\sigma, l} \left(e^{i(k_1 - k'_1)a} c_{k\sigma, l}^\dagger c_{k'\sigma, l} c_{k_2\uparrow, \bar{l}}^\dagger c_{k_2\downarrow, \bar{l}} c_{k_1\downarrow, \bar{l}}^\dagger c_{k'_1\uparrow, \bar{l}} \right. \\ &\quad \left. + \text{h.c.} \right). \quad (46) \end{aligned}$$

Such a three particle interaction term was also obtained for the NFL phase of the 2D Hubbard model from a URG treatment (see Appendix B of Ref.[70]). The channel indices in Eq. 46 can be mapped to the normal directions in [70]. The 2 particle-1 hole interaction in Eq. 46 has a diagonal component which can be obtained by setting $k = k'$, $k_1 = k'_2$ and $k_2 = k'_1$:

$$\begin{aligned} H_{\text{MFL}} &= \sum_{\substack{k, k_1, \\ k_2, \sigma, l}} \frac{2t^2 e^{i(k_1 - k_2)a}}{J} \hat{n}_{k\sigma, l} \hat{n}_{k_2\uparrow, \bar{l}} (1 - \hat{n}_{k_1\downarrow, \bar{l}}) + \text{h.c.} \\ &= \sum_{\substack{k, k_1, \\ k_2, \sigma, l}} \frac{4t^2}{J} \cos a (k_1 - k_2) \hat{n}_{k\sigma, l} \hat{n}_{k_2\uparrow, \bar{l}} (1 - \hat{n}_{k_1\downarrow, \bar{l}}). \quad (47) \end{aligned}$$

The most dominant contribution comes from $k_1 = k_2 = k'$, revealing the non-Fermi liquid metal [94, 45]:

$$H_{\text{MFL}}^* = \frac{4t^2}{J} \sum_{\sigma, k, k', l} \hat{n}_{k\sigma, l} \hat{n}_{k'\uparrow, \bar{l}} (1 - \hat{n}_{k'\downarrow, \bar{l}}) . \quad (48)$$

A non-local version of this effective Hamiltonian was found to describe the normal phase of the Mott insulator of the 2D Hubbard model, as seen from a URG analysis [70, 71]. Following [70], one can track the RG evolution of the dual coupling $R_j = \frac{4t^2}{J_j}$ at the j^{th} RG step, in the form of the URG equation

$$\Delta R_j = -\frac{R_j^2}{\omega - \epsilon_j/2 - R_j/8} . \quad (49)$$

In the RG equation, ϵ_j represents the energy of the j^{th} isoenergetic shell. It is seen from the RG equation that R is relevant in the range of $\omega < \frac{1}{2}\epsilon_j$ that has been used throughout, leading to a fixed-point at $R^*/8 = \omega - \frac{1}{2}\epsilon^*$. The relevance of R is expected because the strong coupling J is irrelevant and $R \sim 1/J$.

The renormalisation in R leads to a renormalisation in the single-particle self-energy [70]. The k -space-averaged self-energy renormalisation is

$$\Delta\Sigma(\omega) = \rho R^{*2} \int_0^{\epsilon^*} \frac{d\epsilon_j}{\omega - \epsilon_j/2 + R_j/8} . \quad (50)$$

The density of states can be approximated to be N^*/R^* , where N^* is the total number of states over the interval R^* . As suggested by the fixed point value of R_j , we can approximate its behaviour near the fixed point by a linear dependence on the dispersion ϵ_j . The two limits of the integration are the starting and ending points of the RG. We start the RG very close to the Fermi surface and move towards the fixed point ϵ^* . Near the starting point, we substitute $\epsilon = 0$ and $R = \omega$, following the fixed point condition. From the fixed point condition, we also substitute $R^*/8 = \omega - \frac{1}{2}\epsilon^*$. On defining $\bar{\omega} = N^*(\omega - \frac{1}{2}\epsilon^*)$, we can write

$$\Delta\Sigma(\omega) \sim \bar{\omega} \ln \frac{N^*\omega}{\bar{\omega}} . \quad (51)$$

The self-energy also provides the quasiparticle residue for each channel[70]:

$$Z(\bar{\omega}) = \left(2 - \ln \frac{2\bar{\omega}}{N^*\omega}\right)^{-1} . \quad (52)$$

As $\omega \rightarrow 0$, the Z vanishes, implying that the ground state is *not adiabatically connected* to the Fermi gas in the presence of the NFL terms. This is the orthogonality catastrophe [102, 103, 104, 105] in the two-channel Kondo problem, and it is brought about by the presence of the channel-non diagonal terms in Eq. 48. Such terms were absent in the single-channel Kondo model, because there was no multiply-degenerate ground state manifold that allowed scattering. This line of argument shows that

the extra degeneracy of the ground state subspace and the frustration of the singlet order that comes about when one upgrades from the single-channel Kondo model to the MCK models is at the heart of the NFL behaviour, and the orthogonality catastrophe is expected to be a general feature of all such frustrated MCK models. A local NFL term and a similar self-energy was also obtained by Coleman, et al. [49] in terms of Majorana fermions at the strong-coupling fixed point of the $\sigma - \tau$ model, which they claimed was equivalent to the intermediate-coupling fixed point of the two-channel Kondo model. Indeed, along with Ref.[39], this demonstration shows the universality between the two-channel Kondo and the $\sigma - \tau$ models.

5. Entanglement properties

5.1. Entanglement properties of the star graph

We will now present the results of our study of various entanglement measures in each of the K degenerate ground states $|J^z\rangle$ of a K channel star graph, labeled by the eigenvalues of J^z .

5.1.1. Entanglement entropy between impurity and the rest In a 1-channel Kondo model, the ground state is unique and a singlet [86]: $|J, J_z = 0, 0\rangle = \frac{1}{\sqrt{2}}(|\uparrow_d, \downarrow_0\rangle - |\downarrow_d, \uparrow_0\rangle)$, and the impurity entanglement entropy (EE_d) is at the maximum possible value of $\log 2$. We will now calculate the same quantity for the ground states

$$|J^z\rangle \equiv |S_d = 1/2, S = K/2; J = (S - 1/2), J^z\rangle \quad (53)$$

of the K channel model.

In order to compute EE_d for a particular state $|J^z\rangle$, we calculate the von-Neumann entropy of the reduced density matrix ρ_d obtained by partially tracing the density matrix $\rho = |J^z\rangle\langle J^z|$ associated with the state $|J^z\rangle$ over the impurity states $|S_d^z = \pm \frac{1}{2}\rangle$:

$$\rho_d = \text{Tr}_d \rho = \sum_{S_d^z} \langle S_d^z | \rho | S_d^z \rangle , \quad EE_d = -\rho_d \log \rho_d . \quad (54)$$

The EE_d for states of various J^z are shown as functions of the number of channels K in fig. 6. At any given value of K on the x -axis, all the points directly above it represent values of EE_d for various values of J^z in the ground state manifold of the MCK problem defined by that particular value of K . The minimum entanglement entropy occurs in the maximum J^z state ($|J^z = J\rangle$), and this minimum value decreases with increase in K . On the other hand, the maximum entanglement entropy is attained in the state with minimum J^z . This maximum value of the entanglement entropy is always $\ln 2$ for odd values of K ; for even values of K , the value asymptotically reaches $\log 2$ at large K .




Figure 6. Variation of impurity entanglement entropy (EE_d) for multiple values of K . The maximum EE_d at large K is $\log 2$.

A very similar computation gives the entanglement entropy between one outer spin and the rest of the spins, and it is shown in Fig. 7. We again find that the minimum entanglement entropy is associated with the state $|J^z = J\rangle$ and it falls with increasing K , approaching zero asymptotically; the maximum entanglement entropy is, as before, associated with the state $|J^z\rangle_{\min}$ and it rises to $\log 2$ in the limit $K \gg 1$.




Figure 7. Variation of the entanglement entropy of an outer spin (EE_o) with the rest for different channels (K). The maximum entanglement entropy in the large K limit is $\log 2$.

Two interesting features emerge from the study:

- For the odd K models, the impurity is maximally entangled with the other spins in the minimum J^z , $|J^z = 0\rangle$, member of the ground state manifold. At large K , the even K models also acquire this property, the minimum J^z there being $\pm \frac{1}{2}$.
- Both EE_d and EE_o are individually equal for the states $|\pm J^z\rangle$, which shows that both types of entanglement entropy are invariant under the transformation $|J^z\rangle \rightarrow \pi |J^z\rangle$. This reflects a parity symmetry in the state space in terms of entanglement.




Figure 8. Variation of mutual information $I^2(d : o)$ of the impurity with the rest, for different values of K . The maximum $I^2(d : o)$ is 0.37.

5.1.2. Mutual Information Mutual information is a measure which captures the correlations present between two subsystems (A,B), in a particular state. It is defined as

$$I^2(A : B) = S_A + S_B - S_{A \cup B}, \quad (55)$$

where $S_{A(B)}$ is the entanglement entropy of the subsystem $A(B)$ with the rest, and $S_{A \cup B}$ is the entanglement entropy of A and B with the rest. We are interested in two types of mutual information: firstly, the mutual information $I^2(d : o)$ between the impurity and one of the other spins, and secondly, the mutual information $I^2(o : o)$ between any two of the outer spins. Both of these have been computed for various channel numbers K , and plotted in Figs. 8 and 9.




Figure 9. Variation of mutual information $I^2(o : o)$ between 2 outer spins, for multiple K . $I^2(o : o)$ vanishes for large K .

In both cases we find that the maximum and minimum mutual information are associated with the $|J^z\rangle_{\min}$ and $|J^z = J\rangle$ states respectively. We also find that the mutual information is the same in the states $|J^z\rangle$ and $\pi^x |J^z\rangle$, indicating the parity symmetry in the mutual information measure. In the large channel limit ($K \gg 1$), the maximum $I^2(d : o)$ and $I^2(o : o)$ saturate to the common value of 0.375.




Figure 10. Variation of multipartite information I^N among N outer spins as a function of N , for $K = 8$. The inset shows I^N vs N for an all-to-all model. Both show power law behavior.

5.1.3. Multipartite information Similar to mutual information, one can calculate higher order multipartite information to study the nature of correlations present in different ground states. We define the tripartite information among three subsystems A, B, C as $I_{ABC}^3 = (S_A + S_B + S_C) - (S_{AB} + S_{BC} + S_{CA}) + S_{ABC}$. I^3 shows behavior similar to the mutual information - the highest I^2 value is associated with the $(J^z)_{\min}$ state and it saturates in the limit of $K \gg 1$. The N -partite information for a collection of subsystems (CSS) $\{\mathcal{A}_N\} \equiv \{A_1, A_2, \dots, A_N\}$ can also be defined as [106]

$$I_{\{\mathcal{A}_N\}}^N = \left[\sum_{m=1}^N (-1)^{m-1} \sum_{Q \in \mathcal{B}_m(\{\mathcal{A}\})} S_{V_{\cup}(Q)} \right] - S_{V_{\cap}(\{\mathcal{A}_N\})} \quad (56)$$

where $\mathcal{B}_m(\{\mathcal{A}_N\}) \equiv \{Q \mid Q \subset \mathcal{P}(\{\mathcal{A}_N\}), |Q| = m\}$, and $\mathcal{P}(\{\mathcal{A}_N\})$ is the powerset of $\{\mathcal{A}_N\}$. For simplicity, we also define the union and intersection of all the subsystems present in Q as $V_{\cup}(Q) \equiv \bigcup_{A \in Q} A$ and $V_{\cap}(Q) \equiv \bigcap_{A \in Q} A$ respectively.

The study of such measures of multipartite information reveals the nature of the multi-party correlations present among the outer spins of the star graph. Our study, as presented in fig. (10), shows that the higher order multi-partite correlations (I^N) decrease as you increase the order (N), and the behaviour follows a power law. In the inset of the same figure, we have shown the I^N vs N plot that is observed in the ground state of the all-to-all effective Hamiltonian (Eq.36), and it shows a similar power-law behaviour. The similarities in the behavior suggest that one can capture entanglement properties either from the star graph or from the corresponding all-to-all model.

5.2. Entanglement properties at the fixed point of the MCK model




Figure 11. This is a schematic diagram of (a) single channel and (b) two channel problem. S_d is the impurity spin. For more details refer to the text.

To study the nature of correlations at the MCK fixed point, we introduce excitations into the star graph ground state. We will work with the Hamiltonian Eq.(39) where we consider nearest-neighbor real space hopping t from the zeroth site into the lattice (Fig. 11). Setting $t = 0$ recovers the zero bandwidth version of the MCK. In the $K = 1$ single channel case, the impurity (S_d) is coupled to the spin degree of freedom at the real space origin ($\{1_{\uparrow}, 1_{\downarrow}\}$) of the single conduction bath via a Heisenberg spin-exchange coupling. For $K = 2$, the impurity is interacting with 2 distinct local spins $\{1_{\uparrow}, 1_{\downarrow}\}$ and $\{2_{\uparrow}, 2_{\downarrow}\}$ belonging to zeroth sites of different conduction channels, and the real space hopping connects these zeroth sites to their nearest neighbor sites. As we increase the hopping strength t from zero to non-zero, the lattice sites start interacting with each other.

5.2.1. Impurity entanglement entropy The impurity entanglement entropy $EE_d(t)$ in the ground state is shown as a function of t in Fig. 12 for three values of channel number K . At low hopping strength ($0 < t \ll 1$), EE_d is independent of t and achieves a constant value. We also find that in this range of t , EE_d decreases with increasing K . This behaviour is reversed at high t , and EE_d is seen to increase with increasing K . Though the values of the impurity entanglement entropy for various values of K are quite similar at low hopping strength, as the single channel EE_d varies smoothly at $t \rightarrow 0^+$, whereas the EE_d for $K > 1$ show a discontinuity at $t \rightarrow 0^+$.

5.2.2. Intra-channel Mutual information Here, we will calculate the mutual information between two electronic states in the ground state wavefunction, as a function of the hopping strength t .

Case 1: We first study the mutual information $MI(d, 1 \uparrow)$ between the impurity spin S_d and the state 1_{\uparrow} (fig. 11), where 1 represents real-space origin of the 1st conduction channel. For a single channel




Figure 12. Variation of impurity entanglement entropy EE_{imp} against t , for $K = 1, 2, 3$.

model, this site is unique because there is just one channel, but in the presence of K channels, there are K possible choices corresponding to each of the channels. However, because of the symmetry of the Hamiltonian under the exchange of the channel indices, all such choices will show identical mutual information signatures. Similarly, due to the $SU(2)$ spin-rotation symmetry of the Hamiltonian, $MI(d, 1 \downarrow)$ will be identical to $MI(d, 1 \uparrow)$. We have numerically computed and plotted $MI(d, 1 \uparrow)$ as a function of t in fig. 13. The inset shows that at low hopping strength, the mutual information for the single channel and the two-channel models saturate to different values. In the single-channel case, the saturation value at $t = 0$ is $\log 2$, and the mutual information changes smoothly as t is turned on. This is similar to the behaviour in the impurity entanglement entropy studied previously. The value of $\log 2$ shows the maximal entanglement between the zeroth site and the impurity, and the perfect screening of the impurity spin. For the two channel problem, we find that the mutual information at $t = 0$ is $MI(d, 1 \uparrow) = 0.2401$. The reduction of the value from $\ln 2$ shows the breakdown of screening. Also note that unlike the single channel case, there is a discontinuity in $MI(d, 1 \uparrow)$ as t is increased from 0.

Case 2: We have also computed the mutual information $MI(1 \uparrow, 1 \downarrow)$ between the two electronic states on the zeroth site of the same conduction channel, and plotted them in fig. 14. We find that it is a smooth function of t for $K = 1$, and for $K = 2$ there is a discontinuity at $t = 0^+$. Note that for small values of t , we find $MI(1 \uparrow, 1 \downarrow) > MI(d, 1 \uparrow)$ for the $K = 2$ model, showing the presence of strong intra-site correlations.

Case 3: Next, we measure the mutual information $MI(d, l_{1 \uparrow})$ between the impurity site and an electronic state of the site that is nearest-neighbour to the real-space origin of one of the conduction channels (see




Figure 13. This figure show the variation of mutual information between the impurity spin and the $1 \uparrow$ state against t . For more details please refer to the text. The error-bar σ_2 of the two channel data at low hopping strength $\sigma_2 \approx 0.002$.




Figure 14. Variation of mutual information ($I^2(1 \uparrow, 1 \downarrow)$) among the two states present on the real-space origin (1) of a conduction channel, against t . The error-bar σ_2 for the two channel at low hopping strength is $\sigma_2 \approx 0.013$.

fig. 11). The results are shown in fig. 15. Vanishing mutual information for the single channel case shows the perfect screening of the impurity spin. The non-zero value of mutual information at small values of t in the $K = 2$ model is again a result of the imperfect screening. As in the previous cases, we find a discontinuity in the mutual information for $K = 2$ as $t \rightarrow 0^+$.

Case 4: A complementary study can be made by calculating the mutual information between the origin of a particular conduction channel and its nearest neighbour site. They are plotted in fig. 16, and we find that at low hopping strength the single channel mutual information vanishes, showing the decoupling of those two states. On the other hand, in the two channel case, there is a non-zero mutual information which indicates the presence of scattering between the local Mott liquid and the local non-Fermi liquid states.




Figure 15. Variation of the mutual information $MI(d, l_{1\uparrow})$ between the impurity spin and the $l_{1\uparrow}$ state, against t . The error-bar σ_2 for the two channel case is $\sigma_2 \approx 0.001$.




Figure 16. Variation of the mutual information $MI(1_\sigma, l_{1\sigma})$ between two nearest neighbor sites against t . The error-bar for the two channel case is $\sigma_2 \approx 0.005$.

Apart from these, we have also computed an intra-channel tripartite information $I^3(1_\uparrow, l_{1\uparrow}, l_{1\downarrow})$ of the two-channel ground state (blue curve in Fig. 17) - it shows that there is a discontinuity in the tripartite information at $t = 0^+$, and at low hopping strength the tripartite information is independent of t with a value 0.084. This non-zero value of the tripartite information is consistent with the presence of a non-Fermi liquid effective Hamiltonian having more than just two-particle interactions within it. We have also studied inter-channel mutual information $I^2(1_\uparrow, 2_\uparrow)$ for $K = 2$ (red curve in fig. 17). It reveals the correlation and quantum entanglement between these two channels, and demonstrates the all-to-all nature of the local Mott liquid.

5.2.3. Bures distance and the orthogonality catastrophe

In addition to these entanglement measures, we have also calculated the Bures distance between the states at $t = 0$ and $t > 0$. The Bures distance [107, 108, 109,




Figure 17. Red: Mutual information $I^2(1_\uparrow, 2_\uparrow)$ against t . Blue: Tripartite information $I^3(1_\uparrow, l_{1\uparrow}, l_{1\downarrow})$ against t . The error-bar (σ) for the I^3 and I^2 near the low hopping strength are 0.002 and 0.003 respectively.




Figure 18. Variation of Bures distance between the reduced density matrices of the star graph ground states at $t = 0$ and $t \neq 0$, against t . The error-bars (σ) for $K = 2$ and $K = 3$ are 0.14 and 0.08 respectively, and near $t \approx 1$ the σ are 0.04 and 0.07 respectively.

110, 111, 112] between two density matrices ρ_1, ρ_2 is defined as

$$d_{Bures}(\rho_1, \rho_2) = \sqrt{2} \left[1 - \text{Tr} \left\{ \left(\rho_1^{1/2} \rho_2 \rho_1^{1/2} \right)^{1/2} \right\} \right], \quad (57)$$

and is a measure of the distance, in the Hilbert space, between the states forming the density matrices. One can see from the above definition that the maximum and minimum possible distances are $\sqrt{2}$ and 0 respectively.

We get the ground state $|\psi(t)\rangle$ of the problem by solving the Hamiltonian in Eq. (39). This ground state $|\psi(t)\rangle(t)$ and the associated density matrix $\rho(t) = |\psi(t)\rangle\langle\psi(t)|$ are functions of the hopping strength t . We are interested in the reduced density matrix $\rho_{sg}(t)$ of only the sites that form the star graph (impurity site and the zeroth sites of the conduction channels), so we trace out all the nearest neighbor sites from

the full density matrix: $\rho_{sg}(t) = \text{Tr}_{n-n}(\rho(t))$. The Bures distance $d_{\text{Bures}}(\rho_{sg}(0), \rho_{sg}(t))$ is then calculated between the reduced density matrices at $t = 0$ and $t > 0$, for $K = 1$, $K = 2$ and $K = 3$.

We have shown the variation of the Bures distance as a function of the hopping strength t in Fig. 18. Although the Bures distance is 0 (by definition) at $t = 0$ for all K , a difference arises as t is made slightly non-zero: there is a discontinuity in the Bures distance for the two and three channel models, while the single channel Bures distance is continuous. The smooth variation of the Bures distance in the single channel case shows the adiabatic continuity of the reduced star graph density matrix from the $t = 0$ case to $t \neq 0$ case. On the other hand, the abrupt jump in the Bures distance for higher-channel cases at $t = 0^+$ shows the breakdown of adiabatic continuity between the $t = 0$ and $t = 0^+$ Hamiltonians and signals an *orthogonality catastrophe* between these two ground states. The discontinuity increases as K is increased, indicating that considering the case of more channels will lead to the maximum Bures distance of $d_{\text{Bures}} = \sqrt{2}$ and hence an exact orthogonality.

6. Duality properties of the MCK model

We start from a strong coupling ($J \rightarrow \infty$) spin- S_d impurity MCK Hamiltonian in the over-screened regime ($K > 2S_d$),

$$H(J) = \sum_{k,\sigma,l} \epsilon_{k,l} \hat{n}_{k\sigma,l} + J \vec{S}_d \cdot \vec{S}. \quad (58)$$

Here, \vec{S} is the total spin $\sum_l \sum_{kk'\alpha\beta} \vec{\sigma}_{\alpha\beta} c_{k\alpha,l}^\dagger c_{k'\beta,l}$ of all the zero modes. At strong coupling, the ground states of the star graph eq.(8) act as a good starting point for a perturbative expansion. As argued previously, there are $K - 2S_d + 1$ ground states, labeled by the K values of the total spin angular momentum $J^z = S_d^z + S^z = -\frac{K}{2} + S_d, -\frac{K}{2} + S_d + 1, \dots, \frac{K}{2} - S_d$. Since a general spin- s object is simply a $2s + 1$ level system, the K -fold degenerate ground state manifold can be used to define a new impurity spin \mathbb{S}_d of multiplicity $2\mathbb{S}_d + 1 = K - 2S_d + 1$ which implies that we need $\mathbb{S}_d = \frac{K}{2} - S_d$. In other words, the spin- S_d impurity has a dual described by a spin- $(K - 2S_d + 1)$ impurity. The states of this new spin are defined by

$$\hat{S}_d^z |S^z\rangle = S^z |S^z\rangle, \\ \hat{S}_d^\pm |S^z\rangle = \sqrt{\mathbb{S}_d(\mathbb{S}_d + 1) - S^z(S^z \pm 1)} |S^z \pm 1\rangle. \quad (59)$$

While the ground state subspace gives rise to the new central spin object, the excited states of the star graph can be used to define bosonic operators that mediate interactions between the central spin and the next-nearest neighbour lattice sites [88]. In terms of the zero-bandwidth spectrum, the bosonic operators

represent scattering between the ground state subspace and the excited subspaces. Through a Schrieffer-Wolff transformation in the small coupling $\frac{t^2}{J}$, one can then remove this interaction and generate an exchange-coupling between the new impurity $\vec{\mathbb{S}}_d$ and the new zero modes formed out of the remaining sites in the lattice [88] (by remaining, we mean those real space sites that have not been consumed into forming the new spin). The new Hamiltonian, characterized by the small super-exchange coupling \mathbb{J} of the general form $\gamma t^2/J$, has the form

$$H'(\mathbb{J}) = \sum_{k,\sigma,l} \epsilon_{k,l} \hat{n}_{k\sigma,l} + \mathbb{J} \vec{\mathbb{S}}_d \cdot \vec{\mathbb{S}}. \quad (60)$$

$\vec{\mathbb{S}}$ is the local bath spin formed by the new zero modes. This Hamiltonian is very similar to the one in eq.(58), and that is the essence of the strong-weak duality: One can go from the over-screened strong coupling spin- S_d MCK model to another over-screened weak coupling spin- $(K - 2S_d + 1)$ MCK model. For the case of $K = 4S_d$, we have $\mathbb{S}_d = S_d$, and both S_d and \mathbb{S}_d describe the same objects: the two models are then said to be self-dual. For example, for the case of spin-1/2 MCK model, two-channel model is self-dual.

One important consequence of the duality relationship between the two over-screened models is that the RG equations are also dual; while the strong coupling model has an irrelevant coupling \mathcal{J} that flows down to the intermediate fixed point \mathcal{J}^* , the weak coupling model has a relevant coupling \mathbb{J} that flows up to the same fixed point $\mathbb{J}^* = \mathcal{J}^*$. From the RG equation for the general spin- S MCK model, we know that $\mathbb{J}^* = \frac{2}{K\rho'}$, where ρ' is the DOS for the bath of the weak coupling Hamiltonian. This constrains the form of the scaling factor γ :

$$\mathbb{J}^* = \frac{\gamma 4t^2}{\mathcal{J}^*} = \frac{2}{K\rho'} \implies \gamma = \frac{1}{4t^2} \mathcal{J}^{*2} = \frac{1}{K^2 t^2 \rho \rho'}. \quad (61)$$

Apart from the strong-weak duality, there exists another set of dual pairs in the MCK model. This was hinted at when we looked at the degree of compensation in eq.(13). Since Γ depends only on the magnitude of δ , both $\pm\delta$ will give the same degree of compensation, same ground state energy and same ground state degeneracy ($g_K^{S_d} = |\delta| + 1$). The definition of δ implies the following duality transformation: $K \rightarrow 2S_d, S_d \rightarrow \frac{K}{2}$. That is, we transform from a K -channel MCK model with spin- S_d impurity, to a $2S_d$ -channel MCK with a spin- $\frac{K}{2}$ impurity. The exactly-screened model $K = 2S_d$ maps on to itself and is therefore self-dual under this transformation.

For $K \neq 2S_d$, we transform an over-screened model into an under-screened model and vice versa. This duality relationship allows us to infer the RG scaling behaviour of one of the models if we know that




Figure 19. *Left:* Variation of the ground state degeneracy against the J_3 keeping $J_1 = J_2 = 1$ fixed. *Right:* Variation of ground state degeneracy against $J_1 = J_2$ keeping J_3 fixed. Both plots show the robustness of the ground state degeneracy against channel anisotropy.

of the other. Importantly, we see that the existence of an intermediate coupling fixed point in the MCK model for a given K and S_d leads to the conclusion that the $2S_d$ -channel spin- $\frac{K}{2}$ model has a strong coupling fixed point.

7. Quantum Phase transition in the MCK model under channel anisotropy

For a channel-anisotropic MCK model with K Kondo couplings $\{\mathcal{J}_i\}$ for each of the K conduction channels, the zero bandwidth model is

$$H_K(\vec{\mathcal{J}}) = \sum_{i=1}^K \mathcal{J}_i \vec{S}_d \cdot \vec{S}_i. \quad (62)$$

For the special case $\mathcal{J}_i = \mathcal{J} \forall i$, we get the usual isotropic star graph model. For the case of spin-1/2 impurity, we find that the Hamiltonian with any value of $\mathcal{J}_i > 0$ has K fold ground state degeneracy. This shows that the ground state degeneracy of the star graph model is extremely robust against the channel anisotropy; the ground state degeneracy does not change until at least one \mathcal{J}_i vanishes.

We have demonstrated this numerically for a three channel anisotropic star graph model. In Fig. 19(a), we show that if two couplings are kept constant and equal to one another ($\mathcal{J}_1 = \mathcal{J}_2 = 1$) and the third coupling \mathcal{J}_3 is tuned from some finite value to zero, the degeneracy does not change from 3 to 2 until \mathcal{J}_3 becomes zero. At that point, the model becomes a two channel Kondo problem. We also show in Fig. 19(b) that taking the coupling \mathcal{J}_3 fixed to 1 and varying the common coupling $\mathcal{J}_1 = \mathcal{J}_2$ from non-zero to zero is equivalent to keeping $\mathcal{J}_1 = \mathcal{J}_2 = 1$ fixed and taking \mathcal{J}_3 to infinity. In this case, when the coupling $\mathcal{J}_1 = \mathcal{J}_2 = 0$, the degeneracy becomes one and reveals the single channel nature of the problem.

The above demonstration makes it clear that the ground state degeneracy can change only when at least one of the Kondo couplings vanish. This can be realised

under RG flow if one considers the anisotropic MCK model.

$$H = \sum_{k,\alpha,l} \epsilon_{k,l} \hat{n}_{k\alpha,l} + \sum_{\substack{k,k',l \\ \alpha,\beta,l}} \mathcal{J}_l \vec{S}_d \cdot \frac{1}{2} \vec{\sigma}_{\alpha\beta} c_{k\alpha,l}^\dagger c_{k'\beta,l}. \quad (63)$$

We consider the specific case where $K-1$ channels have the same coupling $\mathcal{J}_1 = \mathcal{J}_2 = \dots = \mathcal{J}_{K-1} = \mathcal{J}_+$ and the remaining channel has a different coupling $\mathcal{J}_K = \mathcal{J}_-$. The RG equations for such a model are

$$\frac{\Delta \mathcal{J}_\pm}{|\Delta D|} = -\frac{\mathcal{J}_\pm^2 \rho}{\mathcal{D}_\pm} + \frac{\rho^2 \mathcal{J}_\pm}{2} \left[\frac{(K-1)\mathcal{J}_+^2}{\mathcal{D}_+} + \frac{\mathcal{J}_-^2}{\mathcal{D}_-} \right], \quad (64)$$

where $\mathcal{D}_\pm = \omega - \frac{D}{2} - \frac{\mathcal{J}_\pm}{4}$ are the denominators of the URG equations. Setting $\mathcal{J}_+ = \mathcal{J}_-$ leads to the critical fixed point at $\mathcal{J}_+^* = \mathcal{J}_-^* = \mathcal{J}_* = \frac{2}{K\rho}$. We now perturb around this fixed point by defining new variables $j_\pm = \mathcal{J}_\pm - \mathcal{J}_*$. We also assume that the bandwidth is large enough so that $\mathcal{D}_\pm \simeq \omega - \frac{D}{2} - \frac{\mathcal{J}_*}{4} = -|\mathcal{D}_*|$. Performing a linear stability analysis about $j_\pm = 0$ then reveals the following two possibilities:

- (a) If $j_- < 0$ and $j_+ > 0$, the deviation j_- is relevant and \mathcal{J}_- flows to zero. The flow of j_+ are constrained such that the remaining couplings \mathcal{J}_+ flow to the stable intermediate fixed point of the $K-1$ channel MCK model: $\mathcal{J}_{+,*} = \frac{2}{(K-1)\rho}$.
- (b) If $j_- > 0$ and $j_+ < 0$, the couplings \mathcal{J}_+ are irrelevant, and this leads to a single channel Kondo model described by the coupling \mathcal{J}_- which flows to strong coupling.

These conclusions show that the K channel intermediate fixed point is unstable under channel anisotropy [14, 45, 32, 37]. If one of the couplings becomes smaller than the rest, then that coupling flows to zero while the other $K-1$ couplings flow to the $K-1$ channel fixed point. On the other hand, if $K-1$ couplings are smaller than a single coupling, then the smaller couplings vanish while the remaining coupling flows to strong-coupling.

8. Conclusions

In summary, we have explored the low-energy behaviour of the MCK models using the unitary renormalization group (URG) method, and obtained insight into the role of ground state degeneracy and quantum-mechanical frustration in shaping the non-Fermi liquid physics and criticality. We have also obtained the zero-temperature phase diagram of the MCK problem and an effective Hamiltonian with a simple zero bandwidth limit (i.e., the star graph), and shown that it determines the ground state energy, wavefunction and degeneracy of the MCK models. The star graph is found to explain much of the physics of over-screening and criticality of the MCK:

this includes the singularities in the susceptibility and magnetisation, as well as a reduction in the degree to which the impurity spin is compensated. This demonstrates the ground state degeneracy of the star graph (which is unity for single-channel, and greater than 1 for the over-screened models) as the key ingredient in leading to the qualitative and quantitative differences of the MCK from the single-channel Kondo model. The presence of ground state degeneracy also allows the construction of non-local twist operators and fractional excitations that span the degenerate ground state manifold. Integrating out the quantum fluctuations of the impurity leads to an all-to-all effective Hamiltonian for the local conduction bath spins of the star graph. This Hamiltonian is found to contain inter-channel quantum fluctuations involving the scattering of electron-hole pairs, thereby creating a local Mott-liquid phase proximate to the impurity.

The low energy effective Hamiltonian (LEH) for the excitations of the MCK problem is then obtained by considering fluctuations of the conduction bath lying above the ground state. The LEH for the 2-channel Kondo confirms the absence of any local Fermi liquid physics, as well as the presence of a local marginal Fermi liquid arising from inter-channel off-diagonal scattering processes. This reinforces the idea that the ground state degeneracy is crucial to the appearance of non-Fermi liquid physics at the MCK fixed point. Further, we confirm that this arises from an orthogonality catastrophe in the form of singularities that are absent at exact screening. Studies of various thermodynamic properties (presented in the Supplementary Material [74]) from this 2-channel Kondo LEH (e.g., specific heat and susceptibility) show logarithmic dependence at low temperature, in agreement with the literature [30, 31, 32, 9, 33, 34, 35, 42, 43, 44, 45, 46].

The non-diagonal nature of the low energy theory is further investigated through several measures of entanglement. The entanglement entropy calculated from (i) the minimum J^z states, (ii) between the impurity and the outer spins as well as (iii) that between an outer spin and the rest, is observed to saturate to $\ln 2$ in the large channel limit. The opposite behaviour is observed in the maximum J^z state, where the entanglement entropy decreases with the increase of channel number. Moreover, the large values of inter-channel mutual information indicate high inter-channel correlation in the MCK ground state. Indeed, the power-law dependence of the multipartite information among the outer spins of the star graph model is similar to the power-law behavior of the all-to-all local Mott-liquid state. Importantly, the discontinuous behaviour in the entanglement entropy computed for the MCK ground state in the limit of

vanishing bath dispersion, as compared to the smooth behaviour observed in its single-channel counterpart, points to the orthogonality catastrophe in the low-energy phase of the multi-channel models. We confirm this by noting a similar discontinuous behaviour in the Bures distance computed between the density matrices computed from the ground states of the star graph (i.e., zero bath dispersion bandwidth) and the MCK (i.e., a non-zero bath dispersion bandwidth). This discontinuity in the Bures distance is observed to grow as the number of channels is increased. Finally, the URG study of the channel anisotropic MCK shows the expected critical nature of the channel isotropic fixed point: under any anisotropic perturbations, the RG flows go towards either the single-channel model or the symmetric MCK with one less channel coupled to the impurity. This is complemented by the study of the star graph ground state degeneracy which shows that the degeneracy changes only if one or more couplings vanishes. Taken together, we conclude that in the presence of channel anisotropy, the degeneracy of the MCK ground state does not change until the RG reaches the stable fixed point with one (or more channels) completely disconnected.

We end by discussing some open directions. Our work presents a template for the study of how a degenerate ground state manifold arising from symmetry and duality properties in a multichannel quantum impurity model can lead to novel multicritical phases at intermediate coupling. Employed within an auxiliary model framework (such as dynamical mean-field theory), such quantum impurity models will likely unveil novel quantum critical points (or phases) in bulk lattice models of strongly correlated electrons and spins that arise from the competition of various quantum (or symmetry broken) orders. Since the degeneracy of the star graph is observed to be robust under anisotropy, it appears relevant to study the case of the MCK with superconducting leads, i.e., conduction baths with a superconducting gap. Specifically, it will be interesting to see whether the gap can separate the ground state manifold of the star graph from any low-energy excitations, providing thereby a protection for the topological quantum numbers associated with the degenerate ground state manifold. The duality transformations then allow us to view the protected ground state as that belonging to an isolated larger quantum spin. This holds potential for applications in quantum information and quantum computation (see [113] for an alternate proposal based on an MCK model). Interesting behaviour may also be obtained by studying the MCK model with an easy-axis anisotropy term $(J^i)^2$ along one of the axes (i). Such a term would, if it is able to survive the RG flows towards low energy, lift the degeneracy

of the ground state and make it possible to achieve perfect screening even with $K > 2S_d$. Further, understanding the nature of the multicriticality in the MCK model displayed upon introducing channel anisotropy is in itself an important problem (see [114] for a recent work). One can also study the multi-channel lattice models [115, 116] to better understand the role played by singlet frustration and ground state degeneracy in the competition between the local moment versus the heavy fermion sea. Finally, we note that experimental realisations of the star graph have been studied recently with regards to aspects of NMR-based quantum information processing such as algorithmic cooling and information scrambling (see [117] for a review). It will be interesting to see whether some of the results obtained by us for the spin-1/2 Heisenberg star graph can be tested experimentally in a similar manner. Finally, it is tempting to speculate whether a similar study can be conducted for a MCK system of 1D Tomonaga-Luttinger liquid conduction leads [118].

Acknowledgments

S. Patra and Anirban Mukherjee thank the CSIR, Govt. of India and IISER Kolkata for funding through a research fellowship. Abhirup Mukherjee thanks IISER Kolkata for funding through a junior and a senior research fellowship. S. Lal thanks the SERB, Govt. of India for funding through MATRICS grant MTR/2021/000141 and Core Research Grant CRG/2021/000852. We thank three anonymous referees for their valuable questions and comments that greatly helped improve the presentation of the work.

References

- [1] Jun Kondo. Resistance minimum in dilute magnetic alloys. *Progress of theoretical physics*, 32(1):37–49, 1964.
- [2] A. C. Hewson. *The Kondo Problem to Heavy Fermions*. Cambridge University Press, 1993.
- [3] Philip W Anderson and Gideon Yuval. Exact results in the kondo problem: equivalence to a classical one-dimensional coulomb gas. *Physical Review Letters*, 23(2):89, 1969.
- [4] Philip W Anderson, G Yuval, and DR Hamann. Exact results in the kondo problem. ii. scaling theory, qualitatively correct solution, and some new results on one-dimensional classical statistical models. *Physical Review B*, 1(11):4464, 1970.
- [5] P W Anderson. A poor man's derivation of scaling laws for the kondo problem. *Journal of Physics C: Solid State Physics*, 3(12):2436–2441, dec 1970.
- [6] Kenneth G. Wilson. The renormalization group: Critical phenomena and the kondo problem. *Rev. Mod. Phys.*, 47:773–840, Oct 1975.
- [7] N. Andrei, K. Furuya, and J. H. Lowenstein. Solution of the kondo problem. *Rev. Mod. Phys.*, 55:331–402, Apr 1983.
- [8] A M Tsvelick and P B Wiegmann. Exact results in the theory of magnetic alloys. *Adv. in Phys.*, 32:453, 1983.
- [9] Ian Affleck and Andreas WW Ludwig. Exact conformal-field-theory results on the multichannel kondo effect: Single-fermion green's function, self-energy, and resistivity. *Physical Review B*, 48(10):7297, 1993.
- [10] Ian Affleck. Conformal field theory approach to the kondo effect. *Acta Phys.Polon.B*, 26(1869), 1995.
- [11] P Nozières. A “fermi-liquid” description of the kondo problem at low temperatures. *Journal of Low Temperature Physics*, 17:31, 1974.
- [12] A. C. Hewson. Renormalization group and fermi liquid theory. *Advances in Physics*, 43, 1994.
- [13] Y.-H. Zhang, Steffen Kahle, Tobias Herden, Christophe Stroh, Marcel Mayor, Uta Schlickum, Markus Ternes, Peter Wahl, and Klaus Kern. Temperature and magnetic field dependence of a kondo system in the weak coupling regime. *Nature Communications*, 4(1):2110, July 2013.
- [14] Nozières, Ph. and Blandin, A. Kondo effect in real metals. *J. Phys. France*, 41(3):193–211, 1980.
- [15] Ian Affleck and Andreas W. W. Ludwig. Exact critical theory of the two-impurity kondo model. *Phys. Rev. Lett.*, 68:1046–1049, Feb 1992.
- [16] Ian Affleck, Andreas W. W. Ludwig, and Barbara A. Jones. Conformal-field-theory approach to the two-impurity kondo problem: Comparison with numerical renormalization-group results. *Phys. Rev. B*, 52:9528–9546, Oct 1995.
- [17] Antoine Georges and Yigal Meir. Electronic correlations in transport through coupled quantum dots. *Phys. Rev. Lett.*, 82:3508–3511, Apr 1999.
- [18] Gergely Zaránd, Chung-Hou Chung, Pascal Simon, and Matthias Vojta. Quantum criticality in a double-quantum-dot system. *Phys. Rev. Lett.*, 97:166802, Oct 2006.
- [19] Andrew K. Mitchell, David E. Logan, and H. R. Krishnamurthy. Two-channel kondo physics in odd impurity chains. *Phys. Rev. B*, 84:035119, Jul 2011.
- [20] Andrew K. Mitchell, Eran Sela, and David E. Logan. Two-channel kondo physics in two-impurity kondo models. *Phys. Rev. Lett.*, 108:086405, Feb 2012.
- [21] Elio J. König, Piers Coleman, and Yashar Komijani. Frustrated kondo impurity triangle: A simple model of deconfinement. *Phys. Rev. B*, 104:115103, Sep 2021.
- [22] Piers Coleman. Mixed valence as an almost broken symmetry. *Phys. Rev. B*, 35:5072–5116, Apr 1987.
- [23] A. J. Millis and P. A. Lee. Large-orbital-degeneracy expansion for the lattice anderson model. *Phys. Rev. B*, 35:3394–3414, Mar 1987.
- [24] Assa Auerbach and K. Levin. Kondo bosons and the kondo lattice: Microscopic basis for the heavy fermi liquid. *Phys. Rev. Lett.*, 57:877–880, Aug 1986.
- [25] T. M. Rice and K. Ueda. Gutzwiller method for heavy electrons. *Phys. Rev. B*, 34:6420–6427, Nov 1986.
- [26] G. R. Stewart. Heavy-fermion systems. *Rev. Mod. Phys.*, 56:755–787, Oct 1984.
- [27] Mats Granath and Henrik Johannesson. Multichannel kondo effect in an interacting electron system: Exact results for the low-temperature thermodynamics. *Phys. Rev. B*, 57:987–992, Jan 1998.
- [28] A. Zawadowski. Kondo-like state in a simple model for metallic glasses. *Phys. Rev. Lett.*, 45:211–214, Jul 1980.
- [29] D. L. Cox and A. Zawadowski. Exotic kondo effects in metals: Magnetic ions in a crystalline electric field and tunnelling centres. *Advances in Physics*, 47(5):599–942, 1998.
- [30] Ian Affleck and Andreas W.W. Ludwig. Critical theory of overscreened kondo fixed points. *Nuclear Physics B*, 360(2):641–696, 1991.

- [31] Andreas W. W. Ludwig and Ian Affleck. Exact, asymptotic, three-dimensional, space- and time-dependent, green's functions in the multichannel kondo effect. *Phys. Rev. Lett.*, 67:3160–3163, Nov 1991.
- [32] Ian Affleck, Andreas W. W. Ludwig, H.-B. Pang, and D. L. Cox. Relevance of anisotropy in the multichannel kondo effect: Comparison of conformal field theory and numerical renormalization-group results. *Phys. Rev. B*, 45:7918–7935, Apr 1992.
- [33] Olivier Parcollet and Antoine Georges. Transition from overscreening to underscreening in the multichannel kondo model: Exact solution at large N . *Phys. Rev. Lett.*, 79:4665–4668, Dec 1997.
- [34] Ian Affleck. Non-fermi liquid behavior in kondo models. *Journal of the Physical Society of Japan*, 74(1):59–66, 2005.
- [35] V. J. Emery and S. Kivelson. Mapping of the two-channel kondo problem to a resonant-level model. *Phys. Rev. B*, 46:10812–10817, Nov 1992.
- [36] David G. Clarke, Thierry Giamarchi, and Boris I. Shraiman. Curie and non-curie behavior of impurity spins in quantum antiferromagnets. *Phys. Rev. B*, 48:7070–7076, Sep 1993.
- [37] Gergely Zaránd and Jan von Delft. Analytical calculation of the finite-size crossover spectrum of the anisotropic two-channel kondo model. *Phys. Rev. B*, 61:6918–6933, Mar 2000.
- [38] Jan von Delft, Gergely Zaránd, and Michele Fabrizio. Finite-size bosonization of 2-channel kondo model: A bridge between numerical renormalization group and conformal field theory. *Phys. Rev. Lett.*, 81:196–199, Jul 1998.
- [39] A. J. Schofield. Bosonization in the two-channel kondo model. *Phys. Rev. B*, 55:5627–5630, Mar 1997.
- [40] Ralf Bulla, Theo A. Costi, and Thomas Pruschke. Numerical renormalization group method for quantum impurity systems. *Rev. Mod. Phys.*, 80:395–450, Apr 2008.
- [41] H. B. Pang and D. L. Cox. Stability of the fixed point of the two-channel kondo hamiltonian. *Phys. Rev. B*, 44:9454–9457, Nov 1991.
- [42] N. Andrei and C. Destri. Solution of the multichannel kondo problem. *Phys. Rev. Lett.*, 52:364–367, Jan 1984.
- [43] A. M. Tselick and P. B. Wiegmann. Solution of then-channel kondo problem (scaling and integrability). *Zeitschrift für Physik B Condensed Matter*, 54(3):201–206, Sep 1984.
- [44] A M Tselick. The thermodynamics of multichannel kondo problem. *Journal of Physics C: Solid State Physics*, 18(1):159–170, jan 1985.
- [45] Natan Andrei and Andrés Jerez. Fermi- and non-fermi-liquid behavior in the anisotropic multichannel kondo model: Bethe ansatz solution. *Phys. Rev. Lett.*, 74:4507–4510, May 1995.
- [46] Gergely Zaránd, Theo Costi, Andres Jerez, and Natan Andrei. Thermodynamics of the anisotropic two-channel kondo problem. *Phys. Rev. B*, 65:134416, Mar 2002.
- [47] Anirvan M. Sengupta and Antoine Georges. Emery-kivelson solution of the two-channel kondo problem. *Phys. Rev. B*, 49:10020–10022, Apr 1994.
- [48] M. Fabrizio, Alexander O. Gogolin, and Ph. Nozières. Crossover from non-fermi-liquid to fermi-liquid behavior in the two channel kondo model with channel anisotropy. *Phys. Rev. Lett.*, 74:4503–4506, May 1995.
- [49] P. Coleman, L. B. Ioffe, and A. M. Tselick. Simple formulation of the two-channel kondo model. *Phys. Rev. B*, 52:6611–6627, Sep 1995.
- [50] M. Fabrizio, Alexander O. Gogolin, and Ph. Nozières. Anderson-yuval approach to the multichannel kondo problem. *Phys. Rev. B*, 51:16088–16097, Jun 1995.
- [51] A. V. Rozhkov. Impurity entropy for the two-channel kondo model. *International Journal of Modern Physics B*, 12(32):3457–3463, 1998.
- [52] Ru Zheng, Rong-Qiang He, and Zhong-Yi Lu. Order parameter for the multichannel kondo model at quantum criticality. *Phys. Rev. B*, 103:045111, Jan 2021.
- [53] K. Vladár and A. Zawadowski. Theory of the interaction between electrons and the two-level system in amorphous metals. i. noncommutative model hamiltonian and scaling of first order. *Phys. Rev. B*, 28:1564–1581, Aug 1983.
- [54] T. Cichorek, A. Sanchez, P. Gegenwart, F. Weickert, A. Wojakowski, Z. Henkie, G. Auffermann, S. Paschen, R. Kniep, and F. Steglich. Two-channel kondo effect in glasslike thasse. *Phys. Rev. Lett.*, 94:236603, Jun 2005.
- [55] D. C. Ralph and R. A. Buhrman. Observations of kondo scattering without magnetic impurities: A point contact study of two-level tunneling systems in metals. *Phys. Rev. Lett.*, 69:2118–2121, Oct 1992.
- [56] D. C. Ralph, A. W. W. Ludwig, Jan von Delft, and R. A. Buhrman. 2-channel kondo scaling in conductance signals from 2 level tunneling systems. *Phys. Rev. Lett.*, 72:1064–1067, Feb 1994.
- [57] Z. Iftikhar, S. Jezouin, A. Anthore, U. Gennser, F. D. Parmentier, A. Cavanna, and F. Pierre. Two-channel kondo effect and renormalization flow with macroscopic quantum charge states. *Nature*, 526(7572):233–236, Oct 2015.
- [58] L. J. Zhu, S. H. Nie, P. Xiong, P. Schlottmann, and J. H. Zhao. Orbital two-channel kondo effect in epitaxial ferromagnetic 110-mn films. *Nature Communications*, 7(1):10817, Feb 2016.
- [59] R. M. Potok, I. G. Rau, Hadas Shtrikman, Yuval Oreg, and D. Goldhaber-Gordon. Observation of the two-channel kondo effect. *Nature*, 446(7132):167–171, Mar 2007.
- [60] A. J. Keller, L. Peeters, C. P. Moca, I. Weymann, D. Mahalu, V. Umansky, G. Zaránd, and D. Goldhaber-Gordon. Universal fermi liquid crossover and quantum criticality in a mesoscopic system. *Nature*, 526(7572):237–240, Oct 2015.
- [61] VJ Emery and SA Kivelson. Importance of phase fluctuations in superconductors with small superfluid density. *Nature*, 374(6521):434, 1995.
- [62] H. T. Mebrahtu, I. V. Borzenets, H. Zheng, Y. V. Bomze, A. I. Smirnov, S. Florens, H. U. Baranger, and G. Finkelstein. Observation of majorana quantum critical behaviour in a resonant level coupled to a dissipative environment. *Nature Physics*, 9(11):732–737, Nov 2013.
- [63] Bedoor Alkurtass, Abolfazl Bayat, Ian Affleck, Sougato Bose, Henrik Johannesson, Pasquale Sodano, Erik S. Sørensen, and Karyn Le Hur. Entanglement structure of the two-channel kondo model. *Phys. Rev. B*, 93:081106, Feb 2016.
- [64] Donghoon Kim, Jeongmin Shim, and H.-S. Sim. Universal thermal entanglement of multichannel kondo effects. *Phys. Rev. Lett.*, 127:226801, Nov 2021.
- [65] Anirban Mukherjee and Siddhartha Lal. Unitary renormalisation group for correlated electrons-i: a tensor network approach. *Nuclear Physics B*, 960:115170, 2020.
- [66] Anirban Mukherjee and Siddhartha Lal. Unitary renormalisation group for correlated electrons-ii: insights on fermionic criticality. *Nuclear Physics B*, 960:115163, 2020.
- [67] Anirban Mukherjee, Abhirup Mukherjee, N. S. Vidhyadhiraja, A. Taraphder, and Siddhartha Lal. Unveiling the kondo cloud: Unitary renormalization-group study of the kondo model. *Phys. Rev. B*, 105:085119, 2022.

- [68] Santanu Pal, Anirban Mukherjee, and Siddhartha Lal. Correlated spin liquids in the quantum kagome antiferromagnet at finite field: a renormalization group analysis. *New J. Phys.*, 21(2):023019, feb 2019.
- [69] Anirban Mukherjee, Siddhartha Patra, and Siddhartha Lal. Fermionic criticality is shaped by fermi surface topology: a case study of the tomonaga-luttinger liquid. *Journal of High Energy Physics*, 04:148, 2021.
- [70] Anirban Mukherjee and Siddhartha Lal. Scaling theory for mott-hubbard transitions: I. $t = 0$ phase diagram of the $1/2$ -filled hubbard model. *New Journal of Physics*, 22(6):063007, jun 2020.
- [71] Anirban Mukherjee and Siddhartha Lal. Scaling theory for mott-hubbard transitions-II: quantum criticality of the doped mott insulator. *New Journal of Physics*, 22(6):063008, jun 2020.
- [72] Anirban Mukherjee and Siddhartha Lal. Superconductivity from repulsion in the doped 2d electronic hubbard model: an entanglement perspective. *J. Phys.: Condens. Matter*, 34:275601, 2022.
- [73] Siddhartha Patra and Siddhartha Lal. Origin of topological order in a cooper-pair insulator. *Phys. Rev. B*, 104:144514, Oct 2021.
- [74] See the supplementary materials available online for this work.
- [75] Franz Wegner. Flow-equations for hamiltonians. *Annalen der Physik*, 506(2):77–91, 1994.
- [76] Stanisław D. Głazek and Kenneth G. Wilson. Renormalization of hamiltonians. *Phys. Rev. D*, 48:5863–5872, Dec 1993.
- [77] J. Gan. On the multichannel kondo model. *J. Phys.: Condens. Matter*, 6(24):4547–4568, jun 1994.
- [78] Eugene Kogan. Poor man's scaling: anisotropic kondo and coqblin-schrieffer models. *Journal of Physics Communications*, 2(8):085001, aug 2018.
- [79] Y. Kuramoto. Perturbative renormalization of multi-channel kondo-type models. *The European Physical Journal B - Condensed Matter and Complex Systems*, 5:457–464, 1998.
- [80] A. M. Tsvelick and P. B. Wiegmann. Solution of then-channel kondo problem (scaling and integrability). *Zeitschrift für Physik B Condensed Matter*, 54(3):201–206, Sep 1984.
- [81] Andrew K. Mitchell, Martin R. Galpin, Samuel Wilson-Fletcher, David E. Logan, and Ralf Bulla. Generalized wilson chain for solving multichannel quantum impurity problems. *Phys. Rev. B*, 89:121105, Mar 2014.
- [82] V. Tripathi. *Landau Fermi Liquids and Beyond*. CRC Press, 2018.
- [83] Eduardo Fradkin. *Field theories of condensed matter physics*. Cambridge University Press, 2013.
- [84] C. M. Varma and Y. Yafet. Magnetic susceptibility of mixed-valence rare-earth compounds. *Phys. Rev. B*, 13:2950–2954, Apr 1976.
- [85] Kei Yosida. Bound state due to the $s - d$ exchange interaction. *Phys. Rev.*, 147:223–227, Jul 1966.
- [86] Kenneth G Wilson. The renormalization group: Critical phenomena and the kondo problem. *Reviews of Modern Physics*, 47(4):773, 1975.
- [87] Cătălin Pașcu Moca, Ireneusz Weymann, Miklós Antal Werner, and Gergely Zaránd. Kondo cloud in a superconductor. *Phys. Rev. Lett.*, 127:186804, Oct 2021.
- [88] Christian Kolf and Johann Kroha. Strong versus weak coupling duality and coupling dependence of the kondo temperature in the two-channel kondo model. *Phys. Rev. B*, 75:045129, Jan 2007.
- [89] Rok Žitko and Michele Fabrizio. Non-fermi-liquid behavior in quantum impurity models with superconducting channels. *Phys. Rev. B*, 95:085121, Feb 2017.
- [90] J. Gan, N. Andrei, and P. Coleman. Perturbative approach to the non-fermi-liquid fixed point of the overscreened kondo problem. *Phys. Rev. Lett.*, 70:686–689, Feb 1993.
- [91] A. M. Tsvelick and P. B. Wiegmann. Exact solution of the multichannel kondo problem, scaling, and integrability. *Journal of Statistical Physics*, 38(1):125–147, Jan 1985.
- [92] Taro Kimura and Sho Ozaki. Fermi/non-fermi mixing in $su(n)$ kondo effect. *Journal of the Physical Society of Japan*, 86(8):084703, 2017.
- [93] Damien Bensimon, Andrés Jerez, and Mireille Lavagna. Intermediate coupling fixed point study in the overscreened regime of generalized multichannel $SU(n)$ kondo models. *Phys. Rev. B*, 73:224445, Jun 2006.
- [94] D L Cox and M Jarrell. The two-channel kondo route to non-fermi-liquid metals. *J. Phys.: Condens. Matter*, 8(48):9825–9853, nov 1996.
- [95] P. Coleman and C. Pépin. Singular fermi liquid behavior in the underscreened kondo model. *Phys. Rev. B*, 68:220405, Dec 2003.
- [96] Nicolas Roch, Serge Florens, Theo A. Costi, Wolfgang Wernsdorfer, and Franck Balestro. Observation of the underscreened kondo effect in a molecular transistor. *Phys. Rev. Lett.*, 103:197202, Nov 2009.
- [97] Avraham Schiller and Lorenzo De Leo. Phase diagram of the anisotropic multichannel kondo hamiltonian revisited. *Phys. Rev. B*, 77:075114, Feb 2008.
- [98] P. Durganandini. Marginal fermi liquid behaviour in a coupled quantum dot system. *EPL (Europhysics Letters)*, 95(4):47003, jul 2011.
- [99] Santanu Pal and Siddhartha Lal. Magnetization plateaus of the quantum pyrochlore heisenberg antiferromagnet. *Physical Review B*, 100(10):104421, 2019.
- [100] Santanu Pal, Anirban Mukherjee, and Siddhartha Lal. Topological approach to quantum liquid ground states on geometrically frustrated heisenberg antiferromagnets. *Journal of Physics: Condensed Matter*, 32(16):165805, 2020.
- [101] Kunj Tandon, Siddhartha Lal, Swapan K. Pati, S. Ramasesha, and Diptiman Sen. Magnetization properties of some quantum spin ladders. *Phys. Rev. B*, 59:396–410, Jan 1999.
- [102] CM Varma, Z Nussinov, and Wim Van Saarloos. Singular or non-fermi liquids. *Physics Reports*, 361(5-6):267–417, 2002.
- [103] P. W. Anderson. Infrared catastrophe in fermi gases with local scattering potentials. *Phys. Rev. Lett.*, 18:1049–1051, Jun 1967.
- [104] Kōsaku Yamada and Kei Yosida. Orthogonality Catastrophe Due to Local Electron Interaction. *Progress of Theoretical Physics*, 59(4):1061–1071, 04 1978.
- [105] Kōsaku Yamada and Kei Yosida. Orthogonality catastrophe for a system of interacting electrons. ii. *Progress of Theoretical Physics*, 62(2):363–369, 1979.
- [106] Siddhartha Patra, Somnath Basu, and Siddhartha Lal. Unveiling topological order through multipartite entanglement. *Phys. Rev. A*, 105:052428, May 2022.
- [107] Armin Uhlmann. The “transition probability” in the state space of a*-algebra. *Reports on Mathematical Physics*, 9(2):273–279, 1976.
- [108] Donald Bures. An extension of kakutani’s theorem on infinite product measures to the tensor product of semifinite w^* -algebras. *Transactions of the American Mathematical Society*, 135:199–212, 1969.
- [109] Matthias Hübner. Explicit computation of the bures distance for density matrices. *Physics Letters A*, 163(4):239–242, 1992.
- [110] Matthias Hübner. Computation of uhlmann’s parallel transport for density matrices and the bures metric on three-dimensional hilbert space. *Physics Letters A*,

- 179(4-5):226–230, 1993.
- [111] Jochen Dittmann. Some properties of the riemannian bures metric on mixed states. *Journal of Geometry and Physics*, 13(2):203–206, 1994.
 - [112] Paulina Marian, Tudor A Marian, and Horia Scutaru. Bures distance as a measure of entanglement for two-mode squeezed thermal states. *Physical Review A*, 68(6):062309, 2003.
 - [113] Pedro L. S. Lopes, I. Affleck, and E. Sela. Anyons in multichannel kondo systems. *Phys. Rev. B*, 101:085141, Feb 2020.
 - [114] Ru Zheng, Rong-Qiang He, and Zhong-Yi Lu. Order parameter for the multichannel kondo model at quantum criticality. *Phys. Rev. B*, 103:045111, Jan 2021.
 - [115] C. A. Piguett, D. F. Wang, and C. Gruber. Exact solutions of the multichannel kondo-lattice model with infinite range hopping. *Journal of Low Temperature Physics*, 106(1):3–19, Jan 1997.
 - [116] Ming Shaw, Xing-wei Zhang, and Duo Jin. Scaling solution of the overscreened multichannel kondo model. *Phys. Rev. B*, 57:8381–8393, Apr 1998.
 - [117] T.S. Mahesh, D. Khurana, V.R. Krithika, G.J. Sreejith, and C.S. Sudheer Kumar. Star-topology registers: Nmr and quantum information perspectives. *J. Phys.: Condens. Matter*, page 383002, 2021.
 - [118] Siddhartha Lal, Sarang Gopalakrishnan, and Paul M. Goldbart. Approaching multichannel kondo physics using correlated bosons: Quantum phases and how to realize them. *Phys. Rev. B*, 81:245314, Jun 2010.