

A New Auxiliary Model Approach To Fermionic Criticality: Insights on Mottness in Strongly Correlated Systems

*A thesis submitted in partial fulfilment
for the requirements of the degree of*

Doctor of Philosophy

by

ABHIRUP MUKHERJEE
(18IP014)

under the supervision of

Prof. Siddhartha Lal

to the

Department of Physical Sciences,
Indian Institute of Science Education and Research Kolkata

April, 2026



Declaration

Certificate by supervisor

From You, 5 Years Ago

Acknowledgements

List Of Publications

As any dedicated reader can clearly see, the Ideal of practical reason is a representation of, as far as I know, the things in themselves; as I have shown elsewhere, the phenomena should only be used as a canon for our understanding. The paralogisms of practical reason are what first give rise to the architectonic of practical reason. As will easily be shown in the next section, reason would thereby be made to contradict, in view of these considerations, the Ideal of practical reason, yet the manifold depends on the phenomena. Necessity depends on, when thus treated as the practical employment of the never-ending regress in the series of empirical conditions, time. Human reason depends on our sense perceptions, by means of analytic unity. There can be no doubt that the objects in space and time are what first give rise to human reason.

Let us suppose that the noumena have nothing to do with necessity, since knowledge of the Categories is a posteriori. Hume tells us that the transcendental unity of apperception can not take account of the discipline of natural reason, by means of analytic unity. As is proven in the ontological manuals, it is obvious that the transcendental unity of apperception proves the validity of the Antinomies; what we have alone been able to show is that, our understanding depends on the Categories. It remains a mystery why the Ideal stands in need of reason. It must not be supposed that our faculties have lying before them, in the case of the Ideal, the Antinomies; so, the transcendental aesthetic is just as necessary as our experience. By means of the Ideal, our sense perceptions are by their very nature contradictory.

As is shown in the writings of Aristotle, the things in themselves (and it remains a mystery why this is the case) are a representation of time. Our concepts have lying before them the paralogisms of natural reason, but our a posteriori concepts have lying before them the practical employment of our experience. Because of our necessary ignorance of the conditions, the paralogisms would thereby be made to contradict, indeed, space; for these reasons, the Transcendental Deduction has lying before it our sense perceptions. (Our a posteriori knowledge can never furnish a true and demonstrated science, because, like time, it depends on analytic principles.) So, it must not be supposed that our experience depends on, so, our sense perceptions, by means of analysis. Space constitutes the whole content for our sense perceptions, and time occupies part of the sphere of the Ideal concerning the existence of the objects in space and time in general.

Abstract

This thesis reports (i) the development of a new auxiliary model method for studying translation-invariant models of strongly-interacting electrons, and (ii) an explicit demonstration of the holographic principle in the form of a simple and tractable exact holographic mapping (EHM) for a free fermion system with emphasis on the signatures at a critical Fermi surface. Together, these results shed light on how a system of electrons behaves as it undergoes a correlation-driven metal-insulator transition on a lattice.

The auxiliary model method allows us to create a mapping between a lattice model and an auxiliary impurity model in terms of the Hamiltonian, eigenstates, Greens functions and entanglement measures, among other things. This mapping ensures that such lattice model quantities can be computed indirectly from similar computations within the auxiliary impurity model. In terms of phenomenology, this mapping connects various renormalisation group (RG) fixed points of the impurity model to those of the lattice model. As examples, local Fermi liquid and local moment phases of the impurity site get mapped to Fermi liquid and Mott insulating phases, respectively, on the lattice model. The mapping makes use of translation operators and a manybody version of Bloch’s theorem to recreate the lattice model via repeated “tiling” operations on the impurity model. In contrast to other auxiliary model methods such as dynamical mean-field theory (DMFT) and its cluster variants (C-DMFT), this approach does not involve a self-consistency loop, making it more transparent. It also provides significantly more k -space resolution which is necessary for studying phenomena such as the pseudogap observed in the copper-oxide high- T_c superconductors.

A short description of the protocol for applying our method is as follows. We first identify the appropriate quantum impurity model corresponding to the lattice model at hand. For example, the impurity model must have a localisation-delocalisation transition on the impurity site in order to capture a metal-insulator transition on the lattice model. As another example, topological features of the band structure can be encoded by working with a topologically non-trivial conduction bath within the impurity model. Having decided upon a quantum impurity model, we solve the model using an impurity solver. In the works described in this thesis, we use the unitary renormalisation group because of its (i) analytical transparency, (ii) ability to deal with weak and strong correlations, (iii) ability to work at zero temperature, and (iv) reduced computational cost. Using the URG, we obtain a hierarchy of Hamiltonians that describe the high-energy and low-energy behaviour of the auxiliary model in various parameter regimes. This requires numerically solving the URG flow equations for various Hamiltonian couplings.

The phases of the impurity model are then characterised by (a) effective Hamiltonians, (b) static equal-time correlations, (c) dynamical correlations such as Greens functions and self-energy, and

(d) entanglement measures. Calculating some of these objects requires carrying out an iterative diagonalisation procedure on the hierarchy of Hamiltonians. Our tiling method then “periodises” these objects into the lattice model, generating required k –dependence in the process. Since the conduction bath of the impurity model already inherits the lattice geometry of the parent model, the appropriate symmetries are baked into the quantities. As an example, the lattice k –space Greens function G_k obtains contributions from the impurity site Greens function as well as impurity-bath off-diagonal Greens functions. While the former captures the local component of G_k , the latter captures the non-local components. With similar mappings existing for the dynamical correlations and entanglement measures, our method allows us to characterise various phases of the lattice model purely through impurity model computations.

Within our work on the holographic principle, we demonstrate the emergence of a holographic dimension in a system of 2D non-interacting Dirac fermions placed on a torus, by studying the scaling of multipartite entanglement measures under a sequence of renormalisation group (RG) transformations applied in momentum space. Geometric measures defined in this emergent space can be related to the RG beta function of the spectral gap, hence establishing a holographic connection between the spatial geometry of the emergent spatial dimension and the entanglement properties of the boundary quantum theory. We prove, analytically, that changing the boundedness of the holographic space involves a topological transition accompanied by a critical Fermi surface in the boundary theory. We go on to show that this results in the formation of a quantum wormhole geometry that connects the UV and the IR of the emergent dimension. The additional conformal symmetry at the transition also supports a relation between the emergent metric and the stress-energy tensor. In the presence of an Aharonov-Bohm flux, the entanglement gains a geometry-independent piece which is shown to be topological, sensitive to changes in boundary conditions, and related to the Luttinger volume of the system. Upon the insertion of a strong transverse magnetic field, we show that the Luttinger volume is linked to the Chern number of the occupied single-particle Landau levels.

Some Important Questions In this thesis, we have used the auxiliary model tiling method to study three fundamental problems in quantum condensed matter physics: (i) Mott metal-insulator transition on the Bethe lattice in infinite dimensions, (ii) Mott metal-insulator transition on the 2D square lattice, and (iii) quantum criticality in heavy fermion materials in 2D. Towards this, we have applied the method to two paradigmatic models: (a) an extended Hubbard model, and (ii) a bilayer extended Hubbard model. In preparation for applying the tiling approach, we have first analysed the corresponding quantum impurity models.

1. The Mott transition on the Bethe lattice is studied through an extended Anderson impurity model (eSIAM) that hosts local electronic correlation in the conduction bath proximate to the impurity site. Due to the lack of non-local correlations in this limit, it is sufficient to treat the conduction electron in the continuum limit.
2. The Mott transition in 2D is highly sensitive to details of the lattice. This requires us to study a lattice-embedded impurity model, where the impurity site is embedded in a 2D lattice of conduction electrons, and its hybridisation with the bath has the symmetries of the underlying lattice.

-
3. The heavy fermions problem necessitates a two layer impurity model, each layer acting as a lattice-embedded impurity model and coupled via inter-layer hybridisation.

The tiling approach then translates results from the auxiliary models into appropriate lattice models. In this thesis, we have attempted to answer a number of important open questions in the field:

- *Is there a minimal but effective quantum impurity model Hamiltonian that describes the Mott MIT of the $1/2$ -filled Hubbard model on the Bethe lattice in infinite dimensions?* Finding such an impurity model would also reduce the considerable computational effort that is presently required in self-consistent approaches. In the same context, is it possible to obtain a low-energy theory for the local gapless excitations precisely at the transition, where the metal is on the brink of destruction?
- For strongly-correlated electronic models, *is it possible to map the renormalisation group fixed points of such lattice models to those of quantum impurity models?* This would provide a firmer theoretical backing to various quantum impurity models, as well as reduce computational costs.
- The phenomenon of Mott insulation involves the localization of itinerant electrons due to strong local repulsion. Upon doping, a pseudogap (PG) phase emerges - marked by selective gapping of the Fermi surface without conventional symmetry breaking in spin or charge channels. *A key challenge is understanding how quasiparticle breakdown in the Fermi liquid gives rise to this enigmatic state, and how it connects to both the Mott insulating and superconducting phases.*

Questions on entanglement and holography become particularly interesting in systems of critical fermions due to the presence of (i) longer-ranged entanglement, (ii) topological features arising from the incompressibility of the Fermi volume and (iii) Fermi surface gapping phase transitions driven by changes in topological quantum numbers. Unlike topologically ordered phases, there is no geometry-independent topological term in the entanglement entropy of free fermionic systems. It is, however, possible to generate such a term even here by placing the system on a multiply connected manifold such as a cylinder, and inserting a magnetic flux through the cylinder. Several questions are therefore pertinent to this discussion:

- Not much is known about the holographic implications of a Fermi surface gapping transition. *What effect does a topological change in a Fermi surface have on the additional dimension generated for the system?* Can the transition be captured by topological invariants defined within the holographic space?
- *What happens to the topological signatures of entanglement entropy across a Fermi surface instability?* If the insulator on the other side has topological features, can we relate topological invariants on either side?

Summary Of The Chapters. For the readers' convenience, we now briefly describe the content of the various chapters.

The *Introduction* introduces the core themes of electronic correlation and Mott transition. After giving some preliminary insights from toy models, we introduce the other major theme – auxiliary models, and describe why quantum impurity models act as auxiliary models for models with strong local correlations. We go on to describe several classes of strongly correlated materials that are likely to be described by such methods, and discuss the phenomenology of some interacting models that will be investigated within the thesis. Near the end, we describe some of the features of the new auxiliary model developed within the thesis and list some of the questions addressed by it. We end by introducing the holographic principle and list some questions addressed by our work.

After the introduction, the *Methods and Preliminaries* chapter describes the various techniques and some preliminary ideas employed throughout the thesis. The chapter begins by deriving the unitary renormalisation group formalism and describes its various properties and a practical protocol for implementing it. That is followed by demonstrations of the unitary RG on three simple models - a star graph model of spins in a magnetic field, the two-channel Kondo model, and the problem of a particle in a periodic potential. Through these problems, various aspects of the method are clarified. In order to provide more insights, the chapter also details how the unitary RG method is related to other RG-based methods that rely on canonical transformations, such as Poor Man's scaling, Schrieffer-Wolff transformation and CUT (continuous unitary transformation) RG. Apart from the URG, the chapter also describes an iterative diagonalisation method that has been used within this thesis to compute various correlation functions and entanglement measures from the hierarchy of Hamiltonians obtained from the URG. Since various entanglement measures have been used throughout the later chapters, we have also added a section on various entanglement measures here - what they mean, what are their inter-relations, why they are important and how one can compute some of them. The chapter concludes by describing some of the other strategies that have been employed to reduce computational costs in various projects.

In Chapter 3, *Kondo Frustration via Charge Fluctuations: A Route to Mott Localisation in Infinite Dimensions*, we study our first impurity model, the extended Anderson model with local correlation U_b in the conduction bath. We demonstrate that this model hosts a local version of the Mott metal-insulator transition on the Bethe lattice in infinite dimensions, as seen from DMFT. At a critical value of the ratio of U_b and the impurity-bath hybridisation, the effective impurity model shows a transition from a Kondo screened phase into an unscreened local moment phase. For the case of attractive local bath correlations, the model sheds new light on several aspects of the DMFT phase diagram. For example, the $T = 0$ metal-to-insulator quantum phase transition (QPT) is preceded by an excited state QPT (ESQPT) where the local moment eigenstates are emergent in the low-lying spectrum. Long-ranged fluctuations are observed near both the QPT and ESQPT, suggesting that they are the origin of the quantum critical scaling observed recently at high temperatures in DMFT simulations. The $T = 0$ gapless excitations at the quantum critical point display particle-hole interconversion processes, and exhibit power-law behaviour in self-energies and two-particle correlations. These are signatures of non-Fermi liquid behaviour that emerge from the partial breakdown of the Kondo screening. A many-body perturbation theoretic treatment of the Hubbard sidebands reveals that they are comprised of holon-doublon excitations created by the hybridisation of the impurity site with the conduction bath. These excitations consist of decoupled local Fermi liquids for

the holons and doublons at the lowest order, which, at higher orders, become coupled via correlated holon-doublon scattering between impurity and bath.

In Chapter 4, *A New Auxiliary Model Approach for Interacting Electrons: Mapping Impurity Models To Lattice Models*, we lay out the details of the auxiliary model method developed to map impurity models to lattice models. We first motivate the general idea behind auxiliary model methods. This is followed by a construction of the manybody translation operators. We use this, along with a manybody Bloch's theorem, to construct lattice models from quantum impurity models and relate their eigenstates. This then allows us to map correlations functions, self-energies and entanglement measures across models. In particular, we demonstrate how the presence of the conduction bath introduces k -space resolution in various quantities.

In Chapter 5, *Mott Criticality as the Confinement Transition of a Pseudogap-Mott Metal: Insights on Mott Transition and Pseudogap in Two Dimensions*, we apply the auxiliary model tiling method to a lattice-embedded extended SIAM. Applying a many-body tiling scheme to the fixed-point impurity model uncovers a lattice model with electron interactions and Kondo physics. At half-filling, the interplay between Kondo screening and bath charge fluctuations in the impurity model leads to Fermi liquid breakdown. This reveals a pseudogap phase characterized by a non-Fermi liquid (the Mott metal) residing on nodal arcs, gapped antinodal regions of the Fermi surface (Luttinger surfaces), and an anomalous scaling of the electronic scattering rate with frequency. The eventual confinement of holon-doublon excitations of this exotic metal obtains a continuous transition into the Mott insulator. Our results identify the pseudogap as a distinct long-range entangled quantum phase, and offer a new route to Mott criticality beyond the paradigm of local quantum criticality. The phase appears to encode multipartite entanglement, as seen from a large value of the quantum Fisher information. The critical point at the pseudogap-Mott insulator boundary is found to be described by the exactly-solvable Hatsugai-Kohmoto model. The emergence of Luttinger surfaces alters the anomaly structure associated with the generalized symmetry of the FS [44–46]. The corresponding topological charge is defined via the Luttinger–Ward functional. This charge is modified by Green's function zeros, signalling a shift in the underlying anomaly. Within our framework, this reorganization grants topological protection to the RG flows terminating in the PG regime, ensuring the stability of its NFL excitations.

In Chapter 8, *Competing tendencies In Heavy Fermion Systems: Kondo screening vs. Mott localisation*, we ...

In Chapter 7, *Holography of Entanglement in 2D Free Fermions: Emergent Geometry and Fermionic Criticality*, we provide an explicit demonstration of the holographic principle in the form of a simple and tractable exact holographic mapping (EHM) for a free fermion system with and without a mass gap. Our program yields analytic relations between the RG flow parameters and their dual bulk quantities such as distances and curvature. This also allows us to study the consequences of a critical Fermi surface (lying precisely at the quantum phase transition) on the nature of the holographic space.

We conclude in the final chapter by summarising the main results obtained within the thesis, in terms of the methods developed as well as insights obtained into the behaviour of various models. We describe the impact of these results on the field, the broad conclusions emerging from them and some open questions resulting from the work.

Contents

1	Introduction	17
1.1	Metal-insulator transition arising from strong correlation	17
1.2	Features of Mott metal-insulator transition	19
1.3	Preliminary Insights into Mottness From a Toy Model	22
1.4	Strong local correlations: models and materials	23
1.4.1	Introduction to the auxiliary model approach	23
1.4.2	Correspondence between impurity and lattice models in $d = \infty$	24
1.4.3	Examples of materials with strong local correlations	28
1.4.4	Generator of strong local correlations: Quantum impurity models	32
1.5	Mott transition in Hubbard-like models	35
1.5.1	In the limit of infinite dimensions	35
1.5.2	The two-dimensional Hubbard model	37
1.6	Brief introduction to the tiling auxiliary model approach	39
1.6.1	Why do we need another auxiliary model method? Limitations of existing approaches	39
1.6.2	Overview	39
1.6.3	Features	39
1.7	Entanglement holography and fermionic criticality	39
2	Methods and Preliminaries	45
2.1	The Unitary renormalisation group method	45
2.2	Example I: URG analysis of the star graph model	55
2.3	Example II: URG analysis of the two-channel Kondo problem	60
2.4	Example III: Particle in a periodic potential	64
2.5	Comparison of renormalisation group approaches based on canonical transformations	70
2.6	Iterative Diagonalisation of Quantum Impurity Models	72
2.7	Optimising Fermi Surface Calculations	76
2.8	Measures of entanglement and their computation	78
2.8.1	Reduced density matrix elements as equal-time correlation functions	78
2.8.2	Decomposition of entanglement entropy in terms of multipartite information I_n	84
2.9	Wavefunction Renormalisation Group	87
2.9.1	General idea	87

2.9.2	Evolution equation for state tensor coefficients	89
2.10	Kondo model spectral function	91
2.11	Zero temperature Greens function in frequency domain	93
2.11.1	Spectral representation of Greens function	93
2.11.2	Real and imaginary parts - The Sokhotski-Plemelj theorem	94
2.11.3	Zero temperature spectral function	95
2.11.4	Reconstructing full Green function from spectral function: Kramers-Kronig relations	96
3	Kondo Frustration via Charge Fluctuations: A Route to Mott Localisation in Infinite Dimensions	98
3.1	Introduction	98
3.2	Model Hamiltonian and method	101
3.2.1	The extended Anderson impurity model	101
3.2.2	The unitary renormalisation group method	102
3.3	$T = 0$ scaling theory for the e-SIAM	104
3.3.1	URG equations for the e-SIAM	104
3.3.2	Phase diagram	105
3.3.3	Phase transition at r_{c2}	107
3.3.4	The case of repulsive U_b	108
3.4	Descriptors of the local MIT	108
3.4.1	Evolution of the impurity spectral function	108
3.4.2	Using entanglement to track correlations across the transition	110
3.5	Universal theory for the local metal-insulator transition	112
3.5.1	Local pairing and the destruction of Kondo screening	114
3.5.2	Emergent self-consistency in the e-SIAM	114
3.6	Coexistence of local metallic and insulating phases in the e-SIAM	116
3.6.1	Excited state quantum phase transition at r_{c1} : the Mott-Hubbard scenario	116
3.6.2	Theory for the charge excitations in the Hubbard sidebands	118
3.6.3	Hysteresis and the first-order line	120
3.6.4	Zero temperature origin of critical fluctuations above the second order point	121
3.7	Non-Fermi liquid signatures at the MIT	124
3.7.1	Death of the local Fermi liquid: the Brinkman-Rice scenario	124
3.7.2	Emergence of non-Fermi liquid excitations at the transition	126
3.7.3	Impurity magnetisation and entanglement entropy	129
3.7.4	Friedel scattering phase shift and fractional excess charge	130
3.8	Discussions and Outlook	131
3.9	Appendix: Derivation of renormalisation group equations for the extended-SIAM	133
3.9.1	Renormalisation of U_b	133
3.9.2	Renormalisation of the impurity correlation U	134
3.9.3	Renormalisation of the hybridisation V	135
3.9.4	Renormalisation of the spin-exchange coupling J	137
3.9.5	Final URG equations	137

3.10	Appendix: Expressing correlation functions in terms of entanglement	138
3.11	Appendix: One-shot decoupling of impurity from zeroth site	144
3.11.1	Renormalisation from V^*	144
3.11.2	Renormalisation from J^*	146
3.11.3	Total renormalisation	146
3.12	Appendix: Effective Hamiltonian for the excitations within the Hubbard sidebands	146
3.13	Appendix: Behaviour of the Kondo scale T_K close to the transition	149
3.14	Appendix: Effective Hamiltonian for the low-lying excitations at the critical point .	150
3.15	Appendix: Non-Fermi liquid exponents in self-energy and two-particle correlations	156
3.16	Appendix: Entanglement entropy, magnetisation and phase shift, at the QCP	157
3.17	Appendix: Additional correlations near the ESQPT and the QCP	160
4	A New Auxiliary Model Approach for Interacting Electrons: Mapping Impurity Models To Lattice Models	162
4.1	The Philosophy of Auxiliary Models	162
4.2	What Is The Correct Auxiliary Model For Correlated Electrons on a 2D Square Lattice?	164
4.3	Mapping Auxiliary Model To Lattice Model	166
4.3.1	Manybody Translation Operators	166
4.3.2	Periodisation of Auxiliary Model Into Extended Hubbard Model	167
4.3.3	Form Of The Eigenstates: Bloch's Theorem	169
4.4	Properties of the Bloch states	170
4.4.1	Translation invariance	170
4.4.2	Orthonormality	170
4.4.3	Eigenstates	171
4.5	Mapping Greens Functions from Auxiliary Model to Lattice Model	172
4.5.1	One-particle k -space Greens functions	172
4.5.2	One-particle real-space Greens functions	177
4.5.3	Equal-Time Ground State Correlators	179
4.6	Entanglement Measures	180
5	Mott Criticality as the Confinement Transition of a Pseudogap-Mott Metal: Insights on Mott Transition and Pseudogap in Two Dimensions	183
5.1	Introduction	183
5.2	Auxiliary Model and URG analysis	186
5.2.1	Lattice-embedded Impurity Model	187
5.2.2	The unitary renormalisation group method	187
5.3	Derivation of unitary RG equations for the lattice-embedded impurity model	189
5.3.1	RG scheme	189
5.3.2	Renormalisation of the bath correlation term W	191
5.3.3	Renormalisation of the Kondo scattering term J	192
5.4	Reconstructing of lattice model from auxiliary model	194
5.5	Low-energy theory for the lattice-embedded impurity model	198

5.5.1	RG Phase Diagram	198
5.5.2	Unravelling of Kondo Screening	198
5.5.3	Emergence of a two-channel Kondo Model	199
5.5.4	Momentum-resolved Dynamical Spectral Weight Transfer	200
5.6	Non-Fermi liquid excitations within the Pseudogap	201
5.6.1	Pseudogapped spectral function and singular self-energy	201
5.6.2	Finite-Frequency Scattering Rate and Optical Response Through the Pseudogap	202
5.6.3	Non-local correlations in the pseudogap	202
5.7	Exactly solvable nodal Non-Fermi liquid at Mott Criticality	204
5.8	Heisenberg Model as a low-energy description of the tiled Mott insulator	208
5.9	Effective Hamiltonian for excitons in the Mott insulator	210
5.10	Pseudogap as a strongly coupled phase of quantum matter	211
5.11	Conclusions	212
5.12	Appendix: Effective action for the Hubbard-Heisenberg model and the eSIAM	212
5.12.1	Local theory for the Hubbard-Heisenberg model	213
5.12.2	Local theory for the extended SIAM	215
6	Competing tendencies In Heavy Fermion Systems: Kondo screening vs. Mott localisation	217
6.1	Bilayer extended Hubbard model: Impurity model	217
6.2	Tiling Into Bilayer Extended Hubbard Model	218
6.3	Unitary RG Analysis of The Bilayer Lattice-Embedded ESIAM	220
6.4	Renormalisation Group Flows of Kondo Couplings: RG Phase Diagram	223
6.5	Results at half-filling	224
7	Holography of Entanglement in 2D Free Fermions: Emergent Geometry and Fermionic Criticality	226
7.1	Introduction	226
7.2	Preliminaries and definitions	227
7.2.1	The system	227
7.2.2	Entanglement measures	230
7.2.3	Luttinger volume	231
7.3	Entanglement hierarchy in mixed momentum and real space	231
7.4	EHM, the Ryu-Takayanagi bound and holographic inequalities	236
7.5	Holographic geometry of the RG flow	239
7.5.1	RG evolution as the emergent dimension	239
7.5.2	Embedding the emergent dimension on a metric space	242
7.6	Curvature of the emergent space	243
7.6.1	Topological nature of the sign of curvature	244
7.7	Entanglement Holography and Fermionic Criticality	248
7.7.1	Equivalence between a critical Fermi surface and a holographic wormhole	248
7.7.2	Connecting modular energy with the metric	248

7.7.3	Kinematics of the RG flows	249
7.8	Extension to the massive case	251
7.9	Topological content of the entanglement spectrum	252
7.9.1	Luttinger's volume as a winding number: the massless case	252
7.9.2	A relation between the quantum hall Chern numbers and Luttinger's volume	256
7.10	Conclusions and Outlook	258
8	Summary, Conclusions and Future Work	263
9	Appendices	274
9.1	Zero temperature Greens function in frequency domain	274
9.2	Spectrum of the Hubbard dimer	276
9.3	Greens function of Hubbard model in the atomic limit	277
9.4	Local Greens function for the Hubbard dimer	278
9.5	Contributions of various excitations to the site local spectral function	280
9.6	Relation between single-particle Greens function and the Greens operator ($T = 0$) .	280
9.7	Writing single-particle excitations of ground state in terms of $N = 3, S^z = \frac{1}{2}$ eigenstates	282
9.8	Matrix elements of G^{-1} between single-particle momentum excitations, for the Hubbard dimer	283
9.9	Topological interpretation of the Wilson ratio	283

Chapter 1

Introduction

The rich physics of Mott metal-insulator transitions in strongly correlated systems has been an active subject of study for quite some time [Hubbard(1963), Mott(1968), Milligan and Thomas(1985), Imada et al.(1998a)Imada, Fujimori, and Tokura]. Their appearance in various materials such as the nickelates, cuprates, iridates and manganites, among others, is marked by complex phase diagrams that display dramatic changes in properties across changes in filling and/or bandwidth, even at low temperatures. The physics of Mott transitions is driven by the competition between two energy scales in the system: (i) the scale of itineracy, quantified by the bandwidth, and (ii) the scale of electron-electron repulsion. A Mott insulator arises when the electronic correlations are strong enough that low-energy electronic excitations become gapped, and the metallic valence band gets split into lower and upper bands; for an initially half-filled valence band, the lower band is filled at $T = 0$ while the upper band is vacant. Because of strong electronic repulsion, the electrons in the lower band form local moments which can show a variety of behaviour. This includes various symmetry-broken ground states such as an antiferromagnet [Lee et al.(2006)Lee, Nagaosa, and Wen], or symmetry-preserved states such as quantum spin liquids [Pustogow et al.(2018)Pustogow, Bories, Löhle, Rösslhuber, Zhukova, Gorshunov, Tomić, Schlueter, Hübner, Hiramatsu et al.]. The physics of Mottness is also responsible for several very interesting phenomena, such as high-temperature superconductivity [Lee et al.(2006)Lee, Nagaosa, and Wen], non-Fermi liquids [Stewart(2001)] and the various phases in magic-angle twisted bilayer graphene [Cao et al.(2018a)Cao, Fatemi, Demir, Fang, Tomarken, Luo, Sanchez-Yamagishi, Watanabe, Taniguchi, Kaxiras et al., Cao et al.(2018b)Cao, Fatemi, Fang, Watanabe, Taniguchi, Kaxiras, and Jarillo-Herrero].

1.1 Metal-insulator transition arising from strong correlation

Initial attempts at explaining metal-insulator transitions was based on a non-interacting picture of electrons (through the work of Bethe, Sommerfeld, Bloch and Wilson among others); in such a picture, electronic bands can arise only from the scattering of electrons off the periodic lattice potential. One then distinguishes between metals and insulators at zero temperature by the position of the Fermi level E_F with respect to the bands: if E_F lies within a band gap, an integer number of bands will be completely filled and the system will be an insulator, whereas if E_F lies within one of




Figure 1.1

the bands, the system will be a metal because there are vacant gapless excitations within the band.

It was first pointed out by Boer and Verwey [de Boer and Verwey(1937)] that certain insulators and semiconductors (such as nickel oxides) have a particle filling that, purely from band theory, would lead to partially filled bands. This poses a challenge because a partially filled band necessarily leads to a metallic phase (see Fig. 1.1 (left)). The insulating behaviour of these materials therefore signals a fundamental limitation of single-particle band theory and points to a qualitatively different mechanism for electron localization—one driven by strong electron–electron interactions rather than by scattering from the lattice potential alone. This realization led to the concept of a Mott insulator, in which charge localization emerges as a genuine many-body effect despite the absence of a band gap in the non-interacting electronic structure. A very simple mechanism for this was put forth by Mott [Mott(1949), Mott(1956), Mott(1961)]: in the presence of a large local Coulomb repulsion cost U on a lattice site, the quantum states $|\Psi_1\rangle$ in which the site is singly occupied get separated in energy from the states $|\Psi_2\rangle$ where the site is doubly occupied by a gap of size U ; the former class of states $|\Psi_1\rangle$ form the lower band (LB) while the latter $|\Psi_2\rangle$ form the upper band (UB). If we now have a half-filled system (one electron per site), this electron will completely fill LB and leave UB vacant, leading to an insulating state (see Fig. 1.1 (right)).

Given the above discussion, it is clear that a Mott insulator must involve a gap to charge excitations (excitations that add or remove an electron). It does not however determine the fate of the spin, orbital or other degrees of freedom. One scenario is when the spin or orbital degree of freedom undergoes spontaneous symmetry breaking into an ordered state; in the case of spin, this usually results in a superexchange interaction being generated and leads to a long-range ordered antiferromagnetic Neel state coexisting with the Mott state. Examples of such systems include vanadium sesquioxide V_2O_3 and its derivatives and the high- T_c cuprates (such as La_2CuO_4) [Imada et al.(1998a)Imada, Fujimori, and Tokura]. The other scenario is one where the spin degree of freedom retains its symmetry, and we end up with a quantum mechanically non-trivial entangled state (in order to quench the entropy). Examples of such states include quantum spin liquids [Isono et al.(2016)Isono, Terashima, Miyagawa, Kanoda, and Uji]. Mott insulators without long-range order are of particular interest because of the possibility of topological order, fractional excitations, and novel metallic parent states.

A prototypical model for gaining theoretical understanding of Mott metal-insulator transitions on a lattice is the single-band Hubbard model [Anderson(1959), Hubbard(1963), Kanamori(1963), Gutzwiller(1963)]

$$H = -t \sum_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger c_{j,\sigma} + \text{h.c.}) - \frac{U}{2} \sum_i (n_{i,\uparrow} - n_{i,\downarrow})^2 - \mu N. \quad (1.1)$$

The first term incorporates nearest-neighbour hopping on the lattice, while the second term introduces inter-electron repulsion when two electrons site on the same lattice site; in this way, the model incorporates the two competing tendencies of delocalisation and localisation that are responsible for the Mott MIT. The third term fixes the chemical potential and determines the filling of the lattice. Many of the materials that are studied for Mott MIT involve d -electrons that are five-fold degenerate, and these orbitals can hybridise with ligand atoms. The Hubbard model avoids these complexities by focusing on only a single band that lies closest to the Fermi energy, and it is the states of this band that are described by the Hubbard model.

This leaves us with two channels for observing Mott MIT [Imada et al.(1998a)Imada, Fujimori, and Tokura]: (i) tuning the bandwidth, and (ii) tuning the filling. The former can be done, in experiments, by applying pressure. Filling is tuned by doping specific elements of a compound with elements of a different valence. The former is referred to as bandwidth-control metal-insulator transition, while the latter is called filling-control metal-insulator transition.

1.2 Features of Mott metal-insulator transition

Two complementary pictures of the Mott transition: Brinkman-Rice vs Mott-Hubbard. Initial insights into the Mott MIT were obtained by Hubbard [Hubbard(1963)], thinking along lines similar to Mott. Within this picture, one starts from the insulating large U limit of the Hubbard model (eq. 1.1) ($U \gg t$). We stay at half-filling for simplicity ($\mu = 0$). The large on-site repulsion leads to the formation of two bands at each site, the lower Hubbard band (LHB) associated with singly-occupied states and the upper Hubbard band (UHB) associated with doubles and holes (states that incur a cost U). In the large U limit, these bands are well-separated, and the dynamics within the bands is a result of the motion of tightly bound holons and doubles (more on this later) that are generated because of the small but non-zero hopping probability t . As U/t is lowered, the bands come closer and a metal-insulator transition occurs at the critical value of U at which the gap vanishes. For lower U/t , the bands overlap and we get a metal. This picture fails to provide an accurate description of the metal resulting from the melting of such a Mott insulator [Georges et al.(1996)Georges, Kotliar, Krauth, and Rozenberg].

An alternative approach to the Mott MIT is due to Brinkman and Rice [Brinkman and Rice(1970)], where the authors applied Gutzwiller's variational method [Gutzwiller(1965)] to the half-filled Hubbard model. Gutzwiller's approach involves modifying, through variational factors, the hopping matrix elements that start from doubly occupied states, in order to reflect the electronic correlation at each site. By requiring that the number of doubly-occupied states vanish in the Mott insulating ground state, they obtained a critical value of the interaction strength. In the Brinkman-Rice scenario, therefore, the transition from the metal to the insulating phase occurs when the hopping matrix




Figure 1.2

elements from the doubly-occupied to the singly-occupied states strictly vanishes. Note that in contrast to the Mott-Hubbard picture, this approach starts from a correlated Fermi liquid (renormalised due to the Gutzwiller factors). While it gives a better low-energy description of the correlated metal compared to the Mott-Hubbard picture, it fails to account for the high-energy physics, and treats the Mott insulator as a direct product state of local moments at each state (on account of the strictly zero hopping). This of course means that this method cannot explain antiferromagnetic superexchange processes in the Mott insulator [Georges et al.(1996)Georges, Kotliar, Krauth, and Rozenberg].

Holon-doublon unbinding picture of the Mott MIT. In 1949, Mott proposed a zero temperature explanation for a correlation-driven insulator and a mechanism for its transition into a metal in terms of the motion of holes and doubles in the background of a half-filled lattice (see Fig. 1.2). For $U \ll t$ (strong correlation), the ground state predominantly involves lattice sites with a single electron. Hopping processes between two nearest-neighbour sites transfer an electron between them and create a pair of hole and double, but the pair remains tightly bound to each other. This is because moving the hole independently involves scattering processes that cross the Mott gap, and these processes are not allowed at $T = 0$. But without the independent motion of the hole (or double), there is no current, and the system is an insulator (left, Fig. 1.2).

As U/t is lowered, the gap closes and processes that transport the hole without transporting the double become gapless. This essentially means that the pair becomes deconfined, and the hole can be moved far away from the double (right, Fig. 1.2). Current can then be passed by having the hole move through the lattice, and we get a metallic state. This leads to a picture of the Mott transition into a metal in terms of an unbinding of the hole-double bound states (excitons) that were present in the insulator.

There have been more quantitative studies of this mechanism. I list some of the relatively recent ones here. In [Prelovšek et al.(2015)Prelovšek, Kokalj, Lenarčič, and McKenzie], Prelovšek et al. studied the motion of holon-doublon pairs in a spin polarised background by rewriting the repulsive Hubbard model in terms of electrons as an attractive Hubbard model of holes and doubles. The model was treated using a BCS-like treatment, and the Mott transition was obtained when the

binding energy for the hole-double condensate vanishes. To mark the deconfinement, they show that the approach to the transition from the insulating side is concomitant with an increase in the size of the pairs. Following a slave-boson treatment of holon-doublon unbinding, Zhou et al. [Zhou et al.(2014)Zhou, Wang, and Wang] show that the Mott transition from a paramagnetic metal to a Mott insulator on the half-field honeycomb and 2D square lattices occur through an intervening Slater insulator with coherent quasiparticles. In particular, their approach sheds light on the nature of the incoherent excitations in the Hubbard sidebands of the local spectral function. Finally, an older work by Watanabe et al. [Watanabe et al.(2006)Watanabe, Yokoyama, Tanaka, and Inoue] studies the half-filled Hubbard model on an anisotropic triangular lattice using a Monte carlo method with a variational binding parameter between holons and doublons. Their approach reveals a first order Mott transition, along with robust d -wave superconductivity and short-range magnetic order.

Spectral features of the Mott transition. One of the important diagnostics of the Mott metal-insulator transition is the spectrum for one-particle electron-like excitations, as expressed through the one-particle retarded Greens function $G(\mathbf{k}, \sigma; t) = -i\theta(t)\langle\{c_{\mathbf{k},\sigma}(t), c_{\mathbf{k},\sigma}^\dagger\}\rangle$. In general, the presence of poles in the Green function at the Fermi energy and in its neighbourhood imply the existence of gapless excitations and hence a metallic phase. It turns out that this statement is equivalent to Luttinger's theorem,

$$\frac{N}{V} = 2 \int_{G(0,\mathbf{k})>0} \frac{d\mathbf{k}}{(2\pi)^3}, \quad (2.1)$$

which states that the particle density in a metal is equal to the number of states in the Fermi volume. As clarified in the equation, the Fermi volume is defined as the region of k -space where the Greens function is positive at $\omega = 0$.

In a Mott insulator, strong correlations lead to the destruction of electronic excitations around $\omega = 0$. This results in the Greens function poles transmuting into zeros. This is also tracked by analogous changes in the electronic self-energy $\Sigma(\mathbf{k}; \omega)$; Greens function zeros leads to a Mott pole at $\omega = 0$ in $\Sigma(\mathbf{k}; \omega)$. The points in k -space where $\Sigma(\mathbf{k}; 0)$ vanishes is called a Luttinger surface. The Mott metal-insulator transition is therefore about the topological change of a Fermi surface (surface of Greens function poles) into a Luttinger surface (surface of self-energy poles).

As discussed earlier, the appearance of Greens function zeros at $\omega = 0$ leads to the formation of a charge gap that freezes one-particle low-energy excitations. The spectrum gets redistributed into two bands on either side of the gap, the upper and lower Hubbard bands. In the ground state, the lower band is obviously filled while the upper band is empty. Bands of a Mott insulator formed from electronic correlation differ from those formed due to one-particle scattering in a particular way: dynamical spectral weight transfer. Unlike a band insulator, the bands of a Mott insulator are not rigid; tuning the lattice filling results in a continuous transfer of states from one band to another.

The non-rigidity of the bands can be seen most easily from the atomic limit ($t = 0$) of the Hubbard model (eq. 1.1). The local retarded Greens function at an arbitrary site i can be obtained using the equation of motion approach. The final result is [Phillips(2012)]

$$G_R(i; \omega) = \frac{1+x}{\omega + \mu + U/2} + \frac{1-x}{\omega + \mu - U/2}, \quad (2.2)$$




Figure 1.3

where $x = 1 - \sum_{\sigma} \langle n_{i\sigma} \rangle$ is the average number of holes at any site. The spectral weight for the two poles makes it clear that the lower band has $1 + x$ number of states while the upper band has $1 - x$ states. Changing the number of states in the lower band (by tuning μ and hence x) also changes those in the upper band. For example, by adding 0.1 holes per site to the lower band, the spectral weight in the upper band decreases by the same amount. This is the non-rigidity of the bands mentioned previously.

1.3 Preliminary Insights into Mottness From a Toy Model

The Baskaran-Hatsugai-Kohmoto model [Baskaran(1991), Hatsugai and Kohmoto(1992)] introduced independently in 1991 and 1992 is an exactly solvable model of electronic correlation that displays a transition from a metallic phase to a Mott insulator. While its solvability offers an instructive method for studying Mott transition, the infinitely long-ranged nature of its interaction makes it tricky to apply these conclusions to realistic materials.

At half-filling, the model is defined by the following Hamiltonian:

$$H_{\text{BHK}} = \sum_{\mathbf{k}, \sigma} \epsilon(\mathbf{k}) n_{\mathbf{k}, \sigma} - \frac{U}{2} \sum_{\mathbf{k}} (n_{\mathbf{k}, \uparrow} - n_{\mathbf{k}, \downarrow})^2. \quad (3.1)$$

Clearly the model is diagonal in momentum-space; the Hamiltonian splits up into commuting 4×4 blocks for every point $\mathbf{k}$:

$$H_{\text{BHK}} = \sum_{\mathbf{k}} H(\mathbf{k}), \quad H(\mathbf{k}) = \epsilon(\mathbf{k}) \sum_{\sigma} n_{\mathbf{k}, \sigma} - \frac{U}{2} (n_{\mathbf{k}, \uparrow} - n_{\mathbf{k}, \downarrow})^2. \quad (3.2)$$

The vacant eigenstate $|0\rangle$ of $H(\mathbf{k})$ has zero energy ($E_0 = 0$), the two singly-occupied states $|\uparrow\rangle$ and $|\downarrow\rangle$ have energy $E_1 = \epsilon(\mathbf{k}) - U/2$ and the doubly-occupied state $|\uparrow, \downarrow\rangle$ has energy $E_2 = 2\epsilon(\mathbf{k})$. This leads to two possible single-particle excitations – one for adding an electron on the vacant state: ($\Delta_1 = E_1 - E_0 = \epsilon(\mathbf{k}) - U/2$), the other for adding an additional electron on a singly-occupied state: $\Delta_2 = E_2 - E_1 = \epsilon(\mathbf{k}) + U/2$.

Upon combining states from other k -modes, these isolated excitations broaden into bands around $\pm U/2$ because of the continuum of values available to $\epsilon(\mathbf{k})$. For large enough U (compared to the non-interacting bandwidth W), these bands remain well-separated in energy (Fig. 1.3 left), forming lower and upper Hubbard bands. The half-filled system results in the lower band being completely filled and the upper band empty, leading to a Mott insulator. The states within this band are all singly-occupied (per k -state). Upon lowering the correlation strength U , the bands touch each other at the critical value of $U_c = W$ (obtained by equating the top of the lower band to the bottom of the upper band). For lower values of U , the system is a metal due to the presence of gapless excitations within the combined band (Fig. 1.3 right). This therefore provides a simple demonstration of a correlation-driven Mott metal-insulator transition.

1.4 Strong local correlations: models and materials

1.4.1 Introduction to the auxiliary model approach

One of the main results obtained within this thesis is the development of a formalism to study the more realistic case of finite-dimensional interacting models by using appropriately chosen impurity models as auxiliary models. The key to an *auxiliary model method* is to construct an appropriate auxiliary model that while being tractable captures some essential property of the full lattice model [Martin et al.(2016)Martin, Reining, and Ceperley]. This involves separating the complete degrees of freedom into a system part (H_S) which one treats exactly and the rest of the system (H_R) that one might simplify in order to be able to solve the problem. The non-trivial nature of the problem arises of course from the hybridisation H_{SR} between the system and the bath:

$$H = \begin{bmatrix} H_R & H_{RS} \\ H_{RS}^* & H_S \end{bmatrix}. \quad (4.1)$$

This separation can always be done exactly, if one does not care about tractability. For example, one can take the Hubbard model that describes electrons hopping on a lattice and interacting when two electrons reside on the same lattice site:

$$H_{\text{hub}} = -t \sum_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger c_{j,\sigma} + \text{h.c.}) + U \sum_i n_{i\uparrow} n_{i\downarrow}, \quad (4.2)$$

and separate this into parts, the system being the local orbitals of a specific site l and the rest of the system being all the other sites:

$$H_S = U n_{l\uparrow} n_{l\downarrow}, \quad H_R = -t \sum'_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger c_{j,\sigma} + \text{h.c.}) + U \sum_i n_{i\uparrow} n_{i\downarrow}, \quad H_{SR} + H_{RS} = -t \sum_{i,\sigma} (c_{l,\sigma}^\dagger c_{i,\sigma} + \text{h.c.}), \quad (4.3)$$

where the primed summation is carried out over all nearest-neighbour pairs that do not involve l and the summation in $H_{SR} + H_{RS}$ is over all sites adjacent to l .

The smaller system H_S (often called the *cluster*) is typically chosen such that its eigenstates are known exactly. Progress is then made by choosing a simpler form for H_R and its coupling H_{RS} with

the smaller system. This combination of the cluster and the simpler bath is then called the *auxiliary system*. A tractable auxiliary system for the Hubbard model is the single-impurity Anderson model (SIAM); the correlated impurity site represents the set of local orbitals of any given site, while the conduction bath represents the rest of the lattice sites in an approximate fashion. Such a construction is shown in fig. 4.1.




Figure 1.4: Left: Full Hubbard model lattice with onsite repulsion U^H on all sites and hopping between nearest neighbour sites with strength t^H . Right: Extraction of the auxiliary (cluster+bath) system from the full lattice. The central site on left becomes the impurity site (red) on the right (with an onsite repulsion ϵ_d), while the rest of the $N - 1$ sites on the left form a conduction bath (green circles) (with dispersion ϵ_k and correlation modelled by the self-energy $\Sigma_k(\omega)$) that hybridizes with the impurity through the coupling V .

It should be noted that any reasonable choice of the cluster and bath would break the translational symmetry of the full model: To allow computing quantities, one would need to make the bath (which is a much larger system) simpler than the cluster (which is a single site). This distinction breaks the translational symmetry of the Hubbard model. As a result, calculations that make use of the translation symmetry of the full model (such as k -space objects) will require a periodisation operation of an auxiliary model quantity.

1.4.2 Correspondence between impurity and lattice models in $d = \infty$

Dynamical Mean-Field Theory (DMFT) provides a nonperturbative framework for treating strong local electronic correlations in lattice models by mapping them onto a self-consistent quantum impurity problem [Metzner and Vollhardt(1989), Georges et al.(1996)Georges, Kotliar, Krauth, and Rozenberg]. The central idea of DMFT is that in the limit of infinite spatial dimension, or equivalently infinite coordination number, the electronic self-energy becomes purely local while retaining its full dynamical (frequency-dependent) structure. As a result, the many-body lattice problem reduces to an effective single-site problem embedded in a self-consistently determined bath, capturing local quantum fluctuations exactly while neglecting nonlocal correlations. DMFT has become a cornerstone of modern correlated-electron theory, successfully describing Mott metal–insulator transitions, quasiparticle formation, and spectral weight transfer in Hubbard-like models [Georges et al.(1996)Georges, Kotliar, Krauth, and Rozenberg, Georges(2004)]

In order to motivate the impurity model-lattice model mapping inherited by DMFT, we first recall the textbook example of classical mean-field theory. Mean-field theory provides one of the earliest and most instructive examples of controlled approximations in many-body physics. In its classical incarnation, exemplified by the Ising model,

$$H_{\text{Ising}} = - \sum_{i \neq j} J_{ij} S_i S_j - h \sum_i S_i = - \sum_i S_i (h + \sum_{j \neq i} J_{ij} S_j), \quad (4.4)$$

mean-field theory proceeds by replacing the interaction between a given spin and its neighbors by an *effective static field* proportional to the average magnetization. Writing $S_j = \langle S_j \rangle + \delta S_j$ and neglecting quadratic fluctuations and constant parts, one obtains an *auxiliary problem* in the form of a mean-field Hamiltonian

$$H_{\text{Ising}}^{(\text{MF})} = - \sum_i (h + h_{\text{MF}}(i)) S_i, \quad (4.5)$$

where the mean-field h_{MF} comes out to be

$$h_{\text{MF}}(i) = \sum_{j \neq i} 2 \langle S_j \rangle J_{ij}. \quad (4.6)$$

The above approach assumes two things: (i) the presence of a non-zero *order parameter* $m_j = \langle S_j \rangle$ (in this case, a non-zero magnetisation), and (ii) the smallness of the fluctuations $S_j - \langle S_j \rangle$. The mean-field Hamiltonian can be used to compute the partition function (at inverse temperature β) and hence the magnetisation at a site i of the lattice:

$$m_i = \tanh(\beta(h_{\text{MF}}(i) + h)) = \tanh(\beta \sum_{j \neq i} 2m_j J_{ij} + \beta h). \quad (4.7)$$

The problem hasn't yet been solved of course; one needs to determine the appropriate value of m_j . We therefore need an additional equation; this is provided by the so-called *self-consistency condition*. The translation invariance of the lattice ensures that the magnetisation must be uniform across sites i and j . This allows us to replace m_j with m_i in the above equation:

$$m_i = \tanh(\beta m_i \sum_{j \neq i} 2J_{ij} + \beta h). \quad (4.8)$$

This equation can be solved numerically to obtain the uniform magnetisation $m_i = m$. The mean-field approximation becomes exact in the limit of large coordination, where fluctuations are suppressed and spatial correlations beyond a single site vanish.

An equivalent way of framing the self-consistency equation is to equate the mean-fields on different lattice sites. Combining eqs. 4.7 and 4.6, we have

$$h_{\text{MF}}(i) = \sum_{j \neq i} 2m_j J_{ij} = \sum_{j \neq i} 2 \tanh(\beta(h_{\text{MF}}(j) + h)) J_{ij}. \quad (4.9)$$

The equation in terms of $h_{\text{MF}}(i)$ and $h_{\text{MF}}(j)$ can again be solved numerically by claiming that the mean-field has to be uniform across all the sites owing to translation invariance. This form of the self-consistency requirement (in terms of the field) will be useful to us in the upcoming discussion.

The essential feature of classical mean-field theory is therefore the replacement of a many-body problem by an effective *single-site* problem embedded in a self-consistently determined environment. Quantum lattice models admit a closely analogous construction, but with an important conceptual difference: while the classical mean field is static, quantum fluctuations necessarily introduce a tower of excited states that allow for temporal dynamics. This observation underlies the formulation of dynamical mean-field theory (DMFT), which may be viewed as the natural quantum generalization of classical mean-field theory [Metzner and Vollhardt(1989), Georges et al.(1996)Georges, Kotliar, Krauth, and Rozenberg].

Similar to the previous case of the Ising model, we will now derive the self-consistency equation for a quantum model. Consider an interacting fermionic lattice Hamiltonian, such as the Hubbard model:

$$H = - \sum_{ij\sigma} t_{ij} c_{i\sigma}^\dagger c_{j\sigma} + U \sum_i n_{i\uparrow} n_{i\downarrow}, \quad (4.10)$$

defined on a lattice with coordination number z .

The point of this exercise is to calculate the interacting self-energy $\Sigma_k(\omega)$ for the fermions. This therefore acts as an “order parameter” and plays the same role as the local magnetisation in the previous exercise. The self-energy defines the interacting k -space Green’s function for a translation-invariant problem:

$$G_{\mathbf{k}}(\omega) = \frac{1}{\omega - \varepsilon_{\mathbf{k}} - \Sigma_{\mathbf{k}}(\omega)}, \quad (4.11)$$

where $\varepsilon_{\mathbf{k}}$ denotes the noninteracting dispersion, and a real-space local Green’s function, obtained by summing over momentum:

$$G_{\text{loc}}(\omega) = \sum_{\mathbf{k}} \frac{1}{\omega - \varepsilon_{\mathbf{k}} - \Sigma_{\mathbf{k}}(\omega)}. \quad (4.12)$$

Similar to the earlier exercise, we need to apply a mean-field decoupling to drop certain fluctuations and obtain a similar effective problem. In the context of interacting electrons, such a decoupling is obtained in the limit of infinite dimensionality or coordination number, where the self-energy becomes k -independent [Metzner and Vollhardt(1989)]:

$$\Sigma_{\mathbf{k}}(\omega) = \Sigma_{\text{loc}}(\omega), \text{ or (equivalently) } \Sigma_{ij}(\omega) = \delta_{ij} \Sigma_{\text{loc}}(\omega). \quad (4.13)$$

The mean-field decoupling here therefore corresponds to ignoring spatial fluctuations of the one-particle self-energy, and leads to a local Greens function that is directly connected to the local self-energy:

$$G_{\text{loc}}(\omega) = \sum_{\mathbf{k}} \frac{1}{\omega - \varepsilon_{\mathbf{k}} - \Sigma_{\text{loc}}(\omega)}. \quad (4.14)$$

It turns out that a local Greens function connected to a local self-energy also defines a quantum impurity problem describing the dynamics of a single correlated site with the other environment

sites integrated out:

$$S_{\text{imp}} = - \int_0^\beta d\tau d\tau' c^\dagger(\tau) G_{\text{MF}}^{-1}(\tau - \tau') c(\tau') + U \int_0^\beta d\tau n_\uparrow(\tau) n_\downarrow(\tau). \quad (4.15)$$

This acts as an auxiliary problem for the more complete problem locally interacting electrons on a lattice. The non-interacting Greens function G_{MF} is as of now unknown, and defines the environment of the impurity site and comprises the dynamics of the sites that have been integrated out. Such an impurity problem is solvable, and the impurity Green's function is given by

$$G_{\text{imp}}(\omega) = \frac{1}{G_{\text{MF}}^{-1}(\omega) - \Sigma_{\text{imp}}(\omega)}, \quad (4.16)$$

The first term in eq. 4.15 is therefore what makes this problem dynamical – it represents the probability of electrons hopping into and out of the impurity site, and leads to fluctuations of its occupancy. We can make the identification of the Greens function G_{MF} as the *mean-field* encountered in the previous example. More specifically, G_{MF} acts as a “field” because it is obtained by integrating out the other sites on the lattice, and it acts on the impurity site by affecting its dynamics.

The complete set of equations for the solving such a dynamical mean-field problem is finally obtained by equating the impurity local Greens function G_{imp} to the lattice local Greens function G_{loc} . As discussed earlier, such a solution exists in an exact fashion only in the limit of infinite dimensions, as long as the correct G_{MF} can be found. In order to find the appropriate G_{MF} that allows equating the impurity model Greens function to a lattice mode Greens function, we require an additional equation: the self-consistency condition. Similar to the case of the Ising model, we require that G_{MF} be equal to G_{loc} , owing to translation invariance. The full set of coupled DMFT equations is

$$\begin{aligned} G_{\text{MF}} &= G_{\text{loc}}, \\ \Sigma_{\text{imp}}(\omega) &= G_{\text{MF}}^{-1}(\omega) - G_{\text{imp}}^{-1}(\omega), \\ \Sigma_{\text{loc}}(\omega) &= \Sigma_{\text{imp}}(\omega), \\ G_{\text{loc}}(\omega) &= \sum_{\mathbf{k}} \frac{1}{\omega - \varepsilon_{\mathbf{k}} - \Sigma_{\text{loc}}(\omega)}. \end{aligned} \quad (4.17)$$

- The first equation is equivalent to the feature in the previous exercise where we equated the mean-fields on i and j .
- The second equation is equivalent to computing, in the previous exercise, the magnetisation on the j^{th} site from the partition function.
- The third equation is the essential mean-field approximation, where we equate the self-energy of the simpler decoupled problem to that of the more complete lattice problem and is justified in the limit of infinite dimensions.
- The fourth equation is equivalent, in the previous exercise, to using the computed “magnetisation” of site j in the second step to further compute the mean-field at site i .

In direct analogy with classical mean-field theory, DMFT replaces a lattice model by an effective single-site problem embedded in a self-consistent environment. The crucial distinction is that the mean field is now dynamical rather than static, encoding quantum fluctuations exactly while neglecting spatial correlations. Importantly, this demonstrates the importance of quantum impurity models in investigating correlated lattice models, particularly in the presence of strong local interactions. This is because the effects of local correlations are captured exactly by such an auxiliary model mapping.

1.4.3 Examples of materials with strong local correlations

As discussed in the previous section, the route to studying lattice problems via quantum impurity models makes most sense for materials with strong local correlations. Such systems are often referred to as strongly correlated quantum materials, and their phenomenology often lies beyond conventional approaches like density functional theory or perturbative weak-coupling methods. In practice, such materials often involve transition metal elements or rare earth/actinide elements, due to the presence of partially filled $3d$ – or $4f$ –orbitals in such elements. This is because these orbitals are more more localised around the nucleus than the s – or p –orbitals of comparable energy, and hence lead to a reduced itinerant energy scale and amplify the effects of local correlations [Georges(2004)].

The localised nature of these orbitals can be understood as follows. For the $3s$ –orbital, the $1s$ and $2s$ orbitals have same orbital angular momentum l but lower principal quantum number n ; since all such wavefunctions must be orthogonal to each other, the $3s$ –orbital must develop nodes in its radial form and extend farther away from the nucleus in order to be orthogonal to these other orbitals. However, for the $3d$ –orbital, there is no orbital with same l but lower n ; this means that the orthogonality is already enforced by the behaviour of the angular part (the spherical harmonics), and the radial part can remain localised around the nucleus without having to spread out. A similar argument holds for the localised nature of the $4f$ –orbital.

We now discuss the phenomenology of some of these quantum materials in order to identify the challenges in trying to explain them and the kind of models that are likely to be useful in this endeavour.

Transition Metal Oxides (TMOs) Transition Metal Oxides (TMOs) are defined chemically by the presence of partially filled d –shells in transition metals (e.g., Cu^{2+} , Mn^{3+} , V^{4+}) coordinated by oxygen ligands. The physics of TMOs is governed by the competition between the kinetic energy hopping amplitude t and the on-site Coulomb repulsion U ; due to the small overlap between the localised d –orbitals, the d –electrons can delocalise only by hopping onto the ligand atoms. According to the Zaanen-Sawatzky-Allen (ZSA) classification scheme [Zaanen et al.(1985)Zaanen, Sawatzky, and Allen], these materials are categorized as either Mott-Hubbard insulators, where the charge gap is determined by the d – d Coulomb interaction ($E_g \approx U$), or charge-transfer insulators, where the gap is dictated by the energy difference between the oxygen p –band and the metal d –band. These derive from considerations of several factors, such as crystal field splitting, arrangement of atoms and lattice symmetries.

For example, in the case of octahedral symmetry, repulsion from the surrounding atoms leads to the raising of energy of the $d_{x^2-y^2}$ and $d_{3z^2-r^2}$ orbitals forming the e_g doublet, while the other

three orbitals (d_{xy} , d_{yz} and d_{zx}) form the t_{2g} multiplet. For the heavier TMOs (Ti, V, Cr...), the Fermi level lies within e_g . The increased positive nuclear charge (compared to the lighter elements) lowers the energy of the d -orbitals and brings them closer to the p -orbitals, reducing the charge transfer energy $\Delta = \varepsilon_d - \varepsilon_p$ of transporting an electron from a d -state to a p -state, compared to the d -level repulsion U ($|\Delta| < U$). Geometric reasons (the e_g orbitals point towards the p -orbitals) also enhance the hybridisation between d - and p -orbitals in such materials. If Δ becomes larger than the bandwidth, we obtain a *charge-transfer insulator*, where the charge gap is decided by Δ . In the lighter TMOs, the Fermi level passes through t_{2g} , and because they point away from the p -orbitals, the hybridisation is weaker. Moreover, the reduced positive nuclear charge leads to an increased $\Delta > U$; a U greater than the bandwidth leads to what's called a *Mott-Hubbard insulator*.

Experimental studies of transition metal oxides (TMOs) have established metal-insulator transitions (MITs) as a defining consequence of strong electronic correlations in systems with partially filled (d)-electron shells [Imada et al.(1998b)Imada, Fujimori, and Tokura,McWhan et al.(1969)McWhan, Rice, and Remeika]. Early experiments on vanadates such as V_2O_3 characterised pressure-driven MITs through the pressure- and temperature-dependent conductivity as well as an anomalous enhancement of the optical conductivity [Limelette et al.(2003a)Limelette, Georges, Jérôme, Wzietek, Metcalf, and Honig,Rozenberg et al.(1995)Rozenberg, Kotliar, Kajueter, Thomas, Rapkine, Honig, and Metcalf, Qazilbash et al.(2007)Qazilbash, Brehm, Chae, Ho, Andreev, Kim, Yun, Balatsky, Maple, Keilmann et al.]. Copper oxides constitute the most extensively studied class of correlated TMOs and provide a canonical example of a doping-driven MIT. Transport measurements on parent compounds such as La_2CuO_4 establish a robust insulating state, while carrier doping leads to anomalous metallic behavior characterized by linear-in-temperature resistivity and unconventional superconductivity [Takagi et al.(1992)Takagi, Batlogg, Kao, Kwo, Cava, Krajewski, and Peck,Ando et al.(2004)Ando, Komiya, Segawa, Ono, and Kurita, Lemberger et al.(2011)Lemberger, Hetel, Tsukada, Naito, and Randeria]. Such metal-insulator transition is also reported from angle-resolved photoemission spectroscopy (ARPES) measurements in doped La_2CuO_4 and $Bi_2Sr_2CaCu_2O_8$, along with the emergence of pseudogap near the antinodes at high temperatures and a hole-to-electronlike Fermi surface reconstruction [Damascelli et al.(2003)Damascelli, Hussain, and Shen, Sobota et al.(2021)Sobota, He, and Shen,Vishik(2018)]. Nickelates ($LaNiO_2$, $La_3Ni_2O_7$, $La_4Ni_3O_{10}$, etc) exhibit related but distinct experimental signatures. With Ni^{2+} sharing the same $3d^9$ valence configuration as Cu^{2+} in the cuprate superconductors, these materials also display superconductivity, although the mechanism in the nickelates is different due to the strong coupling between the layers [Medarde(1997), Wang et al.(2025)Wang, Jiang, Ying, Wu, Cheng, Hu, and Chen]. The optimal doped regime in thin films shows strange metal behaviour, while infinite-layer nickelates show the absence of long-range antiferromagnetic order [Puphal et al.(2026)Puphal, Schäfer, Keimer, and Hepting].

Transition Metal Dichalcogenides Transition Metal Dichalcogenides (TMDs) are layered van der Waals materials with the stoichiometry MX_2 , where M is a transition metal (e.g., Mo, W) and X is a chalcogen (e.g. S, Se, Te) [Manzeli et al.(2017)Manzeli, Ovchinnikov, Pasquier, Yazyev, and Kis,Joseph et al.(2023)Joseph, Mohan, Lakshmy, Thomas, Chakraborty, Thomas, and Kalarikkal]. Recent interest has focused on the interplay between strong spin-orbit coupling (SOC), electronic correlations and direct bandgap – properties that make them particularly useful for electronic ap-

plications, spintronics and DNA sequencing. Layered TMDs have the structure $X - M - X$, with the transition metal sandwiched between two hexagonal planes of chalcogenides. Different combinations of the metals with the chalcogens lead to widely varying structures and electronic properties in TMDs. Experimental studies show that metal–insulator transitions (MITs) in TMDs are commonly intertwined with charge density wave (CDW) formation rather than arising from purely on-site Coulomb localization [Wilson et al.(1974)Wilson, Di Salvo, and Mahajan, Wilson et al.(2001)Wilson, Salvo, and Mahajan]. In prototypical compounds such as 1T-TaS₂, ARPES and transport measurements reveal fluctuating magnetic order parameters in the Mott state and a transition to superconductivity driven by pressure [Fazekas and Tosatti(1979),Perfetti et al.(2005)Perfetti, Gloor, Mila, Berger, and Grioni, Sipos et al.(2008)Sipos, Kusmartseva, Akrap, Berger, Forró, and Tutiš]. Scanning tunnelling microscope (STM) pulses have been successfully used to manipulate the metal–insulator transition of 1T-TaS₂, demonstrating the macroscopic controllability of the Mott phase [Cho et al.(2016)Cho, Cheon, Kim, Lee, Cho, Cheong, and Yeom].

ARPES measurements in 2H-NbSe₂ and 2H-TaSe₂ partial gapping of the Fermi surface restricted to nested momentum regions and a temperature-dependent pseudogap similar to the cuprates [Borisenko et al.(2008)Borisenko, Kordyuk, Yaresko, Zabolotnyy, Inosov, Schuster, Büchner, Weber, Follath, Patthey et al., Borisenko et al.(2009)Borisenko, Kordyuk, Zabolotnyy, Inosov, Evtushinsky, Büchner, Yaresko, Varykhalov, Follath, Eberhardt et al.]. This is further supported by optical conductivity experiments that show spectral weight in the midinfrared region along with a low-frequency Drude contribution [Vescoli et al.(1998)]. Transport measurements on single crystals of TaSe₂ reveal that disorder enhances superconductivity by suppressing other competing orders such as CDW [Li et al.(2017)Li, Deng, Wang, Liu, Abeykoon, Dooryhee, Tomic, Huang, Warren, Bozin et al.]. Collectively, these experimental results establish TMDs as a distinct class of correlated materials in which reduced dimensionality, band narrowing, and lattice reconstruction cooperate to produce metal–insulator behavior beyond conventional band theory.

Iron-Based Superconductors The iron-based superconductors (FeSCs), discovered in 2008, constitute a broad family of layered materials that share a common structure: a layer of iron atoms joined by tetrahedrally coordinated pnictogen (P, As) or chalcogen (S, Se, Te) anions arranged in a stacked sequence; the layers are separated by alkali, alkaline-earth or rare-earth and oxygen/fluorine ‘blocking layers’ [Kamihara et al.(2008)Kamihara, Watanabe, Hirano, and Hosono, Paglione and Greene(2010), Hirschfeld et al.(2011)Hirschfeld, Korshunov, and Mazin, Stewart(2011)]. Doping or applying external pressure reveals coupled antiferromagnetic and structural transitions along with superconducting and nematic orders [Johnston(2010), Kasahara et al.(2012)Kasahara, Shi, Hashimoto, Tonegawa, Mizukami, Shibauchi, Sugimoto, Fukuda, Terashima, Nevidomskyy et al., Fernandes et al.(2014)Fernandes, Chubukov, and Schmalian]. Electrical resistivity measurements in the neighbourhood of the quantum critical point show a crossover from a bad-metal behavior (T –linear resistivity) to a Fermi liquid behaviour (T^2 –resistivity) [Analytis et al.(2014)Analytis, Kuo, McDonald, Wartenbe, Rourke, Hussey, and Fisher]. Optical conductivity measurements indicate point to intermediate-to-strong electronic correlations [Qazilbash et al.(2009)Qazilbash, Hamlin, Baumbach, Zhang, Singh, Maple, and Basov].

ARPES measurements reveal a k –dependent superconducting gap and de Haas-van Alphen measurements indicate substantial quasiparticle mass enhancements, shedding more light on the

correlated nature of the material [Shishido et al.(2010)Shishido, Bangura, Coldea, Tonegawa, Hashimoto, Kasahara, Rourke, Ikeda, Terashima, Settai et al., Terashima et al.(2009)Terashima, Sekiba, Bowen, Nakayama, Kawahara, Sato, Richard, Xu, Li, Cao et al.]. Strain-tuning measurements of the nematic susceptibility, and differential elastoresistance measurements have provided compelling evidence for the nematic transition and its surrounding fluctuations being the cause of the structural transition as well as the superconductivity [Chu et al.(2012)Chu, Kuo, Analytis, and Fisher, Kuo et al.(2016)Kuo, Chu, Palmstrom, Kivelson, and Fisher]. Neutron scattering and NMR measurements further reveal persistent low-energy spin fluctuations across wide regions of the phase diagram, supporting scenarios in which unconventional superconductivity emerges from the interplay of magnetic, orbital, and nematic degrees of freedom [Inosov et al.(2010)Inosov, Park, Bourges, Sun, Sidis, Schneidewind, Hradil, Haug, Lin, Keimer et al., Chubukov(2012)].

Heavy-Fermion / Kondo-Lattice Systems Heavy-fermion materials are a class of strongly correlated electron systems typically realized in intermetallic compounds of rare earth metals (such as Ce, Sm and Yb) and actinides (such as U, Np and Pu), where localized f electrons hybridize with itinerant conduction bands [Stewart(1984), Wirth and Steglich(2016)]. At high temperatures, these systems exhibit local-moment behavior governed by Curie–Weiss susceptibilities and incoherent transport, while at low temperatures the Kondo effect leads to the formation of a coherent many-body state with extremely large quasiparticle effective masses [Hewson(1993a)]. Experimentally, this crossover is most clearly manifested in transport and thermodynamic measurements: the electrical resistivity shows a characteristic maximum followed by a rapid drop signaling the onset of coherence, while the linear specific heat coefficient $\gamma = C/T$ reaches values orders of magnitude larger than in conventional metals, directly reflecting the heavy quasiparticle mass [Löhneysen et al.(2007)Löhneysen, Rosch, Vojta, and Wölfle].

A defining feature of heavy-fermion materials is their proximity to magnetic quantum phase transitions, which can be accessed experimentally by pressure, chemical substitution, or magnetic field [Gegenwart et al.(2008)Gegenwart, Si, and Steglich]. Near such quantum critical points (QCPs), transport measurements reveal pronounced non-Fermi-liquid behavior (T –linear resistivity) and a vanishing Hall response, along with a divergent effective quasiparticle mass as the transition is approached [Custers et al.(2003)Custers, Gegenwart, Wilhelm, Neumaier, Tokiwa, Trovarelli, Geibel, Steglich, Pépin, and Coleman, Paschen et al.(2004)Paschen, Lühmann, Wirth, Gegenwart, Trovarelli, Geibel, Steglich, Coleman, and Si]; these are all indications of the breakdown of electronic excitations near the transition. Neutron scattering experiments provide direct evidence for spin fluctuations down to very low temperatures, interplaying with the itinerant electrons at the QCP and potentially driving the superconducting transition [Schröder et al.(2000)Schröder, Aeppli, Coldea, Adams, Stockert, Löhneysen, Bucher, Ramazashvili, and Coleman, Stockert et al.(2011)Stockert, Arndt, Faulhaber, Geibel, Jeevan, Kirchner, Loewenhaupt, Schmalzl, Schmidt, Si et al.]. Quantum oscillation measurements through de Haas–van Alphen experiments further demonstrate Fermi surface reconstruction across the quantum critical point, highlighting the drastic change in Fermi volume [Shishido et al.(2005)Shishido, Settai, Harima, and Ōnuki]. Together, these experimental observations establish heavy-fermion compounds as paradigmatic systems in which strong local correlations, hybridization-driven coherence, and quantum critical fluctuations conspire to produce emergent low-energy quasiparticles and unconventional metallic behavior [Si and Steglich(2010)].

1.4.4 Generator of strong local correlations: Quantum impurity models

Within this thesis, we have developed an auxiliary model approach that combines the physics of quantum impurity models with a manybody version of Bloch's theorem to investigate interacting fermionic lattice models. This is motivated by the relative ease in solving impurity models and transparency afforded by such solutions. We now review the phenomenology of some of these impurity models and point out how the various impurity features affect the physics of the lattice models.

The single-impurity Anderson model describes the dynamics of a system of non-interacting itinerant electrons ($c_{k,\sigma}$) hybridising with a localised correlated impurity orbital (d_σ) [Anderson(1959)]:

$$H_{\text{AIM}} = \sum_{k,\sigma} \epsilon_k n_{k,\sigma} + \sum_{k,\sigma} V_k (c_{k,\sigma}^\dagger d_\sigma + \text{h.c.}) + \epsilon_d \sum_{\sigma} n_{d\sigma} + U n_{d\uparrow} n_{d\downarrow} . \quad (4.18)$$

V_k controls the hybridisation strength, and ϵ_d and U control the onsite energy and local correlation respectively of the impurity site. In the limit of large U , the impurity dynamics gets projected onto its spin states, and the model then describes the scattering of non-interacting electrons off magnetic impurities, through the Kondo model [Kondo(1964)]. Formally, the Kondo model is obtained by carrying out a Schrieffer-Wolff transformation on the Anderson impurity model and then using the condition to $V \ll U$ to drop all contributions beyond second order in the ratio V^2/U [Schrieffer and P.A.Wolff(1966)]:

$$H_{\text{Kondo}} = \sum_{k,\sigma} \epsilon_k n_{k,\sigma} + J \sum_{k_1,k_2} \sum_{\alpha,\beta} \vec{S}_d \cdot \vec{\sigma}_{\alpha,\beta} c_{k_1,\alpha}^\dagger c_{k_2,\beta} , \quad (4.19)$$

where $\vec{S}_d$ is the impurity spin formed by projecting the Hilbert space of the impurity to the singly-occupied states.

Even though the conduction bath is non-interacting, the presence of local repulsion on the impurity site makes the problem *many-electron* in nature: it makes it impossible to study the electrons in isolation. This can be seen as follows: imagine two conduction electrons $|k_1 \uparrow\rangle$ and $|k_2 \downarrow\rangle$ trying to hybridise with an empty impurity site. The first electron moves onto the impurity site through a scattering process of the form $c_{d\uparrow}^\dagger c_{k_1\uparrow}$ and makes the impurity site single occupied ($|\uparrow\rangle_d$), but now the second electron k_2 *must pay a cost* of the order of U in order to hybridise with this singly-occupied impurity site through a process of the kind $c_{d\downarrow}^\dagger c_{k_2\downarrow}$. This shows that the conduction electrons *are aware* of the other electrons, and all of them must be treated together. Another way of seeing this collective behaviour is from the Kondo model. Again imagine two spin-up electrons k_1, k_2 that are about to scatter off the impurity spin (which is presently down) via a *spin-flip* process. The first electron scatters and leaves the impurity spin up while it itself becomes down, but now the second conduction electron cannot spin-flip scatter because the spins cannot be exchanged (both are down!). This is another example of how the conduction electrons can *feel* the presence of the other electrons, and this is what makes the problem many-electron in nature.

Kondo screening Despite the interacting nature of the problem, the problem can be solved in an effectively exact fashion, via Wilson's numerical renormalisation group approach [Wilson(1975a),

Krishna-murthy et al.(1980)Krishna-murthy, Wilkins, and Wilson], density-matrix renormalisation group approach [Nishimoto and Jeckelmann(2004)] or the method of *Bethe ansatz* [Andrei(1980), Wiegmann(1981)]. The former solution is more transparent: for any non-zero value of the hybridisation V (or Kondo coupling J), the low energy physics of the impurity is described by a strong-coupling fixed point ($V = \infty$, or $J = \infty$), where the impurity forms a maximally-entangled singlet state with the local spin density of the conduction bath. The low-energy excitations about such a fixed point can be calculated from renormalised perturbation theory, and describe a local Fermi liquid – one particle excitations in the neighbourhood of the singlet renormalised by a local repulsion term [Nozieres(1974), Affleck(1995)]. The crossover from the free local moment at high temperatures to the screened non-magnetic state at low temperatures occurs at an emergent energy scale T_K , the Kondo temperature, which depends exponentially on the strength of the Kondo coupling, and is the only energy scale remaining at low temperatures, so that all thermodynamic quantities must go as a function of T/T_K [Wilson(1975a)].

At half-filling (attained for $\epsilon_d = -U/2$), the only other fixed points are the free orbital fixed point at $V = 0, U = \epsilon_d = 0$ where the impurity site is freely occupied by two independent spins, and the local moment fixed point at $V = 0, U = \infty$, where the impurity site is occupied by a local moment that is decoupled from the bath [Krishna-murthy et al.(1980)Krishna-murthy, Wilkins, and Wilson, Hewson(1993a)]. Away from half-filling, two other fixed points become available [Krishna-murthy et al.(1980)Krishna-murthy, Wilkins, and Wilson, Hewson(1993a)]: a valence fluctuation fixed point at $V = 0, \epsilon_d = 0, U = \infty$ where the doubly-occupied state has been projected out from the symmetric free orbital fixed point, and a frozen impurity fixed point at $V = U = 0, \epsilon_d = \infty$ where the impurity site is left completely empty.

The crossover from local moment physics to Kondo screening as the system is tuned from high to low temperatures can also be seen from the impurity spectral function [Costi and Hewson(1990), Costi and Hewson(1991), Takayama and Sakai(1993), Oliveira and Oliveira(1994)]. At low frequencies, the spectral function exhibits a sharp resonance around $\omega = 0$ that indicates the presence of gapless single-particle local Fermi liquid excitations. At higher frequencies, one sees satellite peaks that involve impurity excitations that cost energy of the order of U . The height of the central Kondo resonance is a topological invariant; it is fixed by the impurity occupancy owing to an application of the Friedel sum rule to impurity models [Langreth(1966)], and remains unchanged as the interaction U is increased adiabatically while remaining at particle-hole symmetry. The topological nature can also be ascertained by identifying such a sum rule as the impurity model analogue of Luttinger’s theorem [Luttinger(1960), Luttinger and Ward(1960)] that relates the total number density of electrons to the number of Greens function poles within the Fermi volume, and which is itself topological [Oshikawa(2000), Seki and Yunoki(2017)].

Breakdown of Kondo screening While the vanilla Anderson model (or the Kondo model derived from it) only display a strongly-coupled screened phase, it is possible to tweak it in order to introduce Kondo frustration which can eventually lead to a breakdown in the Kondo screening process. Such models are interesting because they provide a means of studying zero temperature *continuous* phase transitions and address experimental signatures of quantum criticality in correlated systems. The canonical example of this is the multichannel Kondo model, involving a single impurity hybridising

with multiple independent conduction channels [Nozières, Ph. and Blandin, A.(1980)]:

$$H_{\text{MCK}} = \sum_{k,\sigma,\alpha} \epsilon_{k,\lambda} n_{k,\sigma,\lambda} + \sum_{k_1,k_2,\lambda} J_\lambda \sum_{\alpha,\beta} \vec{S}_d \cdot \vec{\sigma}_{\alpha,\beta} c_{k_1,\alpha,\lambda}^\dagger c_{k_2,\beta,\lambda}, \quad (4.20)$$

here λ is the channel index. In the case of an isotropic Kondo coupling across the channels ($J_\lambda = J$), the strong-coupling $J = \infty$ fixed-point of the single-channel Kondo model is replaced by an intermediate-coupling fixed-point at a finite value J^* [Nozières, Ph. and Blandin, A.(1980), Gan(1994)]. Powerful methods like boundary conformal field theory [Affleck and Ludwig(1991), Ludwig and Affleck(1991), Affleck et al.(1992)Affleck, Ludwig, Pang, and Cox, Affleck and Ludwig(1993a),Parcollet and Georges(1997),Affleck(2005)], bosonization [Emery and Kivelson(1992), Clarke et al.(1993)Clarke, Giamarchi, and Shraiman,Zaránd and von Delft(2000), von Delft et al.(1998a) von Delft, Záránd, and Fabrizio,Schofield(1997)], numerical renormalization group [Bulla et al.(2008)Bulla, Costi, and Pruschke, Affleck et al.(1992)Affleck, Ludwig, Pang, and Cox, Pang and Cox(1991)] and Bethe ansatz [Andrei and Destri(1984), Tsvelick and Wiegmann(1984), Tsvelick(1985), Andrei and Jerez(1995), Záránd et al.(2002)Záránd, Costi, Jerez, and Andrei] have shown that the low energy theory of such models exhibit non-Fermi liquid excitations. Further, the system displays anomalous behaviour in thermodynamic properties near $T = 0$ and a fractional zero temperature entropy in the thermodynamic limit (upon ensuring that temperature is taken to zero after the system size is taken to infinity [Rozhkov(1998), von Delft et al.(1998a) von Delft, Záránd, and Fabrizio]). This anomalous divergent behaviour is a signature of the fact that the MCK system with a single channel-symmetric exchange coupling is quantum critical: the ground state is susceptible to perturbations that introduce channel anisotropy in the exchange couplings [Nozières, Ph. and Blandin, A.(1980), Andrei and Jerez(1995), Affleck et al.(1992)Affleck, Ludwig, Pang, and Cox, Záránd and von Delft(2000), Zheng et al.(2021)Zheng, He, and Lu].

Another class of models that shows Kondo breakdown is that of pseudogap Anderson and Kondo models, where the impurity hybridises with a conduction bath whose density of states (DOS) varies as $\rho(\omega) \sim |\omega|^r$ with a paramter $r > 0$ that controls how fast the DOS vanishes as the Fermi level is approached. Such behaviour can arise in semi-metals and in long-range ordered systems where the order parameter vanishes at certain points on the Fermi surface (such as d -wave superconductors). The pseudogap impurity models exhibit a boundary quantum phase transition between a Kondo-screened strong-coupling phase and an unscreened local-moment phase as a function of interaction strength or exchange coupling [Withoff and Fradkin(1990), Gonzalez-Buxton and Ingersent(1998), Bulla et al.(1997)Bulla, Pruschke, and Hewson, Ingersent(1996)]. At the critical Kondo coupling, the impurity spectral function and spin susceptibility exhibit nontrivial power-law scaling with anomalous exponents determined by r [Gonzalez-Buxton and Ingersent(1998), Ingersent and Si(2002), Fritz and Vojta(2004), Parcollet and Georges(1999), Si et al.(2001)Si, Rabello, Ingersent, and Smith]. For $0 < r < 1/2$ (in the particle-hole symmetric case), both the local-moment and strong-coupling phases are stable, separated by a critical fixed point characterized by fractional residual entropy and nontrivial hyperscaling relations [Gonzalez-Buxton and Ingersent(1998), Bulla et al.(2000)Bulla, Vojta, and Hofstetter]. For larger exponents, the strong-coupling phase may disappear entirely, and the system remains in the local-moment phase for all couplings [Ingersent(1996)].

The multi-impurity Anderson and Kondo models furnish another canonical realization of Kondo

breakdown, driven by the competition between Kondo screening and inter-impurity interaction. The paradigmatic example is the two-impurity Kondo model (TIKM), in which two localized spins couple both to conduction electrons and to each other via an antiferromagnetic exchange K [Jones and Varma(1987), Jones et al.(1988) Jones, Varma, and Wilkins]. In the limit of large $K > 0$, the impurities form a non-magnetic singlet decoupled from the conduction electrons and suppressing the Kondo resonance. This fixed point is separated from the strong-coupling Kondo screened fixed point through a quantum critical point [Jones and Varma(1987), Jones et al.(1988) Jones, Varma, and Wilkins, Affleck et al.(1995) Affleck, Ludwig, and Jones] displaying logarithmic or power-law scaling in thermodynamic quantities and fractional residual entropy [Affleck and Ludwig(1993b)]. A modified version of this model that uses conduction bath-mediated inter-impurity interaction leads to a metallic spin liquid displaying nonlocal impurity-spin entanglement and fractionalized charge excitations [Wójcik and Kroha(2023)]. More intricate geometries can be constructed by arranging antiferromagnetically coupled Kondo impurities along a triangle, each hybridising with its own conduction bath; such a model displays a topological phase transition from a symmetry-preserved gapless spin liquid phase with "topological order" to a screened Fermi liquid phase [König et al.(2021) König, Coleman, and Komijani]. Indeed, recent experiments have validated such phase diagrams for the case of two impurities by studying pairs of copper atoms in the presence of strong magnetic fields [Spinelli et al.(2015) Spinelli, Gerrits, Toskovic, Bryant, Ternes, and Otte].

1.5 Mott transition in Hubbard-like models

1.5.1 In the limit of infinite dimensions

The Hubbard model (introduced in eq. 1.1) presents one of the most fruitful playgrounds for investigating correlation-driven metal-insulator transitions and its various facets. This model, despite its conceptual simplicity, exhibits a rich variety of phenomena, including antiferromagnetism, superconductivity, and, most notably for our purposes, the correlation-driven Mott metal–insulator transition (MIT) [Imada et al.(1998a) Imada, Fujimori, and Tokura, Tasaki(1998), Arovas et al.(2022) Arovas, Berg, Kivelson, and Raghu, Qin et al.(2022) Qin, Schäfer, Andergassen, Corboz, and Gull]. While exact solutions exist in the limits of $d = 1$ [Lieb and Wu(1968)] and $d = \infty$ [Georges and Kotliar(1992)], the problem remains largely open in more realistic situations such as two and three spatial dimensions.

As discussed earlier, the limit of infinite dimensions presents a drastic simplification for the manybody scattering processes within the problem: the one-particle k -space self-energy becomes momentum-independent due to the vanishing of all non-local contributions (they scale as negative powers of the coordination number, which diverges in the limit of $d = \infty$). This simplification lies at the heart of dynamical mean-field theory and its ability to capture the Mott metal–insulator transition at both zero and finite temperatures for this limiting case [Georges and Kotliar(1992), Janis and Vollhardt(1992), Rozenberg et al.(1992) Rozenberg, Zhang, and Kotliar, Georges et al.(1996) Georges, Kotliar, Krauth, and Rozenberg, Rozenberg et al.(1999) Rozenberg, Chitra, and Kotliar, Noack and Gebhard(1999)]. At half-filling and zero temperature, DMFT predicts a continuous transition as a function of U wherein the quasiparticle weight Z tends to zero at a critical interaction strength

U_{c2} , signaling the vanishing of coherent Fermi liquid behavior and the opening of a Mott gap in the single-particle spectral function [Rozenberg et al.(1992)Rozenberg, Zhang, and Kotliar, Georges et al.(1996)Georges, Kotliar, Krauth, and Rozenberg]. The three-peak structure of the interacting spectral function, with a central quasiparticle resonance flanked by incoherent lower and upper Hubbard bands, indicates the presence of highly renormalised Landau quasiparticles [Rozenberg et al.(1992)Rozenberg, Zhang, and Kotliar]. A doping-driven continuous metal-insulator transition also exists for this model at zero temperature, marked by the vanishing of electronic compressibility at a critical value of the chemical potential [Werner and Millis(2007)]. The quantum critical point itself (at $T = 0$) is likely to be an exotic state, and important questions about it remain unanswered. *Is it possible to obtain a low-energy theory for the local gapless excitations precisely at the MIT, where the metal is on the brink of destruction? How do these excitations compare with those of the local Fermi liquid, e.g., in terms of self-energies and two-particle correlation functions?*

While the DMFT solution captures the zero temperature transition accurately, the self-consistently determined nature of the conduction bath renders the solution somewhat opaque and makes it difficult to write down an effective Hamiltonian for the low-energy theories. Since the vanilla Anderson impurity model does not contain the mechanism to stabilise a localised phase over a finite-sized parameter regime, an important question arising from the DMFT solution is: *what are the minimal changes that one must implement into the Anderson impurity model in order to capture such a metal-insulator transition (1/2-filled Hubbard model on the Bethe lattice in $d = \infty$), in sufficient accuracy and precision?* Finding such an impurity model would also reduce the considerable computational effort that is presently required in self-consistent approaches.

At finite temperatures, the interaction-driven as well as doping-driven Mott transition become first order, and is characterized by a region of phase coexistence and hysteresis [Nozières, Ph.(1998), Rozenberg et al.(1999)Rozenberg, Chitra, and Kotliar]. Two spinodal lines, $U_{c1}(T)$ and $U_{c2}(T)$, emerge; within this region metallic and insulating DMFT solutions coexist, and responses such as resistivity show a significant discontinuity [Kotliar et al.(2002)Kotliar, Murthy, and Rozenberg]. These lines meet at a finite-temperature critical endpoint (U_c, T_c), above which there is no sharp distinction between metallic and insulating solutions and only a crossover exists between the Mott insulator and the Fermi liquid through a bad-metal regime, as the chemical potential is tuned [Rozenberg et al.(1999)Rozenberg, Chitra, and Kotliar, Kotliar, G.(1999), Werner and Millis(2007), Deng et al.(2013)Deng, Mravlje, Žitko, Ferrero, Kotliar, and Georges, Žitko et al.(2013)Žitko, Hansen, Perepelitsky, Mravlje, Georges, and Shastry]. The coexistence of metallic and insulating phases shows that the insulating solution is present within the many-body spectrum of the metallic phase. This naturally leads to the following question: *since a quantum impurity lies at the heart of the self-consistency procedure, what does the presence of coexistence region mean for the Hamiltonian dynamics and the spectrum of this impurity model? Can the impurity model Hamiltonian display the emergence of the insulating state prior to the transition, through changes in the eigenstates?* Analytic insight of this kind is particularly important because approaching a coexistence region numerically is often fairly tricky.

The bad metal regime displays anomalous linear-in-temperature resistivity which corresponds to quantum critical ω/T scaling, along the quantum widom line [Vučićević et al.(2015)Vučičević, Tanasković, Rozenberg, and Dobrosavljević, Dasari et al.(2017)Dasari, Vidhyadhiraja, Jarrell, and McKenzie]. This is accompanied by the disappearance of the Drude peak in optical conductivity,

signalling the loss of quasiparticle transport [Vučićević et al.(2015)Vučičević, Tanasković, Rozenberg, and Dobrosavljević]. Similar quantum critical scaling is also obtained from DMFT solutions at very low temperatures, hinting at a "hidden" quantum criticality that is continuously connected to and responsible for the critical behaviour above the finite temperature endpoint [Eisenlohr et al.(2019)Eisenlohr, Lee, and Vojta]. Luttinger's theorem, which relates the poles of the Greens function within the Fermi volume to the number of electrons in the system at zero temperature, is satisfied within the Fermi liquid phase but is violated in the particle-hole asymmetric Mott insulator by a universal magnitude [Logan and Galpin(2015)]. The emergence of a quantum critical regime at high temperatures and its connection to similar signatures at low temperatures suggests the presence of quantum fluctuations in the underlying impurity model that modify the impurity-bath scattering processes and the associated impurity responses. One can therefore ask: *What are the fluctuations that destroy the metal and lead to the insulating phase? Can we obtain a universal theory for these competing tendencies?*

The above studies of single band models describe the Mott transition to be of the Mott-Hubbard kind (described in previous sections), where the size of the gap is determined by the double-occupancy cost U . In order to obtain transition of the charge-transfer kind, one needs to consider at least a two-band model. Indeed, in a non-degenerate two-band Hubbard model with a charge-transfer energy Δ for hopping from a d -orbital to a p -orbital, one sees a Mott-Hubbard transition when $U < \Delta$ and a charge-transfer transition when $U > \Delta$ [Ōno et al.(2001)Ōno, Bulla, and Hewson]. The degenerate two-band model with inter-band local correlation shows successive Mott-Hubbard transitions at low temperatures whenever the filling passes through integer values and the interaction strength is intermediate [Rozenberg(1997)].

1.5.2 The two-dimensional Hubbard model

The 2D Hubbard model is one of the most well-studied models in condensed matter physics, particularly because of its paradigmatic role in Mott metal-insulator transition and its connections to the physics of copper oxide materials and d -wave superconductivity. Photoemission experiments reveal the presence of p and d bands close to the Fermi surface in such materials [Damascelli et al.(2003)Damascelli, Hussain, and Shen]; further work by Zhang and Rice showed that a specific combination of these orbitals governs the low-energy physics, leading to an effective description in terms of a single band Hubbard model [Zhang and Rice(1988)]. The discovery of high-temperature superconductivity in nickelates and organic charge transfer salts has provided more platforms where the Hubbard model and its extensions can prove to be useful [Kitatani et al.(2020)Kitatani, Si, Jansson, Arita, Zhong, and Held, Powell and McKenzie(2006)].

While simple in appearance, the model has proved to be notoriously hard to solve except in highly specific parameter regimes, despite several decades of computational effort. This is due to several reasons [Qin et al.(2022)Qin, Schäfer, Andergassen, Corboz, and Gull]: (i) the presence of closely-lying energy scales arising from competing tendencies, (ii) the inability of several methods to either reach zero temperature or extrapolate to the thermodynamic limit, (iii) the tendency of approximate/finite-size methods to overestimate proclivities towards spontaneous symmetry breaking, and (iv) the presence of differing phases separated by slow crossovers without any sharp boundaries.

At half-filling, the 2D Hubbard model has a spectral gap to charge excitations for any non-zero interaction strength [Hirsch(1985), White et al.(1989)White, Scalapino, Sugar, Loh, Gubernatis, and Scalettar, Schäfer et al.(2015)Schäfer, Geles, Rost, Rohringer, Arrigoni, Held, Blümer, Aichhorn, and Toschi, Vitali et al.(2016)Vitali, Shi, Qin, and Zhang, Šimkovic et al.(2020)Šimkovic, LeBlanc, Kim, Deng, Prokof'ev, Svistunov, and Kozik]. The behaviour below the charge gap is described by a nonlinear sigma model with parameters that depend on the interaction strength U [Borejsza and Dupuis(2004)]. Upon tuning to finite temperatures, the system goes through a sequence of crossovers from the insulating ordered state to a Fermi liquid state with coherent quasiparticles into a non-Fermi liquid state with partial gapping of the Fermi surface [Kim et al.(2020)Kim, Simkovic, and Kozik, Schäfer et al.(2015)Schäfer, Geles, Rost, Rohringer, Arrigoni, Held, Blümer, Aichhorn, and Toschi, Šimkovic et al.(2020)Šimkovic, LeBlanc, Kim, Deng, Prokof'ev, Svistunov, and Kozik]. Away from half-filling but at weak coupling, the Neel order is accompanied by incommensurate magnetic orders (also called spiral magnetic state) with wave vectors away from (π, π) , as seen from mean-field studies [Schulz(1990), Frésard et al.(1991)Frésard, Dzierzawa, and Wölfle]. Functional RG studies indicate that such solutions can exist up to fairly high doping if the superconducting instability is suppressed [Metzner et al.(2012)Metzner, Salmhofer, Honerkamp, Meden, and Schönhammer, Yamase et al.(2016)Yamase, Eberlein, and Metzner]. Superconductivity with d -wave pairing has been achieved from renormalised perturbation studies as well as from diagrammatic Monte Carlo (diagMC) and functional RG (fRG) calculations [Raghu et al.(2010)Raghu, Kivelson, and Scalapino, Šimkovic et al.(2021)Šimkovic, Deng, and Kozik, Wang et al.(2014)Wang, Eberlein, and Metzner].

The phenomena of pseudogap, where the Fermi surface acquires a momentum-dependent suppression of spectral weight, has also been observed above a certain temperature from methods like diagMC and fRG [Šimkovic et al.(2020)Šimkovic, LeBlanc, Kim, Deng, Prokof'ev, Svistunov, and Kozik, Hille et al.(2020a)Hille, Rohe, Honerkamp, and Andergassen]. The gap is seen to open first near the antinodal point $(\pi, 0)$ and spread across the Fermi surface until it reaches the node $(\pi/2, \pi/2)$. The mechanism of the pseudogap at weak to intermediate coupling is found to be long-range antiferromagnetic correlations [Wu et al.(2018)Wu, Scheurer, Chatterjee, Sachdev, Georges, and Ferrero]. The pseudogap disappears at large doping when the Fermi surface undergoes a topological transition from hole-like to electron-like. At strong coupling, the competition between various phases such as d -wave superconductivity, antiferromagnetic order and stripe states (spin/charge order modulating in real space) is an active subject of study. Recent studies using various methods like DMRG and iPEPS found that the period 8 charge stripe state is lower in energy than the superconducting state at strong coupling $U/t = 8$ and no next-nearest-neighbour hopping [Zheng et al.(2017)Zheng, Chung, Corboz, Ehlers, Qin, Noack, Shi, White, Zhang, and Chan]. This was later corroborated by studies from variational Monte Carlo and other DMRG studies [Tocchio et al.(2019)Tocchio, Montorsi, and Becca, Jiang et al.(2020)Jiang, Zaanen, Devereaux, and Jiang]. At weaker coupling, the above methods report a reduced stability for the strip state and increased tendency to realise a d -wave SC state [Qin et al.(2020)Qin, Chung, Shi, Vitali, Hubig, Schollwöck, White, and Zhang, Šimkovic et al.(2021)Šimkovic, Deng, and Kozik].

1.6 Brief introduction to the tiling auxiliary model approach

1.6.1 Why do we need another auxiliary model method? Limitations of existing approaches

1.6.2 Overview

1.6.3 Features

1.7 Entanglement holography and fermionic criticality

Role of entanglement in fermionic systems The present millennium has seen a rapid surge in the use of quantum entanglement towards the study of quantum and condensed matter systems [Terhal et al.(2003)Terhal, Wolf, and Doherty,Horodecki et al.(2009)Horodecki, Horodecki, Horodecki, and Horodecki,Eisert et al.(2010)Eisert, Cramer, and Plenio,Laflorie(2016a),Zeng et al.(2019)Zeng, Chen, Zhou, and Wen,Erhard et al.(2020)Erhard, Krenn, and Zeilinger]. Being non-local by nature, it is helpful in tracking correlations among particles across both small and large distances, giving valuable insight into the nature of the ground state and the low-energy excitations. Entanglement, in the form of various measures like entanglement entropy and mutual information, is the engine that gives rise to new low-energy degrees of freedom from a renormalisation group (RG) perspective, and is therefore key to understanding and characterising phase transitions and criticality in quantum matter. For instance, entanglement entropy is at the heart of understanding the so-called topological states of matter that can be described in terms of a purely topological effective Lagrangian and have a robust ground state degeneracy and gapless edge states [Wen(1990), Wen and Niu(1990), Wen and Zee(1990), Wen(1991), Wen(1995), Wen(2013), Wen(2017), Tao and Wu(1984), Niu et al.(1985)Niu, Thouless, and Wu, Wen(1989), Li and Haldane(2008), Qi et al.(2012a)Qi, Katsura, and Ludwig, Pollmann et al.(2010)Pollmann, Turner, Berg, and Oshikawa, Yao and Qi(2010), Liu et al.(2011a)Liu, Guo, Vedral, and Fan, Calabrese and Lefevre(2008), Läuchli et al.(2010)Läuchli, Bergholtz, Suorsa, and Haque, Schliemann(2011), Halperin(1982), Qi et al.(2012b)Qi, Katsura, and Ludwig, Laflorie(2016b)]. In contrast to systems with local interactions whose subsystem entanglement entropy scales with the area of the subsystem [Das and Shankaranarayanan(2006), Cramer et al.(2006)Cramer, Eisert, Plenio, and Dreißig, Eisert et al.(2010)Eisert, Cramer, and Plenio], topologically ordered systems have an additional subdominant topological term dubbed the topological entanglement entropy [Blok and Wen(1990), Read(1990), Rokhsar and Kivelson(1988), Read and Chakraborty(1989), Moessner and Sondhi(2001), Ardonne et al.(2004)Ardonne, Fendley, and Fradkin, Levin and Wen(2005), Kitaev and Preskill(2006a), Jiang et al.(2012)Jiang, Wang, and Balents, Hamma et al.(2005)Hamma, Ionicioiu, and Zanardi, Levin and Wen(2006), Kitaev and Preskill(2006b), Zhang et al.(2012)Zhang, Grover, Turner, Oshikawa, and Vishwanath, Grover et al.(2011)Grover, Turner, and Vishwanath, Hamma and Lidar(2008), Lan and Wen(2014), Fendley et al.(2013)Fendley, Isakov, and Troyer, Gu et al.(2009)Gu, Levin, Swingle, and Wen, Slagle et al.(2019)Slagle, Aasen, and Williamson] that quantifies the long-ranged nature of the entanglement in such systems. The non-local nature arises from the fact that the ground state of such a system cannot be made separable (non-entangled) by the application of purely local unitary transformations [Chen et al.(2010)Chen, Gu, and Wen, Gu

and Wen(2009), Zeng et al.(2019)Zeng, Chen, Zhou, and Wen].

More pertinent to the work at hand, however, is the entanglement in free fermionic systems. In this thesis, we will consider relativistic Dirac fermions in two dimensions, both massless and massive. They emerge in several condensed matter systems [Vafeek and Vishwanath(2014a)], appearing at the surfaces of three-dimensional topological insulators [Bernevig and Hughes(2013), Cayssol(2013), Moessner and Moore(2021), Rachel(2018)], quasi-2D organic materials [Wang et al.(2015)Wang, Deng, Liu, and Liu, Wu et al.(2016)Wu, Wang, Liu, Li, Fu, Sun, Liu, Pan, Weng, Dincă et al., Ni et al.(2022)Ni, Huang, and Brédas] and in artificial materials like cold atoms, molecular and microwave crystals [Zhu et al.(2007)Zhu, Wang, and Duan, Lepori et al.(2010)Lepori, Mussardo, and Trombettoni, Boada et al.(2011)Boada, Celi, Latorre, and Lewenstein, Kuno et al.(2018)Kuno, Ichinose, and Takahashi, Gomes et al.(2012)Gomes, Mar, Ko, Guinea, and Manoharan, Yuan et al.(2016)Yuan, Zhang, Liu, Narayan, Song, Shen, Sui, Xu, Yu, An et al., Bellec et al.(2013)Bellec, Kuhl, Montambaux, and Mortessagne]. For massless critical systems (or conformal field theories (CFTs)) living in one spatial dimension, the entanglement entropy of a single subsystem of length l scales as $\sim \frac{c}{3} \ln l$ [Vidal et al.(2003)Vidal, Latorre, Rico, and Kitaev, Latorre et al.(2003)Latorre, Rico, and Vidal, Latorre et al.(2005)Latorre, Lütken, Rico, and Vidal, Jin and Korepin(2004), Lambert et al.(2004)Lambert, Emary, and Brandes, Holzhey et al.(1994)Holzhey, Larsen, and Wilczek, Casini and Huerta(2004), Calabrese and Cardy(2004), Osborne and Nielsen(2002), Osterloh et al.(2002)Osterloh, Amico, Falci, and Fazio], where c is the conformal central charge of the CFT. The case of multiple subsystems is more involved, but has been solved [Casini et al.(2005)Casini, Fosco, and Huerta, Calabrese and Cardy(2004), Korepin(2004)]. For massive fermions, the entanglement entropy (or rather the entropic c -function, which is the derivative of the entropy along the distance) can be expressed in the form of differential equations which need to be solved numerically [Casini et al.(2005)Casini, Fosco, and Huerta, Cardy et al.(2008)Cardy, Castro-Alvaredo, and Doyon, Casini and Huerta(2009a), Doyon(2009)]. The solutions lend themselves to small and large mass expansions, reducing to $\frac{1}{3} \ln \frac{l}{\epsilon} - \frac{1}{6} (ml \ln ml)^2 + O((ml)^2)$ for the case of small ml , and to $\sim (ml)^{-1/2} e^{-2ml}$ for that of large ml , both for a single interval. The $\ln l$ behaviour is a specific case of the more general $l^{d-1} \ln l$ violation of the area law, as observed in d -dimensional fermionic critical systems [Casini et al.(2005)Casini, Fosco, and Huerta, Casini and Huerta(2009b), Gioev and Klich(2006), Wolf(2006), Li et al.(2006)Li, Ding, Yu, Roscilde, and Haas, Barthel et al.(2006)Barthel, Chung, and Schollwöck, Farkas and Zimborás(2007), Wolf et al.(2008)Wolf, Verstraete, Hastings, and Cirac, Castro-Alvaredo and Doyon(2009), Swingle(2010), Helling et al.(2010)Helling, Leschke, and Spitzer, Calabrese et al.(2012a)Calabrese, Mintchev, and Vicari, Swingle(2012a), Calabrese et al.(2012b)Calabrese, Mintchev, and Vicari, Tan and Ryu(2020), Murciano et al.(2020)Murciano, Ruggiero, and Calabrese]. This violation arises from the non-local nature of the quantum fluctuations that exist in quantum critical fermionic systems.

While there is no geometry-independent subdominant term in the entanglement entropy of free fermionic systems [Chen et al.(2017)Chen, Witczak-Krempa, Faulkner, and Fradkin, Bueno and Witczak-Krempa(2017)], it is possible to generate such a term by placing the system on a manifold with a non-trivial topology, say a cylinder, and inserting a magnetic flux through the cylinder [Metlitski et al.(2009)Metlitski, Fuertes, and Sachdev, Arias et al.(2015)Arias, Blanco, and Casini, Whitsitt et al.(2017)Whitsitt, Witczak-Krempa, and Sachdev, Chen et al.(2017)Chen, Witczak-Krempa, Faulkner, and Fradkin, Zhu et al.(2018)Zhu, Chen, He, and Witczak-Krempa]. For a massless

Dirac fermion, if there are ϕ number of electronic flux quanta piercing the system, the additional term that is generated in the entanglement entropy of a subsystem of sufficiently (as defined in Ref. [Chen et al.(2017)Chen, Witczak-Krempa, Faulkner, and Fradkin]) large length is of the form $\frac{1}{6} \ln \left| 2 \sin \frac{\phi}{2} \right|$ [Chen et al.(2017)Chen, Witczak-Krempa, Faulkner, and Fradkin, Arias et al.(2015)Arias, Blanco, and Casini]. There is evidence from numerical computations on discrete models that such a term is a signature of the specific field theory and encodes universality [Zhu et al.(2018)Zhu, Chen, He, and Witczak-Krempa, Hu et al.(2019)Hu, Zhu, Eggert, and He, Crépel et al.(2021)Crépel, Hackenbroich, Regnault, and Estienne]. In this thesis, we put this idea on a stronger analytical footing by relating the flux-dependent term to a topological invariant of the field theory, its Luttinger volume [Luttinger(1960), Luttinger and Ward(1960), Oshikawa(2000), Seki and Yunoki(2017), Heath and Bedell(2020a), Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)]. Given by the number of k -states enclosed by the Fermi surface, Luttinger volume V_L has proved to be one of the most important indicators that distinguish metals from insulating states of matter. In metals, V_L is constrained by the number of particles in the system, following Luttinger’s theorem [Luttinger(1960), Luttinger and Ward(1960), Oshikawa(2000)]. As shown most prominently in Ref. [Oshikawa(2000)], Luttinger’s theorem depends crucially on the presence of extended states in the metal that can respond to flux insertion experiments. We demonstrate that indeed it is this subdominant term in the entanglement entropy that carries this dependence and hence encodes Luttinger’s theorem. It is known that for strongly-interacting states like Mott insulators or quantum hall states, V_L is modified due to changes in the topology of the Fermi surface [Heath and Bedell(2020a), Seki and Yunoki(2017), Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)]. In this context, we relate the topological invariant on the metallic side (Luttinger volume) to the topological invariants (the Chern numbers) in the quantum hall system that the metal would lead to when placed in a perpendicular magnetic field. Such a relation makes explicit the transmutation of the extended states into the localised states of the insulating phase.

Entanglement hierarchy and the holographic principle The leading geometry-dependent piece is also found to have intriguing properties of its own. The fact that the free fermion Lagrangian is diagonal in momentum space means that the entanglement arises purely from real space quantum fluctuations, and this allows us to study the entanglement of subsystems of varying sizes in momentum space. We find that the entanglement entropy hence obtained has a Russian doll-like nested structure, such that increasing the size of the subsystem in momentum space gradually reveals layers of entanglement. In fact, we argue that such transformations between the subsystems amount to a renormalisation group (RG) flow in the space of the field theories with varying spectral gaps, and the above-mentioned hierarchy in the entanglement entropy then stretches across the RG transformations. The precise prescription for choosing the subsystems is described in Sec. 7.3. We also show, simultaneously, that the hierarchy holds across multipartite entanglement measures, leading to an onion-like structure where increasing the number of parties in the entanglement measure (mutual information, tripartite information, 4-partite information and so on) leads to additional entanglement that did not exist before. More importantly, we interpret the hierarchy of entanglement along the RG flow as the emergence of an additional dimension [Hooft(1993), Verlinde(2011), Chiang et al.(2016)Chiang, Hu, and Chen], allowing us to define a measure of distance using some non-negative measure of entanglement, like the mutual information [Van Raamsdonk(2010), Raams-

donk(2017), Qi(2013), Lee and Qi(2016)]: the more the entanglement, the less will be the distance between those points. Generation of an additional dimension is, of course, the essence of the AdS/CFT conjecture or, more generally, the holographic principle.

The holographic principle [Bousso(2002), Katz and Sheng(2010), Green(2013), Hartnoll(2009), Sachdev(2012), Zaanen et al.(2015)Zaanen, Liu, Sun, and Schalm, Joseph(2015), Năstase(2015), Erdmenger(2018)], also known as the gauge-gravity duality, posits that there exists a duality relation between a quantum field theory with a gauge field, and a gravity theory having one additional spatial dimension [Hooft(1974), Polyakov(1978), Polyakov(1987), Gross and Taylor(1993)]. The additional dimension can be visualised as a renormalisation group flow that starts from the boundary field theory and extends into the bulk [Akhmedov(1998), Álvarez and Gómez(1999), Freedman et al.(1999)Freedman, Gubser, Pilch, and Warner, Porrati and Starinets(1999), Kachru et al.(2000)Kachru, Schulz, and Silverstein, Girardello et al.(2000)Girardello, Petrini, Porrati, and Zaffaroni, De Boer et al.(2000)De Boer, Verlinde, and Verlinde, de Boer(2001), Johnson et al.(2001)Johnson, Lovis, and Page, Yamaguchi(2003)]. The most popular realisation of the holographic principle is the AdS/CFT correspondence that links a super-symmetric gauge theory with a string theory [Maldacena(1999), Gubser et al.(1998)Gubser, Klebanov, and Polyakov, Witten(1998)]. The essence of holography is that all the information about the bulk gravity theory is already contained within the boundary gauge theory; this idea is very similar to the discovery by Bekenstein and Hawking [Bekenstein(1973), Hawking(1974), Hawking(1975)] that the entropy of a black hole is determined by its surface area. In fact, Ryu and Takayanagi proposed a generalisation of this result where the entanglement entropy of subsystem region a CFT is determined by the area of the minimal surface that bounds the region in the bulk [Ryu and Takayanagi(2006a), Ryu and Takayanagi(2006b)]. The AdS/CFT correspondence happens to be a strong-weak duality - it relates a strongly coupled CFT to a theory of weak classical gravity. This allows physicists to use the holographic principle and convert a problem of strong quantum correlation to one of classical gravity which can be solved using well-known methods [Hartnoll(2009), Sachdev(2012)]. This has led to new insights in various topics of high energy and condensed matter physics, such as the viscosity of the quark-gluon plasma [Kovtun et al.(2005)Kovtun, Son, and Starinets], the response functions of the Bose-Hubbard model [Damle and Sachdev(1997), Hartnoll et al.(2007)Hartnoll, Kovtun, Müller, and Sachdev], and the phenomenology of non-Fermi liquids [Lee(2009), Liu et al.(2011b)Liu, McGreevy, and Vegh, Faulkner et al.(2011)Faulkner, Liu, McGreevy, and Vegh, Čubrović et al.(2009)Čubrović, Zaanen, and Schalm], holographic superconductors [Gubser et al.(2010)Gubser, Pufu, and Rocha, Gauntlett et al.(2009)Gauntlett, Sonner, and Wiseman, Hartnoll et al.(2008a)Hartnoll, Herzog, and Horowitz, Hartnoll et al.(2008b)Hartnoll, Herzog, and Horowitz, Cai et al.(2015)Cai, Li, Li, and Yang, Wang et al.(2020)Wang, Zhang, Fu, and Wu], striped phases [Donos and Gauntlett(2011a), Donos(2013), Cremonini et al.(2018)Cremonini, Li, and Ren] and holographic Fermi liquids [Kachru et al.(2008)Kachru, Liu, and Mulligan, Čubrović et al.(2011)Čubrović, Liu, Schalm, Sun, and Zaanen, Senthil et al.(2003)Senthil, Sachdev, and Vojta, Sachdev(2011), "Hartnoll et al.('2011)"Hartnoll, Hofman, and Vegh]. An overview of the metallic states (phases that satisfy Luttinger's theorem) and their connections to the gauge-gravity duality have been provided in Refs. [Huijse and Sachdev(2011), Aprile et al.(2011)Aprile, Roest, and Russo, Donos and Gauntlett(2011b)].

Of course, on the flip side, the holographic principle also implies that studying the boundary conformal theory can provide insights into the nature of the bulk quantum gravity theory. This

however requires constructing, *ab initio*, the emergent dimension by starting from the field theory, and has proved to be more difficult than the converse, mostly because the field theories are often strongly coupled. Nevertheless, there exist some examples of constructive attempts towards a bulk gravity theory [Matsueda(2012), Matsueda et al.(2013) Matsueda, Ishihara, and Hashizume, Matsueda(2013), Matsueda(2014), Czech(2018), Faulkner et al.(2017) Faulkner, Haehl, Hijano, Parrikar, Rabideau, and Van Raamsdonk, Faulkner et al.(2014) Faulkner, Guica, Hartman, Myers, and Van Raamsdonk, Gu et al.(2016) Gu, Lee, Wen, Cho, Ryu, and Qi, Wen et al.(2016) Wen, Cho, Lopes, Gu, Qi, and Ryu, Leigh et al.(2015) Leigh, Parrikar, and Weiss, Leigh et al.(2014) Leigh, Parrikar, and Weiss, Douglas et al.(2011) Douglas, Mazzucato, and Razamat, de Mello Koch et al.(2011) de Mello Koch, Jevicki, Jin, and Rodrigues, Lee(2010a), Lee(2010b), Lee(2012), Lee(2014), Vidal(2007), Vidal(2008), Swingle(2012b), Evenbly and Vidal(2014), Milsted and Vidal(2018), Evenbly and Vidal(2011), Reyes and Stoudenmire(2020), McMahon et al.(2020) McMahon, Singh, and Brennen, Qi(2013), Lee and Qi(2016), Mukherjee and Lal(2020a), Mukherjee and Lal(2020b), Mukherjee and Lal(2022a), Kim et al.(2019) Kim, Chung, Park, and Han, Kim et al.(2017) Kim, Park, Cho, and Park, Kim and Park(2016), Kim(2020a), Kim(2020b), Kim(2020c), Kim(2020d), Kim and Ryu(2021)]. Some well-known approaches include the momentum shell renormalisation approach of Sung-Sik Lee [Lee(2010a), Lee(2010b), Lee(2012), Lee(2014)] and the multi-scale entanglement renormalisation ansatz (MERA) of Vidal [Vidal(2007), Vidal(2008), Swingle(2012b), Evenbly and Vidal(2014), Milsted and Vidal(2018)]; both use certain forms of RG flow to generate the extra spatial dimension. The former integrates out high momentum degrees of freedom in favour of generating dynamics for the boundary gauge field, with the RG trajectory variable giving rise to the additional spatial dimension [Green(2013)]. On the other hand, the MERA works by disentangling degrees of freedom in real space, creating a tensor network in the process [Evenbly and Vidal(2011), Reyes and Stoudenmire(2020), McMahon et al.(2020) McMahon, Singh, and Brennen] and leading to a renormalisation in the entanglement. The renormalised entanglement is interpreted as the additional dimension [Swingle(2012b)]. There exist other approaches that also proceed via entanglement renormalisation, such as the exact holographic mapping [Qi(2013), Lee and Qi(2016)] (EHM) and the unitary renormalisation group [Mukherjee and Lal(2020a), Mukherjee and Lal(2020b), Mukherjee and Lal(2022a)] (URG) that operate purely through unitary transformations and have been explicitly shown to lead to a holographic tensor network along the RG flow. Apart from these, there is a body of work due largely to Ki-Seok Kim that provides a geometric description of Wilson's numerical renormalisation group [Wilson(1975b), Krishna-murthy et al.(1980) Krishna-murthy, Wilkins, and Wilson, Kim et al.(2019) Kim, Chung, Park, and Han, Kim et al.(2017) Kim, Park, Cho, and Park, Kim and Park(2016)], and creates a more general RG framework in terms of an order-parameter field that extends the Ads/CFT correspondence to non-critical theories [Kim(2020a), Kim(2020b), Kim(2020c), Kim(2020d), Kim and Ryu(2021)].

Questions on entanglement and holography become particularly interesting in systems of critical fermions. As mentioned earlier, such systems differ from bosonic and gapped systems through the presence of longer-ranged entanglement that follows a modified area-law [Wolf et al.(2008) Wolf, Verstraete, Hastings, and Cirac, Swingle(2010)] as well as topological features arising from the incompressibility of the Fermi volume [Oshikawa(2000), Seki and Yunoki(2017)]. Fermi surface gapping phase transitions in such systems are often driven by changes in topological quantum numbers and the nature of the inter-particle entanglement [Mukherjee and Lal(2020a), Mukherjee and

Lal(2020b), Gegenwart et al.(2008)Gegenwart, Si, and Steglich]. Constructing a framework for fermionic criticality is therefore an active area of research with consequences for important systems such as unconventional superconductors, strange metals and heavy fermions [Mukherjee and Lal(2020c), Mukherjee and Lal(2020d), Mukherjee and Lal(2022b), Custers et al.(2003)Custers, Gegenwart, Wilhelm, Neumaier, Tokiwa, Trovarelli, Geibel, Steglich, Pépin, and Coleman]. Still less is known about the holographic implications of such systems. We extend this body of work by providing an explicit demonstration of the holographic principle in the form of a simple and tractable exact holographic mapping (EHM) for a free fermion system with and without a mass gap. Our program yields analytic relations between the RG flow parameters and their dual bulk quantities such as distances and curvature. This also allows us to study the consequences of a critical Fermi surface (lying precisely at the quantum phase transition) on the nature of the holographic space.

Some open questions

Chapter 2

Methods and Preliminaries

2.1 The Unitary renormalisation group method

The URG method was introduced and formalised in refs. [Mukherjee and Lal(2020a), Mukherjee and Lal(2020b), Mukherjee and Lal(2020c), Mukherjee and Lal(2020d)]. This section is adapted from those references and expanded wherever required.

Description of the problem

We are given a Hamiltonian $\mathcal{H}$ which is not completely diagonal in the occupation number basis of the electrons, $\hat{n}_k: [\mathcal{H}, n_k] \neq 0$. k labels any set of quantum numbers depending on the system. For spin-less Fermions it can be the momentum of the particle, while for spin-full Fermions it can be the set of momentum and spin. There are terms that scatter electrons from one quantum number k to another quantum number k' .

We take a general Hamiltonian,

$$\mathcal{H} = H_e \hat{n}_{q\beta} + H_h (1 - \hat{n}_{q\beta}) + c_{q\beta}^\dagger T + T^\dagger c_{q\beta} \quad (1.1)$$

Formally, we can decompose the entire Hamiltonian in the subspace of the electron we want to decouple ($q\beta$).

$$\mathcal{H} = \begin{pmatrix} |1\rangle & |0\rangle \\ H_1 & T \\ T^\dagger & H_0 \end{pmatrix} \quad (1.2)$$

The basis in which this matrix is written is $\{|1\rangle, |0\rangle\}$ where $|i\rangle$ is the set of all states where $\hat{n}_{q\beta} = i$. The aim of one step of the URG is to find a unitary transformation U such that the new Hamiltonian $U\mathcal{H}U^\dagger$ is diagonal in this already-chosen basis.

$$\tilde{\mathcal{H}} \equiv U\mathcal{H}U^\dagger = \begin{pmatrix} |1\rangle & |0\rangle \\ \tilde{H}_1 & 0 \\ 0 & \tilde{H}_0 \end{pmatrix} \quad (1.3)$$

U_q is defined by

$$\tilde{\mathcal{H}} = U_q \mathcal{H} U_q^\dagger \text{ such that } [\tilde{\mathcal{H}}, n_q] = 0 \quad (1.4)$$

It is clear that U is the diagonalizing matrix for $\mathcal{H}$. Hence we can frame this problem as an eigenvalue equation as well. Let $|\psi_1\rangle, |\psi_0\rangle$ be the basis in which the original Hamiltonian $\mathcal{H}$ has no off-diagonal terms corresponding to $q\beta$. Hence, we can write

$$\mathcal{H} |\psi_i\rangle = \tilde{H}_i |\psi_i\rangle, i \in \{0, 1\} \quad (1.5)$$

Since $|\psi_i\rangle$ is the set of eigenstates of $\mathcal{H}$ and $|i\rangle$ is the set of eigenstates in which $U\mathcal{H}U^\dagger$ has no off-diagonal terms corresponding to $q\beta$, we can relate $|\psi_i\rangle$ and $|i\rangle$ by the same transformation : $|\psi_i\rangle = U^\dagger |i\rangle$. We can expand the state $|\psi_i\rangle$ in the subspace of $q\beta$:

$$|\psi_i\rangle = \sum_{j=0,1} |j\rangle \langle j | \psi_i \rangle \equiv |1\rangle |\phi_1^i\rangle + |0\rangle |\phi_0^i\rangle \quad (1.6)$$

where $|\phi_j^i\rangle = \langle j | \psi_i \rangle$. If we substitute the expansion 1.2 into the eigenvalue equation 1.5, we get

$$\left[H_e \hat{n}_{q\beta} + H_h (1 - \hat{n}_{q\beta}) + c_{q\beta}^\dagger T + T^\dagger c_{q\beta} \right] |\psi_i\rangle = \tilde{H}_i |\psi_i\rangle \quad (1.7)$$

The diagonal parts $H_e = \text{tr} [\mathcal{H} \hat{n}_{q\beta}]$ and $H_h = \text{tr} [\mathcal{H} (1 - \hat{n}_{q\beta})]$ can be separated into a purely diagonal part $\mathcal{H}^d$ that contains the single-particle energies and the multi-particle correlation energies or Hartree-like contributions, and an off-diagonal part $\mathcal{H}^i$ that scatters between the remaining degrees of freedom $k\sigma \neq q\beta$. That is,

$$H_e \hat{n}_{q\beta} + H_h (1 - \hat{n}_{q\beta}) = \mathcal{H}^d + \mathcal{H}^i$$

This gives

$$\left[c_{q\beta}^\dagger T + T^\dagger c_{q\beta} \right] |\psi_i\rangle = \left(\tilde{H}_i - \mathcal{H}^i - \mathcal{H}^d \right) |\psi_i\rangle \quad (1.8)$$

Obtaining the decoupling transformation

We now define a new operator $\hat{\omega}_i = \tilde{H}_i - \mathcal{H}^i$, such that

$$\left[c_{q\beta}^\dagger T + T^\dagger c_{q\beta} \right] |\psi_i\rangle = (\hat{\omega}_i - \mathcal{H}^d) |\psi_i\rangle \quad (1.9)$$

From the definition of $\hat{\omega}_i$, we can see that it is Hermitian and has no term that scatters in the subspace of $q\beta$, so it is diagonal in $q\beta$ and we can expand it as $\hat{\omega}_i = \hat{\omega}_i^1 \hat{n}_{q\beta} + \hat{\omega}_i^0 (1 - \hat{n}_{q\beta})$. Using the expansion 1.6, we can write

$$\hat{\omega}_i |\psi_i\rangle = \hat{\omega}_i^1 |1\rangle |\phi_1^i\rangle + \hat{\omega}_i^0 |0\rangle |\phi_0^i\rangle \quad (1.10)$$

Since the only requirement on $|\psi_i\rangle$ is that it diagonalize the Hamiltonian in the subspace of $q\beta$, there is freedom in the choice of this state. We can exploit this freedom and choose the $|\phi_0^i\rangle$ to be an eigenstates of $\hat{\omega}_i^{1,0}$ corresponding to real eigenvalues $\omega_i^{1,0}$:

$$\left[\mathcal{H}^d + c_{q\beta}^\dagger T + T^\dagger c_{q\beta} \right] |\psi_i(\omega_i)\rangle = (\omega_i^1 - \mathcal{H}^d) |1\rangle |\phi_1^i\rangle + (\omega_i^0 - \mathcal{H}^d) |0\rangle |\phi_0^i\rangle \quad (1.11)$$

If we now substitute the expansion 1.6 and gather the terms that result in $\hat{n}_{q\beta} = 1$, we get

$$c_{q\beta}^\dagger T |0\rangle |\phi_0^i\rangle = (\omega_i^1 - \mathcal{H}^d) |1\rangle |\phi_1^i\rangle \quad (1.12)$$

Similarly, gathering the terms that result in $\hat{n}_{q\beta} = 0$ gives

$$T^\dagger c_{q\beta} |1\rangle |\phi_1^i\rangle = (\omega_i^0 - \mathcal{H}^d) |0\rangle |\phi_0^i\rangle \quad (1.13)$$

We now define two many-particle transition operators:

$$\begin{aligned} \eta^\dagger(\omega_i^1) &= \frac{1}{\omega_i^1 - \mathcal{H}^d} c_{q\beta}^\dagger T \equiv G_1 c_{q\beta}^\dagger T \\ \eta(\omega_i^0) &= \frac{1}{\omega_i^0 - \mathcal{H}^d} T^\dagger c_{q\beta} \equiv G_0 T^\dagger c_{q\beta} \end{aligned} \quad (1.14)$$

where G_j is the propagator $\frac{1}{\omega_i^j - \mathcal{H}^d}$. We can write this compactly as

$$\eta(\hat{\omega}) = G T^\dagger c_{q\beta} = \frac{1}{\hat{\omega}_i - \mathcal{H}^d} T^\dagger c_{q\beta} \quad (1.15)$$

where $\hat{\omega}_i = \omega_i^0 (1 - \hat{n}_{q\beta}) + \omega_i^1 \hat{n}_{q\beta} = \begin{pmatrix} \omega_i^1 & \\ & \omega_i^0 \end{pmatrix}$ is a 2x2 matrix and $\mathcal{H}^d = \mathcal{H}_0^d (1 - \hat{n}_{q\beta}) + \mathcal{H}_1^d \hat{n}_{q\beta}$ and $G = (\hat{\omega} - \mathcal{H}^d)^{-1}$. It is easy to check that this reproduces the previous forms of η_0 and $\eta_1^\dagger$. We will later find that it is important to demand that these two be Hermitian conjugates of each other; that constraint is imposed on the denominators:

$$\eta^\dagger(\omega_i^0) = \eta^\dagger(\omega_i^1) \implies \frac{1}{\omega_i^1 - \mathcal{H}^d} c_{q\beta}^\dagger T = c_{q\beta}^\dagger T \frac{1}{\omega_i^0 - \mathcal{H}^d} \quad (1.16)$$

Henceforth we will assume that this constraint has been imposed.

In terms of these operators, eq. 1.13 becomes

$$|1\rangle |\phi_1^i\rangle = \eta^\dagger |0\rangle |\phi_0^i\rangle, \quad |0\rangle |\phi_0^i\rangle = \eta |1\rangle |\phi_1^i\rangle \quad (1.17)$$

These allow us to write

$$|\psi_1\rangle = |1\rangle |\phi_1^i\rangle + |0\rangle |\phi_0^i\rangle = (1 + \eta) |1\rangle |\phi_1^i\rangle, \quad |\psi_0\rangle = (1 + \eta^\dagger) |0\rangle |\phi_0^i\rangle \quad (1.18)$$

Recalling that $|\psi_i\rangle = U^\dagger |i\rangle$, we can read off the required transformation:

$$U_1 = 1 + \eta \quad (1.19)$$




Figure 2.1: Three steps of the URG: Decompose the Hamiltonian in a 2×2 matrix, apply the unitary operator to rotate it, then repeat these steps with one of the rotated blocks.

Properties of the many-body transition operators

The operators η have some important properties. First is the Fermionic nature:

$$\eta^2 = \eta^{\dagger 2} = 0 \quad [c^{\dagger 2} = c^2 = 0] \quad (1.20)$$

Second is:

$$\begin{aligned} |1\rangle |\phi_1^i\rangle &= \eta^\dagger |0\rangle |\phi_0^i\rangle = \eta^\dagger \eta |1\rangle |\phi_1^i\rangle \implies \eta^\dagger \eta = \hat{n}_{q\beta} \\ |0\rangle |\phi_0^i\rangle &= \eta |1\rangle |\phi_1^i\rangle = \eta \eta^\dagger |0\rangle |\phi_0^i\rangle \implies \eta \eta^\dagger = 1 - \hat{n}_{q\beta} \end{aligned} \quad (1.21)$$

and hence the anticommutator

$$\implies \{\eta, \eta^\dagger\} = 1 \quad (1.22)$$

Note that the three equations in 1.21 work only when applied on the eigenstate $|\psi_i\rangle$ and not any arbitrary state.

$$\begin{aligned} \eta^\dagger \eta |\psi_i\rangle &= |1\rangle |\phi_1^i\rangle = \hat{n}_{q\beta} |\psi_i\rangle \\ \eta \eta^\dagger |\psi_i\rangle &= |0\rangle |\phi_0^i\rangle = (1 - \hat{n}_{q\beta}) |\psi_i\rangle \\ \{\eta^\dagger, \eta\} |\psi_i\rangle &= |\psi_i\rangle \end{aligned}$$

Form of the unitary operators

Although we have found the correct similarity transformations U_i (eqs. 1.19), we need to convert them into a unitary transformation. Say we are trying to rotate the eigenstate $|\psi_1\rangle$ into the state $|1\rangle$. We can then work with the transformation

$$U_1 = 1 + \eta \quad (1.23)$$

In this form, this transformation is not unitary. It can however be written in an exponential form:

$$U_1 = e^\eta \quad (1.24)$$

using the fact that $\eta^2 = 0$. It is shown in ref. [Suzuki and Okamoto(1984)] that corresponding to a similarity transformation e^ω , there exists a unitary transformation e^G where

$$G = \tanh^{-1} (\omega - \omega^\dagger) \quad (1.25)$$

Applying that to the problem at hand gives

$$U_1^\dagger = \exp \{ \tanh^{-1} (\eta - \eta^\dagger) \} \quad (1.26)$$

Let $x = \tanh y$. Then,

$$x = \frac{e^{2y} + 1}{e^{2y} - 1} \implies y = \frac{1}{2} \log \frac{1+x}{1-x} \implies e^y = e^{\tanh^{-1} x} = \sqrt{\frac{1+x}{1-x}} \quad (1.27)$$

Therefore,

$$\exp \{ \tanh^{-1} (\eta - \eta^\dagger) \} = \frac{1 + \eta - \eta^\dagger}{\sqrt{(1 + \eta^\dagger - \eta)(1 - \eta^\dagger + \eta)}} = \frac{1 + \eta - \eta^\dagger}{\sqrt{1 + \{\eta, \eta^\dagger\}}} = \frac{1}{\sqrt{2}} (1 + \eta - \eta^\dagger) \quad (1.28)$$

The *unitary* operator that transforms the entangled eigenstate $|\psi_1\rangle$ to the state $|1\rangle$ is thus

$$U_1 = \frac{1}{\sqrt{2}} (1 + \eta^\dagger - \eta) \quad (1.29)$$

It can also be written as $\exp \{ \frac{\pi}{4} (\eta^\dagger - \eta) \}$ because

$$\begin{aligned} \exp \left\{ \frac{\pi}{4} (\eta^\dagger - \eta) \right\} &= 1 + (\eta^\dagger - \eta) \frac{\pi}{4} + \frac{1}{2!} (\eta^\dagger - \eta)^2 \left(\frac{\pi}{4} \right)^2 + \frac{1}{3!} (\eta^\dagger - \eta)^3 \left(\frac{\pi}{4} \right)^3 + \dots \\ &= 1 + (\eta^\dagger - \eta) \frac{\pi}{4} - \frac{1}{2!} \left(\frac{\pi}{4} \right)^2 - \frac{1}{3!} (\eta^\dagger - \eta) \left(\frac{\pi}{4} \right)^3 + \frac{1}{4!} \left(\frac{\pi}{4} \right)^4 + \dots \\ &= \cos \frac{\pi}{4} + (\eta^\dagger - \eta) \sin \frac{\pi}{4} \\ &= \frac{1}{\sqrt{2}} (1 + \eta^\dagger - \eta) \end{aligned} \quad (1.30)$$

There we used

$$(\eta^\dagger - \eta)^2 = \eta^{\dagger 2} + \eta^2 - \{\eta^\dagger, \eta\} = -1 \quad [\because \eta^2 = \eta^{\dagger 2} = 0] \quad (1.31)$$

and hence

$$(\eta^\dagger - \eta)^3 = -1 (\eta^\dagger - \eta) \quad (1.32)$$

and so on.

Effective Hamiltonian

We can now compute the form of the effective Hamiltonian that comes about when we apply U_1 - that is - when we rotate one exact eigenstate $|\psi_1\rangle$ into the occupied Fock space basis $|1\rangle$. From eq. 1.29,

$$\begin{aligned}
 U_1 \mathcal{H} U_1^\dagger &= \frac{1}{2} (1 + \eta^\dagger - \eta) \mathcal{H} (1 + \eta - \eta^\dagger) \\
 &= \frac{1}{2} (1 + \eta^\dagger - \eta) (\mathcal{H} + \mathcal{H}\eta - \mathcal{H}\eta^\dagger) \\
 &= \frac{1}{2} (\mathcal{H} + \mathcal{H}\eta - \mathcal{H}\eta^\dagger + \eta^\dagger \mathcal{H} + \eta^\dagger \mathcal{H}\eta - \eta^\dagger \mathcal{H}\eta^\dagger - \eta \mathcal{H} - \eta \mathcal{H}\eta + \eta \mathcal{H}\eta^\dagger) \\
 &= \frac{1}{2} (\mathcal{H}^d + \mathcal{H}^i + \mathcal{H}^I + \mathcal{H}\eta - \mathcal{H}\eta^\dagger + \eta^\dagger \mathcal{H} + \eta^\dagger \mathcal{H}\eta - \eta^\dagger \mathcal{H}\eta^\dagger - \eta \mathcal{H} - \eta \mathcal{H}\eta + \eta \mathcal{H}\eta^\dagger) \\
 &= \frac{1}{2} (\mathcal{H}^d + \mathcal{H}^i + \mathcal{H}^I + [\eta^\dagger - \eta, \mathcal{H}] + \eta^\dagger \mathcal{H}\eta - \eta^\dagger \mathcal{H}\eta^\dagger - \eta \mathcal{H}\eta + \eta \mathcal{H}\eta^\dagger)
 \end{aligned} \tag{1.33}$$

In the last two lines, we expanded the Hamiltonian into the three parts $\mathcal{H}^d$, $\mathcal{H}^i$ and a third piece $\mathcal{H}^I \equiv c_{q\beta}^\dagger T + T^\dagger c_{q\beta}$.

For reasons that will become apparent, we will split the terms into two groups:

$$\tilde{\mathcal{H}} = \frac{1}{2} \left(\underbrace{\mathcal{H}^d + \mathcal{H}^i + [\eta^\dagger - \eta, \mathcal{H}] + \eta^\dagger \mathcal{H}\eta + \eta \mathcal{H}\eta^\dagger}_{\text{group 1}} + \underbrace{\mathcal{H}^I - \eta^\dagger \mathcal{H}\eta^\dagger - \eta \mathcal{H}\eta}_{\text{group 2}} \right) \tag{1.34}$$

Group 2 can be easily shown to be 0. Note that terms that have two η or two $\eta^\dagger$ sandwiching a $\mathcal{H}$ can only be nonzero if the intervening $\mathcal{H}$ has an odd number of creation or destruction operators.

$$\eta \mathcal{H} \eta = \eta c_q^\dagger T \eta \tag{1.35}$$

and

$$\eta^\dagger \mathcal{H} \eta^\dagger = \eta^\dagger T^\dagger c_q \eta^\dagger \tag{1.36}$$

Group 2 becomes

$$\text{group 2} = \mathcal{H}^I - \eta^\dagger T^\dagger c_q \eta^\dagger - \eta c_q^\dagger T \eta = c_q^\dagger T + T^\dagger c_q - \eta^\dagger T^\dagger c_q \eta^\dagger - \eta c_q^\dagger T \eta \tag{1.37}$$

To simplify this, we use the relation

$$\begin{aligned}
 \eta c_q^\dagger T \eta &= \frac{1}{\omega_i^0 - \mathcal{H}^d} T^\dagger c_q c_q^\dagger T \eta \\
 &= T^\dagger c_q \frac{1}{\omega_i^1 - \mathcal{H}^d} c_q^\dagger T \eta \quad [\text{eq. 1.16}] \\
 &= T^\dagger c_q \eta^\dagger \eta \quad [\text{eq. 1.15}] \\
 &= T^\dagger c_q \hat{n}_q \quad [\text{eq. 1.21}]
 \end{aligned} \tag{1.38}$$

which gives

$$\eta c_q^\dagger T \eta = T^\dagger c_q \quad (1.39)$$

Taking the Hermitian conjugate of eq. 1.39 gives

$$\eta^\dagger T^\dagger c_q \eta^\dagger = c_q^\dagger T \quad (1.40)$$

Substituting the expressions 1.39 and 1.40 into the expression for group 2, 1.37, shows that it vanishes. This leaves us only with group 1:

$$\tilde{\mathcal{H}} = \frac{1}{2} \left(\mathcal{H}^d + \mathcal{H}^i + \overbrace{\eta^\dagger \mathcal{H} \eta + \eta \mathcal{H} \eta^\dagger}^{\text{group A}} + \underbrace{[\eta^\dagger - \eta, \mathcal{H}]}_{\text{group B}} \right) \quad (1.41)$$

Group A simplifies in the following way. First note that $\eta^\dagger \mathcal{H}^I \eta = \eta^\dagger \mathcal{H}^I \eta = 0$ must be 0 because it will involve consecutive $c_{q\beta}$ or consecutive $c_{q\beta}^\dagger$. We are therefore left with the diagonal part of $\mathcal{H}$, which is $H_e \hat{n}_{q\beta} + H_h (1 - \hat{n}_{q\beta})$.

$$\eta^\dagger [H_e \hat{n}_{q\beta} + H_h (1 - \hat{n}_{q\beta})] \eta + \eta [H_e \hat{n}_{q\beta} + H_h (1 - \hat{n}_{q\beta})] \eta^\dagger = \eta^\dagger H_h \eta + \eta H_e \eta^\dagger \quad (1.42)$$

This can be shown to be equal to the diagonal part:

$$\text{group A} = \eta^\dagger H_h \eta + \eta H_e \eta^\dagger = H_e \hat{n}_{q\beta} + H_h (1 - \hat{n}_{q\beta}) = \mathcal{H}^d + \mathcal{H}^i \quad (1.43)$$

It can also be shown that

$$\text{group B} = [\eta^\dagger - \eta, \mathcal{H}] = 2 [c_{q\beta}^\dagger T, \eta] \quad (1.44)$$

Putting it all together,

$$\tilde{\mathcal{H}} = \mathcal{H}^d + \mathcal{H}^i + [c_{q\beta}^\dagger T, \eta] \quad (1.45)$$

The renormalizing in the Hamiltonian is

$$\Delta \mathcal{H} = \tilde{\mathcal{H}} - \mathcal{H}^d - \mathcal{H}^i = [c_{q\beta}^\dagger T, \eta] \quad (1.46)$$

Because of eq. 1.44, it can also be written as

$$\Delta \mathcal{H} = \frac{1}{2} [\eta^\dagger - \eta, \mathcal{H}_X] = \frac{1}{2} [\eta^\dagger - \eta, \mathcal{H}] \quad (1.47)$$

This form will be useful later when we make the connection with one-shot Schrieffer-Wolff transformation and CUT RG.

To check that the renormalised Hamiltonian indeed commutes with $\hat{n}_{q\beta}$,

$$\begin{aligned} [\tilde{\mathcal{H}}, \hat{n}_{q\beta}] &= [[c_{q\beta}^\dagger T, \eta], \hat{n}_{q\beta}] = [c_{q\beta}^\dagger T \eta, \hat{n}_{q\beta}] - [\eta c_{q\beta}^\dagger T, \hat{n}_{q\beta}] \\ &= c_{q\beta}^\dagger T \eta \hat{n}_{q\beta} - \hat{n}_{q\beta} c_{q\beta}^\dagger T \eta \quad \left[2^{\text{nd}} [\cdot] \text{ is 0, } \because c_{q\beta}^\dagger \hat{n}_{q\beta} = \hat{n}_{q\beta} \eta = 0 \right] \\ &= c_{q\beta}^\dagger T \eta - c_{q\beta}^\dagger T \eta = 0 \end{aligned} \quad (1.48)$$

Fixed point condition

Within the URG, it is a prescription that the fixed point is reached when the denominator of the RG equation vanishes. This is equivalent to either $\omega_i^1 = \mathcal{H}_1^d$ or $\omega_i^0 = \mathcal{H}_0^d$. This shows that at the fixed point, one of the eigenvalues of $\hat{\omega}_i$ matches the corresponding eigenvalue of the diagonal blocks. This also leads to the vanishing of the off-diagonal block, because eqs. 1.12 and 1.13 gives

$$c_{q\beta}^\dagger T |0\rangle |\phi_0^i\rangle = (\omega_i^1 - \mathcal{H}_1^d) |1\rangle |\phi_1^i\rangle = 0 \implies c_{q\beta}^\dagger T = 0 \quad (1.49)$$

Multiple off-diagonal terms

There is a subtle assumption in the definitions eq. 1.14. In order for η to be the Hermitian conjugate of $\eta^\dagger$, $\mathcal{H}_d$ cannot have any information that relates to the structure of T . To see why, say the total off-diagonal term is composed of two parts: $T = T_1 + T_2$.

$$\begin{aligned} \eta &= \frac{1}{\omega_0 - \mathcal{H}_d} (T_1^\dagger + T_2^\dagger) c = \left[\frac{1}{\omega^0 - E_1^0} T_1^\dagger c + \frac{1}{\omega^0 - E_2^0} T_2^\dagger c \right] \\ \eta^\dagger &= \frac{1}{\omega^1 - \mathcal{H}_d} c^\dagger (T_1 + T_2) = \left[\frac{1}{\omega^1 - E_1^1} c^\dagger T_1 + \frac{1}{\omega^1 - E_2^1} c^\dagger T_2 \right] \end{aligned} \quad (1.50)$$

where $\mathcal{H}_d T_i^\dagger c = E_i^0 T_i^\dagger c$ and $\mathcal{H}_d c^\dagger T_i = E_i^1 c^\dagger T_i$. We can now see that in order for $\eta = (\eta^\dagger)^\dagger$ to hold, two conditions must be met:

$$\omega^0 - E_1^0 = \omega^1 - E_1^1, \quad \omega^0 - E_2^0 = \omega^1 - E_2^1 \quad (1.51)$$

This will not hold generally. The correct solution is to realize that each such off-diagonal term T_i will come with its own quantum fluctuation scale ω_i .

$$\begin{aligned} \eta &= \sum_i \frac{1}{\omega_i^0 - E_i^0} T_i^\dagger c \\ \eta^\dagger &= \sum_i \frac{1}{\omega_i^1 - E_i^1} c^\dagger T_i \end{aligned} \quad (1.52)$$

If we now impose the condition that $\eta = (\eta^\dagger)^\dagger$, we get the relations

$$\omega_i^0 - \omega_i^1 = E_i^0 - E_i^1 \quad (1.53)$$

and so

$$\eta^\dagger - \eta = \sum_i \frac{1}{\omega_i^0 - E_i^0} (c^\dagger T_i - T_i^\dagger c) \quad (1.54)$$

The expression for the renormalization will not be just $[c^\dagger T, \eta]$ in this case. That form will be non-Hermitian. The correct form is obtained from the more general form $[\eta^\dagger - \eta, \mathcal{H}_X]$:

$$\begin{aligned}\Delta\mathcal{H} &= \frac{1}{2} [\eta^\dagger - \eta, c^\dagger T + T^\dagger c] = \frac{1}{2} \sum_{ij} \frac{1}{\omega_i^0 - E_i^0} [c^\dagger T_i - T_i^\dagger c, c^\dagger T_j + T_j^\dagger c] \\ &= \frac{1}{2} \sum_{ij} \frac{1}{\omega_i^0 - E_i^0} \left[\hat{n} (T_i T_j^\dagger + T_j T_i^\dagger) - (1 - \hat{n}) (T_i^\dagger T_j + T_j^\dagger T_i) \right] \\ &= \frac{1}{2} \sum_{ij} \left(\frac{1}{\omega_i^0 - E_i^0} + \frac{1}{\omega_j^0 - E_j^0} \right) [\hat{n} T_i T_j^\dagger - (1 - \hat{n}) T_i^\dagger T_j]\end{aligned}\quad (1.55)$$

Prescription

Given a Hamiltonian

$$\mathcal{H} = \mathcal{H}_1 + \mathcal{H}_0 + c^\dagger T + T^\dagger c \quad (1.56)$$

the goal is to look at the renormalization of the various couplings in the Hamiltonian as we decouple high energy electron states. Typically we have a shell of electrons at some energy D . During the process, we make one simplification. We assume that there is only one electron on that shell at a time, say with quantum numbers q, σ , and calculate the renormalization of the various couplings due to this electron. We then sum the momentum q over the shell and the spin β , and this gives the total renormalization due to decoupling the entire shell.

From eq. 1.45, the first two terms in the rotated Hamiltonian are just the diagonal parts of the bare Hamiltonian; they are unchanged in that part. The renormalization comes from the third term. For one electron $q\beta$ on the shell, the renormalization is

$$\Delta\mathcal{H} = [c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}), \eta] = c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}) \eta - \eta c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}) \quad (1.57)$$

Since this assumes we have obtained this from U_1 , it is fair to tag the η with a suitable label:

$$\Delta\mathcal{H} = c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}) \eta_1 - \eta_1 c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}) \quad (1.58)$$

It is clear that the first term takes into account virtual excitations that start from a filled state ($\hat{n}_{q\beta} = 1$ initially) - such a term is said to be a part of the *particle sector*.

$$\Delta_1\mathcal{H} = c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}) \eta_1 \quad (1.59)$$

The second term, on the other hand, considers excitations that start from an empty state. They constitute the *hole sector*.

$$\Delta_0\mathcal{H} = -\eta_1 c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}) \quad (1.60)$$

To write the total renormalization in a particle-hole symmetric form, we can use the relation $\eta_0 = -\eta_1$, such that both the terms will now come with a positive sign:

$$\Delta\mathcal{H} = c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}) \eta_1 + \eta_0 c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}) \quad (1.61)$$

We can make one more manipulation: using eq. 1.16, we get

$$\Delta\mathcal{H} = c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}) \eta_1 + \text{Tr}(c_{q\beta}^\dagger \mathcal{H}) c_{q\beta} \eta_0^\dagger \quad (1.62)$$

This form of the total renormalization is identical to the one we use in the "Poor Man's scaling"-type of renormalization that was used to get the scaling equations in the Kondo and Anderson models [Anderson(1970), Haldane(1978)]. Writing down the forms of η and $\eta^\dagger$ explicitly, we get

$$\Delta\mathcal{H} = c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}) \frac{1}{\omega_1^0 - \mathcal{H}_0^d} \text{Tr}(c_{q\beta}^\dagger \mathcal{H}) c_{q\beta} + \text{Tr}(c_{q\beta}^\dagger \mathcal{H}) c_{q\beta} \frac{1}{\omega_0^1 - \mathcal{H}_1^d} c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}) \quad (1.63)$$

The renormalization due to the entire shell is obtained by summing over all states on the shell.

$$\Delta\mathcal{H} = \sum_{q\beta} \left[c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}) \frac{1}{\omega_1^0 - \mathcal{H}_0^d} \text{Tr}(c_{q\beta}^\dagger \mathcal{H}) c_{q\beta} + \text{Tr}(c_{q\beta}^\dagger \mathcal{H}) c_{q\beta} \frac{1}{\omega_0^1 - \mathcal{H}_1^d} c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta}) \right] \quad (1.64)$$

These equations will now need to be simplified. For example, in the particle sector, we can set $\hat{n}_{q\beta} = 0$ in the numerator, because there is no such excitation in the initial state. Similarly, in the hole sector, we can set $\hat{n}_{q\beta} = 1$ because that state was occupied in the initial state. Another simplification we typically employ is that $\mathcal{H}_{0,1}^d$ will, in general, have the energies of all the electrons. But we consider only the energy of the on-shell electrons in the denominator. After integrating out these electrons, we can rearrange the remaining operators to determine which term in the Hamiltonian it renormalizes and what is the renormalization.

At first sight, one might think that we must evaluate lots of traces to obtain the terms in $\Delta\mathcal{H}$. A little thought reveals that the terms in the numerator are simply the off-diagonal terms in the Hamiltonian; $\text{Tr}(c_{q\beta}^\dagger \mathcal{H}) c_{q\beta}$ is the off-diagonal term that has $c_{q\beta}$ in it, and $c_{q\beta}^\dagger \text{Tr}(\mathcal{H}c_{q\beta})$ is the off-diagonal term that has $c_{q\beta}^\dagger$ in it. $\mathcal{H}^D$ is just the diagonal part of the Hamiltonian.

2.2 Example I: URG analysis of the star graph model

The star graph problem has already been analyzed using URG and an extensive study of its entanglement properties has already been carried out, in ref. [Patra et al.(2023a)Patra, Mukherjee, Mukherjee, Vidhyadhiraja, Taraphder, and Lal]. Here we focus on just deriving the RG equations. The system consists of N spin-like degrees of freedom (labeled 1 through N) individually talking to a spin at the center (labeled 0). Each spin i ($\in [0, N]$) has an on-site energy ϵ_i . The coupling strength between 0 and i ($\in [1, N]$) is J_i . We choose the on-site energies such that $\epsilon_{i+1} > \epsilon_i, i \in [N-1, 1]$. In this way, ϵ_1 is the infrared limit and ϵ_N is the ultraviolet limit.




Figure 2.2: Star Graph model

$$\mathcal{H} = \epsilon_0 S_0^z + \sum_{i=1}^N \left[\epsilon_i S_i^z + J_i \vec{S}_0 \cdot \vec{S}_i \right] \quad (2.1)$$

By converting the last term into S^z and $S^\pm$, we can write the Hamiltonian as

$$\mathcal{H} = \epsilon_0 S_0^z + \sum_{i=1}^N \left[\epsilon_i S_i^z + J_i \left(S_0^z S_i^z + \frac{1}{2} (S_0^+ S_i^- + S_0^- S_i^+) \right) \right] \quad (2.2)$$

Calculation of Renormalization

The RG involves decoupling the nodes N through 1, and looking at the resultant renormalization in ϵ_i and J_i . As a simplification, we will ignore the lower nodes in the denominator and keep only the node currently being decoupled, ie node N . Since node 0 is connected to node N , we will keep node 0 in the denominator as well. Making this simplification gives

$$\mathcal{H}^D = \epsilon_0 S_0^z + \epsilon_N S_N^z + J_N S_0^z S_N^z \quad (2.3)$$

The off-diagonal part in the subspace of the node N is

$$\mathcal{H}_X = S_N^+ T + T^\dagger S_N^- = \frac{1}{2} J_N (S_N^+ S_0^- + S_N^- S_0^+) \quad (2.4)$$

The renormalization on doing one step of the URG is given by

$$\begin{aligned} \Delta \mathcal{H} &= S_N^+ T \frac{1}{\omega_1^0 - \mathcal{H}^D} T^\dagger S_N^- + T^\dagger S_N^- \frac{1}{\omega_1^0 - \mathcal{H}^D} S_N^+ T \\ &= \frac{J_N^2}{4} S_N^+ S_0^- \frac{1}{\omega_1^0 - \epsilon_0 S_0^z - \epsilon_N S_N^z - J_N S_0^z S_N^z} S_0^+ S_N^- + \frac{J_N^2}{4} S_0^+ S_N^- \frac{1}{\omega_1^0 - \epsilon_0 S_0^z - \epsilon_N S_N^z - J_N S_0^z S_N^z} S_N^+ S_0^- \end{aligned} \quad (2.5)$$

There, N refers to the spin being decoupled. The first Greens function has S_0^+ and S_N^- in front of it, so we substitute $S_0^z = \frac{1}{2}$, $S_N^z = -\frac{1}{2}$ in that Greens function. For the other Greens function we do the opposite.

$$\Delta\mathcal{H} = \frac{J_N^2}{4} S_N^+ S_0^- \frac{1}{\omega_1^0 - \frac{1}{2}\epsilon_0 + \frac{1}{2}\epsilon_N + \frac{1}{4}J_N} S_0^+ S_N^- + \frac{J_N^2}{4} S_0^+ S_N^- \frac{1}{\omega_0^1 + \frac{1}{2}\epsilon_0 - \frac{1}{2}\epsilon_N + \frac{1}{4}J_N} S_N^+ S_0^- \quad (2.6)$$

To relate ω_1^0 and ω_0^1 , we use eq. ??:

$$\omega_1^0 + \omega_0^1 = \mathcal{H}_0^D + \mathcal{H}_1^D = -\frac{1}{2}J_N \implies \omega_1^0 \equiv \omega, \quad \omega_0^1 \equiv \omega' = -\frac{1}{2}J_N - \omega \quad (2.7)$$

So, the renormalization becomes

$$\Delta\mathcal{H} = \frac{J_N^2}{4} \frac{1}{\omega - \frac{1}{2}\epsilon_0 + \frac{1}{2}\epsilon_N + \frac{1}{4}J_N} (S_N^+ S_0^- S_0^+ S_N^- - S_0^+ S_N^- S_N^+ S_0^-) = \frac{J_N^2}{4} \frac{1}{\omega - \frac{1}{2}\epsilon_0 + \frac{1}{2}\epsilon_N + \frac{1}{4}J_N} (S_N^z - S_0^z) \quad (2.8)$$

There we used $S^+ S^- = \frac{1}{2} + S^z$ and $S^- S^+ = \frac{1}{2} - S^z$.

We can now read off the renormalizations in ϵ_N and ϵ_0 .

$$\begin{aligned} \Delta\epsilon_N &= \frac{1}{4} J_N^2 \frac{1}{\omega - \frac{1}{2}\epsilon_0 + \frac{1}{2}\epsilon_N + \frac{1}{4}J_N} \\ \Delta\epsilon_0 &= -\frac{1}{4} J_N^2 \frac{1}{\omega - \frac{1}{2}\epsilon_0 + \frac{1}{2}\epsilon_N + \frac{1}{4}J_N} \end{aligned} \quad (2.9)$$

Nature of flows

We are interested in looking at the renormalization of the central node energy ϵ_0 , upon removing the nodes N through 1. We will hence concentrate on the second RG equation. We first make some simplifying assumptions: $J_i = J, \epsilon_i = \epsilon$ for all $i \in \{1, N\}$.

$$\Delta\epsilon_0 = -\frac{1}{4} J^2 \frac{1}{\omega - \frac{1}{2}\epsilon_0 + \frac{1}{2}\epsilon + \frac{1}{4}J} \quad (2.10)$$

Define $\tilde{\omega} = \omega + \frac{1}{2}\epsilon + \frac{1}{4}J$.

$$\Delta\epsilon_0 = -\frac{1}{4} J^2 \frac{1}{\tilde{\omega} - \frac{1}{2}\epsilon_0} \quad (2.11)$$

Our goal here is to look for a fixed-point condition such that the denominator vanishes at some point of the RG. If we start with a bare of ϵ_0 such that $\tilde{\omega} - \frac{1}{2}\epsilon_0 > 0$, the denominator will be positive and the RG equation will be irrelevant. This means that ϵ_0 will keep on decreasing, and the denominator will keep on becoming more and more positive, meaning there cannot be a fixed point in this situation.

If, on other hand, we start with a bare of ϵ_0 such that $\tilde{\omega} - \frac{1}{2}\epsilon_0 < 0$, the denominator will be negative and the RG equation will be relevant. This means that ϵ_0 will keep on increasing, and the denominator will keep on becoming more and more negative, meaning there cannot be a fixed point in this situation either. These situations are depicted in figure 2.3.




Figure 2.3: Left: RG flow for the two cases. The green line is the distance between the bare values of the two couplings, and hence also the magnitude of the denominator. The red arrow denotes the direction in which ϵ_0 will flow. Upward flow is increase. In both cases, the flow is such that the distance between the two quantities (and hence the magnitude of the denominator) increases. The RG fixed point occurs when the magnitude of the denominator goes to 0. This happens if the distance vanishes. Since the distance necessarily increases, we cannot get a fixed point in this way. Right: RG flow for the two cases with the new $-\tilde{\omega} = \omega' - \frac{1}{2}\epsilon + \frac{1}{4}J$. Now we can see that in both cases, the flow is such that the distance (green dotted line) between the couplings decreases. A fixed point is reached when this distance vanishes.

Since we cannot find a fixed point, we will use a different ω . Instead of ω_1^0 , we will use ω_1^1 . From eq. ??, we have

$$\omega_1^0 - \omega_1^1 = \mathcal{H}_0^D - \mathcal{H}_1^D = \epsilon_0 - \epsilon_N = \epsilon_0 - \epsilon \quad (2.12)$$

Defining $\omega_1^1 = \omega'$ and substituting this in eq. 2.10 gives

$$\Delta\epsilon_0 = -\frac{1}{4}J^2 \frac{1}{\omega' - \frac{1}{2}\epsilon + \frac{1}{2}\epsilon_0 + \frac{1}{4}J} \quad (2.13)$$

We again define $-\tilde{\omega} = \omega' - \frac{1}{2}\epsilon + \frac{1}{4}J$.

$$\Delta\epsilon_0 = \frac{1}{4}J^2 \frac{1}{\tilde{\omega} - \frac{1}{2}\epsilon_0} \quad (2.14)$$

We now repeat the exercise of determining the relevance of the flows under various regime. If we start with a bare ϵ_0 such that $\tilde{\omega} + \frac{1}{2}\epsilon_0 > 0$, then the denominator is positive so the renormalization will be irrelevant. ϵ_0 will decrease until we reach $\tilde{\omega} + \frac{1}{2}\epsilon_0 = 0$. This will be a fixed point. However, if we start with a bare ϵ_0 such that $\tilde{\omega} + \frac{1}{2}\epsilon_0 < 0$, then the denominator is negative so the renormalization will be relevant. ϵ_0 will increase until we reach $\tilde{\omega} + \frac{1}{2}\epsilon_0 = 0$. This will again be a fixed point. This new situation is depicted in right panel of figure 2.3.

Effective Hamiltonians

If $\tilde{\omega}$ and ϵ_0 are of the same sign at the bare level, then it is easy to see that since the fixed point is defined by $\tilde{\omega} = \frac{1}{2}\epsilon_0^*$ (* denotes value at fixed point), the effective Hamiltonian at the fixed point will be

$$\mathcal{H}^* = 2\tilde{\omega}S_0^z + \epsilon \sum_i S_i^z + J \sum_i \vec{S}_i \cdot \vec{S}_0, \quad \text{if } \tilde{\omega}\epsilon_0 > 0 \quad (2.15)$$

If, at the bare level, ϵ_0 and $\tilde{\omega}$ are of opposite signs, then ϵ_0 would undergo a change in sign at some point as it flows towards $\tilde{\omega}$. Since we do not expect a coupling to change sign under RG, we will restrict it to 0 in such cases.

$$\mathcal{H}^* = \epsilon \sum_i S_i^z + J \sum_i \vec{S}_i \cdot \vec{S}_0, \quad \text{if } \tilde{\omega}\epsilon_0 < 0 \quad (2.16)$$

Things get much more simpler if we assume the onsite energies of the surrounding nodes are zero.

$$\begin{aligned} \mathcal{H}^* &= 2\tilde{\omega}S_0^z + J \sum_i \vec{S}_i \cdot \vec{S}_0, \quad \text{if } \tilde{\omega}\epsilon_0 > 0 \\ \mathcal{H}^* &= J \sum_i \vec{S}_i \cdot \vec{S}_0, \quad \text{if } \tilde{\omega}\epsilon_0 < 0 \end{aligned} \quad (2.17)$$

Fixed points

The fixed points are obtained numerically by solving the RG equation. As mentioned before, there are two types of solutions: The first kind is those in which ϵ_0 and $\tilde{\omega}$ are of the same sign, and the

former flows to the latter without crossing the 0 axis. These flows are shown (obtained numerically) in fig. 2.4. The second kind are those where the two couplings have different signs, and so ϵ_0 flows to 0. These are shown in fig. 2.5.




Figure 2.4: Flows where ϵ_0 and $\tilde{\omega}$ have same sign. The left and right panels show flows starting from negative and positive values respectively. The two plots in each panel correspond to different values of $\tilde{\omega}$, one greater than the bare ϵ_0 , the other less than that. The fixed point value is $2\tilde{\omega}$.




Figure 2.5: Flows where ϵ_0 and $\tilde{\omega}$ have opposite sign. The left and right panels show flows starting from negative and positive values respectively. The two plots in each panel correspond to different values of $\tilde{\omega}$, one greater than the bare ϵ_0 , the other less than that. The fixed point value is 0.

2.3 Example II: URG analysis of the two-channel Kondo problem

The Hamiltonian for the channel-isotropic MCK model is:

$$H = \sum_l \left[\sum_{\substack{k \\ \alpha=\uparrow,\downarrow}} \epsilon_{k,l} \hat{n}_{k\alpha,l} + \frac{\mathcal{J}}{2} \sum_{\substack{k,k' \\ \alpha,\alpha'=\uparrow,\downarrow}} \vec{S}_d \cdot \vec{\sigma}_{\alpha\alpha'} c_{k\alpha,l}^\dagger c_{k'\alpha',l} \right]. \quad (3.1)$$

The URG equation for the single-channel Kondo model [?] shows a stable strong coupling fixed point. Ferromagnetic interactions are irrelevant. Strictly speaking, that RG equation already encodes, in principle, the multi-channel behaviour, through a modified $\hat{\omega}$. To extract this information, we consider the strong coupling fixed-point $J \gg D$ as a fixed point and analyze its stability from the star graph perspective. For the exactly-screened case, the star graph decouples from the conduction bath, leaving behind a local Fermi liquid interaction on the first site. Similarly, in the under-screened regime, the ground state is composed of states where the impurity spin is only partially screened by the conduction channels. If a particular configuration of the bath-impurity system has the total conduction bath spin down, the impurity will have a residual up spin. This induces a ferromagnetic super-exchange coupling that is irrelevant under RG, so this fixed point is stable as well.

We now come to the over-screened case, where there is a residual spin on the conduction channel site. The neighbouring electrons will now hop in with spins opposite to that of the impurity, so an antiferromagnetic interaction will be induced, and such an interaction is relevant under the RG. This shows that the over-screened regime cannot have a stable strong coupling fixed point, and we need to search for an intermediate coupling fixed point. We therefore need the generator of the unitary transformation that incorporates third order scattering processes explicitly. We should take account of all possible processes that render the set of states $\{|\hat{n}_{q\beta} = 1\rangle, |\hat{n}_{q\beta} = 0\rangle\}$ diagonal. The higher order generator itself has two scattering processes, such that the entire renormalisation term $c_{q\beta}^\dagger T \eta$ has in total three coherent processes. The complete generator up to third order can be written as

$$\eta = \frac{1}{\hat{\omega} - H_D} T^\dagger c \simeq \frac{1}{\omega' - H_D} T^\dagger c + \frac{1}{\omega' - H_D} H_X \frac{1}{\omega' - H_D} T^\dagger c + \frac{1}{\omega' - H_D} T^\dagger c \frac{1}{\omega' - H_D} H_X, \quad (3.2)$$

where $H_X = J \sum_{k,k' < \Lambda_j, \alpha, \alpha'} \vec{S}_d \cdot \vec{\sigma}_{\alpha\alpha'} c_{k\alpha}^\dagger c_{k'\alpha'}$ is scattering between the entangled electrons. There are two third order terms in the above equation corresponding to the two possible sequences in which the processes can occur while keeping the total renormalisation $c_{q\beta}^\dagger T \eta$ diagonal in $q\beta$. The second order processes remain unchanged. The total renormalisation takes the form:

$$\begin{aligned} \Delta H_{(j)} = & \underbrace{c^\dagger T \frac{1}{\omega' - H_D} T^\dagger c + (c^\dagger \leftrightarrow c)}_{\Delta H_{(j)}^{(2)}} \\ & + \underbrace{c^\dagger T \frac{1}{\omega' - H_D} H_X \frac{1}{\omega' - H_D} T^\dagger c + c^\dagger T \frac{1}{\omega' - H_D} T^\dagger c \frac{1}{\omega' - H_D} H_X + (c^\dagger \leftrightarrow c)}_{\Delta H_{(j)}^{(3)}}. \end{aligned} \quad (3.3)$$

$\Delta H_{(j)}^{(2)}$ and $\Delta H_{(j)}^{(3)}$ are the renormalisation arising from the second and third order processes respectively.

It is easier to see the RG flow of the couplings if we write the Hamiltonian in terms of the eigenstates of S_d^z . These eigenstates are defined by $S_d^z |m_d\rangle = m_d |m_d\rangle$, $m_d \in [-S, S]$. In terms of these eigenstates, the Hamiltonian becomes

$$\mathcal{H} = \sum_{k\sigma} \epsilon_k \tau_{k\sigma} + \sum_{m_d=-S}^S \sum_{\substack{kl, \\ \sigma=\uparrow, \downarrow}} J_{m_d}^\sigma |m_d\rangle \langle m_d| c_{k\sigma}^\dagger c_{l\sigma} + \sum_{kl} \sum_{m_d=-S}^{S-1} J_{m_d}^t (|m_d+1\rangle \langle m_d| s_{kl}^- + \text{h.c.}) \quad (3.4)$$

where k, l sum over the momentum states, σ sums over the spin indices, $J_m^\sigma = \frac{1}{2} \sigma m J$ in the UV Hamiltonian, and $J_m^t = J \frac{1}{2} \sqrt{S(S+1) - m(m+1)}$ is the coupling that connects $|m\rangle$ and $|m+1\rangle$. We first calculate $\Delta H_{(j)}^{(2)}$. There will be two types of processes - those processes that start from an occupied state (particle sector) and those that start from a vacant state (hole sector). Due to particle hole symmetry of the Hamiltonian, they will be equal to each other and we will only calculate the particle sector contribution.

In the particle sector, we have ($\hat{n}_{q\beta} = 1$), so we will work at a negative energy shell $\epsilon_q = -D$. The renormalisation can schematically represented as $H_0^I \frac{1}{\omega - H_{q\beta}^D} H_1^I$. Both H_0^I and H_1^I have all three operators $S_d^z, S_d^\pm$. We first consider specifically the case of spin- $\frac{1}{2}$ impurity. Those terms that have identical operators on both sides can be ignored because $S_d^{z2} = \text{constant}$ and $S^\pm{}^2 = 0$. All the six terms that *will* renormalise the Hamiltonian have a spin flip operator on at least one side of the Greens function. This means that in the denominator of the Greens function, S_d^z and s_{qq}^z have to be anti-parallel in order to produce a non-zero result for that term. This means we can identically replace $S_d^z s_{qq}^z = -\frac{1}{4}$. Also, in the particle sector, the Greens function always has $c_{q\beta}$ in front of it, so $\epsilon_q \tau_{q\beta} = \frac{D}{2}$. The upshot of all this is that the denominator of all scattering processes for the spin- $\frac{1}{2}$ impurity Hamiltonian will be $\omega - \frac{D}{2} + \frac{J}{4}$.

We now come to the general case of spin- S impurity. The various terms that renormalise the Hamiltonian can be described in terms of the bath spin operators that come into them. For example, the term that has s^z on both sides of the intervening Greens function can be represented as $z|z$. There are 7 such terms: $z|z, \pm|\mp, z|\pm, \pm|z$. Each of these terms occurs both in the particle and the hole sectors. We will demonstrate the calculation of two of these terms. The $z|z + |-\$ terms evaluate in the following manner.

$$\begin{aligned} z|z : & \sum_{kk', m, \sigma} c_{q\sigma}^\dagger c_{k'\sigma} |m\rangle \langle m| \frac{J_m^{\sigma 2}}{\omega - \frac{D}{2} + \frac{J}{2} \sigma S_d^z} |m\rangle \langle m| c_{k\sigma}^\dagger c_{q\sigma} = - \sum_{kk', m, \sigma} n_{q\sigma} \frac{J_m^{\sigma 2} c_{k\sigma}^\dagger c_{k'\sigma} |m\rangle \langle m|}{\omega_{m, \sigma} - \frac{D}{2} + \frac{J}{2} \sigma m}, \\ +| - : & \sum_{kk', m} c_{q\uparrow}^\dagger c_{k'\downarrow} |m\rangle \langle m+1| \frac{J_m^2}{\omega - \frac{D}{2} + \frac{J}{2} S_d^z} |m+1\rangle \langle m| c_{k\downarrow}^\dagger c_{q\uparrow} = -n_{q\uparrow} \sum_{kk', m} \frac{J_m^2 c_{k\downarrow}^\dagger c_{k'\downarrow} |m\rangle \langle m|}{\omega_{m+1, \uparrow} - \frac{D}{2} + \frac{J}{2} (m+1)}. \end{aligned} \quad (3.5)$$

We similarly compute the rest of the terms. We again define $\sum_q \hat{n}_{q\sigma} = n(D)$. To compare with the spin- $\frac{1}{2}$ RG equations, we will transform the general spin- S ω to the spin- $\frac{1}{2}$ ω , using $\omega_{m, \sigma} \rightarrow$

$\omega - \frac{J}{2} (m\sigma - \frac{1}{2})$. The renormalisation in J_m^σ is

$$\Delta J_m^\sigma = -n(D) \frac{(J_m^\sigma)^2 + \left(J_{m-\frac{1+\sigma}{2}}^t\right)^2}{\omega - \frac{D}{2} + \frac{J}{4}}. \quad (3.6)$$

Here, we have defined $J_m^t = 0$ for $|m| > S$. Two relations can be obtained from this RG equation, the RG equations for the sum and difference of the couplings: $J_m^\pm = \frac{1}{2} (J_m^\uparrow \pm J_m^\downarrow)$. The RG equation for the sum of the couplings is

$$\Delta J_m^+ = -n(D) \frac{\sum_\sigma (J_m^\sigma)^2 + \sum_\sigma \left(J_{m-\frac{1+\sigma}{2}}^t\right)^2}{2 \left(\omega - \frac{D}{2} + \frac{J}{4}\right)} = -n(D) \frac{J^2}{4} \frac{S(S+1)}{\omega - \frac{D}{2} + \frac{J}{4}}. \quad (3.7)$$

This is an m -independent piece, so it can be summed over to produce an impurity-independent potential scattering term, which we ignore.

The second is the RG equation for the difference of the couplings:

$$\Delta J_m^- = -n(D) \frac{1}{2} \frac{(J_{m-1}^t)^2 - (J_m^t)^2}{\omega - \frac{D}{2} + \frac{J}{4}} = -\frac{1}{4} \frac{n(D) m J^2}{\omega - \frac{D}{2} + \frac{J}{4}}. \quad (3.8)$$

The usual J Kondo coupling is recovered through $J = 2J_m^-/m$. Substituting this gives

$$\Delta_{\text{p sector}} J = -\frac{1}{2} n(D) \frac{J^2}{\omega - \frac{D}{2} + \frac{J}{4}}. \quad (3.9)$$

We can also obtain the RG equation for J from the transverse renormalisation:

$$\Delta J_m^t = -\frac{n(D) J_m^t (J_m^\downarrow + J_{m+1}^\uparrow)}{\omega - \frac{D}{2} + \frac{J}{4}} = -\frac{1}{2} \frac{n(D) J_m^t J}{\omega - \frac{D}{2} + \frac{J}{4}}. \quad (3.10)$$

Since $J_m^t \propto J$, we have

$$\Delta_{\text{p sector}} J = -\frac{1}{2} \frac{n(D) J^2}{\omega - \frac{D}{2} + \frac{J}{4}}. \quad (3.11)$$

The total renormalisation from both particle and hole sectors, at this order, is

$$\Delta J^{(2)} = -\frac{n(D) J^2}{\omega - \frac{D}{2} + \frac{J}{4}}. \quad (3.12)$$

We now come to the third-order renormalisation. Following eq. (3.3), the next order renormalisation is

$$\Delta H_j^{(3)} = c^\dagger T \frac{1}{\omega' - H_D} H_X \frac{1}{\omega' - H_D} T^\dagger c + c^\dagger T \frac{1}{\omega' - H_D} T^\dagger c \frac{1}{\omega' - H_D} H_X. \quad (3.13)$$

The first term will be of the form

$$\sum_{q,k,l_1} c_{q\beta,l_1}^\dagger c_{k\alpha,l_1} |m_1\rangle \langle m_2| \frac{1}{\omega - \frac{D}{2} - \frac{\epsilon_k}{2} + \frac{\beta J}{4} S_d^z} |m_2\rangle c_{k_1\sigma_1,l_2}^\dagger c_{k_2\sigma_2,l_2} \langle m_3| \frac{1}{\omega - \frac{D}{2} - \frac{\epsilon_k}{2} + \frac{\beta J}{4} S_d^z} |m_3\rangle \langle m_4| c_{k\alpha,l_1}^\dagger c_{q\beta,l_1}. \quad (3.14)$$

We have not bothered to write all the summations and the couplings correctly, because we will only simplify the denominator here. Evaluating the inner products gives

$$\sum_{qk,l_1} \frac{|m_1\rangle \langle m_4| c_{q\beta}^\dagger c_{k\alpha} c_{k_1\sigma_1,l_2}^\dagger}{\omega_{m_2,\beta} - \frac{D}{2} - \frac{\epsilon_k}{2} + \frac{\beta J m_2}{4}} \frac{c_{k_2\sigma_2,l_2} c_{k\alpha}^\dagger c_{q\beta}}{\omega_{m_3,\beta} - \frac{D}{2} - \frac{\epsilon_k}{2} + \frac{\beta J m_3}{4}}. \quad (3.15)$$

We again use $\omega_{m,\sigma} \rightarrow \omega - \frac{J}{2} (m\sigma - \frac{1}{2})$.

$$|m_1\rangle \langle m_4| c_{k_1\sigma_1,l_2}^\dagger c_{k_2\sigma_2,l_2} \sum_{qk,l_1} \frac{\hat{n}_{q\beta} (1 - \hat{n}_{k\alpha})}{\left(\omega - \frac{D}{2} - \frac{\epsilon_k}{2} + \frac{J}{4}\right)^2}. \quad (3.16)$$

We define $\sum_q \hat{n}_{q\beta} = n(D)$. Performing the sums over k and l_1 gives

$$-\frac{1}{2} |m_1\rangle \langle m_4| c_{k_1\sigma_1,l_2}^\dagger c_{k_2\sigma_2,l_2} \frac{\rho n(D) K}{\omega - \frac{D}{2} + \frac{J}{4}}. \quad (3.17)$$

ρ is the density of states which we have taken to be constant. Reinstating the complete summation and the couplings gives

$$-\frac{1}{2} \frac{\rho n(D) K}{\omega - \frac{D}{2} + \frac{J}{4}} \sum_{\substack{m_1, m_4, k_1, k_2, \\ l_2, \sigma_1 \sigma_2}} \lambda_1 \lambda_2 \lambda_3 |m_1\rangle \langle m_4| c_{k_1\sigma_1,l_2}^\dagger c_{k_2\sigma_2,l_2}. \quad (3.18)$$

There is no sum over m_2 and m_3 because they are constrained by m_1 and m_4 respectively. λ_i represent the couplings present at the three interaction vertices. $k_{1,2}$ sum over the momenta, $\sigma_{1,2}$ sum over the spin indices and l_2 sums over the channels.

The second term in eq. (3.13) can be evaluated in an almost identical fashion. The integral here will be negative of the first term, because of an exchange in the scattering processes.

$$\frac{1}{2} \frac{\rho n(D) K}{\omega - \frac{D}{2} + \frac{J}{4}} \sum_{\substack{m_1, m_4, k_1, k_2, \\ l_2, \sigma_1 \sigma_2}} \lambda_1 \lambda_3 \lambda_2 |m_1\rangle \langle m_4| c_{k_1\sigma_1,l_2}^\dagger c_{k_2\sigma_2,l_2}. \quad (3.19)$$

The first group of terms (those that appear in (3.18)) in the particle sector can be represented as $a|b\rangle|c\rangle$, where $a, b, c \in \{z, +, -\}$ and represent the operator for the conduction electrons in the three connected processes. The $\prime$ on b indicates that it is the state of the electrons *not being decoupled*. The second group of terms (those that appear in (3.19)) are therefore represented as $a|b\rangle|c\rangle$, because in this group, the interaction H_X between the electrons that are not being decoupled occur at the very end. We will only calculate the terms in the particle sector, the ones in hole sector will be equal to these because of particle-hole symmetry. The full list of terms is:

$$\begin{array}{cccccccccccc} z|z'|z & z|z|z' & -|z'|+ & -|+|z' & +|z'|- & +|-|z' & z|+|z & z|-|z & +|+|z & +|+|z & -|z'|+ & -|z'|+ \\ -|+|z & z|z|+ & z|z|- & +|-|+ & +|-|- & -|+|+ & z|z'|z & z|z|z' & -|z'|+ & -|z'|+ & -|z'|+ & -|z'|+ \\ -|+|z' & +|z'|- & +|-|z' & z|+|z & z|-|z & +|+|z & -|+|+ & -|+|+ & z|z|+ & z|z|+ & z|z|+ & z|z|+ \\ +|-|+ & +|-|- & -|+|+ & -|+|+ & -|+|+ & -|+|+ & -|+|+ & -|+|+ & -|+|+ & -|+|+ & -|+|+ & -|+|+ \end{array}.$$

The total renormalisation in J_m^σ is

$$\begin{aligned}\Delta J_m^\sigma &= \frac{1}{2} \frac{\rho n(D)K}{\omega - \frac{D}{2} + \frac{J}{4}} \left[(J_{m-1}^t)^2 J_m^\sigma + (J_m^t)^2 J_m^\sigma - (J_{m-1}^t)^2 J_{m-1}^\sigma - (J_m^t)^2 J_{m+1}^\sigma \right] \\ &= \frac{1}{2} \frac{\rho n(D)K}{\omega - \frac{D}{2} + \frac{J}{4}} J^3 m \sigma.\end{aligned}\quad (3.20)$$

Since we had defined $J_m^\sigma \equiv \frac{1}{2} J m \sigma$, we have $\Delta J = \frac{2}{m\sigma} \Delta J_m^\sigma$, and we get $\Delta_{\text{p sector}} J = \frac{1}{4} \frac{\rho n(D)K}{\omega - \frac{D}{2} + \frac{J}{4}} J^3$. Combining with the hole sector renormalisation, we get

$$\Delta J^{(3)} = \frac{1}{2} \frac{\rho n(D)K}{\omega - \frac{D}{2} + \frac{J}{4}} J^3. \quad (3.21)$$

The total renormalisation in J after combining all orders is

$$\Delta J = -\frac{n(D)J^2}{\omega - \frac{D}{2} + \frac{J}{4}} + \frac{1}{2} \frac{\rho n(D)K}{\omega - \frac{D}{2} + \frac{J}{4}} J^3. \quad (3.22)$$

2.4 Example III: Particle in a periodic potential

We consider electrons moving in one dimension in the presence of a periodic potential $V(x)$ satisfying $V(x+a) = V(x)$. The full Hamiltonian can be represented as

$$\mathcal{H} = \int_{-\infty}^{\infty} dx \, c^\dagger(x) \left[\hat{p}^2/2m + V(x) \right] c(x), \quad (4.1)$$

where $c(x)$ destroys a fermion at the position x .

Physically, the model represents a set of N_{well} number of wells (located at the minima of the potential) linked to each other through hopping (brought about by the kinetic energy term). Each well is described a potential $V(x)$, whose spectrum consists of N_{levs} single particle levels. These levels are localised in real space (around each minima of width a), so the corresponding momentum states are widely separated; each of these momentum states eventually give rise to an energy band, we therefore represent them using the band index: $k_n = nG$ ($n = 0, 1, \dots, N_{\text{levs}} - 1$), where G is the spacing between the momentum states arising from a single potential well, and is equal to the reciprocal lattice vector $2\pi/a$.

These are of course not the only momentum states in the problem; since each potential well is spaced far apart (at distances greater than a), the multiplicity of the wells leads to closely spaced momentum states, N_{well} in number (the number of potential wells). Since these are closely spaced, they lead to the dense set of states within each band: $k_{n,j} = nG + \frac{2\pi j}{N_{\text{well}}}$ ($j = 0, 1, \dots, N_{\text{well}} - 1$). The spacing $2\pi/N_{\text{well}}$ between the states within the band is imposed by applying periodic boundary conditions on the system: $\psi(x=0) = \psi(x=aN_{\text{well}})$. We take the thermodynamic limit by allowing $N_{\text{well}} \rightarrow \infty$, keeping all other independent parameters unchanged. Because of the periodicity of the real space potential, it can only scatter states that are separated by a reciprocal lattice vector G ; this means that states within a band (that are within a reciprocal lattice vector of each other) do not scatter off each other: $\langle k_{n+m,j+l} | \hat{V} | k_{n,j} \rangle = \delta_{l,0} \langle k_{n+m,j} | \hat{V} | k_{n,j} \rangle$.




Figure 2.6: The complete set of momentum states in the system. Each row represents a band of states. Each column contains the set of momenta that are connected by a reciprocal lattice vector nG . The states in the lowest band give rise to the crystal momenta. All the momentum states in a column are connected to each other through the potential term (green dotted line). Any two states within a band, on the other hand, are decoupled from each other.

Decomposition into sum of zero-dimensional problems

We begin by writing the Hamiltonian completely in momentum space. A general momentum is represented as $k + nG$, where n is the band index goes from 0 to $N - 1$, and k is the crystal momentum that spans through the lowest band. With this notation, we have

$$\begin{aligned} \mathcal{H} &= \frac{\hbar^2}{2m} \int dk \sum_n c^\dagger(k + nG) (k + nG)^2 c(k + nG) \\ &\quad + \int dk \sum_{m,n} [\mathcal{V}(nG) c^\dagger(k + nG + mG) c(k + mG) + \text{h.c.}] \\ &= \frac{2\pi}{a} \int dk \mathcal{H}(k) \end{aligned} \quad (4.2)$$

where

$$\begin{aligned} \mathcal{H}(k) &= \frac{a}{2\pi} \frac{\hbar^2}{2m} \sum_n c^\dagger(k + nG) (k + nG)^2 c(k + nG) \\ &\quad + \frac{a}{2\pi} \sum_{m,n} [v(nG) c^\dagger(k + nG + mG) c(k + mG) + \text{h.c.}] \\ &= \frac{a}{2\pi} \int dx' c^\dagger(x') \left[\frac{1}{2m} (-i\hbar \partial_{x'} + \hbar k)^2 + V(x') \right] c(x'). \end{aligned} \quad (4.3)$$

We have used x' to symbolise the position variable conjugate to the new set of momenta $\{nG\} = -i\partial_{x'}$ that span the restricted Hilbert space of $\mathcal{H}(k)$. Also, $\mathcal{V}(k)$ is the Fourier transform of the real space potential $V(x)$.

This decoupling of the full Hamiltonian $\mathcal{H}$ into a block-diagonal form (each block indexed by the crystal momentum k) can be attributed to the discrete translation symmetry present in the problem and could have been anticipated very generally, as we now show. The Hamiltonian satisfies




Figure 2.7: Because of the periodicity of the potential, the problem splits into a sum of particle on a circle (POC) problems, each formed by a specific crystal momentum and other momentum states that are related to it by a reciprocal vector.

$[\mathcal{H}, T(a)] = 0$, where $T(x) = e^{i\hat{p}x}$ is the translation operator. By choosing a starting location $x = 0$, applying the translation operator cycles through the set of states $|x = na\rangle$; $n = 0, 1, \dots, N_{\text{well}}$; this set is closed under the application of $T(a)$. It is easy enough to guess the eigenstates of $T(a)$. Since the application of $T(a)$ increases the position monotonically, the only way to create an eigenstate is to involve all states. With that in mind, the simplest ansatz is of the form $|\Phi\rangle = \sum_n e^{-i\Phi n} |an\rangle$. Applying the translation operator on this state gives

$$T(a) |\Phi\rangle = \sum_{n=0}^{N_{\text{well}}-2} e^{-i\Phi n} |a(n+1)\rangle + e^{-i\Phi(N_{\text{well}}-1)} |aN_{\text{well}}\rangle = e^{i\Phi} \left[e^{-i\Phi N_{\text{well}}} |aN_{\text{well}}\rangle + \sum_{n=1}^{N_{\text{well}}-1} e^{-i\Phi n} |an\rangle \right] \quad (4.4)$$

Since we have applied PBC on the system, we used $|aN_{\text{well}}\rangle$ for $|0\rangle$. If the values of Φ are chosen to be some integer multiple of $2\pi/N_{\text{well}}$, we would have $e^{-i\Phi N_{\text{well}}} = 1$, and the entire state can be resummed into the form

$$T(a) |\Phi\rangle = e^{i\Phi} |\Phi\rangle \quad (4.5)$$

This shows that $|\Phi\rangle$ is an eigenstate of the translation operator. Since the set of states has a dimension of N_{well} , the number of eigenstates and the number of possible values of Φ will also be N_{well} . This fixes the eigenvalues to $\Phi = 2\pi m/N_{\text{well}}$, $m = 0, 1, \dots, N_{\text{well}} - 1$. These will also be eigenstates of the Hamiltonian, because they are indexed by different values of Φ and the Hamiltonian cannot change Φ (because it commutes with T). As will be shown later, the eigenvalues Φ are actually related to the crystal momentum, through $\Phi = ka$. The set of allowed values of Φ form the first Brillouin zone.

However, $|\Phi\rangle$ as defined above is not the only eigenstate with eigenvalue $e^{i\Phi}$. One can simply shift the starting location $x = 0$ by any distance smaller than a , and repeat the entire procedure. The eigenstates hence obtained will have the same eigenvalues as above:

$$|\Phi, \delta\rangle = \sum_n e^{-i\Phi n} |an + \delta\rangle, \quad 0 \leq \delta < a; \quad (4.6)$$

$$T(a) |\Phi, \delta\rangle = \sum_{n=0}^{N_{\text{well}}-2} e^{-i\Phi n} |a(n+1) + \delta\rangle + e^{-i\Phi(N_{\text{well}}-1)+\delta} |aN_{\text{well}} + \delta\rangle = e^{i\Phi} |\Phi, \delta\rangle \quad (4.7)$$

The translation operator is therefore able to at most classify the complete set of states into N_{well} number of blocks, each labelled by a specific value of Φ . Each block has a large number of states (N_{levs}), and the Hamiltonian will general scatter these states into each other. The Hamiltonian for each of these blocks can be represented as $\mathcal{H}(\Phi) = \mathcal{P}_\Phi \mathcal{H} \mathcal{P}_\Phi$ (where $\mathcal{P}_\Phi$ projects onto the block of states with eigenvalue $e^{i\Phi}$), such that the total Hamiltonian can be represented as $\mathcal{H} = \sum_\Phi \mathcal{H}(\Phi)$. Upon rescaling by $2\pi/a$ and taking the limit of $N_{\text{well}} \rightarrow \infty$, this is the same Hamiltonian decoupling relation as that obtained in eq. 4.2.

Since the spacing between two adjacent momentum is G , the maximum value of x' has to be $2\pi/G = a$, with the ends identified with each other: $\psi(x' = 0) = \psi(x' = a)$. We can then define a dimensionless variable $\theta = 2\pi x'/a$ that goes from 0 to 2π , and is conjugate to the angular wave number $Q = -i\partial_\theta$. This definition ensures $[\theta, Q] = i$. For reasons that will become clear very soon, we will also define a dimensionless crystal momentum $\Phi = ka$. With these modifications, the Hamiltonian takes the simpler form

$$\mathcal{H}(k) = \int_0^{2\pi} d\theta \, c^\dagger(\theta) \left[\frac{\hbar^2}{2ma^2} \left(\hat{Q} + \Phi/2\pi \right)^2 + U(\theta) \right] c(\theta). \quad (4.8)$$

By noting that we have the boundary condition $\psi(\theta = 0) = \psi(\theta = 2\pi)$, we can recognise this Hamiltonian as that describing the dynamics of a particle moving on a circle (POC) with a potential $U(\theta) = \frac{a}{2\pi} V(a\theta/2\pi)$, in the presence of an Aharonov-Bohm flux Φ . Note that $U(\theta)$ has a period of 2π .




Figure 2.8: The full problem reduces to that of a particle on a circle, in the presence of potential with period 2π , and a flux Φ .

Even when reduced to the simpler POC, the problem cannot be simply integrated to obtain the eigenstates. This is because the complete Hamiltonian does not commute with either the position or the momentum operator; it contains quantum fluctuations that need to be resolved in order to obtain the ground state energy. One way of obtaining the energy is to calculate the propagator for transition from $\theta = 0$ to $\theta = 2\pi$, using the method of path integrals [?, ?]. This is shown in Appendix ?? . We, however, will attack the problem using a renormalisation group method, called the unitary renormalisation group in order to track the emergence of the tight-binding metal in the limit when the effective amplitude of the single-particle periodic potential at low-energies has become much larger than the typical kinetic energy of the electron.

URG analysis of the particle on a circle problem

A summary of the unitary RG method is provided in Appendix ?? . For the purposes of studying the model under renormalisation group, we write the particle on a circle Hamiltonian in momentum(Q –)space. For that, we introduce the Fourier modes

$$c(\theta) = \frac{1}{\sqrt{2\pi}} \int dQ e^{iQ\theta} c(Q), \text{ and } U(\theta) = \int dQ e^{iQ\theta} \mathcal{U}(Q), \quad (4.9)$$

so that the Hamiltonian becomes

$$\mathcal{H}(k) = \int dQ \varepsilon(Q + \Phi/2\pi) \hat{n}(Q) + \int dQ_1 dQ_2 c^\dagger(Q_1) c(Q_2) \mathcal{U}(Q_1 - Q_2). \quad (4.10)$$

where we have defined the dispersion $\varepsilon(Q) = \hbar^2 Q^2 / (2ma^2)$. The second term involves scattering between momentum states Q_1 and Q_2 . The quantum fluctuations that connect the low-energy states to the high-energy states can be integrated out using the URG method [?]. This proceeds through a reduction in the momentum cutoff, ultimately leading to an IR theory. As the large momenta are integrated out, the potential term for scattering between the lower momenta undergo renormalisation. Let, at the l^{th} RG step, the momentum cutoff be Q_l . At this step, fluctuations between all momentum states more energetic than Q_l have been integrated out. The RG equations for the renormalisation of the potential $\mathcal{U}_{ij}^{(l)}$ that scatters any two lower momenta Q_i and Q_j ($Q_i, Q_j < Q_l$) is given by [?]

$$\Delta \mathcal{U}_{ij}^{(l)}(\omega) = \frac{\mathcal{U}_{il} \mathcal{U}_{lj}}{\omega - \varepsilon(Q_l + \Phi/2\pi)}, \quad (4.11)$$

where ω is an energy scale that tracks the amount of quantum fluctuations present in the system, and is characteristic of the URG method. The dispersion also undergoes renormalisation::

$$\Delta \varepsilon^{(l)}(Q_i, \omega) = \frac{|\mathcal{U}_{il}|^2}{\omega - \varepsilon(Q_l + \Phi/2\pi)}, \quad (4.12)$$

Effective description at low energies

Appearance of bands and band gaps

As the URG proceeds, the quantum fluctuations between the momentum states are resolved, leading to a simpler theory for the IR. We focus on the final two states $|Q_0 + \Phi/2\pi\rangle, |Q_1 + \Phi/2\pi\rangle$, for $\Phi = 0$ (Brillouin zone edge). The Hamiltonian for their dynamics, just prior to the decoupling of Q_j , is

$$H_{01}^* = \varepsilon^*(Q_0) |Q_0\rangle \langle Q_0| + \varepsilon^*(Q_1) |Q_1\rangle \langle Q_1| + (\mathcal{U}_{01}^* |Q_1\rangle \langle Q_0| + \text{h.c.}) \quad (4.13)$$

Since both the states are at the edge of the band, we can assume $\varepsilon^*(Q_0) \simeq \varepsilon^*(Q_1)$, in which case the problem becomes that of a symmetric double well with tunnelling. The eigenstates are $|Q_0\rangle \pm |Q_1\rangle$, with energies $\varepsilon^*(Q_0) \pm |\mathcal{U}_{01}^*|$. We can substitute this estimate of the energy eigenvalues into the RG

equations in the form of the quantum fluctuation scale ω . At the last decoupling, the shifts in the kinetic energy are therefore

$$\Delta\varepsilon^*(Q_i, \omega^\pm) \simeq \frac{|\mathcal{U}_{01}^*|^2}{\varepsilon^*(Q_0) \pm |\mathcal{U}_{01}^*| - \varepsilon(Q_0)} = \pm |\mathcal{U}_{01}^*| = \pm |\mathcal{U}^*(G)|, \quad (4.14)$$

where we used $\mathcal{U}_{01} = \mathcal{U}(Q_1 - Q_0) = \mathcal{U}(G)$. The difference between the two shifts gives the energy difference between the two lowest states of the POC. In this way, the above RG equation shows the emergence of the single-particle band gap and the formation of the band insulator (if the chemical potential is placed within the gap). If we now reintroduce the flux degree of freedom, the lowest energy eigenstates of all the POCs can be combined to form a band of closely spaced energy levels $\{\varepsilon^*(Q_0 + \Phi_j/2\pi)\}$, $\Phi = k_{0,j}a$ (see Fig. 2.9). These levels are indexed by $k_{0,j}$ with j ranging through the first Brillouin zone; $k_{0,j}$ are the crystal momenta of the system.

The second band of states is formed similarly by considering the first excited of the POCs: $\{\varepsilon^*(Q_1 + \Phi_j/2\pi)\}$, $\Phi = k_{0,j}a$. The gap between the first and second band is set by the energy difference between the ground state and first excited state of the POC at $\Phi = 0$. This difference, and hence the gap, is equal to $2|\mathcal{U}_{01}^*|$.




Figure 2.9: Spectral flow transformations on the POC by tuning the flux Φ generates the full band of delocalised single-particle electronic states whose crystal momenta correspond to $\hbar\Phi/a$.

Dispersion relation of the lowest band

In order to derive the dispersion for the lowest band, we note that the final fixed point Hamiltonian (obtained upon decoupling all higher momentum states) is described by the single momentum state $|Q_0\rangle$. In terms of real space hopping, this means that all short wavelength processes have been removed, and the only process that remains is the longest hopping, that from 0 to 2π . The fixed point Hamiltonian can be written in terms of this process:

$$\begin{aligned} \varepsilon^*(Q_0) |Q_0\rangle \langle Q_0| &= \frac{1}{2} \varepsilon^*(2\pi) (|\theta = 0\rangle + |\theta = 2\pi\rangle) (\langle\theta = 0| + \langle\theta = 2\pi|) = \frac{1}{2} \varepsilon^*(2\pi) (\hat{n}(0) + \hat{n}(2\pi)) \\ &\quad + \frac{1}{2} \varepsilon^*(2\pi) (c^\dagger(0)c(2\pi) + \text{h.c.}) \end{aligned} \quad (4.15)$$

We now recall that we had mapped the spatial point $x = 0$ ($x = a$) to the angular point $\theta = 0$ ($\theta = 2\pi$) point. We now transform back to spatial coordinates to get

$$\frac{1}{2}\varepsilon^*(2\pi) (\hat{n}(0) + \hat{n}(a)) + \frac{1}{2}\varepsilon^*(a) (c^\dagger(0)c(a) + \text{h.c.}) . \quad (4.16)$$

This is computed for a specific value $\Phi = 0$ for the flux (or the crystal momentum). The POC corresponding to a particular flux is built from the states within a particular well. Changing the flux is then equivalent to studying the POC corresponding to the states within a different well. What this means is that tuning the flux by one interval amounts to translating the fixed point Hamiltonian by one period a of the potential. Adding the contributions from all the values of the flux leads to the following tight-binding Hamiltonian:

$$H_{TB} = \varepsilon^*(2\pi) \sum_{j=0}^{N_{\text{well}}-1} \hat{n}(ja) + \frac{1}{2}\varepsilon^*(a) \sum_{j=0}^{N_{\text{well}}-1} (c^\dagger(ja)c((j+1)a) + \text{h.c.}) , \quad (4.17)$$

where the first term corresponds to a constant chemical potential applied to the tight-binding metal in one dimension with a single-particle dispersion $\varepsilon(k) = -\varepsilon^*(a) \cos(ka)$ where k corresponds to the set of crystal momenta $k \in [-\frac{\pi}{a}, \frac{\pi}{a}]$. The exercise, therefore, shows the emergence of nearest-neighbour hopping in the lowest band by starting from a more microscopic model of a particle in a periodic potential. (An alternate derivation of the tight-binding metal has also been shown from path-integral methods in Appendix ??).

2.5 Comparison of renormalisation group approaches based on canonical transformations

We have considered three canonical transformations in this section: the Poor Man's scaling (PMS), the Schrieffer-Wolff transformation (SWT) and the continuous unitary transformation renormalization group (CUT-RG). PMS and SWT are more or less identical; they differ in the context in which they are used. PMS is used when there is an entire spectrum of energies in the model, ranging from a low energy limit to a high energy; it is then employed to decouple the highest energy modes in an iterative fashion. SWT is used when the Hamiltonian can be split into one high and one low energy part, and we need to decouple these two modes. It is clear when seen from this perspective that SWT is like a one-shot PMS; it decouples the UV from the IR in one step, compared to the iterative approach of PMS. However, as has been shown in the previous subsections, both PMS and SWT can be formalized in an identical fashion, so that one can be switched for the other in both contexts.

Now that we have established that PMS and SWT are formally identical, we can relate them to URG. URG is similar in philosophy to PMS in the fact that URG also successively decouples high energy modes from the low energy ones. The difference is that URG takes care of the quantum fluctuations, at least in principle, by introducing a new set of energy scales ω . These ω then parameterise the RG flows of URG, compared to the single RG flow of PMS. Since SWT is formally the same as PMS, there is also a comparison between SWT and URG. Both PMS and SWT trivialize the quantum fluctuation scales of URG by replacing it with the bare energy values.

CUT-RG is philosophically different from the other transformations. Its goal is to gradually reduce the contributions of the off-diagonal terms by making them decay along a certain scale l . Off-diagonal terms that connect states with large energy differences decay faster. In this sense, there is no sequential dropping out of off-diagonal terms; all off-diagonal terms disappear at $l = \infty$. In this sense, it is like a continuous version of SWT. While SWT strives to remove an entire off-diagonal term in one-shot, CUT RG does this gradually by introducing a scale l . This separation of scales is absent in SWT. It does exist in URG and PMS, albeit in a different fashion. There, the separation comes in when we decouple single electron states starting from the Brillouin zone boundary Λ_N and come down to the Fermi surface Λ_0 . Each Λ provides a natural energy scale for separating the high and low energy physics.

If one integrates the continuum generator η over all the scales, one should recover something analogous to the SWT generator. This generator is however necessarily perturbative in the off-diagonal term, since by definition η will only be of first order in $\mathcal{H}_X$. This is in contrast to the non-perturbative generators in URG and, at least in principle, PMS. This non-perturbative form is encapsulated in the presence of a second completely different set of energy scales ω (or E in PMS). This second energy scale is absent in CUT RG because it takes care of at most the second order term.

Another point to note is that since SWT keeps the entire off-diagonal piece in the generator, new terms will almost certainly be generated at every step, and they have to be truncated. This is not the case with URG or PMS, because in those methods, we decouple just a single-electron at each step, and so those electrons become integrals of motion at that step, leading to their removal from the off-diagonal piece, and very often the Hamiltonian retains the same form as the bare model. This makes the philosophy of URG and PMS easier to work with. Tied to this is the fact that CUT RG often takes a certain type of interaction in the generator part, and not the entire off-diagonal piece. Hence, at the limit of the flow parameter going to ∞ , the chosen off-diagonal piece goes to zero but the remaining off-diagonal pieces still remain. As a result, the Hamiltonian is at most block diagonal at this stage. On the other hand, URG progressively decouples single electrons, meaning all scattering terms corresponding to that electron vanish at each step.

One last thing to note is that URG, being unitarily implemented with a well-defined generator, does not accommodate for spontaneous symmetry breaking (SSB). In order to see SSB, the symmetry-breaking term has to be added to the bare model explicitly; if this term grows under the RG, then the ground state will be symmetry-broken. CUT RG, however, does allow for SSB through the idea that the generator is not uniquely defined. If the generator commutes with a particular symmetry, then the family of Hamiltonians will also have the symmetry [Kehrein(2007)]. However, if the generator is replaced by something that is normal ordered w.r.t a particular expectation value, then the CUT RG flow will usually be towards either the symmetry-preserved state or the symmetry-broken state, depending on whichever is stable [Wegner(1994)].




Figure 2.10: Comparison of the various unitary transformations and their relationships to each other.

2.6 Iterative Diagonalisation of Quantum Impurity Models

The broad idea is the following. We have a Hamiltonian describing an interacting set of states (in real or momentum space). We can express the total Hamiltonian in an incremental fashion: $H = \sum_i H_i$, where H_{i+1} involves more number of states than H_i . For example, a tight-binding model can be written in that form, with the definition $H_i = c_{i+1}^\dagger c_i + \text{h.c.}$. The iterative diagonalisation method obtains the low-energy spectrum of this problem in the following manner: We first diagonalise the Hamiltonian H for a smaller value of i , small enough such that this can be done exactly. We

then truncate the spectrum to a predefined size, and rotate all existing operators to this truncated basis, including the Hamiltonian H . We then consider the "bonding Hamiltonian" ΔH between the existing sites and the new sites, and rotate the same into the truncated basis. Adding the previous rotated Hamiltonian H and the rotated increment Hamiltonian ΔH gives us a truncated but effective Hamiltonian for the increased number of sites. We again diagonalise this, and again retain only a fixed number of states in the spectrum. We keep repeating this until we reach the required number of sites.

Jordan-Wigner Matrix Representation of Fermionic Operators

For a single qubit, the creation/annihilation matrices are

$$c = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, c^\dagger = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad (6.1)$$

given the convention that $|1\rangle = (0 \ 1)$ is the occupied state. For a many-body system, these must be replaced with field operators that have the canonical fermionic algebra. For a system of N 1-particle levels, this can be accomplished through a *Jordan-Wigner-like* transformation

$$c_j = (\otimes_1^j \sigma_z) \otimes c \otimes (\otimes_1^{N-j} \mathbb{I}), j \in [0, N-1], \quad (6.2)$$

where $\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and $\mathbb{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. For example, for spinless electrons on a three-site lattice, we can have three fermionic operators: c_1, c_2 , and c_3 . Following the above expression, their matrices are

$$c_1 = \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}, c_2 = \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}, c_3 = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix},$$

For later use, we define the *antisymmetriser matrix* $S(j)$ and identity matrix $\mathbb{I}(j)$,

$$\begin{aligned} S(j) &= \otimes_1^j \sigma_z = \sigma_z \otimes \dots j \text{ times } \dots \otimes \sigma_z, \\ \mathbb{I}(j) &= \otimes_1^{N-j} \mathbb{I} = \mathbb{I} \otimes \dots j \text{ times } \dots \otimes \mathbb{I}, \end{aligned} \quad (6.3)$$

to express the fermionic representation compactly:

$$c_j = S(j) \otimes c \otimes \mathbb{I}(N-j). \quad (6.4)$$

Structure of Hamiltonian

We consider a general impurity model, of L number of sites (excluding the impurity site):

$$H_{\text{sys}}(L) = H_{\text{imp}} + H_{\text{imp-bath}} + -t \sum_{\sigma, j=1}^{L-1} \left(c_{j,\sigma}^\dagger c_{j+1,\sigma} + \text{h.c.} \right), \quad (6.5)$$

where H_{imp} is the Hamiltonian for the decoupled impurity site and $H_{\text{imp-bath}}$ is the impurity-bath coupling. Such a system has $2L + 2$ single-particle levels in the full system (2 spin levels for each of the $L + 1$ sites).

Iteration scheme (the initial number of states is L_0):

$$\begin{aligned} H_0(L_0) &= H_{\text{sys}}(L_0), \\ H_{r+1}(L_0) &= H_r(L_0) + \Delta H_r(L_0), \quad r \geq 0, \\ \Delta H_r(L_0) &= -t \sum_{\sigma} \left(c_{L_0+r,\sigma}^\dagger c_{L_0+r+1,\sigma} + \text{h.c.} \right). \end{aligned} \quad (6.6)$$

H_0 is the initial Hamiltonian, consisting of L_0 lattice sites in the bath, $H_{r+1}(L_0)$ is the Hamiltonian after $r + 1$ iterations having started with L_0 sites, and $\Delta H_r(L_0)$ is the increment term that gets added to $H_r(L_0)$ during the $(r + 1)^{\text{th}}$ iteration to give rise to $H_{r+1}(L_0)$.

Iterative Diagonalization

Let the starting Hamiltonian be $H_0(L_0)$, consisting of L_0 single-particle levels (hence a Hilbert space dimension of 2^{L_0}). We start with a single-particle computational basis and construct the L_0 fermionic operators $c_1, c_2, \dots, c_{L_0}$ in this basis, using eq. 6.4. We also keep track of a large antisymmetriser matrix $S(L_0)$ that will be used to attach new sites when we expand the system. Let M_s be the maximum number of eigenstates we retain in the spectrum at any given step. The value of M_s should be chosen so that a $M_s \times M_s$ matrix can be diagonalised in reasonable time.

- S1. Construct the complete Hamiltonian matrix H_0 in our present basis using the field operators $\{c_j\}$. Diagonalise the Hamiltonian (of size $D_0 \times D_0$) and obtain the eigenvalues E_n and eigenstates X_n . Each X_n is a column vector of size D_0 .
- S2. Retain at most M_s number of eigenstates, preferring the ones with lower energy. The reduced basis for this rotated truncated subspace is constructed by stacking the column vectors X_n horizontally:

$$R = [X_1 X_2 \dots X_{M_s}]_{D_0 \times M_s}. \quad (6.7)$$

This matrix also acts as the transformation to rotate and truncate all operators from the old basis into the new one.

- S3. We rotate our Hamiltonian H_0 , our fermionic operators $\{c_j\}$ and the large antisymmetrizer matrix $S(L_0)$ into the new reduced basis, using the transformation $\mathcal{O} \rightarrow R^\dagger \mathcal{O} R$.

- S4. We now need to expand our system by adding the increment Hamiltonian ΔH_0 . Let the number of *new* 1-particle levels in ΔH_0 be L_Δ . These new levels will be indexed as $L_0 + 1, \dots, L_0 + L_\Delta$. We need to define antisymmetrized fermionic operators for the new sites:

$$\begin{aligned} c_{L_0+1} &= c \otimes \mathbb{I}(2^{L_\Delta-1}), \\ c_{L_0+2} &= \sigma_z \otimes c \otimes \mathbb{I}(2^{L_\Delta-2}), \\ &\dots \\ c_{L_0+L_\Delta} &= (\otimes_1^{L_\Delta-1} \sigma) \otimes c. \end{aligned} \tag{6.8}$$

These have to be calculated in the local computational basis (of size 2^{L_Δ}) of the new levels.

- S5. Combining the new sites with the old sites leads to a combined Hilbert space dimension of $(M_s + 2^{L_\Delta}) \times (M_s + 2^{L_\Delta})$. To allow all operators to act on the enlarged Hilbert space, we expand both sets of operators:

$$\begin{aligned} c_j &= S(L_0) \otimes c_j; \quad j = L_0 + 1, \dots, L_\Delta, \\ c_j &= c_j \otimes \mathbb{I}(2^{L_\Delta}); \quad j = 1, 2, \dots, L_0, \\ H_0 &\rightarrow H_0 \otimes \mathbb{I}(2^{L_\Delta}), \\ S(L_1) &= S(L_0) \otimes \mathbb{I}(2^{L_\Delta}). \end{aligned} \tag{6.9}$$

where $\mathbb{I}(2^{L_\Delta})$ is an identity matrix of dimension $2^{L_\Delta} \times 2^{L_\Delta}$. Note that the operators in the last three equations are the rotated ones (following Step 3).

- S6. Using the transformed operators $c_{L_0+1}, \dots, c_{L_0+L_\Delta}$ for the new sites, construct the difference Hamiltonian matrix ΔH_0 and hence the updated Hamiltonian $H_1 = H_0 + \Delta H_0$ for the next step. Repeat the process starting from step 2 with the new Hamiltonian H_1 , the new operators c_j and the new matrix $S(L_1)$ replacing the old counterparts.

Static correlations

We are in general interested in n -point correlations of the form $\langle \mathcal{O}_1 \mathcal{O}_2 \dots \mathcal{O}_n \rangle$, where $\mathcal{O}_i$ are operators that act on 1-particle Hilbert spaces. Let the earliest step of the iterative diagonalisation procedure at which all these operators have entered the system be r . At this step r , construct the correlation operator $\mathcal{O}_1 \mathcal{O}_2 \dots \mathcal{O}_n$ using the fermionic matrices c_j at that step (recall that these matrices will be highly rotated versions of the matrices we started with). Having constructed the operator O , we expand and rotate it after the completion of every future step: $O_{n+1} = (R_n^\dagger O_n R_n) \otimes \mathbb{I}(2^{L_\Delta})$, where R_n is the rotation matrix for the n^{th} step. The last step n^* of the iterative diagonalisation consists of only a diagonalisation and no expansion, resulting in a final set of eigenstates $\{X_i\}$. The form of the correlation operator in this basis is $O_{n^*} = R_{n^*-1}^\dagger O_{n^*-1} R_{n^*}$. The expectation value can now be calculated using the matrix O_{n^*} and the ground state of $\{X_i\}$.

Reduced density matrices

2.7 Optimising Fermi Surface Calculations

Rotated basis for simplified representation

At the renormalisation group fixed point, we have a renormalised theory for the interaction of the impurity spin S_f with the f -layer Fermi surface:

$$H^* = \sum_{\alpha,\beta} \mathbf{S}_f \cdot \boldsymbol{\sigma}_{\alpha,\beta} \sum_{\mathbf{k},\mathbf{q}} J_f^*(\mathbf{k}, \mathbf{q}) f_{\mathbf{k},\alpha}^\dagger f_{\mathbf{q},\beta} . \quad (7.1)$$

We will now obtain a more minimal representation of this interaction. Each momentum label is summed over all four quadrants $\mathcal{Q}_1$ through $\mathcal{Q}_4$; eq. ?? relates $\mathcal{Q}_1$ with $\mathcal{Q}_3$ and $\mathcal{Q}_2$ with $\mathcal{Q}_4$:

$$\begin{aligned} \sum_{\mathbf{k},\mathbf{q}} J_f^*(\mathbf{k}, \mathbf{q}) f_{\mathbf{k},\alpha}^\dagger f_{\mathbf{q},\beta} &= \sum_{\mathbf{q}} \left[\sum_{\mathbf{k} \in \mathcal{Q}_1, \mathcal{Q}_2} J_f^*(\mathbf{k}, \mathbf{q}) f_{\mathbf{k},\alpha}^\dagger + \sum_{\mathbf{k} \in \mathcal{Q}_3, \mathcal{Q}_4} J_f^*(\mathbf{k}, \mathbf{q}) f_{\mathbf{k},\alpha}^\dagger \right] f_{\mathbf{q},\beta} \\ &= \sum_{\mathbf{q}} \left[\sum_{\mathbf{k} \in \mathcal{Q}_1, \mathcal{Q}_2} J_f^*(\mathbf{k}, \mathbf{q}) f_{\mathbf{k},\alpha}^\dagger + \sum_{\mathbf{k} \in \mathcal{Q}_1} J_f^*(\mathbf{k} - \mathbf{Q}_1, \mathbf{q}) f_{\mathbf{k}-\mathbf{Q}_1,\alpha}^\dagger + \sum_{\mathbf{k} \in \mathcal{Q}_2} J_f^*(\mathbf{k} + \mathbf{Q}_2, \mathbf{q}) f_{\mathbf{k}+\mathbf{Q}_2,\alpha}^\dagger \right] f_{\mathbf{q},\beta} \\ &= \sum_{\mathbf{q}} \left[\sum_{\mathbf{k} \in \mathcal{Q}_1} J_f^*(\mathbf{k}, \mathbf{q}) (f_{\mathbf{k},\alpha}^\dagger - f_{\mathbf{k}-\mathbf{Q}_1,\alpha}^\dagger) + \sum_{\mathbf{k} \in \mathcal{Q}_2} J_f^*(\mathbf{k}, \mathbf{q}) (f_{\mathbf{k},\alpha}^\dagger - f_{\mathbf{k}+\mathbf{Q}_2,\alpha}^\dagger) \right] f_{\mathbf{q},\beta} . \end{aligned} \quad (7.2)$$

For ease of notation, we define new fermionic operators $A_{\mathbf{k},\sigma}$:

$$A_{\mathbf{k},\sigma,\pm} = \begin{cases} \frac{1}{\sqrt{2}} (f_{\mathbf{k},\sigma} \pm f_{\mathbf{k}-\mathbf{Q}_1,\sigma}) , & \mathbf{k} \in \mathcal{Q}_1 , \\ \frac{1}{\sqrt{2}} (f_{\mathbf{k},\sigma} \pm f_{\mathbf{k}+\mathbf{Q}_2,\sigma}) , & \mathbf{k} \in \mathcal{Q}_2 , \end{cases} \quad (7.3)$$

which satisfy the appropriate algebra: $\{A_{\mathbf{k},\sigma,p}, A_{\mathbf{k}',\sigma',p'}\} = \delta_{\mathbf{k},\mathbf{k}'} \delta_{\sigma,\sigma'} \delta_{p,p'}$, with $p = \pm$ denoting the flavours of the A fields. Note that only the $p = -1$ flavour enters the Hamiltonian. Henceforth, we drop the label $\pm$ and it is implied that A refers to the $p = -1$ variant.

Decomposing the sum over $\mathbf{q}$ in a similar fashion, we get

$$\sum_{\mathbf{k},\mathbf{q}} J_f^*(\mathbf{k}, \mathbf{q}) f_{\mathbf{k},\alpha}^\dagger f_{\mathbf{q},\beta} = 2 \sum_{\mathbf{k},\mathbf{q} \in \mathcal{Q}_1, \mathcal{Q}_2} J_f^*(\mathbf{k}, \mathbf{q}) A_{\mathbf{k},\alpha}^\dagger A_{\mathbf{q},\beta} . \quad (7.4)$$

Correlations in truncated representation

In order to calculate equal-time correlations (such as $\langle S_d^+ f_{\mathbf{k}\downarrow}^\dagger f_{\mathbf{k}'\uparrow} \rangle$), we use eq. 7.3 to express the bare fields $f_{\mathbf{k},\sigma}$ in terms of the new fields $A_{\mathbf{k},\sigma,\pm}$ and $B_{\mathbf{k},\sigma,\pm}$:

$$\begin{aligned} f_{\mathbf{k},\sigma} &= \frac{1}{\sqrt{2}} (A_{\mathbf{k},\sigma,+} + A_{\mathbf{k},\sigma,-}) , \mathbf{k} \in Q_1, Q_2 \\ f_{\mathbf{k}-\mathbf{Q}_1,\sigma} &= \frac{1}{\sqrt{2}} (A_{\mathbf{k},\sigma,+} - A_{\mathbf{k},\sigma,-}) , \mathbf{k} \in Q_1 \\ f_{\mathbf{k}+\mathbf{Q}_2,\sigma} &= \frac{1}{\sqrt{2}} (A_{\mathbf{k},\sigma,+} - A_{\mathbf{k},\sigma,-}) , \mathbf{k} \in Q_2 \end{aligned} \quad (7.5)$$

where the four relations act on operators in four quadrants. These relations allow us to express correlations $F_{k_i,k_j} = \langle S_d^+ c_{k_i,\uparrow}^\dagger c_{k_j,\downarrow} \rangle$ for the bare fields f in terms of the rotated fields. Using the fact that the Hamiltonian does not couple the fields B_+ and A_+ , we have

$$\begin{aligned} F_{k_1,k_1} &= F_{k_3,k_3} = -F_{k_1,k_3} = \frac{1}{2} \langle S_d^+ A_{k_1,\downarrow,-}^\dagger A_{k_1,\uparrow,-} \rangle , \\ F_{k_2,k_2} &= F_{k_4,k_4} = -F_{k_2,k_4} = \frac{1}{2} \langle S_d^+ A_{k_2,\downarrow,-}^\dagger A_{k_2,\uparrow,-} \rangle , \\ F_{k_1,k_2} &= F_{k_3,k_4} = -F_{k_1,k_4} = -F_{k_2,k_3} = \frac{1}{2} \langle S_d^+ A_{k_1,\downarrow,-}^\dagger A_{k_1,\uparrow,-} \rangle , \end{aligned} \quad (7.6)$$

where the momentum state k_i belongs to the quadrant Q_i .

Effective Star Graph Representation of Kondo Interaction

To make computation less intensive, we want to make the Hamiltonian in eq. 7.4 sparser, by going to a star graph representation where each node of the graph couples only with the impurity spin (central node) and not to the other nodes of the graph. We therefore consider a rotated basis of excitations

$$\gamma_{i,\sigma}^\dagger = \sum_k \Lambda_{i,k} A_{k,\sigma}^\dagger , \text{ where } \sum_k \Lambda_{i,k}^* \Lambda_{j,k} = \delta_{i,j} . \quad (7.7)$$

These fermions of course satisfy the appropriate algebra:

$$\left\{ \gamma_{i,\sigma}, \gamma_{j,\sigma'}^\dagger \right\} = \sum_{k,q} \Lambda_{i,k}^* \Lambda_{j,q} \left\{ A_{k,\sigma}, A_{q,\sigma'}^\dagger \right\} = \delta_{i,j} \delta_{\sigma,\sigma'} . \quad (7.8)$$

In order to calculate the coefficients $\Lambda_{i,k}$, we impose the star graph simplification condition described above:

$$\sum_{\mathbf{k},\mathbf{q}} J_f^*(\mathbf{k},\mathbf{q}) A_{\mathbf{k},\alpha}^\dagger A_{\mathbf{q},\beta} = \sum_i K_i \gamma_{i,\alpha}^\dagger \gamma_{i,\beta} , \quad (7.9)$$

where K_i are the eigenvalues of the Kondo coupling matrix $J_f^*(\mathbf{k},\mathbf{q})$. Substituting the definition of the γ fields (eq. 7.7) leads to the equation

$$J_f^*(\mathbf{k},\mathbf{q}) = \sum_i \Lambda_{k,i} K_i \Lambda_{i,q}^\dagger \implies \mathcal{J} = \Lambda K \Lambda^\dagger , \quad (7.10)$$

where $\mathcal{J}$ is the Kondo coupling matrix with matrix elements $J_f^*(\mathbf{k}, \mathbf{q})$, K is a diagonal matrix with the eigenvalues (K_i) of $\mathcal{J}$ along its diagonal, and Λ is the matrix $\Lambda_{i,k}$ (the rotated indices i along the row, and the earlier momentum indices along the column). From eq. 7.10, it is clear that since $\mathcal{J}$ is Hermitian, the matrix Λ is simply the unitary matrix which diagonalises the Kondo matrix $\mathcal{J}$:

$$\Lambda^\dagger \mathcal{J} \Lambda = K . \quad (7.11)$$

The prescription is therefore as follows. We compute the unitary transformation Λ that diagonalises $\mathcal{J}$ and obtain the eigenvalues $\{K_i\}$. Using the eigenvalues, we work with the sparser Hamiltonian

$$\vec{S}_f \cdot \sum_{\alpha,\beta} \vec{\sigma}_{\alpha,\beta} \sum_i K_i \gamma_{i,\alpha}^\dagger \gamma_{i,\beta} . \quad (7.12)$$

If we have to calculate a certain correlation $C_{k,q}$, we instead calculate the correlation matrix $\tilde{C}_{i,j}$, and then transform back using the unitary transformation:

$$C = \Lambda \tilde{C} \Lambda^\dagger . \quad (7.13)$$

2.8 Measures of entanglement and their computation

The present millennium has seen a rapid surge in the use of quantum entanglement towards the study of quantum and condensed matter systems [Terhal et al.(2003)Terhal, Wolf, and Doherty, Horodecki et al.(2009)Horodecki, Horodecki, Horodecki, and Horodecki, Eisert et al.(2010)Eisert, Cramer, and Plenio, Laflorencie(2016a), Zeng et al.(2019)Zeng, Chen, Zhou, and Wen, Erhard et al.(2020)Erhard, Krenn, and Zeilinger]. By tracking quantum correlations among particles across both small and large distances, entanglement measures offer valuable insights into the nature of the ground state and low-energy excitations. This, for example, allows us to distinguish strongly-correlated topologically ordered states characterised by long-ranged entanglement (in the form of a sub-leading topological term in their entanglement entropy [Wen(1990), Wen and Niu(1990), Wen(1995), Niu et al.(1985)Niu, Thouless, and Wu]) from weakly-interacting topologically trivial phases that have short-ranged area-law entanglement [Das and Shankaranarayanan(2006), Cramer et al.(2006)Cramer, Eisert, Plenio, and Dreißig, Eisert et al.(2010)Eisert, Cramer, and Plenio].

2.8.1 Reduced density matrix elements as equal-time correlation functions

A density matrix plays the role of a probability density function (PDF) for a quantum system. A classical variable X that can take values $x \in [-X^*, X^*]$ has a PDF $F(X = x)$, which is used, for example, for calculating the average value (expectation value) of the variable:

$$\langle X \rangle = \int_{-X^*}^{X^*} x F(x) dx . \quad (8.1)$$

In quantum mechanics, the variables we are interested in are operators $\mathcal{O}$, and a quantum PDF ρ will then give the expectation value of the operator, in a fashion similar to its classical counterpart:

$$\langle \mathcal{O} \rangle = \sum_{\phi} \langle \phi | \rho \mathcal{O} | \phi \rangle ; \quad (8.2)$$

here, the integral over the classical values x is replaced by a sum over the all possible quantum states $|\phi\rangle$ (complete orthonormal basis) of the system. However, we already know that the expectation value of a quantum operator is given by

$$\langle \mathcal{O} \rangle = \langle \Psi | \mathcal{O} | \Psi \rangle . \quad (8.3)$$

To bring this form closer to the one in terms of ρ (eq. 2), we insert an identity resolution: $1 = \sum_{\phi} |\phi\rangle \langle \phi|$:

$$\langle \mathcal{O} \rangle = \sum_{\phi} \langle \Psi | \mathcal{O} | \phi \rangle \langle \phi | \Psi \rangle = \sum_{\phi} \langle \phi | \Psi \rangle \langle \Psi | \mathcal{O} | \phi \rangle . \quad (8.4)$$

A comparison of this equation to eq. (2) gives

$$\rho = |\Psi\rangle \langle \Psi| ; \quad (8.5)$$

a density matrix (also called density operator) is therefore formally defined as the outer product of the wavefunction with its conjugate. It encodes all information about the state; for example, the probability of measuring the system to be at a position x (hence in a state $|x\rangle$) is given by $P(x) = \langle x | \rho | x \rangle$. From its definition in eq. (5), the DM has the very important property that it has unity trace:

$$\text{Tr}[\rho] = 1 , \quad (8.6)$$

where $\text{Tr}[\cdot]$ is the trace of an operator. This is seen very easily by taking the trace in an orthonormal basis in which the state $|\Psi\rangle$ is one of the basis states. For example, consider the DM

$$\rho = \frac{1}{\sqrt{2}} [|\Psi_1\rangle \langle \Psi_1| + |\Psi_2\rangle \langle \Psi_2|] , \quad (8.7)$$

where $|\Psi_{1,2}\rangle$ are orthogonal normalised states. The trace of ρ^2 for such a state is only $1/2$; such states satisfying $\text{Tr}[\rho^2] < 1$ are termed *mixed states*; the DM describing such states cannot be written in the form $\rho = |\Psi\rangle \langle \Psi|$ for any wavefunction $|\Psi\rangle$. On the other hand, the DMs that *can* be expressed like that are termed *pure states*, and have $\text{Tr}[\rho^2] = 1$.

One way in which a pure state can be realistically converted into a mixed state is through a measurement process. Consider the zero total spin state formed by two spins:

$$|\Psi\rangle = \frac{1}{2} [|\uparrow, \downarrow\rangle + |\downarrow, \uparrow\rangle] , \rho = |\Psi\rangle \langle \Psi| . \quad (8.8)$$

The state $|\uparrow, \downarrow\rangle$ has the first spin pointing up and the second one down, and this is in a superposition with its flipped counterpart $|\downarrow, \uparrow\rangle$. This is clearly a pure state.

If we now measure the state of the second spin, it will collapse into one of its two states; the other spin will collapse into the opposite state (given the antiparallel configurations of the spins). What this means is that until one checks what the result of the measurement is, the first spin remains in a statistical superposition of both outcomes (up and down). This is formally expressed through a partial trace operation: the state of the first spin after a measurement of the second spin is obtained

by taking the full density matrix ρ and computing its trace over the states $|\sigma_B\rangle$ of the second spin, leaving out the states of the first spin from the trace:

$$\rho_1 = \text{Tr}_2[\rho] = \sum_{\sigma_2} \langle \sigma_2 | \rho | \sigma_2 \rangle ; \quad (8.9)$$

where $|\sigma_2\rangle$ sums over the states $|\uparrow\rangle$ and $|\downarrow\rangle$ of the second spin. this partial tracing operation Tr_2 codifies the measurement process, and the partially traced density matrix ρ_1 is termed a reduced density matrix (RDM) of the first spin, because it only captures information about the first spin (the second spin has been traced out).

To see the connection with mixedness, let us compute the RDM for the first spin explicitly:

$$\rho_1 = \langle \uparrow_2 | \rho | \uparrow_2 \rangle + \langle \downarrow_2 | \rho | \downarrow_2 \rangle = \frac{1}{2} [|\uparrow_1\rangle \langle \uparrow_1| + |\downarrow_1\rangle \langle \downarrow_1|] , \quad (8.10)$$

where the numbered subscripts on the kets and bras indicate that they only act on the respective Hilbert spaces. To see that ρ_1 describes a mixed state, we adopt a specific representation:

$$\begin{aligned} |\uparrow_1\rangle &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} , |\downarrow_1\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} , \\ \Rightarrow \rho_1 &= \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} , \rho_1^2 = \frac{1}{4} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} . \end{aligned} \quad (8.11)$$

The trace of ρ_1^2 is half, hence it is a mixed state.

The deviation of trace of ρ^2 from its maximum value of unity can be thought of as a measure of the lost information. In the example above, when we traced out the states of the second spin, we lost some information about the state of the first spin as well. This is because the two spins were *entangled*; they had quantum correlations between them. The entanglement arises from the fact that the states of the two spins are anti-correlated.

If instead we had chosen a non-entangled parent state to start from, the reduced density matrix would not have been mixed. For example, consider the state

$$|\Psi\rangle = \frac{1}{2} (|\uparrow_1\rangle + |\downarrow_1\rangle) \otimes (|\uparrow_2\rangle + |\downarrow_2\rangle) . \quad (8.12)$$

It is easy to see that this state does not contain any correlations between the two spins - no matter the configuration of the second spin, the first spin is always in the state $|\uparrow_1\rangle + |\downarrow_1\rangle$. The RDM for the first spin comes out to be

$$\rho_1 = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} , \rho_1^2 = \frac{1}{4} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} , \quad (8.13)$$

revealing that ρ_1 is not mixed.

The above method of calculating an RDM works for simple cases but can fail if (i) we have fermion signs to account for, or (ii) the full basis cannot be expressed as a direct product of local basis sets. The former arises if we are dealing with fermions, and the latter arises if we are working in a

rotated basis. We now derive an alternative method for calculating the RDM, one that automatically takes care of the fermion signs and works even if the basis is rotated and complicated to write down. We start by defining the RDM more precisely. Given a state $|\Psi\rangle$ and a bipartition into subsystems S_1 and its complement S_2 , the density matrix for the state is $\rho = |\Psi\rangle \langle\Psi|$. A basis state of the full system can be written as $|A, \alpha\rangle = |A\rangle \otimes |\alpha\rangle$, where $\{A\}$ and $\{\alpha\}$ are complete sets of basis states describing S_1 and S_2 respectively. The RDM for S_2 is then defined as

$$\rho_1 = \text{Tr}_\alpha \rho = \sum_\alpha \langle \alpha | \rho | \alpha \rangle . \quad (8.14)$$

The matrix elements of ρ_1 in the basis $\{A\}$ can be expressed as

$$\langle A | \rho_1 | B \rangle = \sum_\alpha \langle A, \alpha | \rho | B, \alpha \rangle . \quad (8.15)$$

We next define an operator $O_{A,B}$ that represents scattering processes between the various states of subsystem S_1 : $O_{A,B} = |A\rangle \langle B| \otimes I_2$, but acts as identity on S_2 . The expectation value of this operator is

$$\langle O_{A,B} \rangle = \text{Tr}(\rho O_{A,B}) = \sum_{C,\beta} \langle C, \beta | \rho | A \rangle \langle B | C, \beta \rangle = \sum_\beta \langle B, \beta | \rho | A, \beta \rangle . \quad (8.16)$$

Comparing with eq. 8.15, we obtain the relation

$$\langle A | \rho_1 | B \rangle = \langle O_{B,A} \rangle . \quad (8.17)$$

This result is significant because it guarantees that in order to calculate the RDM for a subsystem, it is enough to calculate the expectation values of all inter-state off-diagonal transition operators. For example, we take subsystem S_1 to be a two fermion system represented by the operators c_1 and c_2 . Assuming a basis $\{|0, 0\rangle, |0, 1\rangle, |1, 0\rangle, |1, 1\rangle\}$, where the two entries represent the occupancies of the two fermions, the RDM for this subsystem is determined by the following matrix:

$$\rho_1 = \begin{pmatrix} \langle h_1 h_2 \rangle & \langle h_1 c_2^\dagger \rangle & \langle c_1^\dagger h_2 \rangle & \langle c_1^\dagger c_2^\dagger \rangle \\ & \langle h_1 n_2 \rangle & \langle c_1^\dagger c_2 \rangle & \langle c_1^\dagger n_2 \rangle \\ & & \langle n_1 h_2 \rangle & \langle n_1 c_2^\dagger \rangle \\ & & & \langle n_1 n_2 \rangle \end{pmatrix} , \quad (8.18)$$

where the blank entries can be determined from hermiticity. To see how the operators are determined, take the example of the top right element. The two basis states for that matrix element are $|0, 0\rangle$ and $|1, 1\rangle$.

Quantum Fisher information: a witness of multipartite entanglement

The quantum Fisher information (QFI) F_Q [Braunstein and Caves(1994), Hauke et al.(2016)Hauke, Heyl, Tagliacozzo, and Zoller] quantifies how sensitive a state ρ is to unitary transformations generated by an observable $\hat{O}$. For a pure state $\rho = |\psi\rangle\langle\psi|$, $F_Q(\psi, \hat{O})$ is defined in terms of the variance of the operator $\hat{O}$ (i.e., the connected correlation function):

$$F_Q(\psi, \hat{O}) = 4\Delta(\hat{O})^2 = 4 \left(\langle\psi|\hat{O}^2|\psi\rangle - \langle\psi|\hat{O}|\psi\rangle^2 \right), \quad (8.19)$$

and it provides an upper bound on the precision that can be achieved in measuring the parameter θ that is dual to the observable $\hat{O}$: $\mathcal{N}(\Delta\theta)^2 \geq F_Q^{-1}$, $\mathcal{N}$ being the number of independent measurements [Braunstein and Caves(1994), Hauke et al.(2016)Hauke, Heyl, Tagliacozzo, and Zoller]. $F_Q(\psi, \hat{O})$ is thus a measure of the entanglement arising from quantum fluctuation content in $|\psi\rangle$ (considered with respect to an eigenstate of $\hat{O}$). This can be made more manifest by considering a traceless operator $\hat{M}$ with eigenstates $\{|m\rangle\}$. Without loss of generality, we can rescale the operator such that $\sum m^2 = 1/2$. If $\varepsilon_m(\psi) \equiv 1 - |\langle m|\psi\rangle|^2$ is the geometric entanglement between a given state $|\psi\rangle$ and the eigenstates of $\hat{M}$, the QFI corresponding to $\hat{M}$ in the state ψ can be written as

$$F_Q(\psi, \hat{M}) = 4 \left[\frac{1}{2} - \sum_m m^2 \varepsilon_m - \left(\sum_m m \varepsilon_m \right)^2 \right]. \quad (8.20)$$

The total geometric entanglement $\sum_m \varepsilon_m$ is constrained to be $d_M - 1$, where d_M is the dimension of the operator $\hat{M}$. Because each eigenvalue m has a magnitude of at most $1/\sqrt{2}$ (all m^2 must add up to $1/2$), the maximum magnitude of the last two terms in eq. (10.13) (and hence the minimum value of the QFI) is attained for the case when some of the ε_m are zero. However, because the eigenstates are orthogonal, only one of them can have perfect overlap with the state $|\psi\rangle$ and only one ε_m can be zero at a time. The case of minimum QFI therefore corresponds to $\varepsilon_{m^*} = 0, \varepsilon_{m \neq m^*} = 1$, leading to $F_Q = 0$. The opposite situation arises when all the geometric entanglement measures are non-zero and equal, $\varepsilon_m = 1/d_M$, leading to a maximum QFI value of $F_Q = 2/d_M$. A uniformly spread geometric entanglement distribution, therefore, leads to a larger QFI, while a more focused distribution leads to a smaller QFI. This sums up the relation between the quantum Fisher information and geometric entanglement.

If the operator $\hat{M}$ is such that its eigenstates are separable states from the perspective of a certain party, the vanishing QFI arises when $|\psi\rangle$ has zero geometric entanglement with respect to one of the separable states, and the QFI saturates when $|\psi\rangle$ has a finite geometric entanglement with all of the states. A large QFI can thus be associated with more entanglement. Moreover, if the eigenstates of $\hat{M}$ are symmetry-broken states, a vanishing QFI indicates that $|\psi\rangle$ is very close to such a symmetry-broken state, while a large QFI indicates that $|\psi\rangle$ is a uniform superposition of such symmetry-broken states and therefore preserves the symmetry as a whole. A larger QFI points towards the presence of more quantum fluctuations and a reduced susceptibility towards symmetry-breaking.

We will now relate the QFI to some other measures of correlation. The QFI corresponding to an observable can be directly related to the static lesser Greens functions associated with that

observable:

$$F_Q(\psi, \hat{O}_2) = 4 \left(\left\langle \hat{O}_2 \hat{O}_1^\dagger \right\rangle \Big|_{\hat{O}_1^\dagger = \hat{O}_2} - \left\langle \hat{O}_2 \hat{O}_1^\dagger \right\rangle^2 \Big|_{\hat{O}_1 = 1} \right) = 4 \left[G_{\hat{O}_2, \hat{O}_2}^<(t \rightarrow \infty) - (G_{1, \hat{O}_2}^<(t \rightarrow \infty))^2 \right] \quad (8.21)$$

Further, the retarded Greens function defined above in eq. (5.6) can very generally be related to the QFI. By using the simplified forms $\frac{i}{2} G_{\hat{O}_2, \hat{O}_2}^R(t \rightarrow \infty) = \langle (\hat{O}_2)^2 \rangle$ and $\frac{i}{2} G_{1, \hat{O}_2}^R(t \rightarrow \infty) = \langle \hat{O}_2 \rangle$, we get

$$F_Q(\psi, \hat{O}_2) = 2i G_{\hat{O}_2, \hat{O}_2}^R(t \rightarrow \infty) + (G_{1, \hat{O}_2}^R(t \rightarrow \infty))^2. \quad (8.22)$$

For the sake of completeness, we generalise the above relations to finite temperatures by adopting the approach of Hauke et al. [Hauke et al.(2016)Hauke, Heyl, Tagliacozzo, and Zoller]. For a mixed state at inverse temperature β characterised by the density matrix $\rho = \sum_n f_n |n\rangle \langle n|$ where $(E_n, |n\rangle)$ are the eigenvalue-eigenstate pairs and $f_n = e^{-\beta E_n} / \sum_n e^{-\beta E_n}$ are the Boltzmann weights, the FQI can be assigned a spectral representation of the form [Hauke et al.(2016)Hauke, Heyl, Tagliacozzo, and Zoller]

$$F_Q(\rho, \hat{O}) = 2 \sum_{m,n} \frac{(f_m - f_n)^2}{f_m + f_n} |\langle m | \hat{O} | n \rangle|^2. \quad (8.23)$$

By using the identity $\frac{f_m - f_n}{f_m + f_n} = \int_{-\infty}^{\infty} d\omega \delta(\omega + E_m - E_n) \tanh(\beta\omega/2)$, the finite temperature FQI can be written in the form

$$F_Q(\rho, \hat{O}) = 2 \int_{-\infty}^{\infty} d\omega \tanh(\beta\omega/2) \sum_{m,n} (f_m - f_n) \delta(\omega + E_m - E_n) |\langle m | \hat{O} | n \rangle|^2. \quad (8.24)$$

In order to relate this with a correlation function, we now point out that the lesser Greens function $G^<$ and the greater Greens function $G^>$ that we will now define admit spectral representations in terms of the eigenstates $|n\rangle$ of the full problem:

$$G_{\hat{O}^\dagger, \hat{O}}^<(t) \equiv i \langle \hat{O}^\dagger \hat{O}(t) \rangle \implies G_{\hat{O}^\dagger, \hat{O}}^<(\omega) = 2\pi i \sum_{m,n} f_n \delta(\omega + E_m - E_n) |\langle m | \hat{O}^\dagger | n \rangle|^2 \quad (8.25)$$

$$G_{\hat{O}^\dagger, \hat{O}}^>(t) \equiv -i \langle \hat{O}(t) \hat{O}^\dagger \rangle \implies G_{\hat{O}^\dagger, \hat{O}}^>(\omega) = -2\pi i \sum_{m,n} f_m \delta(\omega + E_m - E_n) |\langle m | \hat{O} | n \rangle|^2 \quad (8.26)$$

If $\hat{O}$ is a Hermitian operator, eqs. (10.19) can be used to rewrite the QFI in terms of $G_{\hat{O}^\dagger, \hat{O}}^<(\omega)$. Indeed, by combining eqs. (10.19) and eq. (10.18) with the assumption $\hat{O} = \hat{O}^\dagger$, we get

$$F_Q(\rho, \hat{O}) = -\frac{1}{\pi i} \int_{-\infty}^{\infty} d\omega \tanh(\beta\omega/2) (G_{\hat{O}, \hat{O}}^>(\omega) + G_{\hat{O}, \hat{O}}^<(\omega)). \quad (8.27)$$

This constitutes the general relation between the finite temperature QFI for a general Hermitian many-particle operator $\hat{O}$ and the corresponding many-particle correlations at zero and finite frequencies.

The Kubo linear response function associated with the operator $\hat{O}(t)$ due to a perturbation from the same operator is associated with a correlation function/susceptibility $\mathcal{C}_{\hat{O}}(t, t') = i\theta(t -$

$t') \left\langle \left[\hat{O}(t), \hat{O}(t') \right] \right\rangle$. By going to the frequency domain, the imaginary part of such a function can be related to the sum of the lesser and greater Greens functions:

$$\begin{aligned} \mathcal{C}_{\hat{O}}(\omega) &= i \int_0^\infty d(t-t') e^{i\omega(t-t')} \left(\left\langle \hat{O}(t) \hat{O}(t') \right\rangle - \left\langle \hat{O}(t') \hat{O}(t) \right\rangle \right) \\ \Rightarrow \mathcal{C}_{\hat{O}}(\omega) - \mathcal{C}_{\hat{O}}(\omega)^* &= i \int_0^\infty d(t-t') \left(e^{i\omega(t-t')} - e^{-i\omega(t-t')} \right) \left(\left\langle \hat{O}(t) \hat{O}(t') \right\rangle - \left\langle \hat{O}(t') \hat{O}(t) \right\rangle \right) \\ &= i \int_{-\infty}^\infty d(t-t') e^{i\omega(t-t')} \left(\left\langle \hat{O}(t) \hat{O}(t') \right\rangle - \left\langle \hat{O}(t') \hat{O}(t) \right\rangle \right) \\ &= - \left(G_{\hat{O},\hat{O}}^>(\omega) + G_{\hat{O},\hat{O}}^<(\omega) \right) . \end{aligned}$$

This allows us to connect the QFI with the imaginary part of the Kubo correlation function/susceptibility obtained in Ref. [Hauke et al.(2016)Hauke, Heyl, Tagliacozzo, and Zoller]:

$$F_Q(\rho, \hat{O}) = \frac{1}{\pi i} \int_{-\infty}^\infty d\omega \tanh(\beta\omega/2) (\mathcal{C}_{\hat{O}}(\omega) - \mathcal{C}_{\hat{O}}(\omega)^*) . \quad (8.28)$$

2.8.2 Decomposition of entanglement entropy in terms of multipartite information I_n

We first define a partitioning of the entire system into N disjoint regions, $\mathcal{S}_N = \{\nu_1, \nu_2, \dots, \nu_N\}$. Going beyond bipartite mutual information, one can define the n -partite information I_n between the members of $\mathcal{S}_n = \{\nu_1, \nu_2, \dots, \nu_n\}$, which is itself a subset of the exhaustive set of regions $\mathcal{S}_N$:

$$I_n(\nu_1, \dots, \nu_n) = \sum_{\sigma \in \mathcal{P}(\mathcal{S}_n)} (-1)^{1+|\sigma|} S_{\text{EE}}(\sigma) , \quad (8.29)$$

where σ is a subset drawn from the power set of $\mathcal{S}_n$, and $|\sigma|$ is the number of members in the subset σ . The 4-partite information can be written as

$$\begin{aligned} I_4(\nu_1, \nu_2, \nu_3, \nu_4) &= S_{\text{EE}}(\nu_1) + S_{\text{EE}}(\nu_2) + S_{\text{EE}}(\nu_3) + S_{\text{EE}}(\nu_4) - S_{\text{EE}}(\nu_1 \cup \nu_2) - S_{\text{EE}}(\nu_1 \cup \nu_3) \\ &\quad - S_{\text{EE}}(\nu_1 \cup \nu_4) - S_{\text{EE}}(\nu_2 \cup \nu_3) - S_{\text{EE}}(\nu_2 \cup \nu_4) - S_{\text{EE}}(\nu_3 \cup \nu_4) + S_{\text{EE}}(\nu_1 \cup \nu_2 \cup \nu_3) \\ &\quad + S_{\text{EE}}(\nu_1 \cup \nu_2 \cup \nu_4) + S_{\text{EE}}(\nu_1 \cup \nu_3 \cup \nu_4) + S_{\text{EE}}(\nu_2 \cup \nu_3 \cup \nu_4) + S_{\text{EE}}(\nu_1 \cup \nu_2 \cup \nu_3 \cup \nu_4) . \end{aligned} \quad (8.30)$$

One can also define the average n -partite information $\bar{I}_n$ by considering all sets of n regions:

$$\bar{I}_n = \frac{(N-n)!n!}{N!} \sum_{\mathcal{S}_n \in C_n(\mathcal{S}_N)} I_n(\mathcal{S}_n) = \frac{(N-n)!n!}{N!} \sum_{\mathcal{S}_n \in C_n(\mathcal{S}_N)} \sum_{\sigma \in \mathcal{P}(\mathcal{S}_n)} (-1)^{1+|\sigma|} S_{\text{EE}}(\sigma) , \quad (8.31)$$

where $C_n(\mathcal{S}_N)$ represents all ways of choosing n distinct elements out of the full set $\mathcal{S}_N$ of subsystems.

The expression for $\bar{I}_n$ can be simplified by splitting the summation over the elements of the power set $\mathcal{P}$ into sets of definite number:

$$\bar{I}_n = \frac{(N-n)!n!}{N!} \sum_{S_n \in C_n(S_N)} I_n(S_n) = \frac{(N-n)!n!}{N!} \sum_{S_n \in C_n(S_N)} \sum_{m=1}^n \sum_{\sigma_m \in \mathcal{P}_m(S_n)} (-1)^{1+|\sigma|} S_{EE}(\sigma_m), \quad (8.32)$$

where $\mathcal{P}_m$ represents all subsets of the power set that have m elements. With some combinatorial magic, it can be shown that

$$\sum_{S_n \in C_n(S_N)} \sum_{\sigma_m \in \mathcal{P}_m(S_n)} = \binom{N-m}{n-m} \sum_{\sigma_m \in C_m(S_N)}. \quad (8.33)$$

To see how this comes about, note that what this equation says is that summing over combinations of subsets of length n of the super set S_N and further summing over all subsets of length $m \leq n$ for each set S_n is equivalent to summing just once over subsets of length m of the super set S_N but now each subset of length m appears with a frequency $f_m = \binom{N-m}{n-m}$. The value for the frequency f_m is simply the number of subsets S_n in which the smaller subset σ_m appears. This can be calculated by allowing m of the n slots in S_n to be filled by the members of σ_m , and the remaining $n-m$ slots then have to be filled by choosing from the remaining $N-m$ members of the super set S_N ; there's obviously $\binom{N-m}{n-m}$ ways of carrying this out.

Implementing this into the expression for $\bar{I}_n$ gives

$$\bar{I}_n = \sum_{m=1}^n (-1)^{1+m} \frac{\binom{n}{m}}{\binom{N}{m}} \sum_{\sigma_m \in C_m(S_N)} S_{EE}(\sigma_m) = \sum_{m=1}^n (-1)^{1+m} \binom{n}{m} S_{EE}^{(m)}, \quad (8.34)$$

where we have define the average entanglement entropy $S_{EE}^{(m)}$ for all m -member sets of regions:

$$S_{EE}^{(m)} = \frac{1}{\binom{N}{m}} \sum_{\sigma_m \in C_m(S_N)} S_{EE}(\sigma_m). \quad (8.35)$$

With most of the machinery in place, we would now like to express the entanglement entropy S_{EE} purely in terms multipartite measures I_n , for the reasons stated above. For a system that is invariant under real-space translation (as our lattice model is), the entanglement entropy of any real-space interval should only depend on the extent of the interval. It therefore suffices to calculate the entanglement entropy averaged over all such intervals, and the corresponding measure in our formulation is $S_{EE}^{(1)}$.

We will now express $S_{EE}^{(1)}$ in terms of $\bar{I}_n$, by eliminating $S_{EE}^{(m)} (m > 1)$ from the equations. We adopt the ansatz

$$S_{EE}^{(1)} = \sum_{n=2}^N \lambda_n \bar{I}_n = \sum_{n=2}^N \lambda_n \binom{n}{m} \sum_{m=1}^n (-1)^{1+m} S_{EE}^{(m)}. \quad (8.36)$$

In order to separate $S_{EE}^{(1)}$ from the rest of the $S_{EE}^{(m)}$, we exchange the order of summation:

$$S_{EE}^{(1)} = \sum_{m=1}^N (-1)^{1+m} S_{EE}^{(m)} \sum_{n=m}^N \lambda_n \binom{n}{m} - S_{EE}^{(1)} \lambda_1 = S_{EE}^{(1)} \sum_{n=2}^N n \lambda_n - \sum_{m=2}^{N-1} S_{EE}^{(m)} (-1)^m \sum_{n=m}^N \binom{n}{m} \lambda_n, \quad (8.37)$$

where the constraint of $m \leq n$ in the previous form is now enforced by letting n run from m to N . The $N - 1$ coefficients $\{\lambda_n\}$ will be determined by requiring that the $S_{EE}^{(m)}(m > 1)$ vanish on the right-hand side, as well as by matching the coefficients of $S_{EE}^{(1)}$ on both sides. Note that m runs only upto $N - 1$ in the second term; this is because, $S_{EE}^{(N)}$ is the entanglement entropy of the full system and hence automatically zero. The equations are

$$\sum_{n=2}^N n\lambda_n = 1, \quad (8.38)$$

$$\sum_{n=m}^N \binom{n}{m} \lambda_n = 0; \quad m = 2, \dots, N - 1. \quad (8.39)$$

We focus on the second equation first, since it is very short for m close to N . For $m = N - 1$ and $m = N - 2$, we get

$$\begin{aligned} m = N - 1 : \lambda_{N-1} + N\lambda_N &= 0 \implies \lambda_{N-1} = -N\lambda_N \\ m = N - 2 : \lambda_{N-2} + (N-1)\lambda_{N-1} + \frac{N(N-1)}{2}\lambda_N &= 0 \implies \lambda_{N-2} = \frac{N(N-1)}{2}\lambda_N. \end{aligned} \quad (8.40)$$

We extrapolate this to the general solution,

$$\lambda_m = (-1)^m \binom{N}{m} \lambda_N, \quad m < N, \quad (8.41)$$

and then prove that if this is true for λ_{N-m} , it also solves the equation for λ_{N-m-1} , thus amounting to a proof by mathematical induction. The general equation for λ_{N-m-1} is

$$\sum_{n=N-m-1}^N \binom{n}{N-m-1} \lambda_n = 0 \implies \lambda_{N-m-1} + \sum_{m'=0}^m \binom{N-m'}{N-m-1} \lambda_{N-m'} = 0. \quad (8.42)$$

Substituting the extrapolated solution gives

$$\begin{aligned} \lambda_{N-m-1} &= - \sum_{m'=0}^m (-1)^{N-m'} \binom{N-m'}{N-m-1} \binom{N}{m'} \lambda_N = \binom{N}{m+1} \lambda_N \sum_{m'=0}^m (-1)^{N-m'-1} \binom{m+1}{m'} \\ &= (-1)^{N-m-1} \binom{N}{m+1} \lambda_N, \end{aligned} \quad (8.43)$$

which is consistent with the general solution eq. 8.41. This determines all coefficients λ_n for $n < N$ in terms of λ_N . To determine λ_N itself, we use eq. 8.38. Substituting eq. 8.41 into it gives

$$\lambda_N \sum_{n=2}^N n(-1)^n \binom{N}{n} = 1 \implies \lambda_N = \frac{(-1)^N}{N}. \quad (8.44)$$

The decomposition of entanglement entropy into multipartite measures is now complete:

$$\begin{aligned}
 S_{\text{EE}}^{(1)} &= \frac{1}{N} \sum_{n=2}^N (-1)^n \binom{N}{n} \bar{I}_n, \\
 \bar{I}_n &= \frac{(N-n)!n!}{N!} \sum_{S_n \in C_n(S_N)} I_n(S_n),
 \end{aligned} \tag{8.45}$$

where we have assumed N is even in order to drop the global phase factor $(-1)^N$.

2.9 Wavefunction Renormalisation Group

2.9.1 General idea

A reverse URG analysis involves studying the evolution of wavefunctions from the IR to the UV, by implementing inverse renormalisation group transformations. This is made possible by the fact that the IR fixed point Hamiltonians are often tractable and allow the computation of wavefunctions. Such an analysis gives insight into how states change as high-energy quantum fluctuations are resolved during the Hamiltonian renormalisation from the UV to the IR. It also allows studying the evolution of entanglement measures and correlation functions.

We will start with a very simple IR ground state wavefunction, and then go back towards the UV ground state by applying the inverse unitary operator $U^\dagger$:

$$\begin{aligned}
 U : \underbrace{|1, 2, \dots, N\rangle}_{\text{UV ground state}} &\rightarrow |1, 2, \dots, N-1\rangle |N\rangle \rightarrow \dots \rightarrow \underbrace{|1, 2, \dots, N^*\rangle |N^*+1\rangle \dots |N\rangle}_{\text{IR ground state}} \\
 U^\dagger : \underbrace{|1, 2, \dots, N^*\rangle |N^*+1\rangle \dots |N\rangle}_{\text{IR ground state}} &\rightarrow |1, 2, \dots, N^*+1\rangle |N^*+2\rangle \dots |N\rangle \rightarrow \dots \rightarrow \underbrace{|1, 2, \dots, N\rangle}_{\text{UV ground state}}
 \end{aligned}$$

The first process is the forward RG which we used to obtain the scaling equations. The second process is the reverse RG which we will undertake now. The method has already been applied to study the renormalisation of entanglement in various models of strong correlation. In general, we start with an IR wavefunction that consists a certain number of momentum states n_1 still entangled with the impurity, and the remaining momentum states n_2 disentangled from the impurity. The former are said to be part of the emergent Kondo cloud, while the latter are said to be part of the integrals of motion (IOMs) and appear in direct product with the cloud+impurity system in the ground state wavefunction. Each momentum state is tagged with two conduction bath levels, one above the Fermi surface and one below. These will be represented as $k, \pm$. If we represent the emergent cloud momentum states as $\{k_i\}$ and the IOM states as $\{q_i\}$, the ground state wavefunction can be written as

$$|\Psi\rangle_0 = (\otimes_{i=1}^{n_2} |q_i, -, \uparrow\rangle |q_i, -, \downarrow\rangle) |\Phi\rangle_{\text{cloud}} (\otimes_{i=1}^{n_1} |q_i, +, \uparrow\rangle |q_i, +, \downarrow\rangle) \tag{9.1}$$

Throughout, we have assumed that the IOMs below the Fermi surface are occupied while those above are unoccupied. This means:

$$|\Psi\rangle_0 = (\otimes_{i=1}^{n_2} |\hat{n}_{q_i,-,\uparrow} = 1\rangle |\hat{n}_{q_i,-,\downarrow} = 1\rangle) |\Phi\rangle_{\text{cloud}} (\otimes_{i=1}^{n_2} |\hat{n}_{q_i,+,\uparrow} = 0\rangle |\hat{n}_{q_i,+,\downarrow} = 0\rangle) \quad (9.2)$$

$|\Phi\rangle_{\text{cloud}}$ will be obtained by diagonalising a small system of n_1 conduction bath k states. This completes the construction of the IR ground state $|\Psi\rangle_0$. The algorithm of reverse RG then involves applying unitary transformations on this IR wavefunction; the unitary transformations will be the inverse of those that were applied during the forward RG algorithm. Since the forwards RG transformations decoupled k states and transformed them into the IOMS, the inverse transformations will take the momentum states in the IOMS and re-entangle them with the impurity+cloud system, one at a time. This is shown in fig. 2.11.




Figure 2.11: We start with a Hamiltonian with an impurity site (red) coupled with two conduction electrons (dark green), with four other decoupled electrons (bright green). The dotted rectangle represents the emergent window $(-\Lambda_j, \Lambda_j)$ at each step; the electrons inside that rectangle are still entangled with the impurity, while the ones outside have been decoupled. The next step of reverse RG involves applying the inverse transformation on the Hamiltonian, which will couple two more electrons from the IOMS (hence four dark green circles in the second step), leading to an enlargement of the emergent window. The unitary varies for each step, hence the notation U_j .

The next step is to write down the unitaries that will take us from the IR ground state to the UV ground state. In the forward RG, we used the following unitary for decoupling the shell ϵ_q and spin β is

$$U_{q\beta} = \frac{1}{\sqrt{2}} \left[1 + \eta_{0\beta} + \eta_{1\beta}^\dagger \right] . \quad (9.3)$$

Here, the subscripts 0 and 1 indicate it decouples an electron above and below the Fermi surface respectively. The inverse transformation for re-entangling $q\beta$ is

$$U_{q\beta}^\dagger = \frac{1}{\sqrt{2}} \left[1 + \eta_{0\beta}^\dagger + \eta_{1\beta} \right] . \quad (9.4)$$

The wavefunction after reversing one step of the RG will thus be

$$|\Psi\rangle_1 = U_{q\uparrow}^\dagger U_{q\downarrow}^\dagger |\Psi\rangle_0 \quad (9.5)$$

Here, q is the first momentum state immediately outside the emergent cloud. Re-entangling further momentum states using their respective unitaries lead to the sequence of wavefunctions $|\Psi\rangle_2, |\Psi\rangle_3$ and so on. A straightforward way of calculating entanglement and correlations across the RG flow is to create a second-quantised prototypical form of the IR state $|\Psi_0\rangle$, and then apply the inverse unitary operators $\{U_i\}$ to this wavefunction in order to generate the family of states $|\Psi_i\rangle = U_i |\Psi_{i-1}\rangle, i > 0$. Quantities like entanglement entropy and correlations can then be calculated by using this family of states. If the unitary is of a specific form, there is instead a more efficient method of doing this, which is what we will describe in the next subsection.

2.9.2 Evolution equation for state tensor coefficients

In the following, we restrict ourselves to the reverse RG analysis of an impurity problem hybridising with the local degree of freedom of a conduction bath. At every step, we allow two states from the IOMS to interact with the already-entangled states (one above the Fermi energy and one below, in order to maintain particle-hole symmetry). After the j^{th} application of the reverse RG transformation, $2j$ number of states are entangled with the impurity site, and the state can be represented as

$$|\Psi_j\rangle = \sum_{\nu_d, \nu_1, \dots, \nu_{2j}} C_{\nu_d, \nu_1, \dots, \nu_{2j}, I_{2j+1}}^{(j)} |\nu_d\rangle |\nu_1\rangle |\nu_2\rangle \dots |\nu_{2j}\rangle \otimes |I_{2j+1}\rangle, \quad (9.6)$$

where $|\nu_d\rangle$ and $|\nu_i\rangle$ ($i \in [1, 2j]$) sum over the configurations of the impurity site and the effective zeroth site configuration at the j^{th} step. $|I_{2j+1}\rangle$ is the configuration of the integrals of motion from the $(2j+1)^{\text{th}}$ site and beyond. We define the product state $|\psi_{dz}(j)\rangle = |\nu_d\rangle |\nu_1\rangle \dots |\nu_{2j}\rangle$ for future reference.

We consider a restricted form of RG evolution, where the unitary transformation from IR towards UV can be expressed at any step as

$$U_j^\dagger = 1 + \sum_{i \in [1, 2j]} \sum_{\nu_d, \mu_d, \nu_i, \mu_i} \mathcal{T}_{\nu_d}^{\mu_d} (J_{\nu_d, \nu_i}^{\mu_d, \mu_i} c_{2j+1}^{\nu_i} \mathcal{T}_{\nu_i}^{\mu_i} + K_{\nu_d, \nu_i}^{\mu_d, \mu_i} c_{2j+2}^{\nu_i} \mathcal{T}_{\nu_i}^{\mu_i}), \quad (9.7)$$

where $\mathcal{T}_\alpha^\nu = |\nu\rangle \langle \alpha|$ is a transition operator and sums over all transition operators of the specific subspace (impurity or bath sites). We first obtain the action of the transformation $U_j^\dagger$ on the basis states. We have

$$\begin{aligned} & \mathcal{T}_{\mu_d}^{\nu_d} c_{2j+2}^{\nu_i} \mathcal{T}_{\mu_i}^{\nu_i} |\mu_d\rangle \dots |\mu_i\rangle \dots |2_{2j+1}\rangle |0_{2j+2}\rangle \\ &= \mathcal{T}_{\mu_d}^{\nu_d} c_{2j+2}^{\nu_i} (-1)^{\mu_d + \sum_{k < i} \nu_i} |\mu_d\rangle \dots |\nu_i\rangle \dots |2_{2j+1}\rangle |0_{2j+2}\rangle \\ &= \mathcal{T}_{\mu_d}^{\nu_d} (-1)^{\mu_d + \sum_{k < i} \nu_i} (-1)^{\mu_d + \sum_k \nu_i} |\mu_d\rangle \dots |\nu_i\rangle \dots |2_{2j+1}\rangle |1_{2j+2}\rangle \\ &= (-1)^{\sum_{k \geq i} \nu_i} |\nu_d\rangle \dots |\nu_i\rangle \dots |2_{2j+1}\rangle |1_{2j+2}\rangle, \end{aligned} \quad (9.8)$$

and

$$\mathcal{T}_{\mu_d}^{\nu_d} c_{2j+1}^{\nu_i} \mathcal{T}_{\mu_i}^{\nu_i} |\mu_d\rangle \dots |\mu_i\rangle \dots |2_{2j+1}\rangle |0_{2j+2}\rangle = (-1)^{\sum_{k \geq i} \nu_i} |\nu_d\rangle \dots |\nu_i\rangle \dots |1_{2j+1}\rangle |0_{2j+2}\rangle. \quad (9.9)$$

Using these expressions, we construct the state at the $(j+1)^{\text{th}}$ step:

$$\begin{aligned}
 |\Psi_{j+1}\rangle &= \sum_{\nu_d, \nu_1, \dots, \nu_j} C_{\nu_d, \nu_1, \dots}^{(j)} |\psi_{dz}(j)\rangle |2_{2j+1}\rangle |0_{2j+2}\rangle |I_{2j+3}\rangle \\
 &+ \sum_i \sum_{\mu_d, \mu_i, \nu_d, \nu_i} C_{\mu_d, \dots, \mu_i, \dots}^{(j)} J_{\mu_d, \mu_i}^{\nu_d, \nu_i} (-1)^{\sum_{k \geq i} \nu_k} |\psi_{dz}(j)\rangle |1_{2j+1}\rangle |0_{2j+2}\rangle |I_{2j+3}\rangle \\
 &+ \sum_i \sum_{\mu_d, \mu_i, \nu_d, \nu_i} C_{\mu_d, \dots, \mu_i, \dots}^{(j)} K_{\mu_d, \mu_i}^{\nu_d, \nu_i} (-1)^{\sum_{k \geq i} \nu_k} |\psi_{dz}(j)\rangle |2_{2j+1}\rangle |1_{2j+2}\rangle |I_{2j+3}\rangle
 \end{aligned} \tag{9.10}$$

Gathering similar terms and normalising the state, we can write it as

$$\begin{aligned}
 |\Psi_{j+1}\rangle &= \frac{1}{L_j} \sum_{\nu_d, \{\nu_i\}} |\psi_{dz}(j)\rangle \left[C_{\nu_d, \{\nu_i\}, 1_{2j+1}, 0_{2j+2}, I_{j+1}}^{(j)} |2_{2j+1}\rangle |0_{2j+2}\rangle + C_{+, \nu_d, \{\nu_i\}}^{(j)} |2_{2j+1}\rangle |1_{2j+2}\rangle \right. \\
 &\quad \left. + C_{-, \nu_d, \{\nu_i\}}^{(j)} |1_{2j+1}\rangle |0_{2j+2}\rangle \right] |I_{2j+3}\rangle,
 \end{aligned} \tag{9.11}$$

where $L_j^2 = \sum_{\nu_d, \{\nu_i\}} \left(C_{\nu_d, \{\nu_i\}}^{(j)}{}^2 + C_{+, \nu_d, \{\nu_i\}}^{(j)}{}^2 + C_{-, \nu_d, \{\nu_i\}}^{(j)}{}^2 \right)$ defines the normalisation factor, and

$$\begin{aligned}
 C_{+, \nu_d, \{\nu_i\}} &= \sum_{i, \mu_d, \mu_i} C_{\mu_d, \dots, \mu_i, \dots, \nu_{2j}, 1_{2j+1}, 0_{2j+2}}^{(j)} K_{\mu_d, \mu_i}^{\nu_d, \nu_i} (-1)^{\sum_{k \geq i} \nu_k} \\
 C_{-, \nu_d, \{\nu_i\}} &= \sum_{i, \mu_d, \mu_i} C_{\mu_d, \dots, \mu_i, \dots, \nu_{2j}, 1_{2j+1}, 0_{2j+2}}^{(j)} J_{\mu_d, \mu_i}^{\nu_d, \nu_i} (-1)^{\sum_{k \geq i} \nu_k}.
 \end{aligned} \tag{9.12}$$

We can now read off the renormalisation of the state tensor coefficients:

$$\begin{aligned}
 C_{\nu_d, \nu_1, \dots, \nu_{2j}, 1_{2j+1}, 0_{2j+2}, I_{2j+3}}^{(j+1)} &= \frac{1}{L_j} C_{\nu_d, \nu_1, \dots, \nu_{2j}, 2_{2j+1}, 0_{2j+2}, I_{2j+3}}^{(j)}, \\
 C_{\nu_d, \nu_1, \dots, \nu_{2j}, 2_{2j+1}, 1_{2j+2}, I_{2j+3}}^{(j+1)} &= \frac{1}{L_j} \sum_i (-1)^{\sum_{k \geq i} \nu_k} \sum_{\mu_d, \mu_i} C_{\mu_d, \dots, \mu_i, \dots, \nu_{2j}, 1_{2j+1}, 0_{2j+2}}^{(j)} K_{\mu_d, \mu_i}^{\nu_d, \nu_i}, \\
 C_{\nu_d, \nu_1, \dots, \nu_{2j}, 1_{2j+1}, 0_{2j+2}, I_{2j+3}}^{(j+1)} &= \frac{1}{L_j} \sum_i (-1)^{\sum_{k \geq i} \nu_k} \sum_{\mu_d, \mu_i} C_{\mu_d, \dots, \mu_i, \dots, \nu_{2j}, 1_{2j+1}, 0_{2j+2}}^{(j)} J_{\mu_d, \mu_i}^{\nu_d, \nu_i},
 \end{aligned} \tag{9.13}$$

We now make an important assumption: the coefficients $K_{\mu_d, \mu_i}^{\nu_d, \nu_i}$ and $J_{\mu_d, \mu_i}^{\nu_d, \nu_i}$ are uniform for the set of states $S(\nu_i)$ and zero for other states. That gives

$$\begin{aligned}
 C_{\nu_d, \nu_1, \dots, \nu_{2j}, 1_{2j+1}, 1_{2j+2}, I_{2j+3}}^{(j+1)} &= \frac{1}{L_j} K \sum_i (-1)^{\sum_{k \geq i} \nu_k} \sum_{\substack{\mu_d \in S(\nu_d), \\ \mu_i \in S(\nu_i)}} C_{\mu_d, \dots, \mu_i, \dots, \nu_{2j}, 1_{2j+1}, 0_{2j+2}}^{(j)}, \\
 C_{\nu_d, \nu_1, \dots, \nu_{2j}, 1_{2j+1}, 0_{2j+2}, I_{2j+3}}^{(j+1)} &= \frac{1}{L_j} J \sum_i (-1)^{\sum_{k \geq i} \nu_k} \sum_{\substack{\mu_d \in S(\nu_d), \\ \mu_i \in S(\nu_i)}} C_{\mu_d, \dots, \mu_i, \dots, \nu_{2j}, 1_{2j+1}, 0_{2j+2}}^{(j)},
 \end{aligned} \tag{9.14}$$

One can also look at renormalisation of expectation values of operators; this is useful for tracking the evolution of correlations and entanglement measures. Consider an operator $\mathcal{O}_{\mu_k}^{\nu_k}$ that acts on the

k^{th} bath site. The expectation value of this operator at any step can be written as

$$\begin{aligned}\langle \mathcal{O}_{\nu_k}^{\mu_k} \rangle_j &= \sum_{\nu_d, \nu_1, \dots, \nu_{k-1}, \nu_{k+1}, \dots, \nu_{2j}} C_{\nu_d, \dots, \mu_k, \dots}^{(j)} C_{\nu_d, \dots, \nu_k, \dots}^{(j)}, \\ \langle \mathcal{O}_{\nu_k}^{\mu_k} \rangle_{j+1} &= \sum_{\nu_d, \nu_1, \dots, \nu_{k-1}, \nu_{k+1}, \dots, \nu_{2j+2}} C_{\nu_d, \dots, \mu_k, \dots}^{(j+1)} C_{\nu_d, \dots, \nu_k, \dots}^{(j+1)}\end{aligned}\quad (9.15)$$

2.10 Kondo model spectral function

For the Kondo model defined by the Hamiltonian

$$H_K = \sum_{k, \sigma} \epsilon_k n_{k, \sigma} + J \sum_{k_1, k_2} \mathbf{S}_d \cdot \sigma_{\alpha, \beta} c_{k_1, \alpha}^\dagger c_{k_2, \beta}, \quad (10.1)$$

the impurity spin S_d does not have any fermionic operator. In order to define a spectral function for the impurity, we therefore look at the scattering of conduction electrons off it. This is captured by the conduction bath Greens function $G_{k, k'}$, which can be expressed in terms of a T -matrix when there are two-particle scattering processes present in the system:

$$G_{k, k'}(\omega) = G_k^{(0)}(\omega) + G_k^{(0)}(\omega) T_{k, k'}(\omega) G_{k'}^{(0)}(\omega), \quad (10.2)$$

where $T_{k, k'}$ is non-zero because of the presence of the impurity spin which scatters momentum states into each other. Because of the local nature of the impurity scattering, the T -matrix is also local: $T_{k, k'}(\omega) = T(\omega)$. We therefore use this to define a spectral function:

$$A_{\text{imp}} = -\frac{1}{\pi} \text{Im } T(\omega). \quad (10.3)$$

In order to calculate $T(\omega)$, we evaluate $G_{k, k'}$ using the equation of motion method. It is define as

$$G_{k, k'}(t) = -i\theta(t) \langle \langle c_{k, \sigma}(t), c_{k', \sigma}^\dagger \rangle \rangle. \quad (10.4)$$

Differentiating against time gives

$$\partial_t G_{k, k'}(t) = -i\delta(t) \langle \langle c_{k, \sigma}, c_{k', \sigma}^\dagger \rangle \rangle - \theta(t) \langle \langle [c_{k, \sigma}(t), H_K], c_{k', \sigma}^\dagger \rangle \rangle. \quad (10.5)$$

Evaluating the commutator gives

$$[c_{k, \sigma}(t), H_K] = \epsilon_k c_{k, \sigma} + J \sum_{q, \beta} \mathbf{S}_d \cdot \sigma_{\sigma, \beta} c_{q, \beta}. \quad (10.6)$$

For convenience we define the local bath operator $c_{0, \beta} = \sum_q c_{q, \beta}$. Substituting the commutator back into the expression gives

$$\partial_t G_{k, k'}(t) = -i\delta(t) \delta_{k, k'} - \theta(t) \epsilon_k \langle \langle c_{k, \sigma}(t), c_{k', \sigma}^\dagger \rangle \rangle - \theta(t) J \sum_{\beta} \langle \langle \mathbf{S}_d \cdot \sigma_{\sigma, \beta} c_{0, \beta}, c_{k', \sigma}^\dagger \rangle \rangle(t). \quad (10.7)$$

Defining a new Greens function $\tilde{G}$,

$$\tilde{G}_{k'}(t) = -i\theta(t) \sum_{\beta} \langle \langle \mathbf{S}_d \cdot \sigma_{\sigma,\beta} c_{0,\beta}, c_{k',\sigma}^{\dagger} \rangle \rangle, \quad (10.8)$$

we have the compact form

$$\partial_t G_{k,k'}(t) = -i\delta(t)\delta_{k,k'} - \theta(t)\epsilon_k \langle \langle c_{k,\sigma}(t), c_{k',\sigma}^{\dagger} \rangle \rangle - i\tilde{G}_{k'}(t). \quad (10.9)$$

We now transform to frequency space, using the definition $f(t) = \int d\omega e^{-i\omega t} g(\omega)$:

$$\omega G_{k,k'}(\omega) = \delta_{k,k'} + \epsilon_k G_{k,k'}(\omega) + J\tilde{G}_{k'}(\omega). \quad (10.10)$$

We setup the equation of motion for $\tilde{G}$ in a similar fashion. We use the Hermitian conjugate $-i\theta(t) \sum_{\beta} \langle \langle c_{k',\sigma}(t), \mathbf{S}_d \cdot \sigma_{\sigma,\beta} c_{0,\beta}^{\dagger} \rangle \rangle$, because it is easier to work with:

$$\begin{aligned} \partial_t \tilde{G}_{k'}(t) &= -i\delta(t) \sum_{\beta} \delta_{\sigma,\beta} \langle \mathbf{S}_d \cdot \sigma_{\sigma,\beta} \rangle - \theta(t) \langle \langle [c_{k',\sigma}(t), H_K], \sum_{\beta} \mathbf{S}_d \cdot \sigma_{\sigma,\beta} c_{0,\beta}^{\dagger} \rangle \rangle \\ &= -i\delta(t) \langle \mathbf{S}_d^z \rangle - \theta(t) \langle \langle (\epsilon_{k'} c_{k',\sigma}(t) + J \sum_{\alpha} \mathbf{S}_d \cdot \sigma_{\sigma,\alpha} c_{0,\alpha}), \sum_{\beta} \mathbf{S}_d \cdot \sigma_{\sigma,\beta} c_{0,\beta}^{\dagger} \rangle \rangle \\ &= -i\delta(t) \langle \mathbf{S}_d^z \rangle - i\epsilon_{k'} G_{k'}(t) - J\theta(t) \langle \langle \sum_{\alpha} \mathbf{S}_d \cdot \sigma_{\sigma,\alpha} c_{0,\alpha}, \sum_{\beta} \mathbf{S}_d \cdot \sigma_{\sigma,\beta} c_{0,\beta}^{\dagger} \rangle \rangle. \end{aligned} \quad (10.11)$$

At this point, we define a Kondo excitation operator $O_{\sigma} = J \sum_{\alpha} \mathbf{S}_d \cdot \sigma_{\sigma,\alpha} c_{0,\alpha}$. Transforming to frequency space gives

$$\omega \tilde{G}_{k'}(\omega) = \langle S_d^z \rangle + \epsilon_{k'} G_{k'}(\omega) + \frac{1}{J} \langle \langle O_{\sigma}, O_{\sigma}^{\dagger} \rangle \rangle, \quad (10.12)$$

which gives an expression for $\tilde{G}_{k'}(\omega)$:

$$(\omega - \epsilon_{k'}) \tilde{G}_{k'}(\omega) = \langle S_d^z \rangle + \frac{1}{J} \langle \langle O_{\sigma}, O_{\sigma}^{\dagger} \rangle \rangle. \quad (10.13)$$

Substituting this into eq. 10.10 and noting that the non-interacting Greens function of the conduction bath is $G_k^{(0)} = (\omega - \epsilon_k)^{-1}$, we get

$$(G_k^{(0)})^{-1} G_{k,k'}(\omega) = \delta_{k,k'} + J \left[\langle S_d^z \rangle + \frac{1}{J} \langle \langle O_{\sigma}, O_{\sigma}^{\dagger} \rangle \rangle \right] G_{k'}^{(0)}, \quad (10.14)$$

which can be finally compared with the general expression in eq. 10.2:

$$G_{k,k'}(\omega) = G_k^{(0)} + G_k^{(0)} [J \langle S_d^z \rangle + \langle \langle O_{\sigma}, O_{\sigma}^{\dagger} \rangle \rangle] G_{k'}^{(0)}. \quad (10.15)$$

The T -matrix for scattering of conduction electrons off the impurity can be read off:

$$T_{\sigma}(\omega) = J \langle S_d^z \rangle + \langle \langle O_{\sigma}, O_{\sigma}^{\dagger} \rangle \rangle, \quad (10.16)$$

2.11 Zero temperature Greens function in frequency domain

2.11.1 Spectral representation of Greens function

The impurity retarded Green's function is defined as

$$G_{d\sigma}(t) = -i\theta(t) \langle \{ \mathcal{O}_\sigma(t), \mathcal{O}_\sigma^\dagger \} \rangle \quad (11.1)$$

where the average $\langle \rangle$ is over a canonical ensemble at temperature T , and $\mathcal{O}_\sigma$ is the excitation whose spectral function we are interested in. What follows is a standard calculation where we write the Green's function in the spectral representation. The ensemble average for the operator $\mathcal{O}_\sigma$ can be written in terms of the exact eigenstates of the Hamiltonian:

$$H |n\rangle = E_n |n\rangle, \quad \langle \mathcal{O}_\sigma \rangle \equiv \frac{1}{Z} \sum_n \langle n | \mathcal{O}_\sigma | n \rangle e^{-\beta E_n} \quad (11.2)$$

where $Z = \sum_n e^{-\beta E_n}$ is the partition function and $\{|n\rangle\}$ is the set of eigenfunctions of the Hamiltonian. We can therefore write

$$\begin{aligned} & \langle \{ \mathcal{O}_\sigma(t), \mathcal{O}_\sigma^\dagger \} \rangle \\ &= \frac{1}{Z} \sum_m e^{-\beta E_m} \langle m | \{ \mathcal{O}_\sigma(t), \mathcal{O}_\sigma^\dagger \} | m \rangle \\ &= \frac{1}{Z} \sum_{m,n} e^{-\beta E_m} \langle m | (\mathcal{O}_\sigma(t) | n \rangle \langle n | \mathcal{O}_\sigma^\dagger + \mathcal{O}_\sigma^\dagger | n \rangle \langle n | \mathcal{O}_\sigma(t)) | m \rangle \quad \left[\sum_n |n\rangle \langle n| = 1 \right] \\ &= \frac{1}{Z} \sum_{m,n} e^{-\beta E_m} \langle m | (e^{iH^*t} \mathcal{O}_\sigma e^{-iH^*t} | n \rangle \langle n | \mathcal{O}_\sigma^\dagger + \mathcal{O}_\sigma^\dagger | n \rangle \langle n | e^{iH^*t} \mathcal{O}_\sigma e^{-iH^*t}) | m \rangle \\ &= \frac{1}{Z} \sum_{m,n} e^{-\beta E_m} (e^{i(E_m - E_n)t} \langle m | \mathcal{O}_\sigma | n \rangle \langle n | \mathcal{O}_\sigma^\dagger | m \rangle + e^{i(E_n - E_m)t} \langle m | \mathcal{O}_\sigma^\dagger | n \rangle \langle n | \mathcal{O}_\sigma | m \rangle) \\ &= \frac{1}{Z} \sum_{m,n} e^{i(E_m - E_n)t} || \langle m | \mathcal{O}_\sigma | n \rangle ||^2 (e^{-\beta E_m} + e^{-\beta E_n}) \end{aligned} \quad (11.3)$$

The time-domain impurity Green's function can thus be written as (this is the so-called Lehmann-Kallen representation)

$$G_{d\sigma} = -i\theta(t) \frac{1}{Z} \sum_{m,n} e^{i(E_m - E_n)t} || \langle m | \mathcal{O}_\sigma | n \rangle ||^2 (e^{-\beta E_m} + e^{-\beta E_n}) \quad (11.4)$$

We are interested in the frequency domain form.

$$\begin{aligned} G_{dd}^\sigma(\omega) &= \int_{-\infty}^{\infty} dt e^{i(\omega)t} G_{dd}^\sigma(t) \\ &= \frac{1}{Z} \sum_{m,n} || \langle m | \mathcal{O}_\sigma | n \rangle ||^2 (e^{-\beta E_m} + e^{-\beta E_n}) (-i) \int_{-\infty}^{\infty} dt \theta(t) e^{i(\omega + E_m - E_n)t} \end{aligned} \quad (11.5)$$

To evaluate the time-integral, we will use the integral representation of the Heaviside function:

$$\theta(t) = -\frac{1}{2\pi i} \lim_{\eta \rightarrow 0^+} \int_{-\infty}^{\infty} \frac{1}{x + i\eta} e^{-ixt} dx \quad (11.6)$$

With this definition, the integral in $G_{d\sigma}(\omega)$ becomes

$$\begin{aligned} (-i) \int_{-\infty}^{\infty} dt \theta(t) e^{i(\omega + i0^+ + E_m - E_n)t} &= (-i) \frac{-1}{2\pi i} \lim_{\eta \rightarrow 0^+} \int_{-\infty}^{\infty} dx \frac{1}{x + i\eta} \int_{-\infty}^{\infty} dt e^{i(\omega + i0^+ + E_m - E_n - x)t} \\ &= \frac{1}{2\pi} \lim_{\eta \rightarrow 0^+} \int_{-\infty}^{\infty} dx \frac{1}{x + i\eta} 2\pi \delta(\omega + i0^+ + E_m - E_n - x) \\ &= \frac{1}{\omega + E_m - E_n + i0^+} . \end{aligned} \quad (11.7)$$

The frequency-domain Green's function is thus

$$G_{dd}^{\sigma}(\omega) = \frac{1}{Z} \sum_{m,n} ||\langle m | \mathcal{O}_{\sigma} | n \rangle||^2 (e^{-\beta E_m} + e^{-\beta E_n}) \frac{1}{\omega + E_m - E_n + i0^+} \quad (11.8)$$

The infinitesimally positive imaginary part in the denominator shifts the poles onto the lower half of the complex plane, leaving it analytic in the upper half: this is necessary to make the retarded Greens function causal.

2.11.2 Real and imaginary parts - The Sokhotski-Plemelj theorem

In order to write down the real and imaginary parts of the Greens function, we first prove the *Sokhotski-Plemelj* formula:

$$\lim_{\eta \rightarrow 0^+} \frac{1}{x + i\eta} = P\left(\frac{1}{x}\right) - i\pi\delta(x) . \quad (11.9)$$

where $P(f(x))$ is the Cauchy principal value, defined as

$$P\left[\int_{-\infty}^{\infty} \frac{dx}{x}\right] = \int_{-\infty}^{0^-} \frac{dx}{x} + \int_{0^+}^{\infty} \frac{dx}{x} . \quad (11.10)$$

To prove the above identity, we integrate the left-hand side using a test function $f(x)$:

$$\begin{aligned} \lim_{\eta \rightarrow 0^+} \int_{-\infty}^{\infty} dx \frac{f(x)}{x + i\eta} &= \lim_{\eta \rightarrow 0^+} \int_{-\infty}^{\infty} dx \frac{f(x)(x - i\eta)}{x^2 + \eta^2} \\ &= \lim_{\eta \rightarrow 0^+} \int_{-\infty}^{\infty} dx f(x) \left[\frac{x}{x^2 + \eta^2} - i\eta \frac{1}{x^2 + \eta^2} \right] \end{aligned} \quad (11.11)$$

The first integral can be split into three parts:

$$\lim_{\eta \rightarrow 0^+} \int_{-\infty}^{\infty} dx f(x) \frac{x}{x^2 + \eta^2} = \lim_{\eta \rightarrow 0^+} \lim_{\varepsilon \rightarrow 0^+} \left[\int_{-\infty}^{-\varepsilon} dx \frac{f(x)x}{x^2 + \eta^2} + \int_{\varepsilon}^{\infty} dx \frac{f(x)x}{x^2 + \eta^2} + \int_{-\varepsilon}^{\varepsilon} dx \frac{f(x)x}{x^2 + \eta^2} \right] \quad (11.12)$$

For the first two parts, it is safe to take the limit of η , because x is always non-vanishing there. For the third part, we can approximate $f(x)$ as $f(0)$ in the neighbourhood $x \in [-\varepsilon, \varepsilon], \varepsilon \rightarrow 0$. This leaves an odd function $x/(x^2 + \eta^2)$ as the integrand, being integrated over a symmetric range. The third term therefore vanishes. In total, we get

$$\lim_{\eta \rightarrow 0^+} \int_{-\infty}^{\infty} dx f(x) \frac{x}{x^2 + \eta^2} = \int_{-\infty}^{-\varepsilon} dx \frac{f(x)}{x} + \int_{\varepsilon}^{\infty} dx \frac{f(x)}{x} = P \left(\int_{-\infty}^{\infty} dx \frac{f(x)}{x} \right). \quad (11.13)$$

To evaluate the second integral of eq. 11.11, we note that the only region in which the integrand is non-zero is when $|x| \simeq 0$; there, we again approximate $f(x)$ as $f(0)$. The remaining integral can then be evaluated easily:

$$-i \lim_{\eta \rightarrow 0^+} \eta \int_{-\infty}^{\infty} dx \frac{f(x)}{x^2 + \eta^2} = -i \lim_{\eta \rightarrow 0^+} \eta f(0) \int_{-\infty}^{\infty} \frac{dx}{x^2 + \eta^2} = -i\pi f(0), \quad (11.14)$$

where we used $\int \frac{dx}{x^2 + \eta^2} = \frac{1}{\eta} \arctan(x/\eta)$. Combining eqs. 11.11, 11.13 and 11.14, we get

$$\lim_{\eta \rightarrow 0^+} \int_{-\infty}^{\infty} dx \frac{f(x)}{x + i\eta} = P \left(\int_{-\infty}^{\infty} dx \frac{f(x)}{x} \right) - i\pi \int_{-\infty}^{\infty} dx f(x) \delta(x). \quad (11.15)$$

This proves eq. 11.9.

The Sokhotski-Plemelj formula allows us to split the spectral representation into a real and an imaginary part:

$$\begin{aligned} G'_{d\sigma} &= \frac{1}{Z} \sum_{m,n} \|\langle m | \mathcal{O}_\sigma | n \rangle\|^2 (e^{-\beta E_m} + e^{-\beta E_n}) P \left(\frac{1}{\omega + E_m - E_n} \right) \\ G''_{d\sigma} &= -\pi \frac{1}{Z} \sum_{m,n} \|\langle m | \mathcal{O}_\sigma | n \rangle\|^2 (e^{-\beta E_m} + e^{-\beta E_n}) \delta(\omega + E_m - E_n). \end{aligned} \quad (11.16)$$

The spectral function, specifically, is defined as follows:

$$A_{d\sigma}(\omega) = -\frac{1}{\pi} G'_{d\sigma} = \frac{1}{Z} \sum_{m,n} \|\langle m | \mathcal{O}_\sigma | n \rangle\|^2 (e^{-\beta E_m} + e^{-\beta E_n}) \delta(\omega + E_m - E_n) \quad (11.17)$$

2.11.3 Zero temperature spectral function

We specialise to zero temperature by taking the limit of $\beta \rightarrow \infty$. In both the partition function as well as inside the summation, the only term that will survive is the exponential of the ground state energy E_{GS} .

$$Z \equiv \sum_m e^{-\beta E_m} \implies \lim_{\beta \rightarrow \infty} Z = d_{\text{GS}} e^{-\beta E_{\text{GS}}}, \quad E_{\text{GS}} \equiv \min \{E_n\}$$

where d_{GS} is the degeneracy of the ground state. The spectral function then simplifies to

$$\begin{aligned} A_{d\sigma} &= \frac{1}{d_{\text{GS}} e^{-\beta E_{\text{GS}}}} \sum_{m,n} \|\langle m | \mathcal{O}_\sigma | n \rangle\|^2 [e^{-\beta E_m} \delta_{E_m, E_{\text{GS}}} + e^{-\beta E_n} \delta_{E_n, E_{\text{GS}}}] \delta(\omega + E_m - E_n) \\ &= \frac{1}{d_{\text{GS}}} \sum_{n, n_{\text{GS}}} [\|\langle n_{\text{GS}} | \mathcal{O}_\sigma | n \rangle\|^2 \delta(\omega + E_{\text{GS}} - E_n) + \|\langle n | \mathcal{O}_\sigma | n_{\text{GS}} \rangle\|^2 \delta(\omega - E_{\text{GS}} + E_n)] \end{aligned} \quad (11.18)$$

The label n_{GS} sums over all states $|n_{\text{GS}}\rangle$ with energy E_{GS} . In practise, we evaluate this expression by replacing the formal Dirac delta functions with "nascent" ones, such as a Lorentzian function with a sufficiently sharp peak.

2.11.4 Reconstructing full Green function from spectral function: Kramers-Kronig relations

As mentioned earlier, the retarded Greens function $G_R(\omega + i0^+)$ has a complex pole in the lower half of the complex plane. In the following, the frequency ω is assumed to be complex. Consider the integral

$$I(\omega) = \oint_C d\omega' \frac{G_R(\omega')}{\omega' - \omega}, \quad \omega \in \mathbb{R}. \quad (11.19)$$

The contour C encloses the entirety of the upper half of the complex plane as well as almost all points of the real line but avoids the pole at $\omega' = \omega$ by forming a semicircle about that point that extends into the upper half of the plane. Since the integrand is completely analytic in the region enclosed by the contour (recall again that G_R is analytic in the upper half), the integral evaluates to zero (by Cauchy's theorem):

$$I(\omega) = 0. \quad (11.20)$$

One can also evaluate the integral along each part of the contour. The semicircular part contributes zero: this is because while the length of the semicircular arc goes as $|\omega'|$, the integrand vanishes faster than $1/|\omega'|$ as $|\omega'| \rightarrow \infty$ (this assumes that the Green function vanishes in the limit of $\omega \rightarrow \infty$). The part of the contour along the real axis can now be evaluated. The two parts of the contour on either side of the pole contribute

$$\int_{-\infty}^{\omega-0^+} d\omega' \frac{G_R(\omega')}{\omega' - \omega} + \int_{\omega+0^+}^{\infty} d\omega' \frac{G_R(\omega')}{\omega' - \omega} = P \int_{-\infty}^{\infty} d\omega' \frac{G_R(\omega')}{\omega' - \omega} \quad (11.21)$$

To evaluate the part of the integral along the small semicircular around $\omega' = \omega$ (that extends into the upper half), we argue that this contribution should be equal to that obtained by taking the semicircle that goes instead into the lower half but circulates in the opposite direction. Each of these contributions should, in turn, be equal to that obtained from joining these two contours. But joining these contours leads to a closed integral about the pole at $\omega' = \omega$, which is equal to $2\pi G_R(\omega)$.

$$\int_{\text{upper}} d\omega' \frac{G_R(\omega')}{\omega' - \omega} = \int_{\text{lower}} d\omega' \frac{G_R(\omega')}{\omega' - \omega} = \frac{1}{2} \oint_{C(\omega)} d\omega' \frac{G_R(\omega')}{\omega' - \omega} = i\pi G_R(\omega). \quad (11.22)$$

Combining the last two equations, we get

$$I(\omega) = P \int_{-\infty}^{\infty} d\omega' \frac{G_R(\omega')}{\omega' - \omega} + i\pi G_R(\omega). \quad (11.23)$$

Comparing with eq. 11.20 and separating into real part G'_R and imaginary part G''_R gives

$$P \int_{-\infty}^{\infty} d\omega' \frac{[G'_R(\omega') + iG''_R(\omega')]}{\omega' - \omega} + \pi [-G''_R(\omega) + iG'_R(\omega)] = 0. \quad (11.24)$$

Comparing the imaginary parts gives

$$G'_R(\omega) = \frac{1}{\pi} P \int_{-\infty}^{\infty} d\omega' \frac{G''_R(\omega')}{\omega' - \omega} = -\mathcal{H} [G''_R(\omega)] , \quad (11.25)$$

where $\mathcal{H} [g(\omega)] = \frac{1}{\pi} P \int_{-\infty}^{\infty} d\omega' \frac{g(\omega')}{\omega - \omega'}$ is the Hilbert transform of the function $g(\omega)$. The spectral function was defined above as $-1/\pi$ times the imaginary part of the Greens function. Exchanging G''_R for A gives

$$G'_R(\omega) = \pi \mathcal{H} [A(\omega)] . \quad (11.26)$$

This provides a method for calculating the real part of a Greens function if the spectral function is already known.

Chapter 3

Kondo Frustration via Charge Fluctuations: A Route to Mott Localisation in Infinite Dimensions

3.1 Introduction

The rich physics of metal-insulator transitions in strongly correlated systems has been an active subject of study for quite some time [Mott(1968), Milligan and Thomas(1985), Imada et al.(1998a) Imada, Fujimori, and Tokura], yet much still remains to be understood. It involves diverse aspects such as spin and charge fluctuations, quasiparticle renormalisation effects, anomalous metallic phases and unconventional superconductivity, and has been studied using an equally diverse array of methods like mean-field theory, renormalisation group approaches, numerical techniques like exact diagonalisation, quantum Monte Carlo and dynamical mean-field theory, and many others. In particular, dynamical mean-field theory (DMFT) [Kuramoto and Watanabe(1987), Cox and Grewe(1988), Metzner and Vollhardt(1989), Zhang et al.(1993) Zhang, Rozenberg, and Kotliar, Georges et al.(1996) Georges, Kotliar, Krauth, and Rozenberg, Parcollet et al.(2004) Parcollet, Biroli, and Kotliar, Maier et al.(2005) Maier, Jarrell, Pruschke, and Hettler, Kotliar et al.(2006) Kotliar, Savrasov, Haule, Oudovenko, Parcollet, and Marianetti, Ohashi et al.(2008) Ohashi, Momoi, Tsunetsugu, and Kawakami] obtains an exact solution of the Mott metal-insulator transition (MIT) for the 1/2-filled Hubbard model [Mott(1949), Gutzwiller(1963), Kanamori(1963), Hubbard(1963), Brinkman and Rice(1970)] on the Bethe lattice with infinite coordination number, in terms of an Anderson impurity model with a self-consistently determined bath obtained by requiring, in an iterative manner, that its local Greens function be equal to that of the impurity site. The above-mentioned transition can be captured by the local spectral function through (i) the continuous appearance of a Mott gap, followed by (ii) the sharpening and vanishing of the central Kondo resonance. These two features are often referred to, respectively, as the Mott-Hubbard [Hubbard(1963)] and Brinkman-Rice [Brinkman and Rice(1970)] scenarios of the Mott MIT. The exact nature of the solution arises from the fact that all non-local contributions to the lattice self-energy are observed to vanish upon taking the limit of an infinite coordination number for the lattice model. Thus, the simplification is that the dynamics of any local site on the lattice is de-

terminated completely by a quantum impurity problem [Georges et al.(1996)Georges, Kotliar, Krauth, and Rozenberg, Logan and Galpin(2015)]. It must also be noted that this exact solution precludes any long-range order in the system, and corresponds to the case of a maximally frustrated Hubbard model involving long-range and frustrating inter-site hopping such that both the metallic and insulating phases remain paramagnetic [Vučićević et al.(2013)Vučičević, Terletska, Tanasković, and Dobrosavljević]. Due to the exact and non-perturbative nature of the DMFT solution for the MIT in $d = \infty$, the method has been extended to models of strongly correlated electrons in finite spatial dimensions [Park et al.(2008)Park, Haule, and Kotliar, Rohringer et al.(2018)Rohringer, Hafermann, Toschi, Katanin, Antipov, Katsnelson, Lichtenstein, Rubtsov, and Held, Maier et al.(2005)Maier, Jarrell, Pruschke, and Hettler], as well as the study of the electronic properties of various correlated materials [Lichtenstein and Katsnelson(1998), Kotliar et al.(2006)Kotliar, Savrasov, Haule, Oudovenko, Parcollet, and Marianetti, Held(2007)].

Despite this progress, a key aspect of the DMFT solution for the Hubbard model in $d = \infty$ remains to be understood. During the search for a self-consistent impurity model, the conduction bath is modified drastically in order to become correlated [Held et al.(2008)Held, Katanin, and Toschi]. The numerical implementation of self-consistency, however, precludes a deeper understanding of the precise nature of the correlations present in the conduction bath of the impurity model, and its implications for the electron dynamics of the associated bulk lattice (Hubbard) model. Indeed, as we demonstrate below, our work paves the way for a more insightful exploration of the physics of the Mott Hubbard transition in strongly correlated electronic systems. Further, it has considerable consequences for the development of functional quantum materials and next-generation quantum technologies. Below, we lay out the specific questions addressed by us, and summarise our results at the end of this section.

- i. Is there a minimal but effective quantum impurity model Hamiltonian that describes the Mott MIT of the $1/2$ -filled Hubbard model on the Bethe lattice in $d = \infty$? Finding such an impurity model would also reduce the considerable computational effort that is presently required in self-consistent approaches.
- ii. What are the fluctuations that destroy the metal and lead to the insulating phase? Can we obtain a universal theory for these competing tendencies?
- iii. The coexistence of metallic and insulating phases at $T = 0$ within DMFT shows that the insulating solution is present within the many-body spectrum of the metallic phase. Can an associated impurity model Hamiltonian display the emergence of the insulating state prior to the transition? Analytic insight of this kind is particularly important because approaching a coexistence region numerically is often fairly tricky.
- iv. Is it possible to obtain a low-energy theory for the local gapless excitations precisely at the MIT, where the metal is on the brink of destruction? How do these excitations compare with those of the local Fermi liquid, e.g., in terms of self-energies and two-particle correlation functions?

Obtaining answers to these questions would go a long way towards understanding the dominant fluctuations at the heart of the Mott transition and the physical processes that become important as

one approaches it. It would also help in understanding the mechanism of DMFT's search for the self-consistent impurity model within the space of Anderson impurity models.

The essence of our approach is to model phenomenologically the lattice self-energy obtained from DMFT in the form of additional bath correlations within an extended Anderson impurity model. In addition to the usual on-site repulsion (U) and single-particle hybridisation (V) between the impurity and the conduction bath of the Anderson impurity model (eq. (2.1)), we introduce (i) an additional on-site correlation (U_b) on the bath site with which the impurity couples, and (ii) an antiferromagnetic Kondo coupling (J) between the impurity and the conduction bath (eq. (2.2)). We note that a similar impurity model-based approach was taken towards understanding the physics of the heavy fermions several years ago by Si and Kotliar [Si and Kotliar(1993a), Kotliar and Si(1993a)]. We postpone a comparison of our work with theirs to the discussions section. The rest of the work is structured as follows. Sec. (3.2.1) describes the extended model that we will study, and the unitary renormalisation group (URG) method that we employ to study it is presented in Sec. (3.2.2). In Sec. (3.3) and (3.4), we describe the phase diagram and various characteristics of the impurity phase transition. In Sec. (3.6), we use our extended model to explain various features of the coexistence region observed in DMFT. In Sec. (3.7), we describe the effect of the impurity on the low-lying excitations of the bath, near and at the transition. We conclude in Sec. (3.8) with some discussions and possible future directions. For the convenience of readers, we first present below a brief summary of our main results.

Summary of our main results

- *Presence of a local metal-insulator transition:* At a critical value of the parameter $r = -U_b/J$, the effective impurity model shows a transition from a Kondo screened phase into an unscreened local moment phase. The quantum critical point (QCP) involves a degeneration of the Kondo singlet and the local moment states.
- *The physics of Kondo screening and local pairing drives the transition:* The transition involves the frustration of the Kondo screening of the impurity by enhanced local pairing fluctuations in the bath and can be described by a universal theory written in terms of J and U_b .
- *Emergence of insulating solutions in the metallic phase:* Our analysis reveals that at a certain value of the parameter r prior to the transition, the single-particle hybridisation parameter (V) turns irrelevant (in the RG sense), and this leads to the emergence of the local moment solutions within the many-particle spectrum through an excited state quantum phase transition (ESQPT).
- *Critical fluctuations and the coexistence region:* We observe the appearance of long-ranged quantum fluctuations extending into the conduction bath in the vicinity of both the ESQPT at $r = r_{c1}$ and QPT at $r = r_{c2}$. We believe that these are the likely origin of the critical fluctuations observed above the finite temperature second-order critical point in DMFT [Terletska et al.(2011)Terletska, Vučićević, Tanasković, and Dobrosavljević, Vučićević et al.(2013)Vučičević, Terletska, Tanasković, and Dobrosavljević]. The two-step process at $T = 0$ also provides a

natural explanation for the coexistence of metallic and insulating features in the phase diagram, in the regime $r_{c1} < r < r_{c2}$.

- *Emergence of non-Fermi liquid excitations at the QCP:* Precisely at the QCP, the local Fermi liquid is replaced by a quasi-local non-Fermi liquid (NFL) that spans the impurity, zeroth and first sites of the conduction bath. The NFL results from a degeneracy between the local moment and singlet states and leads to (i) “Andreev scattering” of incoming states into orthogonal outgoing states, (ii) anomalous power-law behaviour in the self-energies and two-particle correlations with universal exponents, and (iii) a fractional entanglement entropy of the impurity.
- *Correlated Fermi liquid excitations in the Hubbard sidebands:* A many-body perturbation theoretic treatment of the Hubbard sidebands reveals that they are comprised of the holon-doublon excitations created by the hybridisation of the impurity site with the conduction bath. These excitations consist of decoupled local Fermi liquids for the holons and doublons at the lowest order, which, at higher orders, become coupled via correlated holon-doublon scattering between impurity and bath.

3.2 Model Hamiltonian and method

3.2.1 The extended Anderson impurity model

The single-impurity Anderson model (SIAM) [Anderson(1961), Anderson(1978)] consists of a single impurity site with local repulsive correlation U hybridising with a non-interacting fermionic conduction bath through a (momentum-independent) single-particle transfer whose coupling is V . For the case of a half-filled impurity site, the Hamiltonian of the SIAM is given by

$$\mathcal{H}_A = -\frac{U}{2} (\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2 + \sum_{\vec{k}, \sigma} \epsilon_{\vec{k}} \tau_{\vec{k}, \sigma} + V \sum_{\sigma} \left(c_{d\sigma}^\dagger c_{0\sigma} + \text{h.c.} \right), \quad (2.1)$$

where $\tau_{\vec{k}, \sigma} \equiv c_{\vec{k}, \sigma}^\dagger c_{\vec{k}, \sigma} - 1/2$ indicate the occupancy of the single-particle momentum state $|\vec{k}\rangle$. Also, $c_{d\sigma}$ and $c_{0\sigma} = \sum_k c_{k\sigma}$ are the fermionic annihilation operators of spin σ for the impurity and conduction bath site to which it couples (henceforth referred to as the *zeroth site*) respectively. The conduction bath is typically considered to possess a constant (i.e., energy-independent) density of states.

The SIAM (along with its $U \rightarrow \infty$ limit, the Kondo model) has been studied using several analytical and numerical techniques [Anderson and Yuval(1969), Anderson et al.(1970)Anderson, Yuval, and Hamann, Anderson(1970), Haldane(1978), Jefferson(1977), Wilson(1975b), Krishna-murthy et al.(1980)Krishna-murthy, Wilkins, and Wilson, Andrei(1980), Andrei et al.(1983)Andrei, Furuya, and Lowenstein, Wiegmann(1981), Tsvelick and Wiegmann(1983), Kotliar and Si(1996), Duki(2011), Borda et al.(2008)Borda, Schiller, and Zawadowski, Streib et al.(2013)Streib, Isidori, and Kopietz, Mukherjee et al.(2022)Mukherjee, Mukherjee, Vidhyadhiraja, Taraphder, and Lal]. The general conclusion for the positive U case at $T = 0$ is that on the particle-hole symmetric (that is,

half-filled) line, the impurity local moment is always screened by the conduction electrons (referred to as the Kondo cloud [Sørensen and Affleck(1996), Affleck and Simon(2001), Simon and Affleck(2003), Busser et al.(2010)Busser, Martins, Ribeiro, Vernek, Anda, and Dagotto, Ribeiro et al.(2019)Ribeiro, Martins, Gomez-Silva, and Anda, Goldhaber-Gordon et al.(1998)Goldhaber-Gordon, Shtrikman, Mahalu, Abusch-Magder, Meirav, and Kastner,Cronenwett et al.(1998)Cronenwett, Oosterkamp, and Kouwenhoven, Schmid et al.(1998)Schmid, Weis, Eberl, and v. Klitzing, Pustilnik and Glazman(2004), V. Borzenets et al.(2020)V. Borzenets, Shim, Chen, Ludwig, Wieck, Tarucha, Sim, and Yamamoto, Néel et al.(2008)Néel, Kröger, Berndt, Wehling, Lichtenstein, and Katsnelson,Zhao et al.(2005)Zhao, Li, Chen, Xiang, Wang, Pan, Wang, Xiao, Yang, Hou et al.]). Enhanced spin-flip scattering at low-energies leads to the formation of a macroscopic singlet ground state and local Fermi liquid gapless excitations [Nozaki et al.(2012)Nozaki, Ryu, and Takayanagi, Mora et al.(2015)Mora, Moca, von Delft, and Zaránd]. In order to enhance the SIAM, we introduce two extra two-particle interaction terms into the Hamiltonian:

- a spin-exchange term $J\vec{S}_d \cdot \vec{S}_0$ between the impurity spin $\vec{S}_d$ and the spin $\vec{S}_0$ of the zeroth site, and
- a local particle-hole symmetric correlation term $-U_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2$ on the bath zeroth site.

With these additional terms, the Hamiltonian of the *extended single-impurity Anderson model* (henceforth referred to as the e-SIAM) is, at particle-hole symmetry, given by

$$\mathcal{H}_{\text{E-A}} = \mathcal{H}_A + J\vec{S}_d \cdot \vec{S}_0 - \frac{1}{2}U_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2 . \quad (2.2)$$

All the terms in the Hamiltonian have been depicted schematically in the left panel of Fig. (3.1). The additional interaction terms J and U_b enjoy particle-hole, SU(2)-spin and U(1)-charge symmetries. The e-SIAM Hamiltonian (eq. (2.2)) therefore preserves all the local symmetries of the half-filled Hubbard model on a lattice, and can potentially serve as an effective auxiliary quantum impurity model (with a correlated bath) describing the local physics of the latter.

3.2.2 The unitary renormalisation group method

In order to obtain the various low-energy phases of the e-SIAM, we perform a scaling analysis of the associated Hamiltonian (eq. (2.2)) using the recently developed unitary renormalisation group (URG) method [Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)]. The method has been applied successfully on a wide variety of problems of correlated fermions [Pal et al.(2019)Pal, Mukherjee, and Lal, Mukherjee et al.(2021)Mukherjee, Patra, and Lal, Mukherjee and Lal(2020d), Mukherjee and Lal(2020c), Patra and Lal(2021), Mukherjee et al.(2022)Mukherjee, Mukherjee, Vidhyadhiraja, Taraphder, and Lal, Patra et al.(2023a)Patra, Mukherjee, Mukherjee, Vidhyadhiraja, Taraphder, and Lal, Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)]. The method proceeds by resolving quantum fluctuations in high-energy degrees of freedom, leading to a low-energy Hamiltonian with renormalised couplings and new emergent degrees of freedom. Typically, for a system with Fermi energy ϵ_F and bandwidth D_0 , the sequence of isoenergetic shells $\{D_{(j)}\}$, $D_{(j)} \in [\epsilon_F, D_0]$ define the states whose quantum fluctuations we sequentially resolve. The momentum states lying on shells

$D_{(j)}$ that are far away from the Fermi surface comprise the UV states, while those on shells near the Fermi surface comprise the IR states. This scheme is shown in the right panel of Fig. (3.1).



Figure 3.1: Left: Schematic 1D representation of the extended SIAM Hamiltonian. The red sphere is the impurity site with on-site Hubbard term U . It is connected via two couplings V and J to the bath zeroth site (large blue sphere) that has its own on-site Hubbard term U_b . The small blue spheres make up the rest of the bath, connected through the single-particle hopping t . Right: High energy - low energy scheme defined and used in the URG method. The states away from the Fermi surface form the UV subspace and are decoupled first, leading to a Hamiltonian which is more block-diagonal and comprised of only the IR states near the Fermi surface.

As a result of the URG transformations, the Hamiltonian $H_{(j)}$ at a given RG step j involves scattering processes between the k -states that have energies lower than $D_{(j+1)}$. The unitary transformation $U_{(j)}$ is then defined so as to remove the number fluctuations of the currently most energetic set of states $D_{(j)}$ [Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)]:

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger, \text{ such that } [H_{(j-1)}, \hat{n}_j] = 0. \quad (2.3)$$

The eigenvalue of $\hat{n}_j$ has, thus, been rendered an integral of motion (IOM) under the RG transformation.

The unitary transformations can be expressed in terms of a generator $\eta_{(j)}$ that has fermionic algebra [Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)]:

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 + \eta_{(j)} - \eta_{(j)}^\dagger \right), \quad \left\{ \eta_{(j)}, \eta_{(j)}^\dagger \right\} = 1, \quad (2.4)$$

where $\{\cdot\}$ is the anticommutator. The unitary operator $U_{(j)}$ that appears in Eq. (2.7) can be cast into the well-known general form $U = e^{\mathcal{S}}$, $\mathcal{S} = \frac{\pi}{4} \left(\eta_{(j)}^\dagger - \eta_{(j)} \right)$ that a unitary operator can take, defined by an anti-Hermitian operator $\mathcal{S}$. The generator $\eta_{(j)}$ is given by the expression [Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)]

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - \text{Tr}(H_{(j)} \hat{n}_j)} c_j^\dagger \text{Tr}(H_{(j)} c_j). \quad (2.5)$$

The operators $\eta_{(j)}, \eta_{(j)}^\dagger$ behave as the many-particle analogues of the single-particle field operators $c_j, c_j^\dagger$ - they change the occupation number of the single-particle Fock space $|n_j\rangle$. The important operator $\hat{\omega}_{(j)}$ originates from the quantum fluctuations that exist in the problem because of the non-commutation of the kinetic energy terms and the interaction terms in the Hamiltonian:

$$\hat{\omega}_{(j)} = H_{(j-1)} - H_{(j)}^i. \quad (2.6)$$

$H_{(j)}^i$ is the part of $H_{(j)}$ that commutes with $\hat{n}_j$ but does *not* commute with at least one $\hat{n}_l$ for $l < j$. The RG flow continues up to energy D^* , where a fixed point is reached from the vanishing of the RG function. Detailed comparisons of the URG with other methods (e.g., the functional RG, spectrum bifurcation RG etc.) can be found in Refs. [Mukherjee and Lal(2020c), Mukherjee and Lal(2020a)]. More information on the unitary RG method is provided in Section 1 of the Supplementary Materials [sup(Supplementary Materials are available online)].

3.3 $T = 0$ scaling theory for the e-SIAM

3.3.1 URG equations for the e-SIAM

The derivation of the RG equations for the extended Anderson impurity model (e-SIAM) Hamiltonian is shown in Section 2 of the Supplementary Materials [sup(Supplementary Materials are available online)]. The bath coupling U_b is marginal. We provide below the RG equations of the remaining couplings for a given quantum fluctuation scale ω :

$$\begin{aligned}\Delta U &= 4V^2 n_j \left(\frac{1}{d_1} - \frac{1}{d_0} \right) - n_j \frac{J^2}{d_2}, \\ \Delta V &= -\frac{3n_j V}{8} \left[J \left(\frac{1}{d_2} + \frac{1}{d_1} \right) + \frac{4U_b}{3} \sum_{i=1}^4 \frac{1}{d_i} \right], \\ \Delta J &= -\frac{n_j J (J + 4U_b)}{d_2},\end{aligned}\tag{3.1}$$

where the denominators d_i are given by

$$d_0 = \omega - \frac{D}{2} + \frac{U_b}{2} - \frac{U}{2}, \quad d_1 = \omega - \frac{D}{2} + \frac{U_b}{2} + \frac{U}{2} + \frac{J}{4},\tag{3.2}$$

$$d_2 = \omega - \frac{D}{2} + \frac{U_b}{2} + \frac{J}{4}, \quad d_3 = \omega - \frac{D}{2} + \frac{U_b}{2}.\tag{3.3}$$

The symbols used in the RG equations have the following meanings: ΔU represents the renormalisation of the coupling U in going from the j^{th} Hamiltonian to the $(j-1)^{\text{th}}$ Hamiltonian by decoupling the isoenergetic shell at energy $D_{(j)}$ (see right panel of Fig.(3.1)). n_j is the number of electronic states on the shell $D_{(j)}$. We note that the labels U_0, J_0, V_0 that appear in various figures (and elsewhere in the text) represent the bare values of the associated couplings U, J and V . The RG equations reduce, in the perturbative regime of the couplings U, V and J , to the well-known “poor man’s” scaling forms obtained for the SIAM [Haldane(1978)] and the single-channel Kondo model [Anderson(1970)] respectively.

The RG fixed point Hamiltonian describes the low-energy phase of the system. In general, if the RG fixed point is reached at an energy scale D^* , the fixed point Hamiltonian $\mathcal{H}^*$ is obtained simply from the fixed point values of the couplings (and by noting that the states above D^* are now part of

the IOMs):

$$\mathcal{H}^* = \sum_{\sigma, \vec{k}}^{\epsilon_{\vec{k}} < D^*} \epsilon_{\vec{k}} \tau_{\vec{k}, \sigma} + V^* \sum_{\sigma} \left(c_{d\sigma}^\dagger c_{0\sigma} + \text{h.c.} \right) + J^* \vec{S}_d \cdot \vec{S}_0 - \frac{1}{2} U^* (\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2 - \frac{1}{2} U_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2.$$

The fixed point values of the couplings are obtained by solving the RG equations numerically.




Figure 3.2: Left: Phase diagram of the *e*-SIAM, in the space of r and J_0/D_0 . U_b is set to $-U_0/10$. The various phases are described in the text. Right: Schematic nature of RG flows for $U_b = 0$ and $U_b = -U/10$. Blue curves represent RG flows towards the strong-coupling (SC) fixed point, while red curves represent RG flows towards the local moment (LM) fixed point.

3.3.2 Phase diagram

We will work in the low-energy regime where the quantum fluctuation scale ω is such that all the denominators are negative: $d_i < 0 \forall i$. Moreover, we constrain the impurity and local bath correlations (U and U_b respectively) through the relation $U_b = -U/10$. While the precise value of the factor of $1/10$ is unimportant, we have chosen a factor considerably smaller than 1 in order to demonstrate that an interesting body of results can be obtained with a value of $|U_b|$ that is much smaller than that of U . Further, the negative sign in the above relation is significant, as we shall see below that the MIT is obtained for the case of U_b being negative (i.e., attractive on-site correlations on the zeroth site of the bath). The relation between U_b and U is motivated on phenomenological grounds such that the MIT occurring in the bulk lattice model (obtained upon increasing the on-site Hubbard repulsion to large values) corresponds, in the auxiliary model mapping within DMFT, to the local MIT observed upon tuning the impurity correlation U (and the related bath correlation U_b) within the proposed effective impurity model Hamiltonian. The physical significance of attractive on-site correlations in the bath lies in providing a mechanism for the frustration of Kondo screening (in the RG equation for an antiferromagnetic Kondo coupling $J(> 0)$, eq. (9.28)) within an Anderson impurity coupled to a single channel of conduction electrons. We provide a detailed discussion of this point in the concluding section of our work.

Regime	Low-energy effective Hamiltonian	impurity ground-state
1. $0 < r < r_{c1}$	$\mathbf{K.E}^* + V^* \sum_{\sigma} \left(c_{d\sigma}^{\dagger} c_{0\sigma} + \text{h.c.} \right) + J^* \vec{S}_d \cdot \vec{S}_0$	$\frac{1}{\sqrt{2}} (\uparrow_d\rangle \downarrow_0\rangle - \downarrow_d\rangle \uparrow_0\rangle + 2_d\rangle 0_0\rangle + 0_d\rangle 2_0\rangle)$
2. $r_{c1} < r < r_{c2}$	$\mathbf{K.E}^* + J^* \vec{S}_d \cdot \vec{S}_0 - \frac{1}{2} U_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2$	$\frac{1}{\sqrt{2}} (\uparrow_d\rangle \downarrow_0\rangle - \downarrow_d\rangle \uparrow_0\rangle)$
3. $r_{c2} < r$	$\mathbf{K.E}^* - \frac{U^*}{2} (\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2 - \frac{U_b}{2} (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2$	$\{ \uparrow_d\rangle, \downarrow_d\rangle\}$

Table 3.1: Effective Hamiltonians and ground-states of the three important parts of the phase diagram in Fig. (3.2). The ground-state is simplified to capture only the configuration of the impurity site and at most the bath zeroth site. $\mathbf{K.E}^*$ represents the kinetic energy of the conduction electron states residing within the fixed point window D^* .

The phase diagram is shown in the left panel of Fig. (3.2) in terms of the parameter $r = |U_b|/J$ (y-axis) and the ratio of the bare Kondo coupling (J_0) to the bare conduction bath bandwidth (D_0) (x-axis), we first define two important points in the space of couplings. These are values of the parameter r where there is a qualitative change in the nature of RG flows, and hence in the low-energy physics of the model

- $r = r_{c1} \left(= -\left(\frac{U_b}{J}\right)_{c1} = \frac{3}{20} = \left(\frac{U}{10J}\right)_{c1} > 0 \right)$: At this point, the coupling V becomes irrelevant,
- $r = r_{c2} \left(= -\left(\frac{U_b}{J}\right)_{c2} = \frac{1}{4} = \left(\frac{U}{10J}\right)_{c2} \right)$: At this point, the coupling J also turns irrelevant. Note that $r_{c2} > r_{c1}$.

We will use these two values of the parameter r as checkpoints around which we can describe the low-energy physics. There are three important parts in Fig. (3.2):

- red region, $0 < r < r_{c1}$: the $J - V$ model; V and J are both relevant, but U is irrelevant; spin and charge delocalisation on the impurity; spin-charge mixing in ground-state
- blue region, $r_{c1} < r < r_{c2}$: the $J - U_b$ model; J is relevant, but V and U are both irrelevant; charge localisation and spin delocalisation on the impurity; singlet ground-state
- violet region, $r_{c2} < r$: the $U - U_b$ model; U is relevant, but V and J are both irrelevant; spin and charge localisation on the impurity; local moment ground-state. This phase describes a local Mott insulator on the impurity site, characterised by vanishing local double occupancy in ground state (black curve in left panel of Fig. 3.4).

Typical RG flows that lead to these phases are shown in Fig. (3.3). We also note that the grey region shown in the top right corner of Fig.(3.2) corresponds to a model in which all three couplings (J , U and V) are RG irrelevant. We find, however, that this phase is an artefact of solving the RG equations for an impurity coupled to a finite-sized conduction bath, and it gradually disappears upon increasing the bath size.




Figure 3.3: Variation of couplings U, V and J along the RG transformations, for three values of the transition-tuning ratio $r = -U_b/J_0$. The x-axis represents the distance of the running cutoff from the Fermi surface; the rightmost point is the first RG step (UV) and the leftmost point is the final RG step (IR). The red curves represent the flows for $r < r_{c1}$, where both V and J are relevant. The blue curves represent the RG flows for $r_{c1} < r < r_{c2}$, where J is relevant but V is irrelevant. The violet curves represent RG flows for $r > r_{c2}$, where both V and J are irrelevant but U flows to a finite value.

3.3.3 Phase transition at r_{c2}

The effective Hamiltonians and corresponding impurity ground-states for various phases have been listed in table (3.1). The variation in the ground state has been checked by numerically solving the fixed point Hamiltonian for various bare values of the couplings and is shown in the left panel of Fig. (3.4). A sharp change in the ground state from spin-singlet to local moment shows that the blue and violet phases are separated by an *impurity delocalisation-localisation transition* (black line in Fig. (3.2)). The transition occurs at finite values of the correlations: $r_{c2} = -(\frac{U_b}{J})_{c2} = \frac{1}{4} = (\frac{U}{10J})_{c2}$. This is a stark contrast from the standard SIAM, where the transition can happen only at on-site correlation $U \rightarrow \infty$. Therefore, the presence of the critical point at finite values of the various couplings transforms the landscape of RG phase diagram shown schematically in the right panel of Fig. (3.2) (right panel): the RG flows split into two classes - those that flow towards the strong-coupling Kondo screened fixed point and those that flow towards the local moment fixed point.

By a simple rewriting of the RG equation for J (eq. (9.28)), the impurity transition can be seen to arise from a competition between the Kondo screening physics of J and the local pairing physics of U_b :

$$\Delta J = \frac{\overbrace{(J + 2U_b)^2 n_j}^{\text{usual Kondo physics}}}{|d_2|} - \frac{\overbrace{(2U_b)^2 n_j}^{\text{competing pairing physics}}}{|d_2|}, \quad (3.4)$$

where $d_2 = \omega - D/2 + U_b/2 + J/4$. The competition between the effective Kondo term $(J + 2U_b)^2$ and the competing pairing term $-4U_b^2$ leads to the presence of two stable phases - one that is Kondo screened and one that remains unscreened.

In the DMFT treatment of the $1/2$ -filled Hubbard model on the Bethe lattice in infinite dimensions, the vanishing of non-local contributions to the lattice self-energy means that the lattice Greens function can be computed self-consistently by solving a local quantum impurity problem [Georges

et al.(1996)Georges, Kotliar, Krauth, and Rozenberg]. In the rest of the work, we provide extensive evidence that the local transition observed in the e-SIAM is very similar to the Mott MIT observed in DMFT. We conclude thereby that the e-SIAM models faithfully the round-trip excursions of an electron on the Bethe lattice, and that the impurity phase transition observed in the e-SIAM offers a local description of the Mott MIT in the bulk Hubbard model. With this evidence in mind, we will henceforth refer to the transition at r_{c2} as a *local metal-insulator transition*.

3.3.4 The case of repulsive U_b

Henceforth, we will only consider the case of attractive U_b ($U_b > 0$), because that is the regime in which an impurity phase transition is realised at finite values of the couplings. However, before moving on, we will comment briefly on the effects of a repulsive U_b ($U_b > 0$). From the RG equation for J in eq. 9.28, we see that ΔJ is always positive for $U_b > 0$. Moreover, something similar happens in the RG equation for V , where the two terms in the square brackets now enforce each other (J and U_b now have the same sign). Together, they indicate that there will not be any Kondo destruction in the case of $U_b > 0$, and the only fixed points are the weak-coupling one ($J^* = 0$) and the strong coupling one ($J^* = \infty$); there is no local moment phase. This case is therefore adiabatically connected to the case of $U_b = 0$.

3.4 Descriptors of the local MIT

The present section gives more clarity on the nature of the impurity phase transition in the form of additional descriptors of the transition such as impurity spectral function and measures of entanglement, computed from the e-SIAM Hamiltonian. Throughout the rest of the work, we have set the value of the coupling V equal to the Kondo coupling J while generating the quantitative plots.

3.4.1 Evolution of the impurity spectral function

The impurity spectral function of the standard SIAM (eq. (2.1)) always displays a central peak at finite values of U (along with Hubbard sidebands at sufficiently large U), indicating the presence of gapless local Fermi liquid excitations on the impurity site [Wilson(1975b), Nozieres(1974), Costi and Hewson(1990)]. To demonstrate the impurity localisation transition, we compute the impurity local spectral function of the e-SIAM (eq. (2.2)). This involves numerically diagonalising the effective Hamiltonian at various energy scales along the RG flow and computing the spectral weight at a range of frequencies from UV to IR. We find that the spectral function (shown in the right panel of Fig. (3.4)) displays three notable features, in agreement with DMFT results [Georges et al.(1996)Georges, Kotliar, Krauth, and Rozenberg]:

- The appearance of a *preformed gap* (flattening of spectral function between the central peak and sidebands) as r crosses r_{c1} ($= -(\frac{U_b}{J})_{c1} = \frac{3}{20} = (\frac{U}{10J})_{c1}$): The preformed optical gap simply indicates the separation of the spin (central peak) and charge (sidebands) degrees of freedom beyond r_{c1} . This separation is brought about by the irrelevance of V . We note that

the consequences of an irrelevant hybridisation V can be accounted for in many-body perturbation theory, leading to “in-gap” processes within the optical gap of the impurity spectral function [Nozières, Ph.(1998)]. However, we expect that such contributions will be small, and cannot fill in the optical gap observed in Fig.3.4 completely.

- The sharpening of the central peak, and concomitant increase in the preformed gap, as r is increased in the range $r_{c1} < r < r_{c2}$.
- The vanishing of the central peak and appearance of a *hard gap* as r crosses r_{c2} : The irrelevance of J beyond r_{c2} destroys the Kondo screening and localises the impurity moment. This involves the destabilisation of the singlet ground state, and the stabilisation of the local moment states in its place.




Figure 3.4: Left: Overlap of the RG fixed point ground state $|\Psi\rangle_{gs}$ with the spin-singlet state $|SS\rangle$, the zero charge member of the charge triplet states $|CT\rangle$ and the local moment states $|LM\rangle$, for all three regimes of the model. The black curve corresponds to the double occupancy on the impurity site and is observed to be strongly suppressed beyond r_{c1} . Beyond the critical point $r_{c2} = 0.25$, the overlaps with the entangled states vanish, indicating the transition to a decoupled local moment. Right: Variation of the impurity spectral function from $r = 0$ to $r > r_{c2}$. At small r , the central peak is broad, but at larger r , it sharpens, and the difference in spectral weight is used in creating the Hubbard sidebands. For $r > r_{c2}$, the central peak vanishes.

These features are also reflected in ground-state correlation measures like spin-flip and charge isospin-flip correlations (see left panel of Fig. (3.5)), which are defined as follows:

$$\frac{1}{2} (\langle S_i^+ S_j^- \rangle + \text{h.c.}) = \frac{1}{2} (\langle c_{i\uparrow}^\dagger c_{i\downarrow} c_{j\downarrow}^\dagger c_{j\uparrow} \rangle + \text{h.c.}) , \quad \frac{1}{2} (\langle C_i^+ C_j^- \rangle + \text{h.c.}) = \frac{1}{2} (\langle c_{i\uparrow}^\dagger c_{i\downarrow}^\dagger c_{j\downarrow} c_{j\uparrow} \rangle + \text{h.c.}) \quad (4.1)$$

At small values of r , both the correlations are large as the ground-state has both spin and charge content (row 1 of table (3.1)). The subsequent decrease in the impurity-bath charge flip correlation (violet points of Fig. (3.5)), and the simultaneous increase in the impurity-bath spin-flip correlation (red points of Fig. (3.5)), can be attributed to the irrelevance of V at $r \simeq r_{c1}$ and the appearance of the preformed gap in the spectral function. Beyond r_{c2} , the spin-flip correlation sharply drops to zero due to the irrelevance of J . They are replaced by intra-bath correlations like the charge

isospin-flip correlation between the zeroth site and the first site (blue points of Fig. (3.5)). Such intra-bath correlations are promoted by the bath on-site term U_b . The sudden change in the nature of the ground state, and resulting correlations, at r_{c2} indicates the presence of a quantum critical point whose nature will be analysed further in subsequent sections.




Figure 3.5: Left: Variation of impurity-bath spin-flip correlation (red) and charge isospin-flip correlation (violet), as well as intra-bath charge isospin-flip correlation (blue), from $r \sim 0$ to $r > r_{c2}$. The imp.-bath isospin correlation vanishes at r_{c1} , indicating the change in RG relevance of V . The imp.-bath spin correlation vanishes at r_{c2} , indicating the marginality of J at that point. The intra-bath correlation picks up after the transition, showing the decoupling of the impurity from the bath. Right: Variation of the geometric entanglement ϵ_{ss} (blue) w.r.t. the singlet state and that w.r.t the charge triplet zero state, ϵ_{ct} (red), with r . The former becomes maximum (unity) at r_{c1} , showing that the true ground state has no charge content. The latter discontinuously jumps to unity at r_{c2} , therefore acting as an order parameter for the transition.

3.4.2 Using entanglement to track correlations across the transition

We will now show that a certain measure of entanglement behaves as an order parameter for the transition. We can define a geometric measure of entanglement in terms of wavefunctions $|\psi_1\rangle$ and $|\psi_2\rangle$ [Shimony(1995), Wei and Goldbart(2003), Horodecki et al.(2009)Horodecki, Horodecki, Horodecki, and Horodecki]:

$$\epsilon(\psi_1, \psi_2) = 1 - |\langle \psi_1 | \psi_2 \rangle|^2. \quad (4.2)$$

From this definition, if $|\psi_1\rangle$ corresponds to a separable state, the entanglement content of the state $|\psi_2\rangle$ is small if it's overlap with $|\psi_1\rangle$ is large. For brevity, we will use the notation $\epsilon_{ss} \equiv \epsilon(\psi_{ss}, \psi_{gs}^{(2)})$, $\epsilon_{ct} \equiv \epsilon(\psi_{ct}, \psi_{gs}^{(2)})$ to represent the geometric entanglement between the e-SIAM ground-state $|\psi_{gs}\rangle$ and the singlet state $|\psi_{ss}\rangle$ or the charge triplet zero state $|\psi_{ct}\rangle$. The latter two states are shown in Table (3.1). The entanglement measures ϵ_{ss} and ϵ_{ct} can be related to the impurity Greens function through the following equation (details can be found in Sec. 3 of the Supplementary

Materials [sup(Supplementary Materials are available online)]:

$$G_d(\omega) = \sum_n \left[(1 - \varepsilon_{ss}) G_{\Phi_{ss}, \Phi_{ss}}(\omega, n) + (1 - \varepsilon_{ct}) G_{\Phi_{ct}, \Phi_{ct}}(\omega, n) + 2\sqrt{(1 - \varepsilon_{ss})}\sqrt{(1 - \varepsilon_{ct})} G_{\Phi_{ss}, \Phi_{ct}}(\omega, n) \right], \quad (4.3)$$

where

$$G_{\psi_1, \psi_2}(\omega, n) = \frac{1}{2} \frac{\langle \psi_1 | c_{d\sigma} | \Psi_n \rangle \langle \Psi_n | c_{d\sigma}^\dagger | \psi_2 \rangle + \text{h.c.}}{\omega + E_{gs} - E_n} + \frac{1}{2} \frac{\langle \psi_1 | c_{d\sigma}^\dagger | \Psi_n \rangle \langle \Psi_n | c_{d\sigma} | \psi_2 \rangle + \text{h.c.}}{\omega - E_{gs} + E_n}. \quad (4.4)$$

This expression displays that the evolution of the impurity Greens function G_d with r (and related correlation functions shown in the left panel of Fig.(3.5)) is dependent on that of the entanglement measures ε_{ss} and ε_{ct} (shown in the right panel of Fig. (3.5)). Indeed, we find that the geometric entanglement ε_{ct} with the charge sector increases towards unity as the transition at r_{c2} is approached, and remains unity after the transition. A more sensitive measure is the singlet entanglement ε_{ss} : it initially decreases to zero at r_{c1} owing to the irrelevance of V , but rises discontinuously to unity at the transition and becomes equal to the charge entanglement in the local moment phase. Therefore, ε_{ss} acts as an order parameter for the local MIT at r_{c2} . We find that the cross-term $\sqrt{(1 - \varepsilon_{ss})}\sqrt{(1 - \varepsilon_{ct})}$ decreases monotonically to zero with increasing r (not shown), displaying a continual decrease in the mixing of the spin and charge sectors and the *immobilisation of the doublons and holons on the impurity site*.

In general, any one-particle or two-particle fluctuation $\langle O_1 O_2^\dagger \rangle$ that acts on the combined Hilbert space of the impurity and the zeroth site can be expressed in terms of these entanglement measures $\varepsilon_{ss}, \varepsilon_{ct}$. The detailed derivations and expressions are given in Section 3 of the Supplementary Materials [sup(Supplementary Materials are available online)]. As a demonstration, consider the spin-spin correlation $\langle S_d^+ S_0^- \rangle$ between the impurity and the zeroth site shown in the left panel of Fig. (3.5). The general expression given in Section 3 of the Supplementary Materials [sup(Supplementary Materials are available online)] is of the form

$$\langle S_d^+ S_0^- \rangle = (1 - \varepsilon_{ss}) \langle \Phi_{ss} | S_d^+ S_0^- | \Phi_{ss} \rangle + (1 - \varepsilon_{ct}) \langle \Phi_{ct} | S_d^+ S_0^- | \Phi_{ct} \rangle + \sqrt{1 - \varepsilon_{ss}}\sqrt{1 - \varepsilon_{ct}} (\langle \Phi_{ss} | S_d^+ S_0^- | \Phi_{ct} \rangle + \langle \Phi_{ct} | S_d^+ S_0^- | \Phi_{ss} \rangle). \quad (4.5)$$

From the expression given, we find that only the singlet overlap is non-zero, such that $\langle S_d^+ S_0^- \rangle$ is directly proportional to the quantity $1 - \varepsilon_{ss}$. As the entanglement measure ε_{ss} increases towards the transition (right panel of Fig. (3.5)), the quantity $1 - \varepsilon_{ss}$ decreases, in turn leading to the decrease in the magnitude of the correlation observed in the left panel of Fig. (3.5). For another explicit connection between correlations and entanglement measures, we present relations between the quantum Fisher information (QFI) [Hauke et al.(2016)Hauke, Heyl, Tagliacozzo, and Zoller] and many-particle Greens functions in Section 3 of the Supplementary Materials [sup(Supplementary Materials are available online)]. There, we also plot the QFI for a number of two-particle operators as a function of r , notably the ones corresponding to the degree of compensation for the impurity ($\langle \vec{S}_d \cdot \vec{S}_0 \rangle$) and the impurity magnetisation ($\langle S_d^z \rangle$). These two quantities (and hence the corresponding QFI) are important because they track the local MIT and act as order parameters for the transition, and the QFI corresponding to these two operators quantify the quantum fluctuations present

in the system corresponding to these order parameters. We show in Sec. 3 of the Supplementary Materials that the two phases on either side of the transition are characterised by distinct values of this pair of QFI: while the QFI corresponding to the degree of compensation is zero in the Kondo screened phase, it becomes non-zero in the local moment phase, and the opposite is true for the QFI arising from the impurity magnetisation. The phase precisely at the transition is distinct from those on either side because it displays a non-zero value for both of the QFI. While it is expected that a critical point would show enhanced fluctuations of multiple kinds (giving rise to universality), it is enlightening to find that this is also reflected in a measure of many-particle entanglement.

3.5 Universal theory for the local metal-insulator transition

In order to identify the competing tendencies near the transition at r_{c2} , we will now obtain the minimal effective Hamiltonian that displays the same transition. This involves integrating out the degrees of freedom that do not affect the low-energy physics near r_{c2} . We note that, due to the irrelevance of V , there is no scattering between the spin and charge states on the impurity at low energies. Further, the impurity charge states $|0\rangle$ and $|\uparrow_d\downarrow_d\rangle$ have been pushed to the high energy Hubbard sidebands because of the large value of U close to r_{c2} . As a result, the impurity charge states can be safely decoupled from the spin states through a Schrieffer-Wolff transformation. Up to second order in V^2/U , this transformation accounts for the effects of U and V by generating an additional (Kondo) spin-exchange term $\delta J \left(\sim \frac{V^2}{U+U_b} \right)$, as well as an additional on-site correlation $\delta U_b \left(\sim \frac{2V^2}{U+U_b+J/2} - \frac{8V^2}{U-U_b} \right)$ on the zeroth site of the conduction bath. In this way, we obtain the following renormalised effective Hamiltonian:

$$H_{\text{MIT}} = \mathcal{J} \vec{S}_d \cdot \vec{S}_0 - \frac{1}{2} \mathcal{U}_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2 + H_{\text{K.E.}}, \quad (5.1)$$

where $\mathcal{J} = J + \delta J$ is the renormalised s-d interaction and $\mathcal{U}_b = U_b + \delta U_b$ is the renormalised local correlation on the bath zeroth site. A schematic 1D construction of the effective Hamiltonian is shown in the left panel of Fig. (3.6). We note that a similar approach towards extracting an effective theory for the Mott MIT has been employed in the past ([Moeller et al.(1995)Moeller, Si, Kotliar, Rozenberg, and Fisher, Held et al.(2013)Held, Peters, and Toschi]). However, those works essentially led to renormalised Kondo models that do not possess any frustration of the Kondo screening. Thus, they cannot display an impurity transition in the absence of the requirement of self-consistency and can describe only the physics of the metallic regime. Importantly, the effective $\mathcal{J}-\mathcal{U}_b$ model (eq. (5.1)) is consistent with the IR fixed point Hamiltonian obtained in the appropriate regime $r_{c1} < r < r_{c2}$ (row 2 of table (3.1)).

The RG equations for $\mathcal{J}$ and $\mathcal{U}_b$ of the simplified effective impurity model eq. (5.1) can be obtained by setting $U = V = 0$ in the RG equations (9.28):

$$\Delta \mathcal{J} = -\frac{n_j \mathcal{J} (\mathcal{J} + 4\mathcal{U}_b)}{d_2}, \quad \Delta \mathcal{U}_b = 0. \quad (5.2)$$

For $\mathcal{J} + 4\mathcal{U}_b > 0$, the Kondo coupling is relevant and the low-energy phase is a paramagnetic local Fermi liquid with gapless excitations. But for $\mathcal{J} + 4\mathcal{U}_b < 0$, the Kondo coupling becomes

irrelevant and the ground state is a decoupled local moment that is isolated from the bath. Such an effective picture of the transition can be understood if one notes that a straightforward way to destroy the Kondo screening is to inhibit the coordinated spin fluctuations between the impurity and the bath. Such frustration of Kondo screening is precisely the effect of the $\mathcal{U}_b$ term: it promotes charge fluctuations on the bath site coupled directly to the impurity, reducing thereby the spectral weight for spin-flip scatterings between the impurity and bath.



Figure 3.6: Left: Schematic 1D construction of universal theory for the metal-insulator transition, obtained by integrating out the charge states of the impurity near the transition. Right: Zero bandwidth limit of the low-energy effective Hamiltonian for r between r_{c1} and r_{c2} (second row in table (3.1)). The bath is reduced to just a single degree of freedom - the zeroth site that is directly coupled to the impurity site.

Indeed, the local correlation $\mathcal{U}_b$ encourages entanglement between the sites of the bath and makes the formation of the impurity-bath singlet difficult. In other words, the $\mathcal{J} - \mathcal{U}_b$ model displays the destabilisation of the singlet by redistributing the entanglement from the impurity+bath system to purely within the bath (see left panel of Fig. (3.7)). Given the simplicity of these arguments, our analysis makes the case that local pairing fluctuations of the bath in eq. (5.1) offer a universal mechanism by which to frustrate the Kondo effect and lead thereby to an impurity transition that is the local counterpart of the Mott MIT obtained by auxiliary model approaches such as DMFT. We note that similar conclusions were reached in Refs. [Si and Kotliar(1993a), Kotliar and Si(1993a), Kotliar and Si(1996)] for the emergence of non-Fermi liquid phases at critical points in the mixed valence regime of the periodic Anderson model.

Further insight into the destabilisation of the singlet can be obtained from a zero-bandwidth approximation of the bath. Under such an approximation, the IR effective Hamiltonian obtained from the RG flow (row 2 of Fig. (3.1)) takes the form of a two-site model with modified couplings (shown in the right panel of Fig. (3.6)):

$$\tilde{\mathcal{J}} \vec{S}_d \cdot \vec{S}_0 - \frac{1}{2} \tilde{\mathcal{U}}_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2. \quad (5.3)$$

The spectrum of this model is shown in the right panel of Fig. (3.7). For $|\tilde{\mathcal{U}}_b|/\tilde{\mathcal{J}} < 3/2$, the singlet ground-state (red) is separated from the excited local moment states (blue) by a gap of $\frac{-3\tilde{\mathcal{J}}}{4} - \frac{\tilde{\mathcal{U}}_b}{2}$. As we tune r towards r_{c2} , $|\tilde{\mathcal{U}}_b|$ increases and the fixed point value $\tilde{\mathcal{J}}$ decreases, leading to an overall reduction in the gap. At the critical point, the gap closes and the states become degenerate at zero energy; this is the equivalent of the quantum critical point at r_{c2} of the full impurity model that was discussed earlier. We note that the vanishing of the energy of the metallic state was also observed by Brinkman and Rice from a Gutzwiller-type variational calculation of the Hubbard model at half-filling [Brinkman and Rice(1970)].




Figure 3.7: Left: Very close to the transition, the mutual information $I_2(d : 0)$ (red curve) between impurity and the zeroth site reduces, while that between the zeroth and the first site ($I_2(0 : 1)$, blue curve) increases, showing the redistribution of entanglement. Right: Spectrum of the zero bandwidth Hamiltonian of the right panel of Fig. (3.6). The blue line represents the local moment states at zero energy, while the red line represents the singlet state at an energy that continuously increases upon increasing r .

3.5.1 Local pairing and the destruction of Kondo screening

Very close to the transition at r_{c2} , we find signatures of the breakdown of the Kondo cloud in terms of decaying impurity-zeroth site correlations (blue curve in the left panel of Fig. (3.8)) and enhanced intra-bath correlations. Remarkably, we find an *increase in pairing correlations* between the bath zeroth site and first site (red curve in the left panel of Fig. (3.8)). As discussed above, these are observed to be responsible for the destruction of the Kondo screening: spin and charge degrees of freedom are mutually exclusive, and the increased charge fluctuations on the bath zeroth site suppress the spin-flip scattering processes between the impurity and the zeroth site. We will show later that these fluctuations lead to the destruction of the local Fermi liquid excitations in our model, and replace them with non-Fermi local excitations in the neighbourhood of r_{c2} . We believe that the observed growth in non-local pairing correlations near the transition at r_{c2} is likely tied to a putative low-energy divergence of the local pairing susceptibility. A similar observation was made in Ref. [Si and Kotliar(1993a)] from an auxiliary model-based analysis of a hole-doped extended Hubbard model in infinite dimensions that is pertinent to the physics of the heavy fermions. Similar to the conclusions of Ref. [Si and Kotliar(1993a)], we expect the pairing fluctuations of the bath to become dominant upon tuning the e-SIAM away from half-filling, signalling an instability towards a superconducting state.

3.5.2 Emergent self-consistency in the e-SIAM

As is well-established, the DMFT self-consistency equation is equivalent to requiring that the impurity Greens function become equal to the local Greens function in the bath [Georges et al.(1996)Georges, Kotliar, Krauth, and Rozenberg]. Such a condition is also used in the projective self-consistent technique of Moeller et al. [Moeller et al.(1995)Moeller, Si, Kotliar, Rozenberg, and Fisher] for the states within the central Kondo resonance. We will now show that our model displays a qualitatively similar emergent feature. To proceed, by employing a one-step URG transformation, we




Figure 3.8: Left: Spin-flip correlation between the impurity and the bath (blue), and charge isospin-flip correlation between the bath zeroth and first sites (red), both as a function of the tuning parameter r , very close to the transition. The operators are defined as $S_i^+ = c_{i\uparrow}^\dagger c_{i\downarrow}$ and $C_i^+ = c_{i\uparrow}^\dagger c_{i\downarrow}^\dagger$. Right: New effective SIAM Hamiltonian generated upon integrating out the coupling between the impurity and the bath zeroth site. In the new SIAM, the zeroth site acts as the new impurity site with a renormalised on-site correlation $U_Z^* = U_b + \delta U_Z^*$.

integrate out the impurity site from the rest of the fixed-point Hamiltonian of eq. (3.4). The details are shown in Sec. 4 of the Supplementary Materials [sup(Supplementary Materials are available online)]. The essential idea is similar to that of the Schrieffer-Wolff transformation: removing the impurity-bath couplings J and V generates an additional repulsive correlation δU_Z^* on the zeroth site. The new low-energy model H_Z is therefore an Anderson impurity model with a net local correlation $U_Z^* = \delta U_Z^* + U_b$ on the zeroth site (which is now the “new impurity site”), and a single-particle hybridisation $V_Z^* = -t$ coupling with a conduction bath formed by the remaining sites. This new impurity model is depicted schematically in the right panel of Fig. (3.8).

$$H_Z^* = \underbrace{-\frac{1}{2}U_Z^* (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2}_{\text{new impurity} = 0^{\text{th}} \text{ site}} + \underbrace{V_Z^* \sum_{\langle j,0 \rangle} (c_{0\sigma}^\dagger c_{j\sigma} + \text{h.c.})}_{\text{hopping between } 0^{\text{th}} \text{ site \& new bath}} + \underbrace{(-t) \sum_{\langle i,j \rangle} (c_{i\sigma}^\dagger c_{j\sigma} + \text{h.c.})}_{\text{K.E. of new bath}}. \quad (5.4)$$

Since, for $r < r_{c2}$, the model always flows to strong coupling at low energies, the largest energy scales are J^* and V^* . Using this, the effective correlation U_Z^* on the zeroth site can be expressed to leading order as

$$U_Z^* \simeq \frac{J^*}{4} \frac{1}{1 + 2\gamma} - V^* \frac{\gamma}{\gamma^2 - \frac{1}{4}}, \quad \text{where } \gamma \equiv V^*/J^*. \quad (5.5)$$

From $r = 0$ to $r = r_{c1}$, the factor γ decreases due to the gradual removal of single-particle hopping from the impurity site (irrelevance of V), leading to an increase in the correlation U_Z^* . This is simply a restatement of the fact that the scattering processes that create the central impurity resonance induce a repulsive correlation on the bath zeroth site. The fact that we end up with a standard SIAM on the zeroth site once the impurity site has been integrated out means that the spectral function of this new impurity will again go through a sharpening of the central peak (and the appearance of the Hubbard sidebands) upon increasing the parameter r of the original e-SIAM. This ensures

that the impurity and zeroth site spectral functions look qualitatively similar up to r_{c1} . One can now repeat iteratively this process - decoupling the zeroth site generates a standard SIAM with a repulsive correlation on the first site, and so on. The fact that excursions starting from any point along the bath can be described by a positive U SIAM is, therefore, the emergent self-consistency in our model. Similar indications of a correlated spectral function on lattice sites far away from the impurity were also observed in Ref. [Ribeiro et al.(2019)Ribeiro, Martins, Gomez-Silva, and Anda] from finite- U slave boson calculations of the SIAM. Beyond r_{c1} , the irrelevance of V means that U_Z^* reduces to just $J^*/4$. As r is now increased towards r_{c2} , the correlation decreases because J is moving towards its critical point, indicating that the bath zeroth site is moving away from its local moment regime. This is another reflection of the increase in pairing fluctuations of the bath, as well as the lowering of spin-flip fluctuations between the impurity and the bath.

For $r > r_{c2}$, the impurity site decouples from the bath in the impurity model. This impurity model can be promoted to a bulk model as follows. Recall that in the auxiliary model mapping, any lattice site $\vec{r}_i$ of the bulk lattice can act as the impurity, and one can think of the impurity model with the impurity at $\vec{r}_i$ as a representation of the excursion of an electron from any such site $\vec{r}_i$ into the rest of the bath. When all such impurity models undergo the transition at $r = r_{c2}$ simultaneously, the result is the decoupling of all sites from their respective baths and a paramagnetic bulk insulator is obtained.

3.6 Coexistence of local metallic and insulating phases in the e-SIAM

The DMFT solution of the Hubbard model on the Bethe lattice exhibits a coexistence region of metallic and insulating solutions between two spinodal lines $U_{c1}(T)$ and $U_{c2}(T)$ [Georges et al.(1996)Georges, Kotliar, Krauth, and Rozenberg], with the metallic solution having lower internal energy and the insulating solution being a metastable state at a higher energy [Moeller et al.(1995)Moeller, Si, Kotliar, Rozenberg, and Fisher, Georges et al.(1996)Georges, Kotliar, Krauth, and Rozenberg, Bulla(1999), Georges(2004), Georges and Krauth(1993)]. The $T = 0$ transition is observed to be second order in nature and occurs through a merging of the metallic and insulating solutions at U_{c2} [Rozenberg et al.(1994)Rozenberg, Moeller, and Kotliar, Moeller et al.(1995)Moeller, Si, Kotliar, Rozenberg, and Fisher]. On the other hand, at $T > 0$, the MIT happens along the first-order line $U_c(T)$ ($U_{c1} < U_c < U_{c2}$) where the free energies of the two solutions become equal, and the two spinodal lines merge into a second order critical point at a sufficiently high temperature T_c . We will now show that our Hamiltonian-based approach gives a clear zero temperature picture of various aspects that can lead to the emergence of the metal-insulator coexistence at $T > 0$.

3.6.1 Excited state quantum phase transition at r_{c1} : the Mott-Hubbard scenario

We have already discussed in detail the nature of the phase transition at r_{c2} : the zero-bandwidth picture provided (around eq. (5.3)) shows the merging of the metallic and insulating solutions at

that point, enabling identification of r_{c2} with the $T = 0$ continuous phase transition observed at U_{c2} in the DMFT phase diagram. The focus of this subsection is the physics of the other important point r_{c1} . As has been mentioned before, this point marks the RG irrelevance of the single-particle hybridisation amplitude V , and leads to the exclusion of the charge states from the ground-state (see left panel of Fig. (3.4)). This exclusion means that there is now one fewer scattering channel by which the impurity electron can hybridise with the bath. In turn, this leads to a *partial localisation of the impurity*, and can be thought of as the first step towards the more complete localisation that occurs at r_{c2} .

Apart from the change in the ground state, the physics at r_{c1} also involves a phase transition in certain excited states of the spectrum. To expose this, we consider the following states in the zero-bandwidth spectrum of the impurity model given in eq. (2.2):

$$\begin{aligned}
 |1, \sigma, \pm\rangle &= \alpha_{\pm} |\sigma_d\rangle |0_0\rangle \mp \sqrt{1 - \alpha_{\pm}^2} |0_d\rangle |\sigma_0\rangle, \quad |3, \sigma, \pm\rangle = \alpha_{\pm} |\sigma_d\rangle |2_0\rangle \mp \sqrt{1 - \alpha_{\pm}^2} |2_d\rangle |\sigma_0\rangle, \\
 E_+ &= -\frac{U_0}{4} + \sqrt{V_0^2 + \frac{U_0^2}{16}}, \quad E_- = -\frac{U_{\text{im}}}{4} - \sqrt{V_{\text{im}}^2 + \frac{U_{\text{im}}^2}{16}}, \\
 \alpha_+(U_0, V_0) &= \frac{V_0}{\sqrt{V_0^2 + (E_+ + \frac{U_0}{2})^2}}, \quad \alpha_- = \frac{E_- + \frac{U_{\text{im}}}{2}}{\sqrt{V_{\text{im}}^2 + (E_- + \frac{U_{\text{im}}}{2})^2}}.
 \end{aligned} \tag{6.1}$$

where $(\sigma = \uparrow, \downarrow)$ and $E_{\pm}$ is the energy of the states $|1(3), \sigma, \pm\rangle$ and the subscripts d and 0 refer to the impurity and bath zeroth site respectively. The states $|1(3), \sigma, +\rangle$ are high-energy states, so their energy E_+ and coefficient α_+ involve the bare single-particle hybridisation V_0 and impurity on-site repulsion U_0 . On the other hand, the other states $|1(3), \sigma, -\rangle$ are closer to the IR energy scale and involve renormalised intermediate-scale couplings V_{im} and U_{im} . Both E_+ and E_- are four-fold degenerate because of the SU(2) spin ($\uparrow \leftrightarrow \downarrow$) and particle-hole ($|0\rangle \leftrightarrow |2\rangle$) symmetries. For example, the degenerate subspace corresponding to E_+ is the set of states $\{|1, \uparrow, +\rangle, |1, \downarrow, +\rangle, |3, \uparrow, +\rangle, |3, \downarrow, +\rangle\}$.

All these states are in general delocalised - they involve the impurity hybridising with the bath via V . At $r = r_{c1}$, however, the coupling V becomes irrelevant and $V_{\text{im}} \rightarrow 0$, so that the coefficient α_- of the low-energy state becomes unity beyond that point. The low-energy states $|1(3), \sigma, -\rangle$ thus become localised at r_{c1} and give rise to a set of excited and *degenerate local moment states*, i.e., $\alpha_- \rightarrow 0$ as $r \rightarrow r_{c1}$ leads to

$$|1(3), \sigma, -\rangle \rightarrow \begin{cases} |\uparrow_d\rangle |0_0\rangle, |\downarrow_d\rangle |0_0\rangle \\ |\uparrow_d\rangle |2_0\rangle, |\downarrow_d\rangle |2_0\rangle \end{cases}. \tag{6.2}$$

The other coefficient α_+ remains non-zero because, as mentioned earlier, V is non-zero in the UV scales of the RG flow. As a result, the high-energy states $|1(3), \sigma, +\rangle$ remain delocalised, and are separated from the localised states by an energy-scale

$$\lim_{r \rightarrow r_{c1}^-} (E_+ - E_-) \simeq \sqrt{V_0^2 + \left(\frac{U_0}{4}\right)^2} + \sqrt{V_{\text{im}}^2 + \left(\frac{U_{\text{im}}}{4}\right)^2} + \frac{U_{\text{im}} - U_0}{4}. \tag{6.3}$$

This is shown schematically in the left panel of Fig. (3.9), and corresponds to the preformed gap in the zero mode spectrum. The point r_{c1} therefore represents a *delocalisation-localisation excited*

state quantum phase transition (ESQPT) where degenerate local moment states are emergent as excited states in the many-body spectrum. This localisation of the impurity at r_{c1} is shown in the form of excited state mutual information in the right panel of Fig. (3.9). This ESQPT acts as a precursor to the QPT at r_{c2} , where the local moment states become degenerate with the spin-singlet ground state. The local moment states are stabilised as ground states in the insulating phase for $r > r_{c2}$.




Figure 3.9: Left: Evolution of the excited states mentioned in Eqs. (6.1) as r is tuned through r_{c1} . The high-energy states (red) retain their impurity-bath coupling, while their low-energy counterparts (blue) become disentangled and form a degenerate set of local moment states. The high-energy states allow delocalisation into the bath and lead to the broad Hubbard sidebands. Right: Variation of mutual information (MI) between various parties, across the ESQPT at r_{c1} . The MI (red) between the impurity and the zeroth site vanishes at r_{c1} , showing the localisation of the impurity. The same between the bath zeroth site and the next site becomes maximum at r_{c1} . The MI between the impurity and the other sites (purple and gray) of the bath show an initial rise near r_{c1} , indicating that entanglement is becoming long-ranged near the ESQPT.

3.6.2 Theory for the charge excitations in the Hubbard sidebands

Beyond r_{c1} , excitations into the local moment states $|1, \sigma, -\rangle$ and $|3, \sigma, -\rangle$ reside at the edge of the central peak in the impurity spectral function (purple curve in the right panel of Fig.(3.4)) but provide no spectral weight due to the lack of electron mobility. This explains the development of a preformed gap in the impurity spectral function. That these local moment states have to reside at the edge of the central peak becomes clear when we note that as the width of the central peak shrinks continuously, the local moment states must also recede towards zero frequency and finally replace the zero frequency peak at $r = r_{c2}$ in order to give rise to the insulating local moment phase for $r > r_{c2}$. On the other hand, the still-delocalised high-energy states $|1, \sigma, +\rangle$ and $|3, \sigma, +\rangle$ are pushed into the Hubbard sidebands, and their hybridisation with the bath through V and t is responsible for the broadening of the sidebands.

This isolation of the delocalised states into the sidebands means that charge delocalisation processes are now excluded from the physics at low energies. As accessing the sidebands involves excitations at exorbitantly high energy scales, such processes can only happen virtually and involve

very short time scales. The central Kondo resonance observed at low energies, therefore, does not support any charge delocalisation, and metallic excitations propagate only through spin-flip scattering processes of the impurity. This reveals that the Mott insulator comes about through a local binding of doublons and holons [Mott(1949), Kohn(1964), Castellani et al.(1979) Castellani, Castro, Feinberg, and Ranninger]. Closely related to this is the Mott-Hubbard scenario of the MIT, which is equivalent to our observation of the appearance of an optical gap in the spectrum after r_{c1} .

The fact that the sidebands are broad indicates that there are gapless excitations propagating from the impurity site into the conduction bath, but whose energy lies outside the Mott gap. These can also be viewed as metallic excitations of the bath that are induced by the coupled impurity. In order to expose the nature of these excitations, we perform a calculation similar to the local Fermi liquid calculation of Nozières [Nozieres(1974)]. This involves considering the states at $\omega \sim \pm U/2$ as the ground-states of the Hubbard sidebands, and then computing how excitations into the bath renormalise the ground-state subspace. Up to the second order in the hopping strength t , the effective Hamiltonian for the excitations of the bath can be expressed as

$$H_{\text{eff}}^{(2)} = \frac{4t^2(1 - \alpha_+^2)}{E_+ - E_{\text{gs}}} C_{\text{tot}}^z C_1^z - t \sum_{i>0, \sigma} \left(c_{i\sigma}^\dagger c_{i+1\sigma} + \text{h.c.} \right), \quad (6.4)$$

where E_+ and α_+ have been defined in eq. (6.1), $C_i^z |2(0)\rangle = +(-)\frac{1}{2} |2(0)\rangle$ is the z-component of the charge isospin operator at a particular site i , and $\vec{C}_{\text{tot}} = \vec{C}_d + \vec{C}_0$ is the total isospin for the impurity and zeroth sites. $E_{\text{gs}} = U/2$ is the two-site energy of the ground-state subspace within the sideband. These excitations are of the local Fermi liquid kind - the absence of any isospin-flip scattering term promotes the independent delocalisation of the doublon and holon states into the bath. The second-order effective Hamiltonian can therefore be written as the sum of two decoupled parts corresponding to holon and doublon propagation respectively, and suggesting adiabatic continuity with the $U = 0$ Hubbard model [Krishnamurthy et al.(1990) Krishnamurthy, Jayaprakash, Sarker, and Wenzel].

However, fourth-order corrections lead to the appearance of scattering processes that convert holon and doublons into one another, resulting in the local Fermi liquid becoming correlated

$$H_{\text{eff}}^{(4)} = \frac{\gamma^4 \alpha_+^2 \beta^2}{(1 - \alpha_+^2)(E'_+ - E_{\text{gs}})} \left[-C_{\text{tot}}^z C_{\text{tot}}^2 C_1^z + \sqrt{2} \mathcal{P}_{\text{tot}}^4 (C_0^+ - C_d^+) C_1^- + \text{h.c.} \right], \quad (6.5)$$

where $E'_+ = \frac{U}{4} - \frac{3J}{8} + \sqrt{4V^2 + \left(\frac{U}{4} + \frac{3J}{8}\right)^2}$ is the energy of the state containing the charge isospin triplet zero, and $\beta = \frac{2V}{\sqrt{4V^2 + (U + 3J/4)^2}}$. $\mathcal{P}_{\text{tot}}^4$ projects on to the $n_d + n_0 = 4$ subspace. The scattering processes in eq. (6.5) are clearly non-Fermi liquid in nature as they allow the inter-conversion of the charge isospin eigenstates, and therefore reduce the lifetime of the quasiparticle excitations of the local Fermi liquid obtained in eq. (6.5). Additional details pertaining to the calculation of this effective Hamiltonian are present in Sec. 5 of the Supplementary Materials [sup(Supplementary Materials are available online)].

3.6.3 Hysteresis and the first-order line

As discussed above, the localisation of charge in the emergent local moment states $|1, \sigma, -\rangle$ and $|3, \sigma, -\rangle$ for $r > r_{c1}$ means they do not contribute any spectral weight to the impurity spectral function. Instead, they lead to an emergent preformed/optical gap between the central Kondo resonance and the Hubbard sidebands on each side. This *optical gap* finally becomes the true Mott gap at $r = r_{c2}$ when the central peak disappears. On the other hand, upon starting from $r > r_{c2}$ and reducing r , the system is initially in the (doubly degenerate) local moment ground states, and the impurity DOS is of course gapped. In the rest of the subsection, we will show that the presence of these two transitions leads to finite temperature behaviour for the e-SIAM that is qualitatively similar to that of the Hubbard model as seen from DMFT (that was described at the beginning of this section, and can also be seen in Fig. 3.10).

At non-zero temperatures, the system is described by not only the ground-state but also the excitations. The metallic and insulating solutions become temperature dependent, with energies $E_M(r, T)$ and $E_I(r, T)$ respectively. The curve $r_{c1}(T)$ at a given temperature again represents the values of the parameter r at which the local moment states enter the spectrum as excited eigenstates, while $r_{c2}(T)$ marks the point where the metallic ground-state leaves the spectrum: $E_M(r_{c2}(T), T) = E_I(r_{c2}(T), T)$. As a result, within the parameter range $r_{c1}(T) < r(T) < r_{c2}(T)$ at a finite temperature T , the low-energy part of the spectrum (states within the central Kondo resonance) involves both the metallic and insulating solutions. These two low-lying solutions then govern the partition function and hence the free energy:

$$Z(r, T) \simeq Z_M(r, T) + Z_I(r, T), \quad F(r, T) \simeq -k_B T \ln (Z_M(r, T) + Z_I(r, T)), \quad (6.6)$$

where $Z_M(r, T) = \exp(-\beta \mathcal{V} (E_M(r, T) - T S_M(r, T)))$ is the partition function corresponding to the metallic state with energy $E_M(r, T)$ and entropy $S_M(r, T)$ per unit volume $\mathcal{V}$, and

$$Z_I = \exp(-\beta \mathcal{V} (E_I(r, T) - T S_I(r, T))), \quad (6.7)$$

is similarly the partition function corresponding to the local moment states.

In the thermodynamic limit, only the larger of Z_M and Z_I contributes to the partition function $Z(r, T)$. Given a temperature T , there exists a value $r_c(T)$ lying in the range $r_{c1}(T) < r_c(T) < r_{c2}(T)$ at which the partition functions $Z_M(T)$ and $Z_I(T)$ become equal: $Z_M(r_c(T), T) = Z_I(r_c(T), T)$. This condition defines a curve $\{(T, r_c(T))\}$ at which the system undergoes a first-order transition between the metallic and insulating states:

$$\lim_{\mathcal{V} \rightarrow \infty} F(r, T) = \begin{cases} -k_B T \ln Z_M(r, T) & \text{if } r < r_c \text{ } (Z_M > Z_I) \\ -k_B T \ln Z_I(r, T) & \text{if } r > r_c \text{ } (Z_M < Z_I) \end{cases}. \quad (6.8)$$

The first-order curve $r_c(T)$ is determined from the partition function equality (or equivalently, the free energy equality) condition between the metal and the insulator:

$$T = \frac{E_M(r_c, T) - E_I(r_c, T)}{S_M(r_c, T) - S_I(r_c, T)}. \quad (6.9)$$

At $T = 0$, the first-order line $r_c(T)$ becomes identical to the critical point r_{c2} , as $E_M(r_c) = E_I(r_c)$ at $T = 0$. This simply means that the thermal entropy plays no role at zero temperature, and the transition is purely quantum-mechanical in nature. As temperature is now increased, the large entropy S_I of the doubly degenerate local moment states comes into play and the insulating state is able to take over at a smaller value of r . This suggests that a first-order transition can take place in the thermodynamic limit at $r_c(T) < r_{c2}(T)$. This has been shown schematically in Fig. (3.10).




Figure 3.10: Qualitative structure of the finite temperature coexistence region of the $J - U_b$ model. The dotted lines on the left and the right represent the spinodals r_{c1} and r_{c2} where the insulating and metallic solutions become unstable, respectively. The solid red line represents the first-order line where the free energies and the partition functions of the two solutions become equal, $Z_M = Z_I$.

Further, away from the thermodynamic limit, the free energy in eq. (6.6) admits contributions from the stable metallic state as well as the metastable insulating state, and results in the coexistence of the two phases. This is because, depending on whether we are approaching from r_{c2}^+ or r_{c1}^- , the system will remain mostly in the minimum of either Z_I or Z_M respectively. There exist finite activation barriers in the free energy for passage into the other minimum, leading to hysteresis. The transition is observed only at $r_{c1}(T)$ or $r_{c2}(T)$ where the insulating and metallic solutions, respectively, become unstable. These two lines, therefore, mark the spinodals of the coexistence region. At a sufficiently high temperature T_c , the metallic and insulating solutions become indistinguishable at the first spinodal r_{c1} : $E_M(r_{c1}, T_c) = E_I(r_{c1}, T_c)$, $S_M(r_{c1}, T_c) = S_I(r_{c1}, T_c)$. We recall that of these two equalities, the first ($E_M = E_I$) also corresponds to the locus of the second spinodal (corresponding to r_{c2}). Moreover, combining the two equalities (energy and entropy) leads to the free energy equality condition, marking the locus of the first-order line $r_c(T)$. This has the consequence that the two spinodals and the first-order line merge into a second order critical point $r_c(T_c)$ at temperature T_c . This concludes our discussion of the finite temperature behaviour of the extended SIAM model and its qualitative similarities with the DMFT phase diagram for the half-filled Hubbard model on the Bethe lattice.

3.6.4 Zero temperature origin of critical fluctuations above the second order point

Above the second order point $r_c(T_c)$, the DMFT phase diagram shows a rapid crossover from the paramagnetic metallic solution to the paramagnetic insulating solution [Georges et al.(1996)Georges, Kotliar, Krauth, and Rozenberg, Limelette et al.(2003b)Limelette, Georges, Jérôme, Wzietek, Metcalf, and Honig]. Remarkably, signatures of quantum critical scaling in this crossover region have




Figure 3.11: Top: Variation of the one particle correlations in the state $|1, \sigma, -\rangle$ between impurity and bath zeroth site (blue), as well as within the bath (red and violet). The former vanishes, depicting the excited state localisation transition at r_{c1} . Left: Variation of the spin-flip (red) and charge isospin-flip (blue) correlations between impurity and bath, close to r_{c1} . They both increase, indicating that the impurity is now less strongly coupled with the bath. Right: Variation of the spin-flip (red) and charge isospin-flip (blue) correlations between the zeroth and second sites of the bath, close to r_{c1} . Both show an initial increase, leading to the propagation of relatively long-range correlations into the bath.

been recently predicted from theoretical analyses [Terletska et al.(2011)Terletska, Vučičević, Tanasković, and Dobrosavljević, Vučičević et al.(2013)Vučičević, Terletska, Tanasković, and Dobrosavljević], as well as detected experimentally in transport measurements of several organic compounds [Kagawa et al.(2005)Kagawa, Miyagawa, and Kanoda, Furukawa et al.(2015)Furukawa, Miyagawa, Taniguchi, Kato, and Kanoda]. In Refs. [Terletska et al.(2011)Terletska, Vučičević, Tanasković, and Dobrosavljević, Vučičević et al.(2013)Vučičević, Terletska, Tanasković, and Dobrosavljević], this has been ascribed to the existence of a *hidden quantum criticality* in the maximally frustrated 1/2-filled Hubbard model on the Bethe lattice with infinite coordination number. In order to locate the zero temperature origin of these signatures of quantum criticality, we inspect carefully the quantum-mechanical fluctuations near the two important points r_{c1} and r_{c2} in our model.

Close to r_{c1} , we compute correlations in the state $|1, \sigma, -\rangle$. As shown in the top panel of Fig. (3.11), we find that while the one-particle correlation between the impurity and the bath vanishes, the same between the zeroth and first sites of the bath picks up. Importantly, these correlations extend beyond the immediate neighbourhood of the impurity. This is observed in, for example, the one-particle correlation between the zeroth site and the second site (purple curve in the top panel of Fig. (3.11)). Some other signatures of correlations in the bath are shown in Fig. (3.11): while the left panel displays increased spin-flip and charge isospin-flip correlations between the zeroth and first sites of the bath, a similar phenomenon is observed between the zeroth and second sites of the bath in the right panel. Our findings are consistent with the presence of scale-invariant solutions obtained from recent NRG-DMFT calculations [Eisenlohr et al.(2019)Eisenlohr, Lee, and Vojta] that correspond to the metastable insulating solutions in the coexistence region.

The growth of similar long-ranged correlations within the bath (and leading away from the immediate neighbourhood of the impurity) near r_{c2} as well. We recall that correlations between the impurity and the zeroth site are observed to decrease (see left panel of Fig.(3.5)). Instead, as shown in Fig. (3.12), non-trivial two-particle correlations arise between the impurity and bath zeroth sites with bath sites that are farther away. The spreading of the spin-spin correlations shown in the left panel of Fig. (3.12) indicates a “stretching” of the Kondo singlet state prior to its destruction. We will show later that these enhanced correlations also result in a diverging quasiparticle mass at r_{c2} , indicating a breakdown of the local Fermi liquid metal. Further, the right panel of Fig. (3.12) indicates that longer-ranged pairing correlations develop between the bath zeroth site and bath sites farther away upon approaching the transition.

In general, within the metallic phase and away from r_{c1} or r_{c2} , the impurity is strongly coupled to the bath zeroth site and the impurity-bath entanglement follows an area law characteristic of the local nature of the impurity problem. However, near the excited state and ground state transitions at r_{c1} or r_{c2} respectively, these results indicate a spreading of entanglement into the bath: more and more bath sites beyond the zeroth site get correlated with the impurity site as the ESQPT or the QCP is approached. This is corroborated in Fig. 3 of the Supplementary Materials [sup(Supplementary Materials are available online)]. This suggests a significant enhancement of the entanglement beyond the area law and is consistent with the critical quantum fluctuations observed in the various two-particle correlations.

The results of this subsection indicate that it might be possible to experimentally detect the signatures of such critical quantum fluctuations around the spinodals of the DMFT phase diagram even at $T > 0$. We note that signatures of second-order criticality have been indeed observed recently

in vanadium sesquioxide, in the form of critical slowing down and critical opalescence [Kundu et al.(2020)Kundu, Bar, Nayak, and Bansal]. It is, therefore, tempting to speculate that these signatures arise from the presence of critical quantum fluctuations.




Figure 3.12: Left: Variation of the spin-spin correlations between the impurity and bath sites 1 through 4, close to r_{c2} . They show an increase, revealing the distribution of entanglement into the bath. Right: Variation of pairing correlations within the bath. These correlations show an increase, because of the weakening of the Kondo singlet.

3.7 Non-Fermi liquid signatures at the MIT

3.7.1 Death of the local Fermi liquid: the Brinkman-Rice scenario

The low-lying metallic excitations of the bath can be obtained by considering the singlet ground-state of the model (of energy $\sim -3\tilde{\mathcal{J}}/4$) in the metallic phase, and then studying the effect of an electron hopping term (with coupling t) between the singlet and the rest of the bath as a perturbation (see left panel of Fig. (3.13)). Such a strong-coupling expansion in powers of $t^2/\tilde{\mathcal{J}}$ leads to the usual local Fermi liquid effective Hamiltonian in the Kondo model [Nozières(1974), Wilson(1975b), Nozières, Ph. and Blandin, A.(1980), Mukherjee et al.(2022)Mukherjee, Mukherjee, Vidhyadhiraja, Taraphder, and Lal]:

$$H_{\text{LFL}} = \mathcal{F} \hat{n}_{1\uparrow} \hat{n}_{1\downarrow} + H_{\text{KE}}, \quad \mathcal{F} \sim t^4/\tilde{\mathcal{J}}^3, \quad (7.1)$$

where $\hat{n}_{1\sigma}$ are the number operators for the first site (site adjacent to the zeroth site of the conduction bath), H_{KE} is the kinetic energy arising from the nearest-neighbour hopping among all sites in the bath apart from the zeroth site, and $\mathcal{F}$ is the local Fermi liquid correlation strength. The Kondo singlet (formed between the impurity spin and the bath zeroth site) has decoupled from the rest of the lattice, and eq. (7.1) describes the effective Hamiltonian for the rest of the bath beyond the zeroth site.

Following an identical approach, we find that the effective Hamiltonian for the low-lying excitations in the metallic phase is given by

$$H_{\text{LFL}} = \mathcal{F} [\hat{n}_{1\uparrow} \hat{n}_{1\downarrow} + (1 - \hat{n}_{1\uparrow})(1 - \hat{n}_{1\downarrow})] + H_{\text{KE}}, \quad \text{where } \mathcal{F} = \frac{2t^4}{\tilde{\mathcal{J}} (3\tilde{\mathcal{J}}/4 + \tilde{U}_b)^2}. \quad (7.2)$$

We note that for our extended SIAM, the local Fermi liquid correlation $\mathcal{F}$ diverges as the transition is approached at $3\tilde{\mathcal{J}}/4 + \tilde{U}_b \rightarrow 0$. The divergence of $\mathcal{F}$ leads to the divergence of the renormalised mass of the quasiparticles, and can also be obtained from a Gutzwiller variational calculation as shown by Brinkman and Rice [Brinkman and Rice(1970)]. This shows the breakdown of perturbation theory and is indicative of the fact that the ground state is about to change at the metal-insulator transition.

The death of the local Fermi liquid is also seen from the vanishing of the dynamically generated low-energy Kondo screening scale T_K . Towards obtaining T_K close to the transition, we follow the approach used by Moeller et al. 1995 [Moeller et al.(1995)Moeller, Si, Kotliar, Rozenberg, and Fisher] and Held et al. 2013 [Held et al.(2013)Held, Peters, and Toschi]. Near the transition, they obtained a Kondo model from the SIAM by applying a Schrieffer-Wolff transition that removes the charge fluctuations of the impurity site and retains the physics of only the Kondo coupling J . This amounts to removing the side peaks from the impurity spectral function and focusing on the low-energy central peak (right panel of Fig. (3.13)). Held et al. then integrated the RG equation for this Kondo model by using a Lorentzian DOS $\rho(D)$ in the bath: $\rho(D) = \frac{\rho_0 \Gamma^2}{D^2 + \Gamma^2}$. This is motivated by



Figure 3.13: Left: Setup for obtaining the low-energy excitations above the strong-coupling ground state. The effects of the rest of the bath is included by treating the hopping between the zeroth site (blue sphere on the left) and the first site (sphere labelled 1) as a perturbation on top of the singlet ground state formed by the impurity (red sphere) and the zeroth site. Right: At r_{c1} , the Hubbard sidebands have become isolated from the central peak in the local spectral function because of the finite optical gap. Removing the UV excitations that couple the low-energy and high-energy bands then leads to a theory of the Kondo model with a Lorentzian DOS.

the fact that the central peak of the impurity spectral function is a Lorentzian, and the bath becomes equivalent to the impurity site under self-consistency.

We implement the same approach on the e-SIAM, such that the Schrieffer-Wolff transformation leads to a $J - U_b$ model with a Lorentzian electronic DOS in the bath. The expression of the Kondo temperature for such a system has been derived in Sec. 6 of the Supplementary Materials [sup(Supplementary Materials are available online)]. Very close to the transition, we have $r \rightarrow 0.25^-$, and the Kondo temperature is given by

$$T_K = \frac{D_0}{k_B} \exp \left[-\frac{\ln(r_{c2} - r)}{4U_b \rho_0} \right], \quad (7.3)$$

where $r_{c2} = 0.25$ and D_0 is the bare bandwidth. Note that the pre-factor of the logarithm is positive as U_b is negative: $-4U_b \rho_0 = |4U_b \rho_0|$. As we approach the transition, the parameter r takes the limit $r \rightarrow r_{c2}^-$, and the Kondo temperature scale vanishes:

$$\lim_{r \rightarrow r_{c2}^-} T_K = \frac{D_0}{k_B} \lim_{r \rightarrow r_{c2}^-} (r_{c2} - r)^{4|U_b \rho_0|} \rightarrow 0. \quad (7.4)$$

Following the renormalised perturbation theory approach of Hewson [Hewson(1993a), Coleman(2015)], the imaginary part of the self-energy of the local Fermi liquid quasiparticles is given by

$$\text{Im} [\Sigma(\omega)] \sim \frac{\mathcal{F}^2 \omega^2}{D_0 T_K^2}, \quad (7.5)$$

while the quasiparticle residue is given by $Z \sim T_K$. The divergence of $\text{Im} [\Sigma(\omega)]$ at the transition and the vanishing of T_K and the quasiparticle residue are important indicators of the loss of the local Fermi liquid excitations and the breakdown of Kondo screening.

The vanishing of the Kondo temperature scale also leads to the divergence of thermodynamic quantities such as the impurity contribution to local spin susceptibility χ_{imp} and the specific heat coefficient γ_{imp} . As the low-energy theory is a local Fermi liquid, χ_{imp} and γ_{imp} retain their Fermi liquid forms but involve the highly renormalised Kondo temperature scale [Wilson(1975b), Hewson(1993b)]:

$$\begin{aligned} \chi_{\text{imp}} &= \frac{w}{4k_B T_K}, \quad \lim_{r \rightarrow r_{c2}^-} \chi_{\text{imp}} = \frac{w}{4k_B} (r_{c2} - r)^{-4|U_b \rho_0|}, \\ \gamma_{\text{imp}} &= \frac{\pi^2 k_B^2 w}{6T_K}, \quad \lim_{r \rightarrow r_{c2}^-} \gamma_{\text{imp}} = \frac{\pi^2 k_B^2 w}{6} (r_{c2} - r)^{-4|U_b \rho_0|}, \end{aligned} \quad (7.6)$$

where $w \sim 0.4128$ is the Wilson number for the Kondo model [Wilson(1975b), Mukherjee et al.(2022)Mukherjee, Mukherjee, Vidhyadhiraja, Taraphder, and Lal]. These results also show that the ratio of χ_{imp} and γ_{imp} , referred to as the Wilson ratio $R \equiv \frac{4\pi^2 k_B^2 \chi_{\text{imp}}}{3\gamma_{\text{imp}}}$, remains pinned at the single-channel Kondo value of $R_{\text{LFL}} = 2$ for $r \rightarrow r_{c2}^-$. This shows that the low-frequency Landau quasiparticles are able to survive until very close to the QCP. As we will show in the next subsection, the metallic excitations precisely at the QCP acquire non-Fermi liquid character and involve additional correlations between holons and doublons. These additional correlations are expected to lead to an enhancement of the Wilson ratio. Evidence for this can be found in the DMFT calculation of a local Wilson ratio very close to the Brinkman-Rice transition, $R_{\text{loc}} = 2.9 \pm 0.2$ [Georges et al.(1996)Georges, Kotliar, Krauth, and Rozenberg].

3.7.2 Emergence of non-Fermi liquid excitations at the transition

In order to obtain an effective Hamiltonian for the excitations precisely at the critical point $r = 1/4$, we first note that the ground-state subspace of the zero bandwidth $J - U_b$ effective model (eq. (5.3)) becomes degenerate at this point:

$$\frac{1}{\sqrt{2}} (|\uparrow\rangle_d |\downarrow\rangle_0 - |\downarrow\rangle_d |\uparrow\rangle_0), \quad |\sigma\rangle_d |0\rangle_0, \quad |\sigma\rangle_d |2\rangle_0, \quad (\sigma = \uparrow, \downarrow), \quad (7.7)$$

with all the states lying at zero energy. Here, the first ket (with subscript d) represents the impurity spin configuration while the second ket (with subscript 0) represents the configuration of the bath zeroth site). We now diagonalise these states in the presence of the electron hopping t into the rest of the conduction bath, and obtain the effective Hamiltonian for the low-lying excitations mediated

by t . This is described in detail in Sec. 7 of the Supplementary Materials [sup(Supplementary Materials are available online)]. We note that the ground state is four-fold degenerate at energy $-t$ in the presence of the hopping, and comprises the following states:

$$|N_{\text{tot}} = 2, S_{\text{tot}}^z = 0\rangle, |N_{\text{tot}} = 4, S_{\text{tot}}^z = 0\rangle, |N_{\text{tot}} = 3, S_{\text{tot}}^z = \frac{1}{2}\rangle, |N_{\text{tot}} = 3, S_{\text{tot}}^z = -\frac{1}{2}\rangle, \quad (7.8)$$

where $N_{\text{tot}} = \hat{n}_d + \hat{n}_0 + \hat{n}_1$ and $S_{\text{tot}}^z = S_d^z + S_0^z + S_1^z$ are the total number operator and total magnetisation operator respectively for the impurity, bath zeroth and first sites taken together. It is worth observing that

- the presence of the unscreened states $|N_{\text{tot}} = 3, S_{\text{tot}}^z = \pm \frac{1}{2}\rangle$ in the ground-state subspace is a result of the inexact screening at the critical point, and
- the presence of a degenerate ground-state manifold indicates that these excitations of the bath will likely be of the non-Fermi liquid (NFL) kind [Varma et al.(2002)Varma, Nussinov, and Van Saarloos, Si and Kotliar(1993a), Kotliar and Si(1993b)].

The precise form of the effective Hamiltonian for this non-Fermi liquid, as well as the behaviour of various correlation functions, is obtained by considering the ground-state manifold in conjunction with the excited states at energy $-t$.

Since the total Hamiltonian conserves the total spin S_{tot}^z , the effective Hamiltonian separates into two sectors, $S_{\text{tot}}^z = 0$ and $|S_{\text{tot}}^z| = \frac{1}{2}$. Detailed calculations on obtaining this effective Hamiltonian from the RG fixed point theory are shown in Sec. 7 of the Supplementary Materials [sup(Supplementary Materials are available online)]. In order to highlight certain distinct features, we present simplified effective Hamiltonians only for the $S_{\text{tot}}^z = 0$ and $S_{\text{tot}}^z = \frac{1}{2}$ sectors. We first take a look at the $S_{\text{tot}}^z = 0$ sector:

$$H_{\text{eff}}^{S_{\text{tot}}^z=0} = t\vec{S}_d \cdot (\vec{S}_{1,-} + \vec{S}_{3,-}). \quad (7.9)$$

The effective spin-1/2 ladder operators $\vec{S}_{1,\pm}$ act on the positive and negative parity sectors (labelled by $\pm$) of the $\hat{n}_0 + \hat{n}_1 = 1$ subspace spanned by the states

$$|\uparrow\rangle_{1,\pm} = |\uparrow\rangle_0 |0\rangle_1 \pm |0\rangle_0 |\uparrow\rangle_1, \quad |\downarrow\rangle_{1,\pm} = |\downarrow\rangle_0 |0\rangle_1 \pm |0\rangle_0 |\downarrow\rangle_1, \quad (7.10)$$

leading to a Pauli matrix representation of $\vec{S}_{1,\pm}$ in that basis. The Pauli matrix operators $\vec{S}_{3,\pm}$ similarly act on the $\hat{n}_0 + \hat{n}_1 = 3$ subspace spanned by the states $|\uparrow\rangle_{3,\pm} = |\uparrow\rangle_0 |2\rangle_1 \pm |2\rangle_0 |\uparrow\rangle_1$ and $|\downarrow\rangle_{3,\pm} = |\downarrow\rangle_0 |2\rangle_1 \pm |2\rangle_0 |\downarrow\rangle_1$, obtained by applying the transformation $|0\rangle \rightarrow 2$ on the $\hat{n}_0 + \hat{n}_1 = 1$ counterparts. We now point out two interesting features of this simplified effective Hamiltonian in eq. (7.9).

- The fact that the impurity is now interacting with emergent spins $\vec{S}_1$ and $\vec{S}_3$ that span over two lattice sites (0 and 1) instead of just the zeroth site indicates that the entanglement between the impurity and bath is now extended beyond the bath zeroth site, and that *the singlet is being stretched*.

- The presence of two spins that are trying to simultaneously screen the impurity spin introduces frustration in the dynamics of the impurity. This is reminiscent of the 2-channel Kondo effect, and the connection is made more precise in the concluding section of our work.

Both these features are precursors to the ultimate and complete destruction of screening in the local moment regime, and reflect the fact that the impurity-bath system *is now in an over-screened state*.

We now focus on the effective Hamiltonian of the $S_{\text{tot}}^z \neq 0$ subspace:

$$H_{\text{eff}}^{S_{\text{tot}}^z \neq 0} = -\frac{1}{2}t (S_d^- \mathcal{B}_{\uparrow}^+ + \text{h.c.}) . \quad (7.11)$$

The Pauli matrix operators $\mathcal{B}_{\uparrow}^+$ and $\mathcal{B}_{\uparrow}^-$ flip between the doublet of states $|\uparrow\rangle_0 |\uparrow\rangle_1$ and $\frac{1}{\sqrt{2}} (|2\rangle_0 |0\rangle_1 + |0\rangle_0 |2\rangle_1)$:

$$\begin{aligned} \mathcal{B}_{\uparrow}^- |\uparrow\rangle_0 |\uparrow\rangle_1 &= \frac{1}{\sqrt{2}} (|2\rangle_0 |0\rangle_1 + |0\rangle_0 |2\rangle_1) , \\ \mathcal{B}_{\uparrow}^+ \frac{1}{\sqrt{2}} (|2\rangle_0 |0\rangle_1 + |0\rangle_0 |2\rangle_1) &= |\uparrow\rangle_0 |\uparrow\rangle_1 . \end{aligned} \quad (7.12)$$

The structure of the effective Hamiltonian in eq. (7.11) leads to another drastic difference from the behaviour of the local Fermi liquid: it allows for the electrons to be “Andreev scattered” into an orthogonal state by hopping from the first site into the zeroth site. This is easily seen by considering the scattering processes in eq. (7.12): a state $|\uparrow\rangle_0 |\uparrow\rangle_1$ (i.e., with site 1 in $\uparrow$ configuration) has a finite probability of being flipped into the $|\downarrow\rangle_0 |2\rangle_1$ state (i.e., with site 1 in the doublon configuration). *An incident electron $c_{1\uparrow}^\dagger$ can therefore emerge as a doublon $c_{1\uparrow}^\dagger c_{1\downarrow}^\dagger$ or a hole $c_{1\uparrow}$ upon scattering from the zeroth site.* This is a direct consequence of the entanglement between the singlet and the first site at the critical point that was absent in the metallic regime of the e-SIAM. We note that similar orthogonal scattering processes also occur in a two-channel Kondo problem between effective pseudo-particles and pseudo-holes [von Delft et al.(1998b) von Delft, Zaránd, and Fabrizio].

At this point, it is worth noting that the polarised ground-state $|S_{\text{tot}}^z = \frac{1}{2}\rangle$ can be written as an equal superposition of the singlet state $|\text{SS}\rangle_{a0} \otimes |\uparrow\rangle_1$ and the local moment states $\frac{1}{\sqrt{2}} (|\uparrow, 0, 2\rangle - |\uparrow, 2, 0\rangle)$. This symmetry-broken ground state, along with its counterpart $|S_{\text{tot}}^z = -\frac{1}{2}\rangle$, act as a bridge between the ground states of the metallic and insulating phases. They are thus important in displaying the breakdown of the Kondo cloud at the QCP, and the consequences arising from it. We will hence consider these states in the next subsection and demonstrate some exotic non-Fermi liquid properties of these states, e.g., inexact screening of the impurity, fractional impurity magnetisation and fractional impurity entanglement entropy.

Finally, by mapping the quantum impurity problem of the e-SIAM onto that of a classical Coulomb gas [Anderson and Yuval(1969), Si and Kotliar(1993a)], it can be shown that the local self-energies $\Sigma_{dd}(\omega)$ and $\Sigma_{00}(\omega)$ of the impurity and zeroth sites respectively and certain two-particle correlation functions have power-law behaviours in the frequency domain:

$$\text{Re} [\Sigma_{dd}] (\omega) \sim |\omega|^{\gamma_{dd}}, \quad \text{Re} [\Sigma_{00}] (\omega) \sim |\omega|^{\gamma_{00}}, \quad (7.13)$$

$$\langle S_d^+ \rangle \sim |\omega|^{(\alpha_1-1)/2}, \quad \langle c_{0\uparrow}^\dagger c_{0\downarrow}^\dagger \rangle \sim |\omega|^{(\alpha_3-1)/2}. \quad (7.14)$$

The precise forms of the exponents in terms of the conduction electron scattering phase shifts, as well as several other technical details of the calculation, are provided in Sec. 8 of the Supplementary Materials [sup(Supplementary Materials are available online)]. While the various exponents are found to be non-universal functions of the fixed point value of J and the coupling U_b for $r \rightarrow r_{c2}-$, we argue in subsection (3.7.4) that they assume universal values precisely at the QCP ($r = r_{c2}$). The algebraic behaviour of these local correlations is reminiscent of critical behaviour ascribed to the class of local quantum criticality [Coleman et al.(2001a)Coleman, Pépin, Si, and Ramazashvili, Si et al.(2001)Si, Rabello, Ingersent, and Smith].

3.7.3 Impurity magnetisation and entanglement entropy

More indications of non-Fermi liquid behaviour at the QCP is obtained from a calculation of the impurity magnetisation and the impurity entanglement entropy ($S_{EE}(d)$) (shown in Sec. 9 of the Supplementary Materials [sup(Supplementary Materials are available online)]) for the symmetry-broken states $|S_{\text{tot}}^z = \sigma/2\rangle$. This leads to a fractional entanglement entropy (in units of $\log 2$) for each of the two states:

$$\begin{aligned} \rho_{\text{imp}} &= \begin{pmatrix} \frac{1}{2} + m_{\text{imp}}^z & 0 \\ 0 & \frac{1}{2} - m_{\text{imp}}^z \end{pmatrix} = \begin{pmatrix} \frac{3}{4} & 0 \\ 0 & \frac{1}{4} \end{pmatrix}, \\ S_{EE}(d) &= -\text{Tr}(\rho_{\text{imp}} \ln \rho_{\text{imp}}) \simeq 0.81 \log 2, \end{aligned} \quad (7.15)$$

where the impurity magnetisation m_{imp}^z takes the value $m_{\text{imp}}^z = 1/4$ at the QCP (i.e., half the value for a local moment). Further, $S_{EE}(d)$ can be written in terms of an effective impurity degeneracy g_{imp} , which we define using the impurity magnetisation: $g_{\text{imp}} \equiv 1 + 2|m_{\text{imp}}^z|$, such that it takes the expected values of 1 and 2 in the local Fermi liquid and local moment phases (with m_{imp}^z having values 0 and $\frac{1}{2}$ respectively). The effective degeneracy in the polarised subspace $S_{\text{tot}}^z = \pm 1/2$ at the QCP then turns out to be $3/2$, which is half-way between a unique state and a doublet. This corresponds to partial screening of the impurity degrees of freedom at the QCP, in contrast to complete screening in the metallic phase ($g_{\text{imp}} = 1$) and the absence of screening in the insulating phase ($g_{\text{imp}} = 2$).

In Sec. 9 of the Supplementary Materials [sup(Supplementary Materials are available online)], we show that the incomplete magnetisation (and hence the fractional value of S_{EE}) arises from the mixing of the local Fermi liquid ground-state $|\phi\rangle = \otimes_{k \leq k_F} |k \uparrow\rangle |k \downarrow\rangle$ and the gapless excitations above it, $|e\rangle_\sigma = e_\sigma^\dagger |\phi\rangle$, $e \equiv \sum_{k \in FS} c_{k\sigma}^\dagger$, leading to the decay of the local Fermi liquid quasiparticles [Varma et al.(2002)Varma, Nussinov, and Van Saarloos]. This is captured by the modified ground-state $|-\rangle$ and lowest-lying excited states $|+\rangle$ at the QCP:

$$|\pm\rangle = \frac{1}{\sqrt{2}} (|SS\rangle \otimes |e\rangle_\sigma \pm |LM\rangle_\sigma \otimes |\phi\rangle). \quad (7.16)$$

The spin-singlet state $|SS\rangle \sim \sum_\sigma \sigma |\sigma\rangle_d |\bar{\sigma}\rangle_0$ and the local moment state $|LM\rangle_\sigma = |\sigma\rangle_d |2\rangle_0$ represent the configurations of the impurity and the bath zeroth sites. Such a ground-state $|-\rangle$ should be contrasted with the local Fermi liquid ground-state $|SS\rangle \otimes |\phi\rangle$ that is stable in the metallic phase for $r_{c1} < r < r_{c2}$. This change in the ground-state manifests in the vanishing of the quasiparticle residue of the local Fermi liquid excitations: $Z = |\langle + | e_\sigma^\dagger | - \rangle|^2 = |\langle + | (|LM\rangle_\sigma \otimes |e\rangle_\sigma) \rangle|^2 = 0$,

consistent with the orthogonality catastrophe [Anderson(1967)] described below eq. (7.5). As the local Fermi liquid quasiparticles are rendered unstable, they are replaced by stable composite three-site excitations of the form, $S_d^- c_{0\bar{\sigma}} e_{\sigma}^{\dagger}$. This is further reflected in the fact that the corresponding residue for such three-site composite excitation is non-zero at the QCP: $|\langle + | S_d^- c_{0\bar{\sigma}} e_{\sigma}^{\dagger} | - \rangle|^2 > 0$.

3.7.4 Friedel scattering phase shift and fractional excess charge

The orthogonality catastrophe faced by the infrared excitations of the conduction bath at the QCP points towards the presence of an “unrenormalised” phase shift [Anderson(1990)] of the scattering k -states. Indeed, using the Friedel sum rule [Friedel(1956), Langer and Ambegaokar(1961), Langreth(1966)], we obtain a $\pi/2$ phase shift for the extended states at the Fermi surface for the QCP, which is only half the unitary limit. This can be seen by recalling that the ground state in eq. (7.16) is an equal admixture of a decoupled local moment state and a singlet state. Through the singlet state, the impurity contributes an excess charge of unity to the Fermi surface of the conduction bath (because of the presence of gapless spin-flip fluctuations). The local moment state does not contribute any excess charge, because the impurity is decoupled from the bath. The net excess charge then comes out to be $n_{\text{exc}} = \frac{1}{2}(1 + 0) = \frac{1}{2}$. From the Friedel sum rule, the phase shift at the fixed point is $\delta^* = \pi n_{\text{exc}} = \pi/2$, as mentioned above. Expectedly, n_{exc} is zero in the local moment phase. Also, as the exponents of the algebraic form for the local quantities shown in eq. (7.13) are functions of the scattering phase shift, we find that these exponents take universal values precisely at the QCP.

The values of the excess charge at the QCP and in the local moment phase lead to important corollaries. Firstly, the jump in the excess charge by unity in going from $r < r_{c2}$ to $r > r_{c2}$ is in fact a reduction in the Luttinger volume V_L of the conduction bath [Luttinger(1960), Martin(1982a)]. V_L counts the number of extended states present at and below the Fermi volume, and corresponds to a topological quantum number [Oshikawa(2000), Seki and Yunoki(2017), Heath and Bedell(2020b), Mukherjee and Lal(2020a)]. Since the excess charge n_{exc} calculated above is simply the impurity contribution to V_L at the Fermi surface, the difference in the excess charge across the transition implies that the two phases acquire different values for the invariant V_L . Thus, n_{exc} tracks the change (ΔV_L) in the Luttinger volume, and the transition is topological in nature. Secondly, in a recent work, Sen et al. [Sen et al.(2020)Sen, Wong, and Mitchell] have shown that the Mott MIT of the infinite-dimensional Hubbard model proceeds through the dissociation of domain walls in a fictitious Su-Schrieffer-Heeger chain connected to the physical lattice sites. If one couples their observation with the fact that the two dissociated domain walls at the ends of the SSH chain are together known to host a single charge [Heeger et al.(1988)Heeger, Kivelson, Schrieffer, and Su, Stone(1985)], it becomes evident that the fractional excess charge we obtain at the QCP corresponds to the state that is localised in the SSH chain near the physical lattice site as obtained by Sen et al. The well-known bulk-boundary correspondence of the SSH model (see, e.g., [Asbóth et al.(2016)Asbóth, Oroszlány, and Pályi]) suggests that the excess charge n_{exc} and the change ΔV_L in the Luttinger volume are the respective topological invariants of the boundary and bulk in the e-SIAM that are tied to one another. Further, the appearance of a half-quantised n_{exc} at the MIT of the e-SIAM corresponds, in the SSH model, to the dissociation of a domain wall-anti domain wall pair with each carrying a half charge [Heeger et al.(1988)Heeger, Kivelson, Schrieffer, and Su, Stone(1985)].

We show in Sec. 9 of the Supplementary Materials [sup(Supplementary Materials are available

online)] that it is in fact possible to connect the scattering phase shift (δ^*), impurity magnetisation (m_{imp}^z), effective degeneracy ($\tilde{g}$) and excess charge (n_{exc}) in a single relation:

$$\delta^* = \pi n_{\text{exc}} = \left[1 + \left(\frac{2 - \tilde{g}}{\tilde{g} - 1} \right)^2 \right]^{-1} = \frac{4m_{\text{imp}}^z{}^2}{1 - 4|m_{\text{imp}}^z| + 8m_{\text{imp}}^z{}^2}. \quad (7.17)$$

As we now discuss, this provides a unified picture of the effect of frustration on the impurity spin. As we have seen earlier, the presence of U_b in the Hamiltonian of the e-SIAM introduces local moment states into the spectrum and leads to states with non-vanishing impurity magnetisation at the QCP. This non-zero magnetisation can be interpreted as a partial screening of the impurity spin, in turn resulting in an effective impurity degeneracy ($1 < \tilde{g} < 2$ between that of a unique screened state and an unscreened local moment. The partial screening also manifests in only half the excess charge being contributed to the conduction bath. This reduction in the excess charge acts as a change in the boundary conditions felt by the conduction electrons coupled to the impurity and manifests as a phase shift that is less than the unitarity limit of π .

3.8 Discussions and Outlook

In summary, we have shown that an attractive correlation on the bath zeroth site is enough to frustrate the Kondo effect and stabilise the local moment phase. The destruction of the Kondo cloud, and the associated local Fermi liquid, occurs through *pairing fluctuations in the bath and proximate to the impurity* (left panel of Fig. (3.8)). This is reminiscent of a subdominant superconducting tendency that was observed in the half-filled Hubbard model at $T = 0$ from a unitary RG treatment in Refs. [Mukherjee and Lal(2020c), Mukherjee and Lal(2022a)]. We also note the finding of non-local attractive effective interactions in a recent theoretical study of the 2D Hubbard model [Gazizova and LeBlanc(2023)]. We find that the critical point displays non-Fermi liquid behaviour with a vanishing quasiparticle residue Z . The strong agreement of our results with several aspects of DMFT suggests that the local self-energy obtained self-consistently in that method can be represented quite faithfully through the e-SIAM. It will, thus, be important to test the predictions offered by our approach directly within the DMFT method. We note that the non-Fermi liquid at the critical point displays a *partial correlation of doublons and holons* that is distinct from not only the correlated Fermi liquid metal (unbound holons and doublons), but also the paramagnetic insulator (comprised of bound holons and doublons). The excitations that propagate through the two unscreened states in the ground-state subspace (Eq. (7.8)) involve the simultaneous creation of a doublon and a holon: in these channels, the doublons and holons cannot propagate in the absence of each other. Such an incomplete correlation is an intermediate step towards complete confinement in the local moment phase. It appears interesting to experimentally test some of these ideas on a mesoscopic quantum dot [Iftikhar et al.(2015)Iftikhar, Jezouin, Anthore, Gennser, Parmentier, Cavanna, and Pierre, Iftikhar et al.(2018)Iftikhar, Anthore, Mitchell, Parmentier, Gennser, Ouerghi, Cavanna, Mora, Simon, and Pierre] which is, apart from the usual electron tunnel coupling to an electronic reservoir, additionally coupled through the proximity effect to a superconducting lead.

We stress here that the e-SIAM analysed in this work represents an impurity model that (i) has a single correlated channel of conduction electrons, (ii) is consistent with the symmetries of

a half-filled Hubbard model (SU(2)-spin, U(1)-charge and particle-hole), and (iii) shows a local metal-insulator transition. The model, therefore, provides a minimal route towards obtaining a Hamiltonian-based understanding of DMFT. Further, at the level of the renormalisation group flows that involve the frustration of Kondo screening, the effect of the attractive on-site interaction U_b introduced by us is equivalent to that of additional bath correlations that have been introduced in related impurity models. These include a single impurity Anderson model with multiple anisotropic conduction channels studied by Giamarchi et al. [Giamarchi et al.(1993)Giamarchi, Varma, Ruckenstein, and Nozières], and a periodic Anderson model enhanced with explicit s-d coupling (as well as a density-density interaction) between the conduction and impurity bands, studied by Si and Kotliar [Si and Kotliar(1993a), Kotliar and Si(1996), Si and Kotliar(1993b)] as well as by Ruckenstein et al., [Ruckenstein and Varma(1991)]. In the former K -channel model of Ref. [Giamarchi et al.(1993)Giamarchi, Varma, Ruckenstein, and Nozières], the z -component (V_l , $l \in [1, K - 1]$) of the Kondo interaction term in the $K - 1$ additional conduction channels acts as a source of frustration for the spin-flip term γ_0 of the Kondo interaction in the original $l = 0$ channel, leading to a breakdown of Kondo screening. The U_b coupling in the e-SIAM acts as a similar source of frustration: the equivalence lies at the level of the renormalisation group equations, such that both V_l and U_b oppose the RG relevance of γ_0 and J respectively. This can be made more precise by comparing the RG equations of our work (eqs.(9.28)) and that of Giamarchi et al. [Giamarchi et al.(1993)Giamarchi, Varma, Ruckenstein, and Nozières]: the mapping between the two frustration terms U_b and V_l is found to be of the form $U_b \sim \sum_{l=1}^{K-1} V_l^2 \rho$, where ρ is the conduction bath DOS. A similar relationship can be found between the RG equations obtained by us and those for the extended periodic Anderson model studied by Si and Kotliar [Si and Kotliar(1993a)]. In this sense, our analysis applies to a wide variety of models where the Kondo effect is stable in a certain regime of parameters but is destroyed in other parameter regimes by some form of quantum-mechanical frustration of the impurity spin degree of freedom.

Extensions of the present work involve analysing the mixed valence regime of the impurity site (i.e., away from half-filling) within the e-SIAM auxiliary model. This allows the possibility that the finite-temperature critical end-point of the DMFT first-order transition could turn into a quantum critical point from the merging of the $T = 0$ ESQPT and QPT observed in the present work. Recent DMFT calculations [Vučićević et al.(2015)Vučičević, Tanasković, Rozenberg, and Dobrosavljević] show a shrinking of the coexistence region with hole doping but fall short of revealing a QCP. If such a QCP does exist, can it harbour pair fluctuations between the impurity and zeroth bath sites that become dominant upon doping, signalling thereby a putative superconducting instability of a related bulk model? We recall that a superconducting state of matter was indeed observed to be emergent from a QCP in the hole-doped Hubbard model at $T = 0$ via a unitary RG treatment in Refs. [Mukherjee and Lal(2020d), Mukherjee and Lal(2022a)]. The phenomenon of electronic differentiation in k -space can perhaps be captured by considering cluster variants of the e-SIAM, i.e., multiple impurities connected to one another through single-particle hopping and/or RKKY-like interactions [Ferrero et al.(2007)Ferrero, Leo, Lecheminant, and Fabrizio, Sakai et al.(2009a)Sakai, Motome, and Imada]. In general, the presence of multiple and varied classes of correlations in the auxiliary model - localisation from Mott physics, delocalisation from spin and charge fluctuations, and pairing from local attractive correlations - makes this a strong candidate for an auxiliary model that carries the potential for describing the emergence of a variety of novel phases of correlated

quantum matter.

3.9 Appendix: Derivation of renormalisation group equations for the extended-SIAM

The Hamiltonian we are working with is the extended Anderson impurity model (described in Section III of the main manuscript):

$$\mathcal{H} = -\frac{1}{2}U(\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2 + \sum_{\vec{k},\sigma} \epsilon_{\vec{k}} \tau_{\vec{k},\sigma} + J \vec{S}_d \cdot \vec{S}_0 + V \sum_{\sigma} \left(c_{d\sigma}^{\dagger} c_{0\sigma} + \text{h.c.} \right) - \frac{1}{2}U_b(\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2. \quad (9.1)$$

The renormalisation in the Hamiltonian $H_{(j)}$ at the j^{th} RG step, upon decoupling an electronic state $q\beta$, is given by the expression:

$$(\Delta H_{(j)})_{\vec{q},\beta} = H_{(j-1)} - H_{(j)} = c_{\vec{q},\beta}^{\dagger} T_{\vec{q},\beta} \frac{1}{\omega_e - H_D} T_{\vec{q},\beta}^{\dagger} c_{\vec{q},\beta} + T_{\vec{q},\beta}^{\dagger} c_{\vec{q},\beta} \frac{1}{\omega_h - H_D} c_{\vec{q},\beta}^{\dagger} T_{\vec{q},\beta}, \quad (9.2)$$

where $\omega_{e,h}$ are the quantum fluctuation scales for the electron and hole scattering channels, H_D is the part of the Hamiltonian that is diagonal in k -space, and $T_{\vec{q},\beta} + T_{\vec{q},\beta}^{\dagger}$ is the part of the Hamiltonian that does not commute with $\hat{n}_{\vec{q},\beta}$ (in short, it is the part that is off-diagonal with respect to a particular state $\vec{q},\beta$). This off-diagonal part is made up of contributions from V, J and U_b : $T_{\vec{q}\uparrow} = \left[V c_{d\uparrow}^{\dagger} + J \sum_{\vec{k}} S_d^+ c_{\vec{k}\downarrow}^{\dagger} + U_b \sum_{\vec{k}_1, \vec{k}_2, \vec{k}_3} c_{\vec{k}_1\uparrow}^{\dagger} c_{\vec{k}_3\downarrow}^{\dagger} c_{\vec{k}_4\downarrow} \right] c_{\vec{q}\uparrow}$, while the diagonal part is given by $H_D = \epsilon_{\vec{q}} \tau_{\vec{q}\uparrow} + \frac{1}{4} J S_d^z (\hat{n}_{\vec{q}\uparrow} - \hat{n}_{\vec{q}\downarrow}) - \frac{1}{2} U_b (\tau_{\vec{q}\uparrow} - \tau_{\vec{q}\downarrow})^2$. The contribution from U_b is obtained by taking the Hartree contributions corresponding to the states $|q\sigma\rangle$ from the full interacting term in the Hamiltonian. Note that we have ignored potential scattering terms in the off-diagonal part $T_{\vec{q}\beta}$. In order to allow both spin-flip and charge transfer scattering processes, we start from initial states with $\tau_{\vec{q}\uparrow} = -\tau_{\vec{q}\downarrow}$. As a result, the contribution to H_D from the kinetic energy is $\epsilon_{\vec{q}} \tau_{\vec{q}\uparrow} = (\pm D) \times (\pm \frac{1}{2}) = D/2$, while the contribution from U_b is $-\frac{1}{2} U_b (\pm \frac{1}{2} - (\mp \frac{1}{2}))^2 = -U_b/2$.

3.9.1 Renormalisation of U_b

U_b can renormalise only via itself. The relevant renormalisation term in the particle sector is

$$U_b^2 \sum_{q\beta} \sum_{\substack{k_1, k_2, k_3, \\ k'_1, k'_2, k'_3}} c_{q\beta}^{\dagger} c_{k_1\beta} c_{k_3\beta}^{\dagger} c_{k'_1\beta} \frac{1}{\omega - H_D} c_{k'_2\beta}^{\dagger} c_{k'_3\beta} c_{k_2\beta}^{\dagger} c_{q\beta}. \quad (9.3)$$

In order to renormalise U_b , we need to contract one more pair of momenta. There are two choices. The first is by setting $k_3 = k'_3 = q$. The two internal states, then, are $q\beta$ and $\bar{q}\beta$. As discussed above, the intermediate state energy is $-U_b/4$. We therefore have

$$\frac{U_b^2 n_j}{\omega - D/2 + \frac{U_b}{2}} \sum_{\beta} \sum_{k_1, k_2, k'_1, k'_2} c_{k_1\beta} c_{k'_1\beta} c_{k'_2\beta}^{\dagger} c_{k_2\beta}^{\dagger} = \frac{U_b^2 n_j}{\omega - D/2 + \frac{U_b}{2}} \sum_{\beta} \sum_{k_1, k_2, k'_1, k'_2} c_{k'_2\beta}^{\dagger} c_{k'_1\beta} c_{k_2\beta}^{\dagger} c_{k_1\beta} \quad (9.4)$$

Another way to contract the momenta is by setting $k'_1 = k'_2 = q$, which gives a renormalisation of

$$\frac{U_b^2 n_j}{\omega - D/2 + \frac{U_b}{2}} \sum_{\beta} \sum_{k_1, k_2, k_3, k'_3} c_{k_1\beta} c_{k_3\bar{\beta}}^{\dagger} c_{k_3'\bar{\beta}}^{\dagger} c_{k_2\beta} = -\frac{U_b^2 n_j}{\omega - D/2} \sum_{\beta} \sum_{k_1, k_2, k_3, k'_3} c_{k_3\bar{\beta}}^{\dagger} c_{k_3'\bar{\beta}}^{\dagger} c_{k_2\beta} c_{k_1\beta} . \quad (9.5)$$

The two contributions cancel each other. The same cancellation happens in the hole sector as well.

3.9.2 Renormalisation of the impurity correlation U

The coupling U is renormalised by two kinds of vertices: V^2 and J^2 . We will consider these processes one at a time. For convenience, we define $\epsilon_d = -U/2$.

The renormalisation arising from the first kind of terms, in the particle sector, is

$$\begin{aligned} \sum_{q\beta} c_{q\beta}^{\dagger} c_{d\beta} \frac{V^2}{\omega - H_D} c_{d\beta}^{\dagger} c_{q\beta} &= \sum_{q\beta} V^2 \hat{n}_{q\beta} (1 - \hat{n}_{d\beta}) \left(\frac{1 - \hat{n}_{d\bar{\beta}}}{\omega_0 - E_0} + \frac{\hat{n}_{d\bar{\beta}}}{\omega_1 - E_1} \right) \\ &= V^2 n_j \sum_{\beta} (1 - \hat{n}_{d\beta}) \left(\frac{1 - \hat{n}_{d\bar{\beta}}}{\omega_0 - E_0} + \frac{\hat{n}_{d\bar{\beta}}}{\omega_1 - E_1} \right) . \end{aligned} \quad (9.6)$$

q runs over the momentum states that are being decoupled at this RG step: $|q| = \Lambda_j$. $E_{1,0}$ are the diagonal parts of the Hamiltonian at $\hat{n}_{d\bar{\beta}} = 1, 0$ respectively: $E_1 = \frac{D}{2}$ and $E_0 = \frac{D}{2} + \epsilon_d - \frac{J}{4}$. In order to relate ω_0 and ω_1 with the common fluctuation scale ω for the conduction electrons, we will replace these quantum fluctuation scales with the current renormalised values of the single-particle self-energy for the initial state from which we started scattering. For $\hat{n}_{d\bar{\beta}} = 0$, there is no additional self-energy because the impurity does not have any spin: $\omega_0 = \omega$. For $\hat{n}_{d\bar{\beta}} = 1$, we have an additional self-energy of ϵ_d arising from the correlation on the impurity: $\omega_1 = \omega + \epsilon_d$. Substituting the values of $E_{0,1}$ and $\omega_{0,1}$, we get

$$V^2 n_j \sum_{\beta} (1 - \hat{n}_{d\beta}) \left(\frac{1 - \hat{n}_{d\bar{\beta}}}{\omega - \frac{D}{2} - \epsilon_d + \frac{J}{4}} + \frac{\hat{n}_{d\bar{\beta}}}{\omega - \frac{D}{2} + \epsilon_d} \right) . \quad (9.7)$$

Performing a similar calculation for the hole sector gives the contribution:

$$V^2 n_j \sum_{\beta} \hat{n}_{d\beta} \left(\frac{1 - \hat{n}_{d\bar{\beta}}}{\omega - \frac{D}{2} + \epsilon_d} + \frac{\hat{n}_{d\bar{\beta}}}{\omega - \frac{D}{2} - \epsilon_d + \frac{J}{4}} \right) . \quad (9.8)$$

We now come to the second class of terms: spin-spin. We first look at the particle sector:

$$\frac{J^2}{4} \sum_{q\beta} c_{d\bar{\beta}}^{\dagger} c_{d\beta} c_{q\beta}^{\dagger} c_{-q\bar{\beta}} \frac{1}{\omega - H_D} c_{d\beta}^{\dagger} c_{d\bar{\beta}} c_{q\bar{\beta}}^{\dagger} c_{q\beta} = \frac{J^2}{4} n_j \frac{1}{\omega - \frac{D}{2} + \frac{J}{4}} \sum_{\beta} \hat{n}_{d\bar{\beta}} (1 - \hat{n}_{d\beta}) . \quad (9.9)$$

The diagonal part in the denominator was simple to deduce in this case because the nature of the scattering requires the spins S_d^z and $\frac{\beta}{2} (\hat{n}_{q\beta} - \hat{n}_{q\bar{\beta}})$ to be anti-parallel. This ensures that the intermediate state has an energy of $E = \frac{D}{2} + \epsilon_d - \frac{J}{4}$, and the quantum fluctuation scale is $\omega' = \omega + \epsilon_d$, such that $\omega' - E = \omega - \frac{D}{2} + \frac{J}{4}$. In the hole sector, we have

$$\frac{J^2}{4} n_j \frac{1}{\omega - \frac{D}{2} + \frac{J}{4}} \sum_{\beta} \hat{n}_{d\beta} (1 - \hat{n}_{d\bar{\beta}}) . \quad (9.10)$$

From eqs. (9.7), (9.8), (9.9) and (9.10), we write

$$\begin{aligned} \Delta U &= \Delta\epsilon_2 + \Delta\epsilon_0 - 2\Delta\epsilon_1 \\ &= -\frac{4V^2n_j}{\omega + \frac{U_b}{2} - \frac{D}{2} - U/2} + \frac{4V^2n_j}{\omega + \frac{U_b}{2} - \frac{D}{2} + U/2 + \frac{J}{4}} - \frac{J^2n_j}{\omega + \frac{U_b}{2} - \frac{D}{2} + \frac{J}{4}}, \end{aligned} \quad (9.11)$$

where we have restored the contribution from U_b in the denominator.

3.9.3 Renormalisation of the hybridisation V

Renormalisation of V happens through two kinds of processes: VJ and VU_b . Within the first kind, the scattering can be either via S_d^z or through $S_d^\pm$. For the first kind, we have the following contribution in the particle sector:

$$\begin{aligned} &\sum_{q\beta} V c_{q\beta}^\dagger c_{d\beta} \frac{1}{\omega - H_D} \frac{1}{4} J \sum_k (\hat{n}_{d\beta} - \hat{n}_{d\bar{\beta}}) c_{k\beta}^\dagger c_{q\beta} \\ &= \frac{1}{4} V J n_j \frac{1}{2} \left(\frac{1}{\omega'_1 - E} + \frac{1}{\omega'_2 - E} \right) \sum_{k\beta} (1 - \hat{n}_{d\bar{\beta}}) c_{d\beta} c_{k\beta}^\dagger. \end{aligned} \quad (9.12)$$

The transformation from $\frac{1}{\omega - H_D}$ to $\frac{1}{2} \left(\frac{1}{\omega'_1 - E} + \frac{1}{\omega'_2 - E} \right)$ is made so that we can account for both the initial state and the final state energies through the two fluctuation scales ω'_1 and ω'_2 respectively; we calculate the denominators for both the initial and final states and then take the mean of the two (hence the factor of half in front). This was not required previously because, in the earlier scattering processes, the impurity returned to its initial state at the end, at least in terms of $\epsilon_d (\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2$, and so we had $\omega'_1 = \omega'_2 = \omega'$. Substituting the energies and the ω -scales, we get

$$- \left(\frac{\frac{n_j}{4} V J \frac{1}{2}}{\omega - \frac{D}{2} + \frac{J}{4}} + \frac{\frac{n_j}{4} V J \frac{1}{2}}{\omega - \frac{D}{2} - \epsilon_d + \frac{J}{4}} \right) \sum_{k\beta} (1 - \hat{n}_{d\bar{\beta}}) c_{k\beta}^\dagger c_{d\beta}. \quad (9.13)$$

One can generate another such process by exchanging the single-particle process and the spin-exchange process:

$$\sum_{q\beta} \frac{1}{4} J \sum_k (\hat{n}_{d\beta} - \hat{n}_{d\bar{\beta}}) c_{q\beta}^\dagger c_{k\beta} \frac{1}{\omega - H_D} V c_{d\beta}^\dagger c_{q\beta}. \quad (9.14)$$

This is simply the Hermitian conjugate of the previous contribution. Combining this with the previous then gives

$$-\frac{n_j}{8} V J \left(\frac{1}{\omega - \frac{D}{2} + \frac{J}{4}} + \frac{1}{\omega - \frac{D}{2} - \epsilon_d + \frac{J}{4}} \right) \sum_{k\beta} (1 - \hat{n}_{d\bar{\beta}}) \times (c_{d\beta}^\dagger c_{k\beta} + \text{h.c.}). \quad (9.15)$$

If we similarly calculate the contributions from the spin-exchange processes involving $S_d^\pm$, we get

$$-\frac{1}{4} V J n_j \left(\frac{1}{\omega - \frac{D}{2} + \frac{J}{4}} + \frac{1}{\omega - \frac{D}{2} - \epsilon_d + \frac{J}{4}} \right) \sum_{k\beta} (1 - \hat{n}_{d\beta}) (c_{k\bar{\beta}}^\dagger c_{d\bar{\beta}} + \text{h.c.}). \quad (9.16)$$

The contributions from the hole sector are obtained by making the transformation $\hat{n}_{d\bar{\beta}} \rightarrow 1 - \hat{n}_{d\bar{\beta}}$ on the particle sector contributions. The total renormalisation to V from VJ processes are

$$-\frac{3n_j}{8}VJ \left(\frac{1}{\omega + U_b/4 - \frac{D}{2} + \frac{J}{4}} + \frac{1}{\omega + U_b/4 - \frac{D}{2} + U/2 + \frac{J}{4}} \right) \sum_{k\beta} \left(c_{d\beta}^\dagger c_{k\beta} + \text{h.c.} \right). \quad (9.17)$$

The renormalisation in V from U_b arises through terms of VU_b and U_bV kind. The first term gives

$$\begin{aligned} - \sum_{k\beta} c_{d\beta}^\dagger c_{k\beta} & \left[\frac{\hat{n}_{d\bar{\beta}}}{2} \left(\frac{n_j U_b V}{\omega - \frac{D}{2} - \frac{U}{2} + \frac{U_b}{4}} + \frac{n_j U_b V}{\omega - \frac{D}{2} + \frac{U_b}{4}} \right) \right. \\ & \left. + \frac{1 - \hat{n}_{d\bar{\beta}}}{2} \left(\frac{n_j U_b V}{\omega - \frac{D}{2} + \frac{U_b}{4} + \frac{U}{2} + \frac{J}{4}} + \frac{n_j U_b V}{\omega - \frac{D}{2} + \frac{U_b}{4} + \frac{J}{4}} \right) \right]. \end{aligned} \quad (9.18)$$

The second term is of the form $\sum_{q\beta} \sum_k U_b V c_{q\beta}^\dagger c_{d\beta} \frac{1}{\omega - H_D} \hat{n}_{q\bar{\beta}} c_{k\beta}^\dagger c_{q\beta}$, and this is just the Hermitian conjugate of the previous term, so these two terms together lead to

$$\begin{aligned} -n_j U_b V \sum_{k\beta} \left(c_{d\beta}^\dagger c_{k\beta} + \text{h.c.} \right) & \left[\frac{\hat{n}_{d\bar{\beta}}}{2} \left(\frac{1}{\omega - \frac{D}{2} - \frac{U}{2} + \frac{U_b}{4}} + \frac{1}{\omega - \frac{D}{2} + \frac{U_b}{4}} \right) \right. \\ & \left. + \frac{1 - \hat{n}_{d\bar{\beta}}}{2} \left(\frac{1}{\omega - \frac{D}{2} + \frac{U_b}{4} + \frac{U}{2} + \frac{J}{4}} + \frac{1}{\omega - \frac{D}{2} + \frac{U_b}{4} + \frac{J}{4}} \right) \right]. \end{aligned} \quad (9.19)$$

In the hole sector, we have

$$-n_j U_b V \sum_{k\beta} c_{k\beta}^\dagger \left[\frac{\hat{n}_{d\bar{\beta}}}{2} \left(\frac{1}{\omega_1 - E_1} + \frac{1}{\omega'_1 - E_1} \right) + \frac{1 - \hat{n}_{d\bar{\beta}}}{2} \left(\frac{1}{\omega_0 - E_0} + \frac{1}{\omega'_0 - E_0} \right) \right] c_{d\beta}. \quad (9.20)$$

$E_1 = D/2 - U_b/4 - U/2 - J/4$, $E_0 = D/2 - U_b/4 - K/4$. The fluctuation scales are $\omega_1 = \omega = \omega'_0$, $\omega'_1 = \omega - U/2 = \omega_0$. Substituting these gives

$$\begin{aligned} - \sum_{k\beta} c_{d\beta}^\dagger c_{k\beta} & \left[\frac{1 - \hat{n}_{d\bar{\beta}}}{2} \left(\frac{n_j U_b V}{\omega - \frac{D}{2} - \frac{U}{2} + \frac{U_b}{4}} + \frac{n_j U_b V}{\omega - \frac{D}{2} + \frac{U_b}{4}} \right) + \frac{\hat{n}_{d\bar{\beta}}}{2} \left(\frac{n_j U_b V}{\omega - \frac{D}{2} + \frac{U_b}{4} + \frac{U}{2} + \frac{J}{4}} \right) \right. \\ & \left. + \frac{n_j U_b V}{\omega - \frac{D}{2} + \frac{U_b}{4} + \frac{J}{4}} \right]. \end{aligned} \quad (9.21)$$

The other term, obtained by exchanging V and U_b , gives the Hermitian conjugate, so the overall contribution from the hole sector is the same as the total contribution from the particle sector, but with $\hat{n}_{d\bar{\beta}} \rightarrow 1 - \hat{n}_{d\bar{\beta}}$. Combining both the sectors, we get

$$\begin{aligned} - \sum_{k\beta} \left(c_{d\beta}^\dagger c_{k\beta} + \text{h.c.} \right) & \left[\left(\frac{n_j U_b V/2}{\omega - \frac{D}{2} - \frac{U}{2} + \frac{U_b}{4}} + \frac{n_j U_b V/2}{\omega - \frac{D}{2} + \frac{U_b}{4}} \right) + \left(\frac{n_j U_b V/2}{\omega - \frac{D}{2} + \frac{U_b}{4} + \frac{U}{2} + \frac{J}{4}} \right) \right. \\ & \left. + \frac{n_j U_b V/2}{\omega - \frac{D}{2} + \frac{U_b}{4} + \frac{J}{4}} \right]. \end{aligned} \quad (9.22)$$

3.9.4 Renormalisation of the spin-exchange coupling J

The term $J\vec{S}_d \cdot \vec{S}_0$ can be split into three parts: $J^z S_d^z, \frac{1}{2}J^+ S_d^+ S_0^-$ and $\frac{1}{2}J^- S_d^- S_0^+$. We will only calculate the renormalisation in J^+ , which will be equal to that of J^- , J^z due to spin-rotation symmetry. The terms that renormalise J^+ are of the form $S_d^+ S_d^z$ and $S_d^z S_d^+$. In the particle sector, we have

$$\begin{aligned} & - \sum_q \sum_{kk'} \frac{1}{4} J^2 S_d^+ c_{q\downarrow}^\dagger c_{k'\uparrow} \frac{1}{\omega - H_D} S_d^z c_{k\downarrow}^\dagger c_{q\downarrow} = n_j \frac{1}{4} J^2 \left(-\frac{1}{2} S_d^+ \right) \sum_{kk'} c_{k\downarrow}^\dagger c_{k'\uparrow} \frac{1}{\omega + \frac{U_b}{2} - \frac{D}{2} + \frac{J}{4}}, \\ & \sum_q \sum_{kk'} \frac{1}{4} J^2 S_d^z c_{q\uparrow}^\dagger c_{k'\uparrow} \frac{1}{\omega - H_D} S_d^+ c_{k\downarrow}^\dagger c_{q\uparrow} = -n_j \frac{1}{4} J^2 \left(\frac{1}{2} S_d^+ \right) \sum_{kk'} c_{k\downarrow}^\dagger c_{k'\uparrow} \frac{1}{\omega + \frac{U_b}{2} - \frac{D}{2} + \frac{J}{4}}. \end{aligned} \quad (9.23)$$

The denominator is determined using $E = \frac{D}{2} + \epsilon_d - \frac{J}{4}$ and $\omega' = \omega + \epsilon_d$. In the hole sector, we similarly have

$$\begin{aligned} & \sum_q \sum_{kk'} \frac{1}{4} J^2 S_d^+ c_{k\downarrow}^\dagger c_{q\uparrow} \frac{1}{\omega - H_D} S_d^z c_{q\uparrow}^\dagger c_{k'\uparrow} = n_j \frac{1}{4} J^2 \left(-\frac{1}{2} S_d^+ \right) \sum_{kk'} c_{k\downarrow}^\dagger c_{k'\uparrow} \frac{1}{\omega + \frac{U_b}{2} - \frac{D}{2} + \frac{J}{4}}, \\ & - \sum_q \sum_{kk'} \frac{1}{4} J^2 S_d^z c_{k\downarrow}^\dagger c_{q\downarrow} \frac{1}{\omega - H_D} S_d^+ c_{q\downarrow}^\dagger c_{k'\uparrow} = -n_j \frac{1}{4} J^2 \left(\frac{1}{2} S_d^+ \right) \sum_{kk'} c_{k\downarrow}^\dagger c_{k'\uparrow} \frac{1}{\omega + \frac{U_b}{2} - \frac{D}{2} + \frac{J}{4}} \end{aligned} \quad (9.24)$$

The renormalisation due to U_b also happens through multiple terms. One of the terms is

$$\frac{1}{2} J U_b \sum_q \sum_{k,k'} S_d^+ c_{q\downarrow}^\dagger c_{k\uparrow} \frac{1}{\omega - H_D} \hat{n}_{q\uparrow} c_{k'\downarrow}^\dagger c_{q\downarrow} = -\frac{1}{2} \frac{J U_b n_j}{\omega - \frac{D}{2} + \frac{U_b}{2} + \frac{J}{4}} \sum_{k,k'} S_d^+ c_{k'\downarrow}^\dagger c_{k\uparrow}. \quad (9.25)$$

The factor of half in front is the same half factor that appears in front of the $S_1^+ S_2^-, S_1^- S_2^+$ terms when we rewrite $\vec{S}_1 \cdot \vec{S}_2$ in terms of $S^z, S^\pm$. Another term is obtained by switching J and U_b :

$$\frac{1}{2} J U_b \sum_q \sum_{k,k'} \hat{n}_{q\downarrow} c_{q\uparrow}^\dagger c_{k\uparrow} \frac{1}{\omega - H_D} S_d^+ c_{k'\downarrow}^\dagger c_{q\uparrow} = -\frac{1}{2} \frac{J U_b n_j}{\omega - \frac{D}{2} + \frac{U_b}{2} + \frac{J}{4}} \sum_{k,k'} S_d^+ c_{k'\downarrow}^\dagger c_{k\uparrow}. \quad (9.26)$$

The corresponding terms in the hole sector are

$$-\frac{1}{2} \frac{J U_b n_j}{\omega - \frac{D}{2} + \frac{U_b}{2} + \frac{J}{4}} \sum_{k,k'} S_d^+ c_{k'\downarrow}^\dagger c_{k\uparrow}, -\frac{1}{2} \frac{J U_b n_j}{\omega - \frac{D}{2} + \frac{U_b}{2} + \frac{J}{4}} \sum_{k,k'} S_d^+ c_{k'\downarrow}^\dagger c_{k\uparrow}. \quad (9.27)$$

3.9.5 Final URG equations

In summary, the renormalisation in the couplings takes the form:

$$\begin{aligned} \Delta U &= 4V^2 n_j \left(\frac{1}{d_1} - \frac{1}{d_0} \right) - n_j \frac{J^2}{d_2}, & \Delta V &= -\frac{3n_j V}{8} \left[J \left(\frac{1}{d_2} + \frac{1}{d_1} \right) + \frac{4U_b}{3} \sum_{i=1}^4 \frac{1}{d_i} \right], \\ \Delta J &= -\frac{n_j J (J + 4U_b)}{d_2}, & \Delta U_b &= 0, \end{aligned} \quad (9.28)$$

where the denominators d_i are given by

$$d_0 = \omega - \frac{D}{2} + \frac{U_b}{2} - \frac{U}{2}, \quad d_1 = \omega - \frac{D}{2} + \frac{U_b}{2} + \frac{U}{2} + \frac{J}{4}, \quad (9.29)$$

$$d_2 = \omega - \frac{D}{2} + \frac{U_b}{2} + \frac{J}{4}, \quad d_3 = \omega - \frac{D}{2} + \frac{U_b}{2}. \quad (9.30)$$

For the sake of completeness, we present the RG equation for a charge isospin Kondo coupling K between the impurity and the bath:

$$\Delta K = -\frac{n_j K (K + 4U_b)}{\omega - \frac{D}{2} + \frac{U_b}{2} + \frac{K}{4}}. \quad (9.31)$$

Indeed, the RG equation for K is observed to be identical in form to that for the spin Kondo coupling J , obtained through the transformation $J \rightarrow K$. This indicates that charge fluctuations between the bath zeroth and first sites (that are incited by an attractive interaction U_b) lead to the RG irrelevance of $K(> 0)$, and thereby a destabilisation of the charge Kondo effect similar to that presented in the main manuscript for J .

3.10 Appendix: Expressing correlation functions in terms of entanglement

We will first relate the impurity Greens function to the geometric entanglement. Given a ground state $|\Psi\rangle_{\text{gs}}$ and a spectrum of energies $\{E_n\}$, the retarded impurity Greens function (in time domain) is defined as $G_{d\sigma}(t) = -i\theta(t) \langle \{c_{d\sigma}(t), c_{d\sigma}^\dagger\} \rangle$. It can be given a spectral representation in terms of the eigenstates $\{|\Psi\rangle_n\}$ of the e-SIAM:

$$G_{d\sigma}(\omega) = \frac{1}{Z} \sum_n \left[\frac{|\langle \Psi_{\text{gs}} | c_{d\sigma} | \Psi_n \rangle|^2}{\omega + E_{\text{gs}} - E_n} + \frac{|\langle \Psi_n | c_{d\sigma} | \Psi_{\text{gs}} \rangle|^2}{\omega - E_{\text{gs}} + E_n} \right], \quad (10.1)$$

where $Z \equiv \sum_n e^{-\beta E_n}$ is the partition function. We will now insert a complete basis into the expression. The basis will be the set of eigenstates of the Hamiltonian $H = H_1 + H_2$, where H_1 is the two-site $J - U_b$ Hamiltonian formed by the impurity and zeroth sites, and H_2 is a tight-binding Hamiltonian formed by the remaining sites. Since the Hamiltonians are decoupled, the eigenstates $|\Psi\rangle_{m,n}$ will be direct product states formed by combining the eigenstates $|\phi\rangle_m, |\psi\rangle_n$ of the individual Hamiltonians H_1 and H_2 respectively: $|\Phi\rangle_{m,n} = |\phi\rangle_m \otimes |\psi\rangle_n$. Inserting this basis leads to the expression:

$$|\Psi\rangle_{\text{gs}} = \sum_{m,n} |\Phi\rangle_{m,n} (\langle \phi_m | \otimes \langle \psi_n |) |\Psi_{\text{gs}}\rangle. \quad (10.2)$$

The ground state of H_1 is the spin-singlet: $|\phi\rangle_0 = |ss\rangle$. We denote the ground state of the tight-binding Hamiltonian H_2 by $|\psi\rangle_0$. Because of the highly renormalised couplings V and J , the impurity site is almost maximally entangled with the zeroth site, such that the ground state $|\Psi\rangle_{\text{gs}}$ in the $r <$

0.25 regime can be thought of as a direct product of the two-site ground state, $c|ss\rangle + \sqrt{1-c^2}|ct\rangle$, in direct product with the tight-binding ground state:

$$|\Psi\rangle_{\text{gs}} \simeq \left(c|ss\rangle + \sqrt{1-c^2}|ct\rangle \right) \otimes |\psi\rangle_0 = |\Phi\rangle_{\text{ss}} + |\Phi\rangle_{\text{ct}} , \quad (10.3)$$

where $|\Phi\rangle_{\text{ss(ct)}} = |ss(ct)\rangle \otimes |\psi\rangle_0$. This suggests that not all terms in the summation of eq. (10.2) contribute; out of all $\{|\psi\rangle_n\}$, only the ground state $n = 0$ contributes, while only $|ss\rangle$ and $|ct\rangle$ contribute from the set $\{|\phi\rangle_m\}$. The summation then simplifies to

$$|\Psi\rangle_{\text{gs}} = |\Phi\rangle_{\text{ss}} \langle \text{ss} | \Psi_{\text{gs}}^{(2)} \rangle + |\Phi\rangle_{\text{ct}} \langle \text{ct} | \Psi_{\text{gs}}^{(2)} \rangle , \quad (10.4)$$

where $|\Psi_{\text{gs}}^{(2)}\rangle$ is the two-site part of the ground state and can be obtained by starting with the full ground state $|\Psi\rangle_{\text{gs}}$ and tracing over the other lattice sites of the system.

We assume that the global phases of the wavefunctions are real, and since the internal weights of the wavefunctions are real as well, the overlaps $\langle \text{ss} | \Psi_{\text{gs}}^{(2)} \rangle$, $\langle \text{ct} | \Psi_{\text{gs}}^{(2)} \rangle$ are also real. We can use these overlaps to define a geometric measure of entanglement [Shimony(1995), Wei and Goldbart(2003), Horodecki et al.(2009)Horodecki, Horodecki, Horodecki, and Horodecki]:

$$\varepsilon(\psi_1, \psi_2) = 1 - |\langle \psi_1 | \psi_2 \rangle|^2 . \quad (10.5)$$

If $|\psi_1\rangle$ is thought of as a separable state, then $|\psi_2\rangle$ should be less entangled if its overlap with $|\psi_1\rangle$ is large, which is indeed borne out by the definition. The overlaps then become $\langle \text{ss} | \Psi_{\text{gs}}^{(2)} \rangle = \sqrt{1 - \varepsilon(\text{ss}, \Psi_{\text{gs}}^{(2)})}$, and similarly for the state $|ct\rangle$. For brevity, we will use the notation $\varepsilon_{\text{ss}} \equiv \varepsilon(\text{ss}, \Psi_{\text{gs}}^{(2)})$, $\varepsilon_{\text{ct}} \equiv \varepsilon(\text{ct}, \Psi_{\text{gs}}^{(2)})$. The retarded impurity Greens function for spin σ can now be written in terms of these measures:

$$\begin{aligned} G_{d\sigma}(\omega) = \frac{1}{Z} \sum_n & \left[(1 - \varepsilon_{\text{ss}}) \left(\frac{|\langle \Phi_{\text{ss}} | c_{d\sigma} | \Psi_n \rangle|^2}{\omega + E_{\text{gs}} - E_n} + \frac{|\langle \Psi_n | c_{d\sigma} | \Phi_{\text{ss}} \rangle|^2}{\omega - E_{\text{gs}} + E_n} \right) + (1 - \varepsilon_{\text{ct}}) \frac{|\langle \Phi_{\text{ct}} | c_{d\sigma} | \Psi_n \rangle|^2}{\omega + E_{\text{gs}} - E_n} \right. \\ & + (1 - \varepsilon_{\text{ct}}) \frac{|\langle \Psi_n | c_{d\sigma} | \Phi_{\text{ct}} \rangle|^2}{\omega - E_{\text{gs}} + E_n} + \sqrt{(1 - \varepsilon_{\text{ss}})} \sqrt{(1 - \varepsilon_{\text{ct}})} \frac{\langle \Phi_{\text{ss}} | c_{d\sigma} | \Psi_n \rangle \langle \Psi_n | c_{d\sigma}^\dagger | \Phi_{\text{ct}} \rangle + \text{h.c.}}{\omega + E_{\text{gs}} - E_n} \\ & \left. + \sqrt{(1 - \varepsilon_{\text{ss}})} \sqrt{(1 - \varepsilon_{\text{ct}})} \frac{\langle \Phi_{\text{ct}} | c_{d\sigma} | \Psi_n \rangle \langle \Psi_n | c_{d\sigma}^\dagger | \Phi_{\text{ss}} \rangle + \text{h.c.}}{\omega - E_{\text{gs}} + E_n} \right] . \quad (10.6) \end{aligned}$$

We now generalise this to any real-space two-particle correlation involving operators O_1, O_2 that are at most two-particle operators and act on the combined Hilbert space of the impurity site and the zeroth site. The so-called lesser Greens function corresponding to these operators is defined as

$$G_{O_1^\dagger, O_2}^<(t) \equiv i \langle O_1^\dagger O_2(t) \rangle , \quad G_{O_1^\dagger, O_2}^<(t \rightarrow \infty) = i \langle \Psi_{\text{gs}} | O_1^\dagger O_2 | \Psi_{\text{gs}} \rangle . \quad (10.7)$$

We focus on the static case ($t \rightarrow \infty$) here. Following eq. (10.4), we can rewrite the ground states in terms of the entanglement measures mentioned above.

$$|\Psi\rangle_{\text{gs}} = |\Phi\rangle_{\text{ss}} \sqrt{1 - \varepsilon_{\text{ss}}} + |\Phi\rangle_{\text{ct}} \sqrt{1 - \varepsilon_{\text{ct}}} , \quad (10.8)$$

where $|\Phi\rangle_{\text{ss(ct)}}$ are the zero spin and zero charge isospin eigenstates of the Hamiltonian $H = H_1 + H_2$ defined below eq. (10.1). Substituting this in the static lesser Greens function gives

$$\begin{aligned} \frac{1}{i} G_{O_1^\dagger, O_2}^<(t \rightarrow \infty) = & (1 - \varepsilon_{\text{ss}}) \langle \Phi_{\text{ss}} | O_2 O_1^\dagger | \Phi_{\text{ss}} \rangle + (1 - \varepsilon_{\text{ct}}) \langle \Phi_{\text{ct}} | O_2 O_1^\dagger | \Phi_{\text{ct}} \rangle \\ & + \sqrt{1 - \varepsilon_{\text{ss}}} \sqrt{1 - \varepsilon_{\text{ct}}} \left(\langle \Phi_{\text{ss}} | O_2 O_1^\dagger | \Phi_{\text{ct}} \rangle + \langle \Phi_{\text{ct}} | O_2 O_1^\dagger | \Phi_{\text{ss}} \rangle \right). \end{aligned} \quad (10.9)$$

This can be extended to generalised retarded Greens functions $G_{O_1^\dagger, O_2}^R(t) \equiv -i\theta(t) \langle \{O_1^\dagger(t), O_2\} \rangle$, whose spectral representation is of the form

$$G_{O_1^\dagger, O_2}^R(\omega) = \frac{1}{Z} \sum_n \left[\frac{\langle \Psi_{\text{gs}} | O_2 | \Psi_n \rangle \langle \Psi_n | O_1^\dagger | \Psi_{\text{gs}} \rangle}{\omega + E_{\text{gs}} - E_n} + \frac{\langle \Psi_n | O_2 | \Psi_{\text{gs}} \rangle \langle \Psi_{\text{gs}} | O_1^\dagger | \Psi_n \rangle}{\omega - E_{\text{gs}} + E_n} \right]. \quad (10.10)$$

Following an approach very similar to the one that led to eq. (10.6), we can cast the generalised Greens function in terms of the geometric entanglement measures:

$$\begin{aligned} G_{O_1^\dagger, O_2}^R(\omega) = \frac{1}{Z} \sum_n & \left[(1 - \varepsilon_{\text{ss}}) \left(\frac{(O_1)_{\text{ss},n}^* (O_2)_{\text{ss},n}}{\omega + E_{\text{gs}} - E_n} + \frac{(O_1)_{n,\text{ss}}^* (O_2)_{n,\text{ss}}}{\omega - E_{\text{gs}} + E_n} \right) + (1 - \varepsilon_{\text{ct}}) \frac{(O_1)_{\text{ct},n}^* (O_2)_{\text{ct},n}}{\omega + E_{\text{gs}} - E_n} \right. \\ & + (1 - \varepsilon_{\text{ct}}) \frac{(O_1)_{n,\text{ct}}^* (O_2)_{n,\text{ct}}}{\omega - E_{\text{gs}} + E_n} + \sqrt{(1 - \varepsilon_{\text{ss}})} \sqrt{(1 - \varepsilon_{\text{ct}})} \frac{(O_2)_{\text{ss},n} (O_1)_{\text{ct},n}^* + \text{h.c.}}{\omega + E_{\text{gs}} - E_n} \\ & \left. + \sqrt{(1 - \varepsilon_{\text{ss}})} \sqrt{(1 - \varepsilon_{\text{ct}})} \frac{(O_2)_{\text{ct},n} (O_1)_{\text{ss},n}^* + \text{h.c.}}{\omega - E_{\text{gs}} + E_n} \right], \end{aligned} \quad (10.11)$$

where $(O_2)_{\text{ss},n}$ is the matrix element $\langle \Phi_{\text{ss}} | O_2 | \Psi_n \rangle$, and similar definitions exist for $(O_1)_{\text{ss},n}$ and its $|\Phi_{\text{ct}}\rangle$ counterparts.

Closely related to geometric entanglement is the quantum Fisher information (QFI) F_Q [Braunstein and Caves(1994), Hauke et al.(2016)Hauke, Heyl, Tagliacozzo, and Zoller], which quantifies how sensitive a state ρ is to unitary transformations generated by an observable $\hat{O}$. For a pure state $\rho = |\psi\rangle \langle \psi|$, $F_Q(\psi, \hat{O})$ is defined in terms of the variance of the operator $\hat{O}$ (i.e., the connected correlation function):

$$F_Q(\psi, \hat{O}) = 4\Delta(\hat{O})^2 = 4 \left(\langle \psi | \hat{O}^2 | \psi \rangle - \langle \psi | \hat{O} | \psi \rangle^2 \right), \quad (10.12)$$

and it provides an upper bound on the precision that can be achieved in measuring the parameter θ that is dual to the observable $\hat{O}$: $\mathcal{N}(\Delta\theta)^2 \geq F_Q^{-1}$, $\mathcal{N}$ being the number of independent measurements [Braunstein and Caves(1994), Hauke et al.(2016)Hauke, Heyl, Tagliacozzo, and Zoller]. $F_Q(\psi, \hat{O})$ is thus a measure of the entanglement arising from quantum fluctuation content in $|\psi\rangle$ (considered with respect to an eigenstate of $\hat{O}$). This can be made more manifest by considering a traceless operator $\hat{M}$ with eigenstates $\{|m\rangle\}$. Without loss of generality, we can rescale the operator such that $\sum m^2 = 1/2$. There are several examples of such operators in the context of the present work, such as the impurity magnetisation operator (S_d^z), the Kondo spin-flip operator ($S_d^+ S_0^- + \text{h.c.}$) and the Kondo isospin-flip operator ($C_d^+ C_0^- + \text{h.c.}$). If $\varepsilon_m(\psi) \equiv 1 - |\langle m | \psi \rangle|^2$ is the geometric entanglement between a given state $|\psi\rangle$ and the eigenstates of $\hat{M}$, the QFI corresponding to $\hat{M}$ in

the state ψ can be written as

$$F_Q(\psi, \hat{M}) = 4 \left[\frac{1}{2} - \sum_m m^2 \varepsilon_m - \left(\sum_m m \varepsilon_m \right)^2 \right]. \quad (10.13)$$

The total geometric entanglement $\sum_m \varepsilon_m$ is constrained to be $d_M - 1$, where d_M is the dimension of the operator $\hat{M}$. Because each eigenvalue m has a magnitude of at most $1/\sqrt{2}$ (all m^2 must add up to $1/2$), the maximum magnitude of the last two terms in eq. (10.13) (and hence the minimum value of the QFI) is attained for the case when some of the ε_m are zero. However, because the eigenstates are orthogonal, only one of them can have perfect overlap with the state $|\psi\rangle$ and only one ε_m can be zero at a time. The case of minimum QFI therefore corresponds to $\varepsilon_{m^*} = 0, \varepsilon_{m \neq m^*} = 1$, leading to $F_Q = 0$. The opposite situation arises when all the geometric entanglement measures are non-zero and equal, $\varepsilon_m = 1/d_M$, leading to a maximum QFI value of $F_Q = 2/d_M$. A uniformly spread geometric entanglement distribution, therefore, leads to a larger QFI, while a more focused distribution leads to a smaller QFI. This sums up the relation between the quantum Fisher information and geometric entanglement.

If the operator $\hat{M}$ is such that its eigenstates are separable states from the perspective of a certain party, the vanishing QFI arises when $|\psi\rangle$ has zero geometric entanglement with respect to one of the separable states, and the QFI saturates when $|\psi\rangle$ has a finite geometric entanglement with all of the states. A large QFI can thus be associated with more entanglement. Moreover, if the eigenstates of $\hat{M}$ are symmetry-broken states, a vanishing QFI indicates that $|\psi\rangle$ is very close to such a symmetry-broken state, while a large QFI indicates that $|\psi\rangle$ is a uniform superposition of such symmetry-broken states and therefore preserves the symmetry as a whole. A larger QFI points towards the presence of more quantum fluctuations and the lack of susceptibility towards symmetry-breaking.

In order to demonstrate these ideas in the present problem, we computed the QFI in the ground-state of the e-SIAM for a number of Hermitian operators, as a function of the parameter $r = -U_b/J_0$. These are shown in the left panel of Fig. (3.14) (left panel). In order to explain the behaviour depicted in the figure, we point out again that the ground-states for $r \ll r_{c2}$, $r \lesssim r_{c2}$ and $r > r_{c2}$ are $|\text{SS}\rangle + |\text{CT}\rangle$, $|\text{SS}\rangle$ and $|\text{LM}\rangle$ respectively, where $|\text{SS}\rangle$, $|\text{CT}\rangle$ and $|\text{LM}\rangle$ are the spin-singlet state, charge triplet and local moment states respectively. We focus on the spin-flip QFI, corresponding to the operator $\hat{O} = S_d^+ S_0^- + \text{h.c.}$.

- For $r < r_{c1}$, the ground-state involves both $|\text{SS}\rangle$ and $|\text{CT}\rangle$, meaning that neither of the two entanglement measures ε_{SS} or ε_{CT} will be zero. As mentioned in the preceding paragraph, this leads to the maximal QFI, as can be seen in the figure as well.
- At $r = r_{c1}$, the charge triplet content vanishes and the ground-state is purely a singlet beyond that point. As a result, in the range $r_{c1} < r < r_{c2}$, the geometric entanglement corresponding to the singlet state is zero, leading to a minimal and vanishing QFI. This is again shown in the figure.
- Finally, for $r > r_{c2}$, the local moment states become the ground-states, where the bath zeroth site displays all four configurations as a superposition, because of the hopping into the rest of the bath. This state is again not an eigenstate of the spin-flip operator, and the geometric

entanglement will be uniformly distributed across the states, none being zero. This explains the large QFI in the local moment phase.

Similar explanations hold for the single-particle hopping FQI between the impurity and the zeroth site, and the charge isospin FQI between the zeroth site and the first site. In Fig. (3.14) (right panel), we show the QFI for the same three operators very close to the MIT (i.e., for $r \lesssim r_{c2}$). Two important quantities that track the local MIT and act as order parameters for the transition are the degree of compensation for the impurity ($\langle \vec{S}_d \cdot \vec{S}_0 \rangle$) and the impurity magnetisation ($\langle S_d^z \rangle$), and the first two QFI (red and blue curves) in Fig. 3.14 are therefore particularly important because that they quantify the quantum fluctuations present in the system corresponding to these order parameters. Our analysis shows that the two phases on either side of the transition are characterised by distinct values of this pair of QFI: while the QFI corresponding to the degree of compensation zero in the Kondo screened phase, it becomes non-zero in the local moment phase, and the opposite is true for the QFI arising from the impurity magnetisation. The phase precisely at the transition is distinct from those on either side because it displays a non-zero value for both the QFI. While it is expected that a critical point would show enhanced fluctuations of multiple kinds (giving rise to universality), it is enlightening to find that this is also reflected in a measure of many-particle entanglement.

All three QFIs are observed to saturate to finite, non-zero values at the MIT, indicating that the non-Fermi liquid state therein possesses quantum fluctuations of all three varieties.

We will now relate the QFI to some other measures of correlation. The QFI corresponding to an observable can be directly related to the static lesser Greens functions associated with that observable:

$$F_Q(\psi, \hat{O}_2) = 4 \left(\left\langle \hat{O}_2 \hat{O}_1^\dagger \right\rangle \Big|_{\hat{O}_1^\dagger = \hat{O}_2} - \left\langle \hat{O}_2 \hat{O}_1^\dagger \right\rangle^2 \Big|_{\hat{O}_1 = 1} \right) = 4 \left[G_{\hat{O}_2, \hat{O}_2}^<(t \rightarrow \infty) - (G_{1, \hat{O}_2}^<(t \rightarrow \infty))^2 \right] \quad (10.14)$$

In the context of the present work, eq. (10.9) allows relating the QFI to the geometric entanglement measured with respect to the spin-singlet and charge triplet states. By combining eqs. (10.9) and (10.14), we get

$$\begin{aligned} F_Q(\psi, \hat{O}_2) &= 4 \left[(1 - \varepsilon_{ss}) \langle \Phi_{ss} | \hat{O}_2^2 | \Phi_{ss} \rangle + (1 - \varepsilon_{ct}) \langle \Phi_{ct} | \hat{O}_2^2 | \Phi_{ct} \rangle \right. \\ &\quad + \sqrt{1 - \varepsilon_{ss}} \sqrt{1 - \varepsilon_{ct}} \left(\langle \Phi_{ss} | \hat{O}_2^2 | \Phi_{ct} \rangle + \langle \Phi_{ct} | \hat{O}_2^2 | \Phi_{ss} \rangle \right) - \left\{ (1 - \varepsilon_{ss}) \langle \Phi_{ss} | \hat{O}_2 | \Phi_{ss} \rangle \right. \\ &\quad \left. \left. + (1 - \varepsilon_{ct}) \langle \Phi_{ct} | \hat{O}_2 | \Phi_{ct} \rangle + \sqrt{1 - \varepsilon_{ss}} \sqrt{1 - \varepsilon_{ct}} \left(\langle \Phi_{ss} | \hat{O}_2 | \Phi_{ct} \rangle + \langle \Phi_{ct} | \hat{O}_2 | \Phi_{ss} \rangle \right) \right\}^2 \right] \quad (10.15) \end{aligned}$$

Further, the retarded Greens function defined above in eq. (5.6) can very generally be related to the QFI. By using the simplified forms $\frac{i}{2} G_{\hat{O}_2, \hat{O}_2}^R(t \rightarrow \infty) = \langle (\hat{O}_2)^2 \rangle$ and $\frac{i}{2} G_{1, \hat{O}_2}^R(t \rightarrow \infty) = \langle \hat{O}_2 \rangle$, we get

$$F_Q(\psi, \hat{O}_2) = 2i G_{\hat{O}_2, \hat{O}_2}^R(t \rightarrow \infty) + (G_{1, \hat{O}_2}^R(t \rightarrow \infty))^2. \quad (10.16)$$

For the sake of completeness, we generalise the above relations to finite temperatures by adopting the approach of Hauke et al. [Hauke et al.(2016)Hauke, Heyl, Tagliacozzo, and Zoller]. For a mixed




Figure 3.14: Left: Variation of the quantum Fisher information for three separate fluctuations, across the entire range of r . Right: The QFI corresponding to the same three operators, but very close to the QCP at r_{c2} .

state at inverse temperature β characterised by the density matrix $\rho = \sum_n f_n |n\rangle \langle n|$ where $(E_n, |n\rangle)$ are the eigenvalue-eigenstate pairs and $f_n = e^{-\beta E_n} / \sum_n e^{-\beta E_n}$ are the Boltzmann weights, the FQI can be assigned a spectral representation of the form [Hauke et al.(2016)Hauke, Heyl, Tagliacozzo, and Zoller]

$$F_Q(\rho, \hat{O}) = 2 \sum_{m,n} \frac{(f_m - f_n)^2}{f_m + f_n} |\langle m | \hat{O} | n \rangle|^2. \quad (10.17)$$

By using the identity $\frac{f_m - f_n}{f_m + f_n} = \int_{-\infty}^{\infty} d\omega \delta(\omega + E_m - E_n) \tanh(\beta\omega/2)$, the finite temperature FQI can be written in the form

$$F_Q(\rho, \hat{O}) = 2 \int_{-\infty}^{\infty} d\omega \tanh(\beta\omega/2) \sum_{m,n} (f_m - f_n) \delta(\omega + E_m - E_n) |\langle m | \hat{O} | n \rangle|^2. \quad (10.18)$$

In order to relate this with a correlation function, we now point out that the lesser Greens function $G^<$ defined in eq. (10.7), as well as the greater Greens function $G^>$ that we will now define, also admit spectral representations in terms of the eigenstates $|n\rangle$ of the full problem:

$$G_{O^\dagger, O}^<(t) \equiv i \langle \hat{O}^\dagger \hat{O}(t) \rangle \Rightarrow G_{O^\dagger, O}^<(\omega) = 2\pi i \sum_{m,n} f_n \delta(\omega + E_m - E_n) |\langle m | \hat{O}^\dagger | n \rangle|^2 \quad (10.19)$$

$$G_{O^\dagger, O}^>(t) \equiv -i \langle \hat{O}(t) \hat{O}^\dagger \rangle \Rightarrow G_{O^\dagger, O}^>(\omega) = -2\pi i \sum_{m,n} f_m \delta(\omega + E_m - E_n) |\langle m | \hat{O} | n \rangle|^2 \quad (10.20)$$

If $\hat{O}$ is a Hermitian operator, eqs. (10.19) can be used to rewrite the QFI in terms of $G_{O^\dagger, O}^<(\omega)$. Indeed, by combining eqs. (10.19) and eq. (10.18) with the assumption $\hat{O} = \hat{O}^\dagger$, we get

$$F_Q(\rho, \hat{O}) = -\frac{1}{\pi i} \int_{-\infty}^{\infty} d\omega \tanh(\beta\omega/2) (G_{O, O}^>(\omega) + G_{O, O}^<(\omega)). \quad (10.21)$$

This constitutes the general relation between the finite temperature QFI for a general Hermitian many-particle operator $\hat{O}$ and the corresponding many-particle correlations at zero and finite frequencies.

The Kubo linear response function associated with the operator $\hat{O}(t)$ due to a perturbation from the same operator is associated with a correlation function/susceptibility $\mathcal{C}_{\hat{O}}(t, t') = i\theta(t - t') \langle [\hat{O}(t), \hat{O}(t')] \rangle$. By going to the frequency domain, the imaginary part of such a function can be related to the sum of the lesser and greater Greens functions:

$$\begin{aligned} \mathcal{C}_{\hat{O}}(\omega) &= i \int_0^\infty d(t - t') e^{i\omega(t-t')} \left(\langle \hat{O}(t) \hat{O}(t') \rangle - \langle \hat{O}(t') \hat{O}(t) \rangle \right) \\ \Rightarrow \mathcal{C}_{\hat{O}}(\omega) - \mathcal{C}_{\hat{O}}(\omega)^* &= i \int_0^\infty d(t - t') (e^{i\omega(t-t')} - e^{-i\omega(t-t')}) \left(\langle \hat{O}(t) \hat{O}(t') \rangle - \langle \hat{O}(t') \hat{O}(t) \rangle \right) \\ &= i \int_{-\infty}^\infty d(t - t') e^{i\omega(t-t')} \left(\langle \hat{O}(t) \hat{O}(t') \rangle - \langle \hat{O}(t') \hat{O}(t) \rangle \right) \\ &= - (G_{\hat{O}, \hat{O}}^>(\omega) + G_{\hat{O}, \hat{O}}^<(\omega)) \end{aligned} \quad (10.22)$$

This allows us to connect the QFI with the imaginary part of the Kubo correlation function/susceptibility obtained in Ref. [Hauke et al.(2016)Hauke, Heyl, Tagliacozzo, and Zoller]:

$$F_Q(\rho, \hat{O}) = \frac{1}{\pi i} \int_{-\infty}^\infty d\omega \tanh(\beta\omega/2) (\mathcal{C}_{\hat{O}}(\omega) - \mathcal{C}_{\hat{O}}(\omega)^*) . \quad (10.23)$$

3.11 Appendix: One-shot decoupling of impurity from zeroth site

We will decouple the impurity states from the fixed point Hamiltonian using a single URG transformation. This will, of course, generate correlations on the zeroth site. The resulting Hamiltonian will be an AIM with the hopping between the zeroth site and the rest of the chain acting as the effective hybridisation. Schematically, we will have

$$\begin{aligned} H^* = H_{\text{imp}} + V_{\text{imp-0}} + H_0 + H_{0-1} + H_{\text{rest}} &\xrightarrow{\text{decouple imp.}} E_{\text{imp}} + \tilde{H}_0 + H_{0-1} + H_{\text{rest}} \\ &= H_{\text{new imp}} + V_{\text{new imp - rest}} + H_{\text{rest}} . \end{aligned} \quad (11.1)$$

Since the impurity is not coupled with any site beyond the zeroth site, the parts of the Hamiltonian that involve "rest" will not change in the process. This also means that the decoupling can be performed by looking at the smaller Hamiltonian

$$\begin{aligned} H_{\text{imp+0}} = H_{\text{imp}} + H_0 + V_{\text{imp-0}} &= -\frac{U^*}{2} (\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2 - U_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2 + J^* S_d^z S_0^z \\ &+ V^* \sum_{\sigma} (c_{d\sigma}^\dagger c_{0\sigma} + \text{h.c.}) + \frac{J^*}{2} (c_{d\uparrow}^\dagger c_{d\downarrow} c_{0\downarrow}^\dagger c_{0\uparrow} + \text{h.c.}) . \end{aligned} \quad (11.2)$$

3.11.1 Renormalisation from V^*

From the off-diagonal term involving V^* , we generate the following term, in the particle sector for 0:

$$\sum_{\sigma} c_{0\sigma}^\dagger c_{d\sigma} \frac{V^{*2}}{\tilde{\omega} - H_d} c_{d\sigma}^\dagger c_{0\sigma} = \sum_{\sigma} c_{0\sigma}^\dagger c_{d\sigma} \frac{V^{*2}}{\tilde{\omega} + \frac{U^*}{2} (1 - \hat{n}_{d\bar{\sigma}})^2 + U_b \hat{n}_{0\bar{\sigma}} + \frac{J^*}{4} (1 - \hat{n}_{d\bar{\sigma}}) \hat{n}_{0\bar{\sigma}}} c_{d\sigma}^\dagger c_{0\sigma} \quad (11.3)$$

where $\tilde{\omega}$ is the quantum fluctuation operator for the impurity site and $H_d = -\frac{U^*}{2}(\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2 - U_b(\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2 + J^*S_d^z S_0^z$, is the diagonal part of the Hamiltonian for the imp+0 system. In order to resolve the operators in the denominator, we expand the unity in the numerator using the identity $1 = \hat{n}_{d\bar{\sigma}}\hat{n}_{0\bar{\sigma}} + \hat{n}_{d\bar{\sigma}}\hat{h}_{0\bar{\sigma}} + \hat{h}_{d\bar{\sigma}}\hat{n}_{0\bar{\sigma}} + \hat{h}_{d\bar{\sigma}}\hat{h}_{0\bar{\sigma}}$, where $\hat{h} = 1 - \hat{n}$ is hole operator. On Substituting this, we get

$$\begin{aligned} \sum_{\sigma} c_{0\sigma}^{\dagger} c_{d\sigma} \frac{V^{*2}}{\tilde{\omega} - H_d} c_{d\sigma}^{\dagger} c_{0\sigma} &= V^{*2} \sum_{\sigma} c_{0\sigma}^{\dagger} c_{d\sigma} \frac{\hat{h}_{d\bar{\sigma}}\hat{h}_{0\bar{\sigma}} + \hat{h}_{d\bar{\sigma}}\hat{n}_{0\bar{\sigma}} + \hat{n}_{d\bar{\sigma}}\hat{h}_{0\bar{\sigma}} + \hat{n}_{d\bar{\sigma}}\hat{n}_{0\bar{\sigma}}}{\tilde{\omega} + \frac{U^*}{2}\hat{h}_{d\bar{\sigma}} + U_b\hat{n}_{0\bar{\sigma}} + \frac{J^*}{4}\hat{h}_{d\bar{\sigma}}\hat{n}_{0\bar{\sigma}}} c_{d\sigma}^{\dagger} c_{0\sigma} \\ &= V^{*2} \sum_{\sigma} \hat{h}_{d\sigma}\hat{n}_{0\sigma} \left[\frac{\hat{h}_{d\bar{\sigma}}\hat{h}_{0\bar{\sigma}}}{\tilde{\omega}_{00} + \frac{U^*}{2}} + \frac{\hat{h}_{d\bar{\sigma}}\hat{n}_{0\bar{\sigma}}}{\tilde{\omega}_{01} + \frac{U^*}{2} + U_b + \frac{J^*}{4}} + \frac{\hat{n}_{d\bar{\sigma}}\hat{h}_{0\bar{\sigma}}}{\tilde{\omega}_{10}} + \frac{\hat{n}_{d\bar{\sigma}}\hat{n}_{0\bar{\sigma}}}{\tilde{\omega}_{11} + U_b} \right]. \quad (11.4) \end{aligned}$$

$\tilde{\omega}_{(0,1),(0,1)}$ represents the quantum fluctuation scale corresponding to the configuration in the numerator.

In order to "freeze" the impurity dynamics, we will substitute $\hat{n}_{d\sigma} = \hat{n}_{d\bar{\sigma}} = \frac{1}{2}$, because of the Z_2 symmetry and the particle-hole symmetry of the impurity levels. This gives

$$\frac{1}{4} V^{*2} \sum_{\sigma} \hat{n}_{0\sigma} \left[\frac{\hat{h}_{0\bar{\sigma}}}{\tilde{\omega}_{00} + \frac{U^*}{2}} + \frac{\hat{n}_{0\bar{\sigma}}}{\tilde{\omega}_{01} + \frac{U^*}{2} + U_b + \frac{J^*}{4}} + \frac{\hat{h}_{0\bar{\sigma}}}{\tilde{\omega}_{10}} + \frac{\hat{n}_{0\bar{\sigma}}}{\tilde{\omega}_{11} + U_b} \right]. \quad (11.5)$$

The state that most closely represents the metallic ground state is $\tilde{\omega}_{10}$, we take that as the reference quantum fluctuation scale $\bar{\omega}$. It is of the order of $\bar{\omega} \sim -\frac{J^*}{4} - \frac{U^*}{2} - U_b$. We will now relate the other scales to $\bar{\omega}$ by expressing them in terms of the energy of the initial state:

$$\tilde{\omega}_{00} \sim -U_b = \bar{\omega} + \frac{J^*}{4} + \frac{U^*}{2}, \quad \tilde{\omega}_{01} \sim 0 = \bar{\omega} + \frac{J^*}{4} + \frac{U^*}{2} + U_b, \quad \tilde{\omega}_{11} \sim -\frac{U^*}{2} = \bar{\omega} + \frac{J^*}{4} + U_b \quad (11.6)$$

Substituting these gives

$$\frac{1}{4} V^{*2} \sum_{\sigma} \hat{n}_{0\sigma} \left[\hat{h}_{0\bar{\sigma}} \alpha_1 + \hat{n}_{0\bar{\sigma}} \alpha_2 \right], \quad (11.7)$$

where $\alpha_1 = (\bar{\omega} + U^* + \frac{J^*}{4})^{-1} + (\bar{\omega})^{-1}$ and $\alpha_2 = (\bar{\omega} + U^* + 2U_b + \frac{J^*}{2})^{-1} + (\bar{\omega} + 2U_b + \frac{J^*}{4})^{-1}$.

Because of the particle-hole symmetry on the impurity as well as in the bath, the renormalisation from the hole sector is obtained simply by transforming $\hat{n} \leftrightarrow \hat{h}$:

$$\frac{1}{4} V^{*2} \sum_{\sigma} \hat{h}_{0\sigma} \left[\hat{n}_{0\bar{\sigma}} \alpha_1 + \hat{h}_{0\bar{\sigma}} \alpha_2 \right]. \quad (11.8)$$

The total renormalisation arising from V is therefore

$$\frac{1}{4} V^{*2} \sum_{\sigma} \left[\alpha_1 \left(\hat{n}_{0\sigma} \hat{h}_{0\bar{\sigma}} + \hat{h}_{0\sigma} \hat{n}_{0\bar{\sigma}} \right) + \alpha_2 \left(\hat{n}_{0\sigma} \hat{n}_{0\bar{\sigma}} + \hat{h}_{0\sigma} \hat{h}_{0\bar{\sigma}} \right) \right] = \frac{1}{2} V^{*2} (\alpha_1 - \alpha_2) (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2 + \text{constant}. \quad (11.9)$$

3.11.2 Renormalisation from J^*

The renormalisation arising from decoupling the Kondo coupling has two terms. The first term arises when the spin of the zeroth site is initially up:

$$\begin{aligned} \frac{J^{*2}}{4} c_{d\downarrow}^\dagger c_{d\uparrow} c_{0\uparrow}^\dagger c_{0\downarrow} \frac{1}{\tilde{\omega} - H_d} c_{0\downarrow}^\dagger c_{0\uparrow} c_{d\uparrow}^\dagger c_{d\downarrow} &= c_{d\downarrow}^\dagger c_{d\uparrow} c_{0\uparrow}^\dagger c_{0\downarrow} \frac{J^{*2}/4}{\tilde{\omega} + \frac{U^*}{2} + U_b + \frac{J^*}{4}} c_{0\downarrow}^\dagger c_{0\uparrow} c_{d\uparrow}^\dagger c_{d\downarrow} \\ &= \frac{J^{*2}}{4} \frac{\hat{n}_{d\downarrow} \hat{h}_{d\uparrow} \hat{n}_{0\uparrow} \hat{h}_{0\downarrow}}{\tilde{\omega} + \frac{U^*}{2} + U_b + \frac{J^*}{4}}. \end{aligned} \quad (11.10)$$

The quantum fluctuation scale $\tilde{\omega}$ for this process can be similarly related to $\bar{\omega}$: $\tilde{\omega} \sim -\frac{U^*}{2} - U_b - \frac{J^*}{4} = \bar{\omega}$. This gives

$$\frac{J^{*2}}{4} \frac{\hat{n}_{d\downarrow} \hat{h}_{d\uparrow} \hat{n}_{0\uparrow} \hat{h}_{0\downarrow}}{\bar{\omega} + \frac{U^*}{2} + U_b + \frac{J^*}{4}} = \frac{J^{*2}}{16} \frac{\hat{n}_{0\uparrow} \hat{h}_{0\downarrow}}{\bar{\omega} + \frac{U^*}{2} + U_b + \frac{J^*}{4}}, \quad (11.11)$$

as the renormalisation for this configuration. At the final step, we substituted $\hat{n}_{d\sigma} = \hat{h}_{d\sigma} = \frac{1}{2}$. For the other configuration where the spin of the zeroth site is down, we get

$$\frac{J^{*2}}{16} \frac{\hat{n}_{0\downarrow} \hat{h}_{0\uparrow}}{\bar{\omega} + \frac{U^*}{2} + U_b + \frac{J^*}{4}}. \quad (11.12)$$

Adding both sectors, we get

$$\frac{J^{*2}}{16} \frac{1}{\bar{\omega} + \frac{U^*}{2} + U_b + \frac{J^*}{4}} (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2. \quad (11.13)$$

3.11.3 Total renormalisation

Adding the contributions from both V^* and J^* , the net renormalisation is the generation of a local correlation term $-U' (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2$ on the zeroth site, where U' is given by

$$U' = \frac{-J^{*2}/16}{\bar{\omega} + \frac{U^*}{2} + U_b + \frac{J^*}{4}} - \frac{1}{2} V^{*2} \left(\frac{1}{\bar{\omega} + U^* + \frac{J^*}{4}} + \frac{1}{\bar{\omega}} - \frac{1}{\bar{\omega} + U^* + 2U_b + \frac{J^*}{2}} - \frac{1}{\bar{\omega} + 2U_b + \frac{J^*}{4}} \right) \quad (11.14)$$

The effective Hamiltonian for the zeroth site is therefore

$$H_{0+\text{rest}} = \underbrace{-(U' + U_b) (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2}_{\text{new correlated impurity}} - t \underbrace{\sum_{\substack{j \in \text{n.n. of } 0, \\ \sigma}} (c_{0\sigma}^\dagger c_{j\sigma} + \text{h.c.})}_{\text{hopping between new impurity \& new bath}} - t \underbrace{\sum_{\langle i, j \rangle} (c_{i\sigma}^\dagger c_{j\sigma} + \text{h.c.})}_{\text{K.E. of new bath}} \quad (11.15)$$

3.12 Appendix: Effective Hamiltonian for the excitations within the Hubbard sidebands

Beyond r_{c1} , the Hubbard sidebands can be seen to separate from the central Kondo resonance in the impurity spectral function. These sidebands are broad, indicating that the impurity site is hybridising with the bath from within the sidebands, at high energies $\omega \sim U/2$. These processes lead to the

presence of gapless excitations within the rest of the bath, outside the Mott gap. To find the effective Hamiltonian for the rest of the bath (sites 1 and onwards), we will take the eigenstates of the two-site Hamiltonian H that reside within the sideband as our ground-states, and treat the hopping from the zeroth site to the rest of the bath as a perturbation. The two-site Hamiltonian is

$$H_{\text{two site}} = V \sum_{\sigma} \left(c_{d\sigma}^{\dagger} c_{0\sigma} + \text{h.c.} \right) + J \vec{S}_d \cdot \vec{S}_0 - \frac{U}{2} (\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2 + \frac{U}{2}. \quad (12.1)$$

In order to align the doubly-occupied impurity level at the more conventional value of $U/2$, we have shifted the Hamiltonian by the same constant value. The effect of U_b has been ignored because it is small compared to U at the frequencies pertinent to the physics of the sidebands. The two-site states that reside within the Hubbard sidebands are

$$\begin{aligned} |\Psi_{cc}\rangle &\equiv |c\rangle_d |c\rangle_0; \quad c \in \{0, 2\}; \quad |\Psi_{CS}\rangle \equiv \frac{1}{\sqrt{2}} (|2\rangle_d |0\rangle_0 - |0\rangle_d |2\rangle_0); \\ |\Psi_{\sigma,c}\rangle &= \alpha |\sigma\rangle_d |c\rangle_0 + \sqrt{1 - \alpha^2} |c\rangle_d |\sigma\rangle_0; \quad \sigma \in \{\uparrow, \downarrow\}; \\ |\Psi_{CT}\rangle &= \beta \frac{1}{\sqrt{2}} (|\uparrow\rangle_d |\downarrow\rangle_0 - |\downarrow\rangle_d |\uparrow\rangle_0) + \sqrt{1 - \beta^2} \frac{1}{\sqrt{2}} (|2\rangle_d |0\rangle_0 + |0\rangle_d |2\rangle_0), \end{aligned} \quad (12.2)$$

where the coefficients of the eigenstates are given by

$$\alpha \simeq \frac{V}{\sqrt{V^2 + (E_{X1} + \frac{U}{2})^2}}, \quad \beta = \frac{2V}{\sqrt{4V^2 + (E_{CS} + \frac{U}{2} + \frac{3J}{4})^2}}. \quad (12.3)$$

The energies of the states are

$$\begin{aligned} E_{\text{gs}} \equiv E_{00} = E_{22} = E_{CS} &= \frac{U}{2}, \quad E_{X1} \equiv E_{\sigma,c} = \frac{U}{4} + \sqrt{V^2 + \frac{U^2}{16}}, \\ E_{X2} \equiv E_{CS} &= \frac{U}{4} - \frac{3J}{8} + \sqrt{4V^2 + \left(\frac{U}{4} + \frac{3J}{8} \right)^2}. \end{aligned} \quad (12.4)$$

The ground state (comprising the states shown in eq. (12.2)) is triply degenerate. In order to study the dynamics of the bath, we now introduce the perturbation $H_I = -t \sum_{\sigma} (c_{0\sigma}^{\dagger} c_{1\sigma} + \text{h.c.})$. We also take an expanded set of eigenstates $\{|\Psi_n\rangle \otimes |a\rangle\}$, where $\{|\Psi_n\rangle\}$ is the set of eigenstates of $H_{\text{two site}}$, and $|a\rangle \in \{|0\rangle, |\uparrow\rangle, |\downarrow\rangle, |2\rangle\}$ represents the configuration of the first site. The eigenstates of $H_{\text{two site}}$ are also eigenstates the total number operator $\hat{n}_d + \hat{n}_0 + \hat{n}_1$, and since H_I does not conserve the total number, all odd order shifts are zero: $\langle \Psi_n | H_I^b | \Psi_n \rangle$, $b = 1, 3, \dots$, is zero for all the eigenstates.

For the second order shift, we need to calculate the overlaps $\langle \Psi_m | H_I | \Psi_n \rangle$, $m \neq n$ for n in the ground state. Since all three ground-states involve the impurity and zeroth sites in charge configuration (0 or 2), $H_I |\Psi_n\rangle$ will result in states of the form $|0(2)\rangle_d |\sigma\rangle_0 |a\rangle_1$. This resulting state has non-zero overlap only with the part $\sqrt{1 - \alpha^2} |c\rangle_d |\sigma\rangle_0 |a\rangle$ of the expanded state $|\Psi_{\sigma,c}\rangle |a\rangle$. The square of each such overlap is, therefore, $t^2 (1 - \alpha^2)$.

- Excitations from the ground-states $|\Psi_{00}\rangle |0\rangle$ and $|\Psi_{22}\rangle |2\rangle$ have zero overlap with any other state, such that the second order shifts in these states is zero.

- Excitations from the ground-states $|\Psi_{00}\rangle |\sigma\rangle$ and $|\Psi_{22}\rangle |\sigma\rangle$ have non zero overlap with one excited state each, such that the second order shift in these states are $\gamma^2 \equiv t^2 (1 - \alpha^2) / (E_{X1} - E_{gs})$.
- Excitations from the remaining ground states have non-zero overlaps with two excited states each, such that the second order shift in these states are $2\gamma^2$.

Upon dropping the zeroth order part (because it has no dynamics for the rest of the bath), the effective Hamiltonian at second order in t takes the form

$$H_{\text{eff}}^{(2)} = \gamma^2 \left[(|\Psi_{00}\rangle \langle \Psi_{00}| + |\Psi_{22}\rangle \langle \Psi_{22}| + 2 |\Psi_{CS}\rangle \langle \Psi_{CS}|) (\sum_{\sigma} |\sigma\rangle_1 \langle \sigma|_1) + 2 (|\Psi_{22}\rangle \langle \Psi_{22}| + |\Psi_{CS}\rangle \langle \Psi_{CS}|) |0\rangle_1 \langle 0|_1 + 2 (|\Psi_{00}\rangle \langle \Psi_{00}| + |\Psi_{CS}\rangle \langle \Psi_{CS}|) |2\rangle_1 \langle 2|_1 \right] \quad (12.5)$$

By using the completeness relation $1 = \sum_{\sigma} |\sigma\rangle_1 \langle \sigma|_1 + |0\rangle_1 \langle 0|_1 + |2\rangle_1 \langle 2|_1$ of the Hilbert space of the first site, this evaluates to

$$H_{\text{eff}}^{(2)} = \text{constant} + \gamma^2 (|\Psi_{22}\rangle \langle \Psi_{22}| - |\Psi_{00}\rangle \langle \Psi_{00}|) (|2\rangle_1 \langle 2|_1 - |0\rangle_1 \langle 0|_1), \quad (12.6)$$

where the "constant" term refers to operators that do not depend on the dynamics of the rest of the bath. By defining the charge isospin operator $C^z |2(0)\rangle = +(-)\frac{1}{2} |2(0)\rangle$, the effective Hamiltonian can be written as (dropping the non-dynamical part)

$$H_{\text{eff}}^{(2)} = 4\gamma^2 C_{\text{tot}}^z C_1^z, \quad (12.7)$$

where C_i^z is the charge isospin for site i and $C_{\text{tot}}^z \equiv C_d^z + C_0^z$ is the z-component of the total charge isospin for the impurity and the zeroth sites.

The fourth order shift in the energy of the eigenstate $|\Psi_n\rangle$ will be given by

$$E_n^{(4)} = \sum_{j,k,l} \frac{(H_I)_{nl} (H_I)_{lk} (H_I)_{kj} (H_I)_{jn}}{(E_l - E_n) (E_k - E_n) (E_j - E_n)} - E_n^{(2)} \left[\frac{(H_I)_{kn}}{E_k - E_n} \right]^2, \quad (12.8)$$

where the matrix elements $(H_I)_{in}$ are defined as $\langle i|H_I|n\rangle$. This fourth-order shift vanishes for almost all the ground-states. For the remaining states, it becomes important to treat their degeneracy. The "appropriate" combinations of states that provide non-zero fourth order contribution are

$$|+\rangle = \sqrt{\frac{2}{3}} |\Psi_{22}\rangle |0\rangle_1 + \sqrt{\frac{1}{3}} |\Psi_{CS}\rangle |2\rangle_1, \quad |-\rangle = -\sqrt{\frac{2}{3}} |\Psi_{00}\rangle |2\rangle_1 + \sqrt{\frac{1}{3}} |\Psi_{CS}\rangle |0\rangle_1. \quad (12.9)$$

Corresponding to these ground states, the fourth-order shift in the eigenvalue is

$$E^{(4)} = \frac{3\gamma^4 \alpha^2 \beta^2}{(1 - \alpha^2) (E_{X2} - E_{gs})}, \quad (12.10)$$

such that the fourth-order contribution to the effective Hamiltonian is

$$H_{\text{eff}}^{(4)} = E^{(4)} (|+\rangle \langle +| + |-\rangle \langle -|) = \frac{\gamma^4 \alpha^2 \beta^2}{(1 - \alpha^2) (E_{X2} - E_{gs})} [-C_{\text{tot}}^z C_{\text{tot}}^2 C_1^z + \sqrt{2} \mathcal{P}_{\text{tot}}^4 (C_0^+ - C_d^+) C_1^- + \text{h.c.}] . \quad (12.11)$$

The operator $\mathcal{P}_{\text{tot}}^4$ acts on states belonging to the combined $d + 0$ Hilbert space and projects on to the $\hat{n}_d + \hat{n}_0 = 4$ subspace.

3.13 Appendix: Behaviour of the Kondo scale T_K close to the transition

Towards obtaining the emergent Kondo temperature scale T_K close to the transition, we follow the approach used by Moeller et al. 1995 [Moeller et al.(1995)Moeller, Si, Kotliar, Rozenberg, and Fisher] and Held et al. 2013 [Held et al.(2013)Held, Peters, and Toschi]. Near the transition, they obtained a Kondo model from the full AIM by applying a Schrieffer-Wolff transition that removes the charge fluctuations of the impurity site and retains the physics of only the s-d coupling J . This amounts to removing the side peaks from the impurity spectral function and focusing on the low-energy central peak. Held et al. then integrated the RG equation for this Kondo model by using a Lorentzian DOS in the bath. This is motivated by the fact that the central peak of the impurity spectral function is a Lorentzian, and the bath becomes equivalent to the impurity site under self-consistency. We implement the same approach but on our extended impurity model, such that the transformation then leads to a $J - U_b$ model. The relevant RG equations are mentioned in Section IV of the main manuscript:

$$\Delta J = -\frac{n_j J (J + 4U_b)}{\omega - D/2 + U_b/2 + J/4}, \quad \Delta U_b = 0, \quad (13.1)$$

where $n_j = \rho(D)|\Delta D|$ is proportional to the bath density of states. In Section VII of the main manuscript, we show that the spectral function of the zeroth site is tracked by that on the impurity site through the coherent spin-flip fluctuations, and this shows that the appropriate density of states for the bath in this regime is of the Lorentzian kind (same as the shape of the central peak of the impurity spectral function). We therefore set

$$\rho(D) = \frac{\rho_0 \Gamma^2}{D^2 + \Gamma^2}, \quad (13.2)$$

where Γ represents the half-width of the Lorentzian. A similar line of arguments was presented by Held et al. [Held et al.(2013)Held, Peters, and Toschi]. Close to the transition, the RG flow of the Kondo coupling J is stunted by the competing pairing term, and the RG equation can be simplified into the Poor man's scaling 1-loop form:

$$\frac{dJ}{dD} \simeq -2\rho_0 \Gamma^2 \frac{J (J + 4U_b)}{(\Gamma^2 + D^2) D}. \quad (13.3)$$

We now integrate this RG equation over the range $D \in [D_0, D^*]$, where D_0 is the bare bandwidth, and D^* is the fixed-point bandwidth. D^* represents the energy scale for the conduction electrons in the IR.

$$\begin{aligned} \int_{J_0}^{J^*} \frac{dJ}{J (J + 4U_b)} &= - \int_{D_0}^{D^*} \frac{2\rho_0 \Gamma^2 dD}{(\Gamma^2 + D^2) D} \\ \Rightarrow \frac{1}{4U_b \rho_0} \left(\ln \frac{J^*}{J_0} - \ln \frac{J^* + 4U_b}{J_0 + 4U_b} \right) &= \ln \frac{D^* + \Gamma^2}{D_0 + \Gamma^2} - \ln \frac{D^*}{D_0}. \end{aligned} \quad (13.4)$$

We will now take the limit of $J_0 + 4U_b \rightarrow 0^-$ and *equate the singular terms on both sides* of the equation:

$$\ln \frac{D^*}{D_0} = -\frac{1}{4U_b\rho_0} \ln \frac{J_0 + 4U_b}{J_0} = -\frac{1}{4U_b\rho_0} \ln (r_{c2} - r) + \text{non-singular part} . \quad (13.5)$$

We have replaced the couplings J and U_b with the dimensionless parameters $r = -U_b/J_0$ and $r_{c2} = 1/4$. Dropping the non-singular part and solving for D^* then gives the IR energy scale:

$$D^* = D_0 \exp \left[-\frac{\ln (r_{c2} - r)}{4U_b\rho_0} \right] . \quad (13.6)$$

The dynamically generated Kondo temperature scale is then obtained as

$$T_K = \frac{D^*}{k_B} = \frac{D_0}{k_B} \exp \left[-\frac{\ln (r_{c2} - r)}{4U_b\rho_0} \right] . \quad (13.7)$$

Note that the prefactor of the logarithm is positive because U_b is negative: $-4U_b\rho_0 = |4U_b\rho_0|$. As we approach the transition, the parameter r takes the limit $r \rightarrow r_{c2}^-$, and the Kondo temperature scale vanishes:

$$\lim_{r \rightarrow r_{c2}^-} T_K = \frac{D_0}{k_B} \lim_{r \rightarrow r_{c2}^-} (r_{c2} - r)^{4|U_b\rho_0|} \rightarrow 0 . \quad (13.8)$$

3.14 Appendix: Effective Hamiltonian for the low-lying excitations at the critical point

In order to obtain an effective Hamiltonian for the low-lying excitations in the bath, we will introduce a hopping term V between the zeroth site and the first site into the two-site Hamiltonian H_0 , and treat that term perturbatively.

$$H = \underbrace{J\vec{S}_d \cdot \vec{S}_0 - U_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2}_{H_0} - \underbrace{t \sum_{\sigma} (c_{0\sigma}^\dagger c_{1\sigma} + \text{h.c.})}_{V} . \quad (14.1)$$

As mentioned before, H_0 is our zeroth Hamiltonian, while V is the perturbation. At the critical point, the zeroth Hamiltonian has a 20-fold degenerate ground-state manifold, and we will show later that it is this degeneracy that leads to NFL excitations. The ground-states are

$$\{|S = S^z = 0\rangle, |\uparrow, 0\rangle, |\uparrow, 2\rangle, |\downarrow, 0\rangle, |\downarrow, 2\rangle\} \otimes \{|0\rangle, |\uparrow\rangle, |\downarrow\rangle, |\uparrow\downarrow\rangle\} . \quad (14.2)$$

The ket to the left of $\otimes$ describes the configuration of the impurity and zeroth sites; $S = S^z = 0$ represents the singlet, while the remaining four states $|\sigma, 0\rangle, |\sigma, 2\rangle$ are local moment states. The ket to the right of $\otimes$ represents the configuration of the first site. Eight of these states have zero matrix elements within the ground-state subspace, even in the presence of the perturbation, so

we drop these states. Since the full Hamiltonian conserves the total number of particles $N_{\text{tot}} = \sum_{\sigma} (\hat{n}_{d\sigma} + \hat{n}_{0\sigma} + \hat{n}_{1\sigma})$ and the total spin $S_{\text{tot}}^z = S_d^z + S_0^z + S_1^z$, the remaining 12 states can be split into decoupled blocks based on the values of these two operators:

$$\begin{aligned} N_{\text{tot}} = c + 2, S_{\text{tot}}^z = 0 & : |S = S^z = 0\rangle |c\rangle, |\uparrow, c, \downarrow\rangle, |\downarrow, c, \uparrow\rangle; c = 0, 2, \\ N_{\text{tot}} = 3, S_{\text{tot}}^z = \pm \frac{1}{2} & : |S = S^z = 0\rangle |\sigma\rangle, |\sigma, 0, 2\rangle, |\sigma, 2, 0\rangle; \sigma = \uparrow, \downarrow. \end{aligned} \quad (14.3)$$

The perturbation V will lift the degeneracy in each block; to calculate the modified spectrum, we diagonalise each block in the presence of V . We will first write down the action of the perturbation on the singlet ground-states: this will make it easier to identify the true eigenstates.

$$\begin{aligned} N_{\text{tot}} = c + 2, S_{\text{tot}}^z = 0 & : V |\text{SS}_c\rangle = \frac{t}{\sqrt{2}} (|\downarrow, c, \uparrow\rangle - |\uparrow, c, \downarrow\rangle), c = 0, 2, \\ N_{\text{tot}} = 3, S_{\text{tot}}^z = \frac{\sigma}{2} & : V |\text{SS}_{\sigma}\rangle = \frac{t}{\sqrt{2}} |\sigma\rangle \otimes (|0, 2\rangle + |2, 0\rangle), \sigma = \uparrow (1), \downarrow (-1), \end{aligned} \quad (14.4)$$

where we have introduced the notation $|\text{SS}_{\alpha}\rangle = |S = S^z = 0\rangle \otimes |\alpha\rangle, \alpha = 0, \uparrow, \downarrow, 2$. This shows that the action of the perturbation on the singlet state is to create linear combinations of the local moment states in the respective blocks. We therefore define the two possible linear combinations $|\text{LM}_{\pm}^{N_{\text{tot}}, S_{\text{tot}}^z}\rangle$ within each block:

$$\begin{aligned} N_{\text{tot}} = c + 2, S_{\text{tot}}^z = 0 & : |\text{LM}_{\pm}^{c+2,0}\rangle = \frac{1}{\sqrt{2}} (|\uparrow, c, \downarrow\rangle \pm |\downarrow, c, \uparrow\rangle), c = 0, 2, \\ N_{\text{tot}} = 3, S_{\text{tot}}^z = \frac{\sigma}{2} & : |\text{LM}_{\sigma}^{3, \frac{1}{2}}\rangle = \frac{1}{\sqrt{2}} (|\sigma, 0, 2\rangle \pm |\sigma, 2, 0\rangle), \sigma = \uparrow (1), \downarrow (-1). \end{aligned} \quad (14.5)$$

In terms of these linear combinations, we then get

$$\begin{aligned} V |\text{SS}_0\rangle &= t |\text{LM}_{-}^{2,0}\rangle, \quad V |\text{LM}_{-}^{2,0}\rangle = t |\text{SS}_0\rangle, \quad V |\text{LM}_{+}^{2,0}\rangle = 0; \quad V |\text{SS}_{\uparrow}\rangle = t |\text{LM}_{+}^{3,\uparrow}\rangle, \\ V |\text{LM}_{+}^{3,\uparrow}\rangle &= t |\text{SS}_{\uparrow}\rangle, \quad V |\text{LM}_{-}^{3,\uparrow}\rangle = 0, \quad V |\text{SS}_{\downarrow}\rangle = t |\text{LM}_{+}^{3,\downarrow}\rangle, \quad V |\text{LM}_{+}^{3,\downarrow}\rangle = t |\text{SS}_{\downarrow}\rangle, \\ V |\text{LM}_{-}^{3,\downarrow}\rangle &= 0; \quad V |\text{SS}_2\rangle = t |\text{LM}_{-}^{4,0}\rangle, \quad V |\text{LM}_{-}^{4,0}\rangle = t |\text{SS}_2\rangle, \quad V |\text{LM}_{+}^{4,0}\rangle = 0. \end{aligned} \quad (14.6)$$

The zero matrix elements at the far right of each line should be understood as the fact that V takes those states out of the ground-state subspace. For e.g., the state $V |\text{LM}_{-}^{2,0}\rangle$ is not really zero, but the overlap of $V |\text{LM}_{-}^{2,0}\rangle$ with any other state in the ground-state subspace spanned by the 20 degenerate states is zero. We now see that each of the 3×3 blocks decouples into a 2×2 block $\begin{pmatrix} 0 & t \\ t & 0 \end{pmatrix}$ and a 1×1 block (0) when written in terms of the linear combinations $|\text{LM}_{\pm}\rangle$. The state in the final block is therefore already an eigenstate, but with zero energy, so that will drop out from the effective Hamiltonian. The remaining 2×2 block has eigenstates $|\pm N_{\text{tot}}, S_{\text{tot}}^z\rangle$ that are linear combinations of

the singlet states and the surviving local moment combination, with eigenvalues $\pm t$:

$$\begin{aligned}
 N_{\text{tot}} = 2, S_{\text{tot}}^z = 0 : |\pm^{2,0}\rangle &= \frac{1}{\sqrt{2}} (|\text{SS}_0\rangle \pm |\text{LM}_{-}^{2,0}\rangle), \\
 N_{\text{tot}} = 3, S_{\text{tot}}^z = \frac{1}{2} : |\pm^{3,\frac{1}{2}}\rangle &= \frac{1}{\sqrt{2}} (|\text{SS}_{\uparrow}\rangle \pm |\text{LM}_{+}^{3,\frac{1}{2}}\rangle), \\
 N_{\text{tot}} = 3, S_{\text{tot}}^z = -\frac{1}{2} : |\pm^{3,-\frac{1}{2}}\rangle &= \frac{1}{\sqrt{2}} (|\text{SS}_{\downarrow}\rangle \pm |\text{LM}_{+}^{3,-\frac{1}{2}}\rangle), \\
 N_{\text{tot}} = 4, S_{\text{tot}}^z = 0 : |\pm^{4,0}\rangle &= \frac{1}{\sqrt{2}} (|\text{SS}_2\rangle \pm |\text{LM}_{-}^{4,0}\rangle). \tag{14.7}
 \end{aligned}$$

At this point, it is worth noting that the polarised ground-states $|\text{SS}_{\sigma}^z = \sigma \frac{1}{2}\rangle$, $\sigma = \pm 1$ can be written as an equal superposition of the singlet states $|\text{SS}\rangle_{a0} \otimes |\sigma\rangle_1$ and the local moment states $\frac{1}{\sqrt{2}} (|\sigma, 0, 2\rangle - |\sigma, 2, 0\rangle)$. The polarised symmetry-broken states therefore act as a bridge between the ground states of the two phases and are important in displaying the breakdown of the Kondo cloud at the QCP and the consequences arising from it (like inexact screening of the impurity and fractional magnetisation and entanglement entropy, which will be discussed in the next section).

By combining the two eigenstates $|\pm\rangle$ for each block, one can then reconstruct the effective Hamiltonian $\mathcal{H}_{\text{eff}}^{N_{\text{tot}}, S_{\text{tot}}^z}$ that describes the dynamics of the first site from within the ground-state subspace of the zeroth Hamiltonian. Formally, this is written as

$$\begin{aligned}
 \mathcal{H}_{\text{eff}}^{N_{\text{tot}}, S_{\text{tot}}^z} &= t |+\rangle^{N_{\text{tot}}, S_{\text{tot}}^z} \langle +|^{N_{\text{tot}}, S_{\text{tot}}^z} - t |-\rangle^{N_{\text{tot}}, S_{\text{tot}}^z} \langle -|^{N_{\text{tot}}, S_{\text{tot}}^z}, \\
 &= \begin{cases} t \left[|\text{SS}_{\alpha}\rangle \langle \text{LM}_{-}^{N_{\text{tot}}, S_{\text{tot}}^z}| + |\text{LM}_{-}^{N_{\text{tot}}, S_{\text{tot}}^z}\rangle \langle \text{SS}_{\alpha}| \right], & \alpha = 0, 2, \\ t \left[|\text{SS}_{\alpha}\rangle \langle \text{LM}_{+}^{N_{\text{tot}}, S_{\text{tot}}^z}| + |\text{LM}_{+}^{N_{\text{tot}}, S_{\text{tot}}^z}\rangle \langle \text{SS}_{\alpha}| \right], & \alpha = \uparrow, \downarrow. \end{cases} \tag{14.8}
 \end{aligned}$$

We first write down the effective Hamiltonian for $\alpha = 0$ which corresponds to $N_{\text{tot}} = 2, S_{\text{tot}}^z = 0$:

$$\mathcal{H}_{\text{eff}}^{2,0} = t [|\text{SS}_0\rangle \langle \text{LM}_{-}^{2,0}| + |\text{LM}_{-}^{2,0}\rangle \langle \text{SS}_0|]. \tag{14.9}$$

In order to convert this into a second-quantised form, we will use the following identities:

$$|\text{SS}_0\rangle = \frac{1}{\sqrt{2}} (P_d^{\uparrow} c_{0\downarrow}^{\dagger} - P_d^{\downarrow} c_{0\uparrow}^{\dagger}) P_0^{(0)} P_1^{(0)} |0\rangle; \quad |\text{LM}_{-}^{2,0}\rangle = \frac{1}{\sqrt{2}} (P_d^{\uparrow} c_{1\downarrow}^{\dagger} - P_d^{\downarrow} c_{1\uparrow}^{\dagger}) P_0^{(0)} P_1^{(0)} |0\rangle, \tag{14.10}$$

where $P_i^{(m)}$ ($m = 0, 1, 2$) projects onto the $\hat{n}_i = m$ subspace on the i^{th} site, and P_d^{σ} projects the impurity spin into $|\sigma\rangle$; $\sigma = \uparrow, \downarrow$ configuration. We then have

$$\begin{aligned}
 |\text{SS}_0\rangle \langle \text{LM}_{-}^{2,0}| &= \frac{1}{2} (P_d^{\uparrow} c_{0\downarrow}^{\dagger} - P_d^{\downarrow} c_{0\uparrow}^{\dagger}) P_0^{(0)} P_1^{(0)} (P_d^{\uparrow} c_{1\downarrow} - P_d^{\downarrow} c_{1\uparrow}) \\
 &= P_1^{(0)} \frac{1}{2} (P_d^{\uparrow} c_{0\downarrow}^{\dagger} c_{1\downarrow} + P_d^{\downarrow} c_{0\uparrow}^{\dagger} c_{1\uparrow} - S_d^{+} c_{0\downarrow}^{\dagger} c_{1\uparrow} - S_d^{-} c_{0\uparrow}^{\dagger} c_{1\downarrow}) P_0^{(0)}, \tag{14.11}
 \end{aligned}$$

where we have used $P^2 = P$ if P is a projector, and that $P_d^{\uparrow} P_d^{\downarrow} = S_d^{+}$. Substituting these into the effective Hamiltonian and recognising that $P_d^{\uparrow} = \frac{1}{2} + S_d^z$, $P_d^{\downarrow} = \frac{1}{2} - S_d^z$ gives

$$\begin{aligned}
 \mathcal{H}_{\text{eff}}^{2,0} &= \frac{t}{2} \left[\left(\frac{1}{2} + S_d^z \right) c_{0\downarrow}^{\dagger} P_0^{(0)} P_1^{(0)} c_{1\downarrow} - S_d^{+} c_{0\downarrow}^{\dagger} P_0^{(0)} P_1^{(0)} c_{1\uparrow} - S_d^{-} c_{0\uparrow}^{\dagger} P_0^{(0)} P_1^{(0)} c_{1\downarrow} \right. \\
 &\quad \left. + \left(\frac{1}{2} - S_d^z \right) c_{0\uparrow}^{\dagger} P_0^{(0)} P_1^{(0)} c_{1\uparrow} \right] + \text{h.c.} . \tag{14.12}
 \end{aligned}$$

In order to make this more illuminating, we define the holon excitation operators $h_{i\sigma}^\dagger = c_{i\sigma}^\dagger P_i^{(0)}$. In terms of these operators, the effective Hamiltonian becomes

$$\begin{aligned} \mathcal{H}_{\text{eff}}^{2,0} &= \frac{t}{2} \left[\left(\frac{1}{2} + S_d^z \right) \left(h_{0\downarrow}^\dagger h_{1\downarrow} + h_{1\downarrow}^\dagger h_{0\downarrow} \right) - S_d^+ \left(h_{0\downarrow}^\dagger h_{1\uparrow} + h_{1\downarrow}^\dagger h_{0\uparrow} \right) - S_d^- \left(h_{0\uparrow}^\dagger h_{1\downarrow} + h_{1\uparrow}^\dagger h_{0\downarrow} \right) + \right. \\ &\quad \left. \left(\frac{1}{2} - S_d^z \right) \left(h_{0\uparrow}^\dagger h_{1\uparrow} + h_{1\uparrow}^\dagger h_{0\uparrow} \right) \right] \\ &= -t \left[S_d^z \frac{1}{2} \left(h_{0\uparrow}^\dagger h_{1\uparrow} + h_{1\uparrow}^\dagger h_{0\uparrow} - h_{0\downarrow}^\dagger h_{1\downarrow} - h_{1\downarrow}^\dagger h_{0\downarrow} \right) + \frac{1}{2} S_d^+ \left(h_{0\downarrow}^\dagger h_{1\uparrow} + h_{1\downarrow}^\dagger h_{0\uparrow} \right) \right. \\ &\quad \left. + \frac{1}{2} S_d^- \left(h_{0\uparrow}^\dagger h_{1\downarrow} + h_{1\uparrow}^\dagger h_{0\downarrow} \right) \right] + \frac{t}{4} \left(h_{0\uparrow}^\dagger h_{1\uparrow} + h_{1\uparrow}^\dagger h_{0\uparrow} + h_{0\downarrow}^\dagger h_{1\downarrow} + h_{1\downarrow}^\dagger h_{0\downarrow} \right). \end{aligned} \quad (14.13)$$

The first part of the Hamiltonian (with a $-t$ factor in front) is reminiscent of a two-spin Heisenberg Hamiltonian. To investigate this, we write down the potential second spin:

$$\mathcal{S}_1^z \equiv \frac{1}{2} \left(h_{0\uparrow}^\dagger h_{1\uparrow} + h_{1\uparrow}^\dagger h_{0\uparrow} - h_{0\downarrow}^\dagger h_{1\downarrow} - h_{1\downarrow}^\dagger h_{0\downarrow} \right). \quad (14.14)$$

Note that each term that comprises the full operator $\mathcal{S}_1^z$ has a combined projector $P_{01}^{(0)} = P_0^{(0)} P_1^{(0)} = P_1^{(0)} P_0^{(0)}$ in the middle with a single annihilation operator to the right. This means that $\mathcal{S}^z$ projects onto the smaller set of only those states that satisfy $\hat{n}_0 + \hat{n}_1 = 1$. That is also the meaning of the subscript 1 in $\mathcal{S}_1^z$. There are four such states: $|\uparrow, 0\rangle, |\downarrow, 0\rangle, |0, \uparrow\rangle, |0, \downarrow\rangle$. In fact, it is easy to see that the eigenstates of $\mathcal{S}^z$ are

$$|\uparrow\rangle_{1,\pm} = \frac{1}{\sqrt{2}} (|\uparrow, 0\rangle \pm |0, \uparrow\rangle), \quad \mathcal{S}_1^z |\uparrow\rangle_{1,\pm} = \pm \frac{1}{2} |\uparrow\rangle_{1,\pm}, \quad (14.15)$$

$$|\downarrow\rangle_{1,\pm} = \frac{1}{\sqrt{2}} (|\downarrow, 0\rangle \pm |0, \downarrow\rangle), \quad \mathcal{S}_1^z |\downarrow\rangle_{1,\pm} = \mp \frac{1}{2} |\downarrow\rangle_{1,\pm}. \quad (14.16)$$

The label $\pm$ on top of the eigenstates indicates that these are also eigenstates of the parity operator $\mathcal{P}_{01}$ that switches the site labels 0 and 1:

$$\mathcal{P}_{01} |\alpha, \alpha'\rangle = |\alpha', \alpha\rangle; \quad \alpha, \alpha' \in \{0, \uparrow, \downarrow, 2\}. \quad (14.17)$$

In fact, the eigenstates $|\uparrow\rangle_{1,+1}$ and $|\downarrow\rangle_{1,-1}$ both have $\mathcal{S}^z = \frac{1}{2}$, but are orthogonal because they have different parities. Analogous to the $\mathcal{S}_1^z$ operator, we can also read off the potential $\mathcal{S}_1^\pm$ ladder operators from the effective Hamiltonian:

$$\mathcal{S}_1^+ = h_{0\uparrow}^\dagger h_{1\downarrow} + h_{1\uparrow}^\dagger h_{0\downarrow}, \quad \mathcal{S}_1^- = (\mathcal{S}_1^+)^\dagger. \quad (14.18)$$

Applying these ladder operators on the eigenstates of $\mathcal{S}_1^z$ gives

$$\mathcal{S}_1^+ |\downarrow\rangle_{1,\pm} = \pm |\uparrow\rangle_{1,\pm}; \quad \mathcal{S}_1^- |\uparrow\rangle_{1,\pm} = \pm |\downarrow\rangle_{1,\pm}; \quad \mathcal{S}_1^+ |\uparrow\rangle_{1,\pm} = \mathcal{S}_1^- |\downarrow\rangle_{1,\pm} = 0. \quad (14.19)$$

This makes it clear that the pair of states $|\uparrow\rangle_1^+$ and $|\downarrow\rangle_1^+$ form a spin-half doublet, as does the other pair $|\uparrow\rangle_1^-$ and $|\downarrow\rangle_1^-$. We can define new spin operators that act only on a single parity subspace:

$$\mathcal{S}_{1,\pm}^a = \pm \mathcal{S}^a \frac{1}{2} (1 \pm \mathcal{P}_{01}); \quad a = x, y, z. \quad (14.20)$$

The operator $\frac{1}{2}(1 \pm \mathcal{P}_{01})$ projects on to the subspace with ± 1 parity. In the notation $\mathcal{S}_{1,\pm}^a$, the label $\pm$ indicates the parity sector to which the operator applies, while 1 indicates that we are operating in the subspace with $\hat{n}_0 + \hat{n}_1 = 1$. Two operators corresponding to different parity subspaces will commute because they act on disjoint Hilbert spaces: $[\mathcal{S}_{+1}^a, \mathcal{S}_{-1}^b] = 0$. In terms of these projected spin operators, we recover the usual spin-half SU(2) algebra:

$$\begin{aligned} \mathcal{S}_{1,\pm}^z |\uparrow\rangle_{1,\pm} &= \frac{1}{2} |\uparrow\rangle_{1,\pm}; & \mathcal{S}_{1,\pm}^z |\downarrow\rangle_{1,\pm} &= -\frac{1}{2} |\downarrow\rangle_{1,\pm}; \\ \mathcal{S}_{1,\pm}^+ |\downarrow\rangle_{1,\pm} &= |\uparrow\rangle_{1,\pm}; & \mathcal{S}_{1,\pm}^- |\uparrow\rangle_{1,\pm} &= |\downarrow\rangle_{1,\pm}. \end{aligned} \quad (14.21)$$

The part of the effective Hamiltonian that involves the impurity can now be written in terms of these new spin operators:

$$t \left(\vec{S}_d \cdot \vec{S}_{-1} - \vec{S}_d \cdot \vec{S}_{+1} \right); \quad \vec{S}_{\pm} = (\mathcal{S}_{\pm}^x \quad \mathcal{S}_{\pm}^y \quad \mathcal{S}_{\pm}^z). \quad (14.22)$$

The last term in eq. (14.13) (with a prefactor of $\frac{t}{4}$) can now be identified as the parity operator $\mathcal{P}_{01}$ acting within the subspace $\hat{n}_0 + \hat{n}_1 = 1$, because it has eigenstates $|\uparrow, 0\rangle \pm |0, \uparrow\rangle$ and $|\downarrow, 0\rangle \pm |0, \downarrow\rangle$ with eigenvalues ± 1 . The full effective Hamiltonian in the $N_{\text{tot}} = 2$ subspace then takes the compact form

$$\mathcal{H}_{\text{eff}}^{2,0} = t \left(\vec{S}_d \cdot \vec{S}_{-1} - \vec{S}_d \cdot \vec{S}_{+1} \right) + \frac{t}{4} \mathcal{P}_{01} P_{01}^{(1)}, \quad (14.23)$$

where $P_{01}^{(1)} = P_0^{(1)} P_1^{(0)} + P_0^{(0)} P_1^{(1)}$ projects on to the $\hat{n}_0 + \hat{n}_1 = 1$ subspace. In light of these new spin operators, the original eigenstates $|\pm^{2,0}\rangle$ can be written in terms of these emergent spins $|\uparrow\rangle_{1,\pm}$ and $|\downarrow\rangle_{1,\pm}$:

$$\begin{aligned} |\pm^{2,0}\rangle &= \frac{1}{2} (|\uparrow, \downarrow, 0\rangle - |\downarrow, \uparrow, 0\rangle \pm |\uparrow, 0, \downarrow\rangle \mp |\downarrow, 0, \uparrow\rangle) = \frac{1}{2} [|\uparrow\rangle (|\downarrow, 0\rangle \pm |0, \downarrow\rangle) - |\downarrow\rangle (|\uparrow, 0\rangle \pm |0, \uparrow\rangle)] \\ &= \frac{1}{\sqrt{2}} \left(|\uparrow\rangle |\downarrow\rangle_{1,\pm} - |\downarrow\rangle |\uparrow\rangle_{1,\pm} \right). \end{aligned} \quad (14.24)$$

Both the ground and excited states are therefore spin-singlet states, but they belong to the negative and positive parity sectors respectively.

We now proceed to the $N_{\text{tot}} = 4$ sector, which is equivalent to $\alpha = 2$ and $\hat{n}_0 + \hat{n}_1 = 3$. From eq. (14.8), we note that the effective Hamiltonian of the $\alpha = 2$ sector is obtained from that of the $\alpha = 0$ sector by replacing the holons with doublons, $|0\rangle \rightarrow |2\rangle$. This allows us to use the results of the $N_{\text{tot}} = 2$ sector simply by making the above-mentioned holon-doublon transformation. Accordingly, the composite up and down spins of the $0 + 1$ -site system become

$$|\uparrow\rangle_{3,\pm} = \frac{1}{\sqrt{2}} (|\uparrow, 2\rangle \pm |2, \uparrow\rangle), \quad |\downarrow\rangle_{3,\pm} = \frac{1}{\sqrt{2}} (|\downarrow, 2\rangle \pm |2, \downarrow\rangle), \quad (14.25)$$

which then allow us to define spin operators for the positive and negative parity sectors:

$$\mathcal{S}_{\pm}^z = \frac{1}{2} \left(|\uparrow\rangle_{3,\pm} - |\downarrow\rangle_{3,\pm} \right), \quad \mathcal{S}_{\pm}^+ = |\uparrow\rangle_{3,\pm} \langle \downarrow|_{3,\pm}, \quad (14.26)$$

analogous to those of the $N_{\text{tot}} = 2$ sector in eqs. (14.15) and (14.18). These definitions then allow us to rewrite the effective Hamiltonian in terms of the eigenstates of $\mathcal{S}_{\pm}^z$, and hence in terms of the spin operators $\vec{\mathcal{S}}_{3,\pm}$. The eigenstates $|\pm^{4,0}\rangle$ and the effective Hamiltonian $H_{\text{eff}}^{4,0}$ for this sector can be written as

$$|\pm^{4,0}\rangle = \frac{1}{\sqrt{2}} \left(|\uparrow\rangle_d |\downarrow\rangle_{3,\pm} - |\downarrow\rangle_d |\uparrow\rangle_{3,\pm} \right). \quad (14.27)$$

The effective Hamiltonian can then be written in terms of these emergent spins

$$H_{\text{eff}}^{4,0} = \frac{t}{2} \left[\left(|\uparrow\rangle_d |\downarrow\rangle_{3,+} - |\downarrow\rangle_d |\uparrow\rangle_{3,+} \right) \left(\langle\uparrow|_d \langle\downarrow|_{3,+} - \langle\downarrow|_d \langle\uparrow|_{3,+} \right) \right. \\ \left. - \left(|\uparrow\rangle_d |\downarrow\rangle_{3,-} - |\downarrow\rangle_d |\uparrow\rangle_{3,-} \right) \left(\langle\uparrow|_d \langle\downarrow|_{3,-} - \langle\downarrow|_d \langle\uparrow|_{3,-} \right) \right]. \quad (14.28)$$

By replacing the emergent spins with the spin operators $\vec{\mathcal{S}}_{3,\pm}$, we get the final form of the Hamiltonian

$$H_{\text{eff}}^{4,0} = t \left(\vec{S}_d \cdot \vec{\mathcal{S}}_{3,-} - \vec{S}_d \cdot \vec{\mathcal{S}}_{3,+} \right) + \frac{t}{4} \mathcal{P}_{01} P_{01}^{(3)}, \quad (14.29)$$

where $\mathcal{P}_{01}$ is the parity transformation operator defined in eq. (14.17) and $P_{01}^{(3)}$ projects on to the $\hat{n}_0 + \hat{n}_1 = 3$ subspace. This completes the effective Hamiltonian for the $S_{\text{tot}}^z = 0$ sector.

We now come to the $S_{\text{tot}}^z = +1/2$ sector. From eq. (14.8), the effective Hamiltonian for this sector is

$$H_{\text{eff}}^{3,\frac{1}{2}} = t \left[|\text{SS}\uparrow\rangle \langle \text{LM}_-^{3,\frac{1}{2}}| + |\text{LM}_-^{3,\frac{1}{2}}\rangle \langle \text{SS}\uparrow| \right], \quad (14.30)$$

where

$$|\text{SS}\uparrow\rangle = \frac{1}{\sqrt{2}} \left(P_d^\uparrow c_{0\downarrow}^\dagger - P_d^\downarrow c_{0\uparrow}^\dagger \right) c_{1\uparrow}^\dagger |0\rangle; \quad |\text{LM}_-^{3,\frac{1}{2}}\rangle = \frac{1}{\sqrt{2}} P_d^\uparrow \left(c_{0\uparrow}^\dagger c_{0\downarrow}^\dagger + c_{1\uparrow}^\dagger c_{1\downarrow}^\dagger \right) |0\rangle, \quad (14.31)$$

such that

$$|\text{SS}\uparrow\rangle \langle \text{LAM}_-^{3,\frac{1}{2}}| = \frac{1}{2} \left(P_d^\uparrow c_{0\downarrow}^\dagger - P_d^\downarrow c_{0\uparrow}^\dagger \right) c_{1\uparrow}^\dagger (c_{0\downarrow} c_{0\uparrow} + c_{1\downarrow} c_{1\uparrow}) P_d^\uparrow \quad (14.32)$$

$$= -\frac{1}{2} \left[P_d^\uparrow \left(c_{1\uparrow}^\dagger c_{0\uparrow} \hat{n}_{0\downarrow} + c_{0\downarrow}^\dagger c_{1\downarrow} \hat{n}_{1\uparrow} \right) + S_d^- \left(c_{1\uparrow}^\dagger c_{0\downarrow} \hat{n}_{0\uparrow} - c_{0\uparrow}^\dagger c_{1\downarrow} \hat{n}_{1\uparrow} \right) \right] \quad (14.33)$$

The full effective Hamiltonian then becomes

$$H_{\text{eff}}^{3,\frac{1}{2}} = -\frac{t}{2} \left[P_d^\uparrow \left\{ \left(c_{1\uparrow}^\dagger c_{0\uparrow} + c_{0\downarrow}^\dagger c_{1\downarrow} \right) \left(P_0^{(2)} P_1^{(0)} + P_0^{(0)} P_1^{(2)} \right) + \left(c_{0\uparrow}^\dagger c_{1\uparrow} + c_{1\downarrow}^\dagger c_{0\downarrow} \right) P_0^{(1)} P_1^{(1)} \right\} \right. \\ \left. + S_d^- \left(c_{1\uparrow}^\dagger c_{0\downarrow} - c_{0\uparrow}^\dagger c_{1\downarrow} \right) \left(P_0^{(2)} P_1^{(0)} + P_0^{(0)} P_1^{(2)} \right) + S_d^+ \left(c_{0\downarrow}^\dagger c_{1\uparrow} - c_{1\downarrow}^\dagger c_{0\uparrow} \right) P_0^{(1)} P_1^{(1)} \right] \\ = t \left[P_d^\uparrow \sqrt{2} \mathcal{A}_{\frac{1}{2}}^z - \frac{1}{2} S_d^+ \mathcal{B}_{\frac{1}{2}}^- - \frac{1}{2} S_d^- \mathcal{B}_{\frac{1}{2}}^+ \right], \quad (14.34)$$

where we have defined

$$\mathcal{A}_{\frac{1}{2}}^z = -\frac{1}{2\sqrt{2}} \left[\left(c_{1\uparrow}^\dagger c_{0\uparrow} + c_{0\downarrow}^\dagger c_{1\downarrow} \right) \left(P_0^{(2)} P_1^{(0)} + P_0^{(0)} P_1^{(2)} \right) + \left(c_{0\uparrow}^\dagger c_{1\uparrow} + c_{1\downarrow}^\dagger c_{0\downarrow} \right) P_0^{(1)} P_1^{(1)} \right] \quad (14.35)$$

and

$$\mathcal{B}_{\frac{1}{2}}^+ = \left(c_{1\uparrow}^\dagger c_{0\downarrow} - c_{0\uparrow}^\dagger c_{1\downarrow} \right) \left(P_0^{(2)} P_1^{(0)} + P_0^{(0)} P_1^{(2)} \right); \quad \mathcal{B}_{\frac{1}{2}}^- = \left(c_{0\downarrow}^\dagger c_{1\uparrow} - c_{1\downarrow}^\dagger c_{0\uparrow} \right) P_0^{(1)} P_1^{(1)} \quad (14.36)$$

The effective Hamiltonian for the $S_{\text{tot}}^z = -1/2$ sector is obtained by applying the transformations $P_d^\sigma \rightarrow P_d^{\bar{\sigma}}, c_{i\sigma} \rightarrow c_{i\bar{\sigma}}$ on the Hamiltonian. This is motivated by the fact that applying the same transformations on the $S_{\text{tot}}^z = 1/2$ eigenstates give the $S_{\text{tot}}^z = -1/2$ eigenstates.

$$H_{\text{eff}}^{3,-\frac{1}{2}} = t \left[P_d^\dagger \sqrt{2} \mathcal{A}_{-\frac{1}{2}}^z - \frac{1}{2} S_d^+ \mathcal{B}_{-\frac{1}{2}}^- - \frac{1}{2} S_d^- \mathcal{B}_{-\frac{1}{2}}^+ \right]. \quad (14.37)$$

The operators $\mathcal{A}_{-\frac{1}{2}}^z$ and $\mathcal{B}_{-\frac{1}{2}}^\pm$ are obtained by applying the above mentioned transformation on their $S_{\text{tot}}^z = +1/2$ counterparts.

3.15 Appendix: Non-Fermi liquid exponents in self-energy and two-particle correlations

The one-particle self-energies relating to the impurity and zeroth sites, as well as some relevant two-particle correlation functions, can be shown to follow power-law behaviour close to the Brinkman-Rice transition at r_{c2} , by mapping the impurity model to a classical Coulomb gas, similar to the work of Anderson, Yuval and Hamman for the Kondo model [Anderson and Yuval(1969)] and that of Si and Kotliar for a periodic Anderson model [Si and Kotliar(1993a)]. Indeed, the latter work uses an auxiliary that is similar to our extended Anderson impurity model. In the following, we adapt their results to our model. These non-universal power-laws are signatures of the non-Fermi liquid excitations that emerge at r_{c2} .

Near r_{c2} , extended SIAM can be reduced to a $J - U_b$ model (shown in subsection V.A of the main manuscript). The conduction electrons scattering off the impurity suffer phase shifts because of the presence of the potentials J and U_b . These phase shifts are given by $\delta_{\text{sp}} = \arctan(\pi \rho_0 J/2)$ and $\delta_{\text{ch}} = \arctan(\pi \rho_0 U_b/2)$ [Si and Kotliar(1993a), Giamarchi(2004)], where ρ_0 is the electronic density of states in the conduction bath. The low-energy behaviour of the local impurity and bath self-energies can be expressed in terms of these phase shifts [Si and Kotliar(1993a)]:

$$\begin{aligned} \Sigma_{dd}(\omega) &\sim \omega^{\gamma_{dd}}, \quad \Sigma_{d0}(\omega) \sim \omega^{\gamma_{d0}}, \quad \Sigma_{00}(\omega) \sim \omega^{\gamma_{00}}, \\ \gamma_{dd} &= \frac{5}{4} - \left(\frac{\delta_{\text{tot}}^*}{\pi} \right)^2 - \left(\frac{\delta_{\text{ch}}^*}{\pi} \right)^2, \quad \gamma_{d0} = 2 \left[\left(\frac{\delta_{\text{ch}}^*}{\pi} \right)^2 + \left(\frac{\delta_{\text{tot}}^*}{\pi} \right)^2 - \frac{1}{4} \right], \\ \gamma_{00} &= \left(\frac{\delta_{\text{tot}}^*}{\pi} - \frac{1}{2} \right)^2 + \left(\frac{\delta_{\text{ch}}^*}{\pi} \right)^2 - 2, \end{aligned} \quad (15.1)$$

where we have defined a total phase shift parameter $\frac{\delta_{\text{tot}}}{\pi} \equiv \frac{\delta_{\text{sp}} + \delta_{\text{ch}}}{\pi} - \frac{1}{2}$. This can be extended to certain correlation functions as well. The impurity spin-flip correlation, the impurity-bath excitonic correlation and the zeroth-site pairing correlation [Si and Kotliar(1993a)]:

$$\begin{aligned} \langle S_d^+(\tau) S_d^-(0) \rangle &\sim \tau^{-\alpha_1} \implies \langle S_d^+ \rangle(\omega) \sim \omega^{(\alpha_1-1)/2}, \\ \langle (c_{d\uparrow}^\dagger c_{0\uparrow})(\tau) (c_{0\uparrow}^\dagger c_{d\uparrow})(0) \rangle &\sim \tau^{-\alpha_2} \implies \langle c_{d\uparrow}^\dagger c_{0\uparrow} c_{0\uparrow}^\dagger c_{d\uparrow} \rangle(\omega) \sim \omega^{(\alpha_2-1)/2}, \\ \langle (c_{0\uparrow}^\dagger c_{0\downarrow}^\dagger)(\tau) (c_{0\downarrow} c_{0\uparrow})(0) \rangle &\sim \tau^{-\alpha_3} \implies \langle c_{0\uparrow}^\dagger c_{0\downarrow}^\dagger \rangle(\omega) \sim \omega^{(\alpha_3-1)/2}, \end{aligned} \quad (15.2)$$

where the exponents are given by

$$\alpha_1 = 2 \left(1 - \frac{\delta_{\text{sp}}^*}{\pi} \right)^2, \alpha_2 = \left(\frac{\delta_{\text{tot}}^*}{\pi} - \frac{1}{2} \right)^2 + \left(\frac{\delta_{\text{ch}}^*}{\pi} \right)^2, \alpha_3 = \left(\frac{\delta_{\text{tot}}^*}{\pi} + \frac{1}{2} \right)^2 + \left(1 + \frac{\delta_{\text{ch}}^*}{\pi} \right)^2. \quad (15.3)$$

3.16 Appendix: Entanglement entropy, magnetisation and phase shift, at the QCP

As mentioned in the previous section, the polarised ground-states $|-\rangle^{3,\pm\frac{1}{2}}$ of eq. (14.7) are special because they involve the strong-coupling ground-state as well as the local moment ground-state. The overall ground states, therefore, contain both entangled and non-entangled impurity contributions, leading to a departure from maximal entanglement. We will now calculate the impurity entanglement entropy $S_{\text{EE}}(d)$ in this polarised ground-state(s) that emerges exactly at the QCP, and show that $S_{\text{EE}}(d)$ has a fractional value (in units of $\log 2$). This turns out to be linked to the fact that the impurity is screened partially in these states, leading to an impurity magnetisation that is non-vanishing but only half of that in the local moment phase.

The fractional entropy arises because of the singular scattering of the gapless excitations at the Fermi surface leading to the destruction of the local Fermi liquid. In order to access these processes, we will work with a *semi-continuous* conduction bath beyond the zeroth site (total number of lattice sites $N \rightarrow \infty$): $H_{\text{bath}} = \sum_{k,\sigma} \epsilon(k) \psi_\sigma^\dagger(k) \psi_\sigma(k)$. This conduction bath that is composed of the k -states formed from the Hilbert space of the lattice sites beyond the zeroth site will be referred to as the reduced conduction bath. This is in addition to the Hamiltonian for the impurity and zeroth sites: $H_{d0} = J \vec{S}_d \cdot \vec{S}_0 - \frac{U_b}{2} (c_{0\uparrow}^\dagger c_{0\uparrow} - c_{0\downarrow}^\dagger c_{0\downarrow})^2$, where $\vec{S}_d$ is the spin operator for the impurity site, and $c_{0\sigma}^\dagger$ creates an electron of spin σ at the zeroth site. The full Hamiltonian also involves a single-particle hopping term that couples these two Hamiltonians: $H = H_{d0} + H_{\text{bath}} + H_{\text{int}}$, where

$$H_{\text{int}} = -t \sum_{\sigma} \left(c_{0\sigma}^\dagger c_{1\sigma} + \text{h.c.} \right) = -\frac{t}{N} \sum_{k,\sigma} \left[c_{0\sigma}^\dagger \psi_\sigma(k) + \text{h.c.} \right]. \quad (16.1)$$

$c_{1\sigma}^\dagger = \frac{1}{N} \sum_k \psi_\sigma^\dagger(k)$ is the creation operator for the first site of the conduction bath of N sites (but effectively the zeroth site for the reduced conduction bath defined by H_{bath}). We will treat H_{int} perturbatively in t/J , in order to obtain the set of renormalised g -fold degenerate ground-states $\{ |\psi_{\text{gs}}^{(n)}\rangle; n = 1, \dots, g \}$.

As mentioned before, the intention here is to calculate magnetisation and entanglement entropy in the non-Fermi liquid ground states. In the case of a degenerate subspace with $S_{\text{tot}}^z = \pm 1/2$ where $S_{\text{tot}}^z = S_d^z + S_0^z + S_1^z$ is the total spin operator, the magnetisation m_d^z must be calculated in any *one* of the degenerate states for our problem. This is accomplished by solving the problem in the presence of an infinitesimal magnetic field term $-hS_{\text{tot}}^z$ that lifts the degeneracy and chooses one of the symmetry-broken ground-states. This is similar to the "two self-energy description" introduced by David Logan for computing quantities in the local moment phase/Mott insulating phase in terms of different self-energies obtained by inserting positive and negative magnetic fields [Glossop and Logan(2002), Logan et al.(2014) Logan, Tucker, and Galpin, Logan and Galpin(2015)]. Due to the symmetry of the Hamiltonian under $S_{\text{tot}}^z \rightarrow -S_{\text{tot}}^z$, both the limits $h = 0^+$ and $h = 0^-$ give the same results for the density matrix and the entanglement entropy. We will work with $h = 0^+$. Since the total Hamiltonian H and the perturbation H_{int} preserve the total spin $S_{\text{tot}}^z = S_d^z + S_0^z + \int dk S_k^z$, it is sufficient to solve only for the sector with the most positive value of S_{tot}^z , since those are the states that will be selected by the positive magnetic field $h = 0^+$.

In the absence of H_{int} , the set of ground-states with a positive S_{tot}^z will contain, to begin with, the local moment states $|\uparrow\rangle_d |0\rangle_0 |\phi\rangle$ and $|\uparrow\rangle_d |2\rangle_0 |\phi\rangle$, where $|\phi\rangle$ represents the filled Fermi sea of the reduced conduction bath, acting as its spinless ground-state. These states have $S_{\text{tot}}^z = 1/2$. We can label these states as $|n_{\text{tot}}, \mu_d^z\rangle$, using the total number of particles $n_{\text{tot}} = n_d + n_0 + n_{\text{bath}}$ (which is also conserved, and where we set $n_{\text{bath}} = 1$ for the set $|\phi\rangle$), and the average magnetisation μ_d^z of the impurity state: $|2, \frac{1}{2}\rangle \equiv |\uparrow\rangle_d |0\rangle_0 |\phi\rangle$, $|4, \frac{1}{2}\rangle \equiv |\uparrow\rangle_d |2\rangle_0 |\phi\rangle$. Since the singlet state $|SS\rangle_{d0}$ (on the d and zeroth sites) is degenerate with the local moment states $|\uparrow\rangle_d |0(2)\rangle_0$, one can construct other states with $S_{\text{tot}}^z = 1/2$ by creating magnetic excitations at the Fermi surface. The fact that we are working with a thermodynamically large bath ($N \rightarrow \infty$) means that there will be gapless excitations vanishingly close to the Fermi surface. Such states are of the form $|SS\rangle_{d0} |e_\uparrow\rangle$ (labeled as $|4, 0\rangle$ in the $|n_{\text{tot}}, \mu_d^z\rangle$ notation) and $|SS\rangle_{d0} |h_\downarrow\rangle$ (labeled as $|2, 0\rangle$), where $|e_\sigma\rangle$ and $|h_\downarrow\rangle$ are electron and hole states respectively:

$$|e_\sigma\rangle = \sum_k^{\text{FS}} \psi_\sigma^\dagger(k) |\phi\rangle, \quad |h_\sigma\rangle = \sum_k^{\text{FS}} \psi_\sigma(k) |\phi\rangle. \quad (16.2)$$

The sum is over k -states on the Fermi surface. The singlet state was set to have zero spin on the impurity site in the state notation, because $\langle S_d^z \rangle$ vanishes in the singlet state. The full set of relevant degenerate ground-states, classified by the total number of particles, is

$$n_{\text{tot}} = 4 : |4, 0\rangle, |4, \frac{1}{2}\rangle; \quad n_{\text{tot}} = 2 : |2, 0\rangle, |2, \frac{1}{2}\rangle. \quad (16.3)$$

Within each 2×2 block, the interaction matrix H_{int} takes the form (in the basis of the states in eq. (16.3))

$$(H_{\text{int}})_{\text{gs}}^{n_{\text{tot}}=2} = \frac{t}{N} \times \begin{pmatrix} 0 & 1/\sqrt{2} \\ 1/\sqrt{2} & 0 \end{pmatrix} = - (H_{\text{int}})_{\text{gs}}^{n_{\text{tot}}=4}, \quad (16.4)$$

and the $S_{\text{tot}}^z = +\frac{1}{2}$ ground-states are then easily obtained to be of the form $|\psi_{\text{gs}}\rangle^{(n_{\text{tot}})} = \frac{1}{\sqrt{2}} (|n_{\text{tot}}, 0\rangle - |n_{\text{tot}}, \frac{1}{2}\rangle)$, $n_{\text{tot}} = 2$ & 4 , independent of N . Both ground-states involve an equal superposition of a $\mu_d^z = 1/2$ state

and a $\mu_d^z = 0$ state, such that the net impurity magnetisation in either of the renormalised polarised states is

$$m_d^z = \frac{1}{2} \left(0 + \frac{1}{2} \right) = \frac{1}{4}. \quad (16.5)$$

The reduced density matrix $\rho_{\text{imp}} = \text{Tr}' \left(|\psi_{\text{gs}}\rangle^{(n_{\text{tot}})} \langle \psi_{\text{gs}}|^{(n_{\text{tot}})} \right)$ (where Tr' indicates partial trace over all states apart from those of the impurity) for the impurity spin in the polarised states can now be immediately obtained. It is independent of n_{tot} , and actually turns out to be halfway between the maximally mixed density matrix $\rho_{\text{MM}} = \frac{1}{2}I$ and the non-entangled density matrix $\rho_{\text{NE}} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$:

$$\rho_{\text{imp}} = \begin{pmatrix} 3/4 & 0 \\ 0 & 1/4 \end{pmatrix} = \frac{1}{2} (\rho_{\text{MM}} + \rho_{\text{NE}}), \quad S_{\text{EE}}(d) = -\text{Tr}(\rho_{\text{imp}} \ln \rho_{\text{imp}}) \simeq 0.81 \log 2. \quad (16.6)$$

This fractional value of the entropy in units of $\log 2$ is another signature of the non-Fermi liquid. We can express this in terms of an effective degeneracy $\tilde{g}$ of the impurity degree of freedom. We define this effective degeneracy as $\tilde{g} = 1 + 2|m_d^z|$, in order to recover the unique ground-state in the local Fermi liquid phase (that has $m_d^z = 0, \tilde{g} = 1$) and the magnetic doublet in the local moment phase (that has $m_d^z = 1/2, \tilde{g} = 2$). The proposed form also results in a non-integer impurity degeneracy for the NFL at the QCP, $\tilde{g}^* = 3/2$. Using the effective degeneracy $\tilde{g}$, one can also write down a general ground-state

$$|\psi_{\text{gs}}(g)\rangle = \frac{1}{\sqrt{(2-\tilde{g})^2 + (\tilde{g}-1)^2}} [(2-\tilde{g}) |n_{\text{tot}}, 0\rangle - (\tilde{g}-1) |n_{\text{tot}}, 1/2\rangle], \quad (16.7)$$

that interpolates between the strong-coupling ground-state for $r < r_{c2}$ (therefore $\tilde{g} = 1$) and the weak coupling ground-state $r > r_{c2}$ (therefore $\tilde{g} = 2$), also describing the QCP in between ($\tilde{g} = 3/2$). The entanglement entropy, when written in terms of $\tilde{g}$, takes the form

$$\rho_{\text{imp}} = \frac{1}{2} \begin{pmatrix} 2-\tilde{g} & 0 \\ 0 & \tilde{g} \end{pmatrix} = \frac{1}{2} + \begin{pmatrix} m_d^z & 0 \\ 0 & -m_d^z \end{pmatrix}, \quad (16.8)$$

$$S_{\text{EE}}(d) = - \left(1 - \frac{\tilde{g}}{2} \right) \ln \left(1 - \frac{\tilde{g}}{2} \right) - \frac{\tilde{g}}{2} \ln \frac{\tilde{g}}{2}. \quad (16.9)$$

The relation between the reduced density matrix ρ_{imp} and the impurity magnetisation m_d^z is consistent with that obtained by Kopp et al., [Kopp et al.(2007)Kopp, Jia, and Chakravarty].

The general form $|\psi_{\text{gs}}(g)\rangle$ of polarised interpolating ground-state also allows us to calculate the excess charge (n_{exc}) added, by the impurity, to the conduction bath Fermi surface (through gapless excitations). This then allows the scattering phase shift of the conduction electrons (obtained via the Friedel sum rule, [Friedel(1956), Langer and Ambegaokar(1961), Langreth(1966)]) to be linked with the impurity magnetisation m_d^z . From the definitions of the states $|n_{\text{tot}}, 0\rangle$ and $|n_{\text{tot}}, 1/2\rangle$, we note that the former state involves an impurity-bath singlet while the latter involves a decoupled local

moment. The singlet state contributes an excess charge of unity to the conduction bath: $n_{\text{exc}}^{\text{SS}} = 1$, because of the impurity-bath entanglement within it. The local state moment does not contribute any excess charge, because the impurity spin is no longer hybridising with the bath in that state: $n_{\text{exc}}^{\text{LM}} = 0$. Putting these together, the net excess charge contributed by the state in eq. (16.7) is

$$n_{\text{exc}} = \frac{1}{(2 - \tilde{g})^2 + (\tilde{g} - 1)^2} [(2 - \tilde{g})^2 n_{\text{exc}}^{\text{SS}} + (\tilde{g} - 1)^2 n_{\text{exc}}^{\text{LM}}] = \left[1 + \left(\frac{2 - \tilde{g}}{\tilde{g} - 1} \right)^2 \right]^{-1}. \quad (16.10)$$

At the QCP, the excess charge acquires a fractional value of $1/2$, indicating once more that only half of the impurity remains coupled with the bath at the transition. The Friedel sum rule can be used to equate this excess charge with the phase shift δ^* suffered by conduction electrons as they scatter off the impurity, at the fixed point:

$$\delta^* = \pi n_{\text{exc}} = \left[1 + \left(\frac{2 - \tilde{g}}{\tilde{g} - 1} \right)^2 \right]^{-1} = \frac{4m_d^z{}^2}{1 - 4|m_d^z| + 8m_d^z{}^2}. \quad (16.11)$$

Equation (16.11) unifies the effect of several important quantities that lead to the frustration of the impurity and the destruction of the Kondo effect. The presence of U_b introduces local moment states into the spectrum and leads to states with non-vanishing impurity magnetisation m_d^z at the QCP; this non-zero m_d^z can be interpreted as an effective impurity degeneracy $\tilde{g}$ that is only partially screened, resulting in only half the excess charge n_{exc} being contributed to the conduction bath. This reduction in the excess charge acts as a shift in the boundary conditions felt by the conduction electrons and manifests as a phase shift that is less than the unitarity limit of π .

3.17 Appendix: Additional correlations near the ESQPT and the QCP

We have included some additional correlations for the extended SIAM here to highlight the features that are mentioned in the main manuscript. Left panel of Fig. (3.15) shows the impurity and zeroth site double occupancy. Both vanish at r_{c1} , indicating the emergence of the Kondo model. Very close to r_{c2} , the zeroth site doublon occupancy picks up (see inset), showing the destruction of the Kondo cloud. Right panel of Fig. (3.15) shows various measures of entanglement close to r_{c2} . The impurity-bath mutual information (blue) decreases as the transition is approached, showing the breakdown of impurity screening. The impurity and first site MI (violet), the zeroth site and first site MI (red) and the impurity, zeroth site and first site tripartite information (grey) increase, showing the emergence of long-ranged correlations near the transition. Fig. (3.16) shows the emergence of long-ranged intra-bath mutual information near r_{c1} and r_{c2} , signalling critical behaviour near the ESQPT and QCP respectively. Intra-bath CDW correlations and impurity-bath Ising correlations in Fig. (3.17) also show the previously-mentioned emergence of long-ranged correlations.




Figure 3.15: Left: Evolution of the double occupancy of impurity (red) and zeroth sites (blue) ground-state double occupancy, with r . Right: Evolution of various quantum information-theoretic measures, very close to the transition.




Figure 3.16: Variation of intra-bath mutual information between zeroth site and sites 1, 2 and 3, near r_{c1} (left) and r_{c2} (right).




Figure 3.17: Left: Variation of charge isospin Ising correlations between the zeroth site and bath sites further down the chain. Right: Ising correlations between the impurity site and bath sites beyond the zeroth site. Both sets of correlations show an overall increase as $r \rightarrow r_{c2}$.

Chapter 4

A New Auxiliary Model Approach for Interacting Electrons: Mapping Impurity Models To Lattice Models

4.1 The Philosophy of Auxiliary Models

In the previous chapter, we extended the Anderson impurity model to host a correlation-driven delocalisation-localisation transition on the impurity site, from a local Fermi liquid state to a local moment state. We demonstrated that this transition captures very well the phenomenology of the Hubbard dimension in infinite dimensions (as seen from, say, dynamical mean-field theory (DMFT)). In this chapter, we extend this approach by developing a formalism to study the more realistic case of finite-dimensional interacting models by using appropriately chosen impurity models as auxiliary models. The key to an *auxiliary model method* is to construct an appropriate auxiliary model that while being tractable captures some essential property of the full lattice model [Martin et al.(2016)Martin, Reining, and Ceperley]. This involves separating the complete degrees of freedom into a system part (H_S) which one treats exactly and the rest of the system (H_R) that one might simplify in order to be able to solve the problem. The non-trivial nature of the problem arises of course from the hybridisation H_{SR} between the system and the bath:

$$H = \begin{bmatrix} H_R & H_{RS} \\ H_{RS}^* & H_S \end{bmatrix}. \quad (1.1)$$

This separation can always be done exactly, if one does not care about tractability. For example, one can take the Hubbard model that describes electrons hopping on a lattice and interacting when two electrons reside on the same lattice site:

$$H_{\text{hub}} = -t \sum_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger c_{j,\sigma} + \text{h.c.}) + U \sum_i n_{i\uparrow} n_{i\downarrow}, \quad (1.2)$$

and separate this into parts, the system being the local orbitals of a specific site l and the rest of the

system being all the other sites:

$$H_S = U n_{l\uparrow} n_{l\downarrow}, \quad H_R = -t \sum_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger c_{j,\sigma} + \text{h.c.}) + U \sum_i n_{i\uparrow} n_{i\downarrow}, \quad H_{SR} + H_{RS} = -t \sum_{i,\sigma} (c_{l,\sigma}^\dagger c_{i,\sigma} + \text{h.c.}), \quad (1.3)$$

where the primed summation is carried out over all nearest-neighbour pairs that do not involve l and the summation in $H_{SR} + H_{RS}$ is over all sites adjacent to l .

The smaller system H_S (often called the *cluster*) is typically chosen such that its eigenstates are known exactly. Progress is then made by choosing a simpler form for H_R and its coupling H_{RS} with the smaller system. This combination of the cluster and the simpler bath is then called the *auxiliary system*. A tractable auxiliary system for the Hubbard model is the single-impurity Anderson model (SIAM); the correlated impurity site represents the set of local orbitals of any given site, while the conduction bath represents the rest of the lattice sites in an approximate fashion. Such a construction is shown in fig. 4.1.




Figure 4.1: Left: Full Hubbard model lattice with onsite repulsion U^H on all sites and hopping between nearest neighbour sites with strength t^H . Right: Extraction of the auxiliary (cluster+bath) system from the full lattice. The central site on left becomes the impurity site (red) on the right (with an onsite repulsion ϵ_d), while the rest of the $N - 1$ sites on the left form a conduction bath (green circles) (with dispersion ϵ_k and correlation modelled by the self-energy $\Sigma_k(\omega)$) that hybridizes with the impurity through the coupling V .

It should be noted that any reasonable choice of the cluster and bath would break the translational symmetry of the full model: To allow computing quantities, one would need to make the bath (which is a much larger system) simpler than the cluster (which is a single site). This distinction breaks the translational symmetry of the Hubbard model. As a result, calculations that make use of the translation symmetry of the full model (such as k -space objects) will require a periodisation operation of an auxiliary model quantity.

The present work is aimed towards developing a new auxiliary model method to study correlated electronic systems. As described above, the first step is to identify the correct impurity model which can faithfully capture the physics of the lattice model we are trying to solve. The local behaviour of this impurity model should reflect the essential local physics of the lattice model. Typically, we will

consider impurity model geometries where the real-space bath site connected directly to the impurity hosts some local interaction. We will henceforth refer to this site as the *zeroth site*. This model must be solved using an impurity solver. In the present work, we use the recently-developed unitary renormalisation group method [Mukherjee and Lal(2020c), Mukherjee and Lal(2020d), Mukherjee and Lal(2020a), Mukherjee and Lal(2020b), Patra and Lal(2021), Pal et al.(2019) Pal, Mukherjee, and Lal]. The leap to the bulk model is then made by applying **manybody lattice translation operators** on the auxiliary model. This process, referred to as *tiling* here, allows us to reconstruct the lattice model Hamiltonian from that of the auxiliary model and restores the translation invariance of the full lattice model. Once the Hamiltonians can be mapped onto one another, it is possible to relate the eigenstates and correlations functions between the auxiliary model and lattice model, using a manybody version of **Bloch's theorem**.

4.2 What Is The Correct Auxiliary Model For Correlated Electrons on a 2D Square Lattice?

As discussed in more detail in the previous chapter, it was necessary to extend the single-impurity Anderson model (SIAM) in order to obtain a phase transition on the impurity site. Without any modification, the standard Anderson model (at particle-hole symmetry) leads to the emergence of an $\omega = 0$ Kondo resonance in the impurity ($d\sigma$) spectral function at any value of the impurity correlation U_d , at sufficiently low temperatures:

$$H_{\text{SIAM}} = \sum_{k,\sigma} \varepsilon_k n_{k,\sigma} - V \sum_{\sigma} (c_{0,\sigma}^\dagger c_{d,\sigma} + \text{h.c.}) + \varepsilon_d \sum_{\sigma} n_{d,\sigma} + U_d n_{d,\uparrow} n_{d,\downarrow}. \quad (2.1)$$

The operator $c_{0,\sigma}$ acts on the local conduction bath density that couples with the impurity site. The low-energy is one of local Fermi liquid for all parameter regimes. Increasing U_d only serves to reduce the width of the central peak at the cost of the appearance of side bands at energy scales of the order of U_d , but the resonance never disappears. The extended Anderson model that we developed augments the SIAM by allowing for local correlation in the local conduction bath charge density that couples with the impurity spin:

$$H_{\text{ESIAM}} = H_{\text{SIAM}} + J_K \mathbf{S}_d \cdot \mathbf{S}_0 - \frac{U_b}{2} (n_{0,\uparrow}^2 - n_{0,\downarrow}^2); \quad (2.2)$$

tuning this bath correlation to large attractive values (see [Gazizova and LeBlanc(2023)]) for some recent findings of non-local effective attractive interactions within the Hubbard model) *does* lead to a localisation transition on the impurity where the local Fermi liquid excitations are replaced by the physics of a gapped local moment on the impurity. We note that the generation of such correlations in the bath is something that occurs in DMFT as well – the conduction bath acquires a *non-trivial self-energy* during the convergence to self-consistency.

While this modified impurity model captures well the Mott metal-insulator transition of the Hubbard model in infinite dimensions, it *lacks any knowledge of the underlying lattice geometry*, and is therefore ill-suited to address the phenomenology of more realistic low-dimensional models. Examples of such phenomenology include the pseudogap phase of the cuprates (with nodal-antinodal

dichotomy), strange metal (with T -linear resistivity) and high- T_c superconducting phases observed in several families of quantum materials as well as graphene. This leads us to propose the framework of *lattice-embedded quantum impurity models*. Upon embedding the impurity site within the underlying lattice, the coupling between it and its environment must be consistent with the lowered symmetry of the lattice (see Fig. 4.2). Impurity-bath correlations calculated within such an impurity model are sensitive to the lattice geometry, and are much better suited to address the phenomenology mentioned above.




Figure 4.2

As a concrete example, we take the extended SIAM defined in eq. 2.2, and write down its lattice-embedded form on a 2D square lattice, with the impurity placed at the site r_d . This is done by allowing the impurity to hybridise with the four nearest-neighbours, through V and J_K . The same four sites also carry the local bath interaction. The complete auxiliary model is

$$\begin{aligned}
 H = & -t \sum_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger c_{j,\sigma} + \text{h.c.}) + \sum_{\langle i, r_d \rangle} \left[-\frac{V}{Z} \sum_{\sigma} (c_{r_d,\sigma}^\dagger c_{i,\sigma} + \text{h.c.}) + \frac{J_K}{Z} \mathbf{S}_{r_d} \cdot \mathbf{S}_i - \frac{W}{2Z} (n_{i,\uparrow}^2 - n_{i,\downarrow}^2) \right] \\
 & + \varepsilon_d \sum_{\sigma} n_{r_d,\sigma} + U_d n_{r_d,\uparrow} n_{r_d,\downarrow},
 \end{aligned}
 \tag{2.3}$$

where Z is the coordination number of the lattice, and the second summation $\sum_{\langle i, r_d \rangle}$ is carried out over all sites i that are nearest-neighbour to the site r_d . Note that the one-particle hybridisation V and the spin-exchange coupling J_K have their *symmetry lowered* from the s -wave spherical symmetry of the continuum impurity models to a C_4 -symmetry consistent with the underlying lattice. This feature is crucial in explaining the phenomenology of the various low-dimensional materials.

Various enhancements can be applied to such a model. The lattice can be modified to study different materials. More complicated geometries such as a two-layer model can be studied using a two-impurity auxiliary model, where each impurity couples with its own conduction bath as well as to each other.

4.3 Mapping Auxiliary Model To Lattice Model

In this subsection, we provide an explicit example of constructing a lattice model from a lattice-embedded impurity model. This will involve the use of manybody lattice translation operators, so we begin with a brief discussion of them.

4.3.1 Manybody Translation Operators

In order to create the bulk model, we need to translate the local auxiliary model over the entire lattice. For this, we define *many-particle global* translation operators $T(\mathbf{a})$ that translate all positions by a vector $\mathbf{a}$. In terms of manybody states and operators, their action is defined as

$$\begin{aligned} T^\dagger(\mathbf{a}) |\mathbf{r}_1, \mathbf{r}_2, \dots\rangle &= |\mathbf{r}_1 + \mathbf{a}\rangle \otimes |\mathbf{r}_2 + \mathbf{a}\rangle \dots \otimes |\mathbf{r}_n + \mathbf{a}\rangle \\ T^\dagger(\mathbf{a}) \mathcal{O}(\mathbf{r}_1, \mathbf{r}_2, \dots) T(\mathbf{a}) &= \mathcal{O}(\mathbf{r}_1 + \mathbf{a}, \mathbf{r}_2 + \mathbf{a}, \dots), \end{aligned} \quad (3.1)$$

where $|\mathbf{r}_1, \mathbf{r}_2, \dots\rangle$ is a state in the manyparticle Fock-space basis with the particles localised at the specified positions. For example, for a local fermionic creation operator $c^\dagger(\mathbf{r})$, we have

$$T^\dagger(\mathbf{a}) c^\dagger(\mathbf{r}) T(\mathbf{a}) = c^\dagger(\mathbf{r} + \mathbf{a}). \quad (3.2)$$

It acts similarly on the auxiliary model Hamiltonian:

$$T^\dagger(\mathbf{a}) H_{\text{aux}}(\mathbf{r}_d) T(\mathbf{a}) = H_{\text{aux}}(\mathbf{r}_d + \mathbf{a}), \quad (3.3)$$

translating all sites by the vector $\mathbf{a}$. By introducing the Fourier transform to momentum space,

$$|\mathbf{r}_1, \mathbf{r}_2, \dots\rangle = \otimes_{j=1}^N \int d\mathbf{k}_j e^{i\mathbf{r}_j \cdot \mathbf{k}_j} |\mathbf{k}_j\rangle, \quad (3.4)$$

it is easy to see that the total momentum states are eigenstates of the global translation operators:

$$\begin{aligned} T^\dagger(\mathbf{a}) |\mathbf{k}_1, \mathbf{k}_2, \dots\rangle &= \otimes_{j=1}^N \int d\mathbf{r}_j e^{-i\mathbf{r}_j \cdot \mathbf{k}_j} T^\dagger(\mathbf{a}) |\mathbf{r}_j\rangle, \\ &= \otimes_{j=1}^N \int d\mathbf{r}_j e^{-i\mathbf{r}_j \cdot \mathbf{k}_j} |\mathbf{r}_j + \mathbf{a}\rangle \\ &= \otimes_{j=1}^N e^{i\mathbf{a} \cdot \mathbf{k}_j} \int d\mathbf{r}_j e^{-i\mathbf{r}_j \cdot \mathbf{k}_j} |\mathbf{r}_j\rangle \\ &= e^{i\mathbf{a} \cdot \mathbf{k}_{\text{tot}}} |\mathbf{k}_1, \mathbf{k}_2, \dots\rangle, \end{aligned} \quad (3.5)$$

where $\mathbf{k}_{\text{tot}} = \sum_j \mathbf{k}_j$ is the total momentum. This gives a compact form for the translation operator:

$$T^\dagger(\mathbf{a}) = e^{i\mathbf{a} \cdot \hat{\mathbf{k}}_{\text{tot}}}, \quad (3.6)$$

where $\hat{\mathbf{k}}_{\text{tot}}$ is the total momentum operator.

We end this section by calculating the effect of the translation operators on the k -space annihilation operators. We have

$$c(\mathbf{k}) = \frac{1}{\sqrt{N}} \sum_{\mathbf{r}} e^{-i\mathbf{k}\cdot\mathbf{r}} c(\mathbf{r}) . \quad (3.7)$$

Applying the translation operator to this decomposition, we get

$$\begin{aligned} T(\vec{a}) c(\mathbf{k}) T^\dagger(\vec{a}) &= \frac{1}{\sqrt{N}} \sum_{\mathbf{r}} e^{-i\mathbf{k}\cdot\mathbf{r}} T(\vec{a}) c(\mathbf{r}) T^\dagger(\vec{a}) \\ &= \frac{1}{\sqrt{N}} \sum_{\mathbf{r}} e^{-i\mathbf{k}\cdot\mathbf{r}} c(\mathbf{r} - \vec{a}) \\ &= e^{-i\mathbf{k}\cdot\vec{a}} c(\mathbf{k}) . \end{aligned} \quad (3.8)$$

4.3.2 Periodisation of Auxiliary Model Into Extended Hubbard Model

We recall the lattice-embedded impurity model defined in eq. 2.3. As discussed above, the route to restoring translation symmetry and reconstructing the lattice model is to translate the auxiliary model using manybody translation operators. Accordingly, we identify the “unit cell” of translation. Since we expect the impurity site to capture most of the physics arising from the correlated electrons, a good choice for the unit cell is the impurity site and its surrounding correlated environment. In order to retain information of the lattice geometry, we convert the hybridisation and bath interaction terms from real space to k -space. The Hamiltonian for the unit cell is therefore

$$\begin{aligned} H_{\text{clust}}(\mathbf{r}_d) &= - \sum_{\mathbf{k}, \sigma} e^{i\mathbf{k}\cdot\mathbf{r}_d} V(\mathbf{k}) (c_{\mathbf{r}_d, \sigma}^\dagger c_{\mathbf{k}, \sigma} + \text{h.c.}) + \sum_{\mathbf{k}_1, \mathbf{k}_2} e^{i(\mathbf{k}_1 - \mathbf{k}_2)\cdot\mathbf{r}_d} J_K(\mathbf{k}_1, \mathbf{k}_2) \mathbf{S}_{\mathbf{r}_d} \cdot \mathbf{S}_{\mathbf{k}_1, \mathbf{k}_2} + \varepsilon_d n_{\mathbf{r}_d} \\ &\quad + U_d n_{\mathbf{r}_d, \uparrow} n_{\mathbf{r}_d, \downarrow} - \frac{1}{2} \sum_{\{\mathbf{k}_i\}, \sigma, \sigma'} \sigma \sigma' e^{i(\mathbf{k}_1 - \mathbf{k}_2 + \mathbf{k}_3 - \mathbf{k}_4)\cdot\mathbf{r}_d} W(\{\mathbf{k}_i\}) c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \sigma'}^\dagger c_{\mathbf{k}_4, \sigma'} , \end{aligned} \quad (3.9)$$

where $\mathbf{S}_{\mathbf{k}_1, \mathbf{k}_2} = \sum_{\alpha, \beta} \boldsymbol{\sigma}_{\alpha, \beta} c_{\mathbf{k}_1, \alpha}^\dagger c_{\mathbf{k}_2, \beta}$, and the Fourier transforms are defined by the following relations:

$$\begin{aligned} \sum_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}_d} V(\mathbf{k}) c_{\mathbf{r}_d, \sigma}^\dagger c_{\mathbf{k}, \sigma} &= \frac{V}{Z} \sum_{r \rightarrow \text{NN of } \mathbf{r}_d} c_{\mathbf{r}_d, \sigma}^\dagger c_{\mathbf{r}, \sigma} , \\ \sum_{\mathbf{k}_1, \mathbf{k}_2} e^{i(\mathbf{k}_1 - \mathbf{k}_2)\cdot\mathbf{r}_d} J_K(\mathbf{k}_1, \mathbf{k}_2) \vec{S}_{\mathbf{r}_d} \cdot \vec{S}_{\mathbf{k}_1, \mathbf{k}_2} &= \frac{J_K}{Z} \sum_{r \rightarrow \text{NN of } \mathbf{r}_d} \mathbf{S}_{\mathbf{r}_d} \cdot \mathbf{S}_{\mathbf{r}} , \\ \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{q}_1, \mathbf{q}_2} e^{i(\mathbf{k}_1 - \mathbf{k}_2 + \mathbf{k}_3 - \mathbf{k}_4)\cdot\mathbf{r}_d} W(\{\mathbf{k}_i\}) c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \sigma'}^\dagger c_{\mathbf{k}_4, \sigma'} &= \frac{W}{2Z} \sum_{r \rightarrow \text{NN of } \mathbf{r}_d} n_{\mathbf{r}, \sigma} n_{\mathbf{r}, \sigma'} , \end{aligned} \quad (3.10)$$

We now translate this unit cell to all sites of the lattice, and reconstruct a translationally-invariant correlated lattice model:

$$H_{\text{latt}} = \sum_{\mathbf{r}} T^\dagger(\mathbf{r} - \mathbf{r}_d) H_{\text{clust}}(\mathbf{r}_d) T(\mathbf{r} - \mathbf{r}_d) . \quad (3.11)$$

We consider the effect of the translation operations on each part of the Hamiltonian. The purely local terms translate directly into a chemical potential term and a Hubbard interaction term:

$$\sum_{\mathbf{r}} T^\dagger(\mathbf{r} - \mathbf{r}_d) (\varepsilon_d \sum_{\sigma} n_{\mathbf{r}_d, \sigma} + U_d n_{\mathbf{r}_d, \uparrow} n_{\mathbf{r}_d, \downarrow}) T(\mathbf{r} - \mathbf{r}_d) = \varepsilon_d \sum_{\mathbf{r}, \sigma} n_{\mathbf{r}, \sigma} + U_d \sum_{\mathbf{r}} n_{\mathbf{r}, \uparrow} n_{\mathbf{r}, \downarrow}. \quad (3.12)$$

The one-particle hybridisation term maps onto a tight-binding hopping term:

$$\begin{aligned} \sum_{\mathbf{r}} T^\dagger(\mathbf{r} - \mathbf{r}_d) \sum_{\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{r}_d} V(\mathbf{k}) c_{\mathbf{r}_d, \sigma}^\dagger c_{\mathbf{k}, \sigma} T(\mathbf{r} - \mathbf{r}_d) &= \sum_{\mathbf{r}} \sum_{\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{r}} V(\mathbf{k}) c_{\mathbf{r}, \sigma}^\dagger c_{\mathbf{k}, \sigma} \\ &= \frac{V}{Z} \sum_{\mathbf{r}} \sum_{\mathbf{r}' \rightarrow \text{NN of } \mathbf{r}} c_{\mathbf{r}, \sigma}^\dagger c_{\mathbf{r}', \sigma} \\ &= \frac{2V}{Z} \sum_{\langle \mathbf{r}, \mathbf{r}' \rangle} c_{\mathbf{r}, \sigma}^\dagger c_{\mathbf{r}', \sigma}, \end{aligned} \quad (3.13)$$

where we used eq. 3.8 and the first equation from 3.10, and the fact that each nearest-neighbour pair $\mathbf{r}, \mathbf{r}'$ appears twice in the summation.

The Kondo terms map onto a nearest-neighbour spin-exchange term:

$$\begin{aligned} \sum_{\mathbf{r}, \mathbf{k}_1, \mathbf{k}_2} T^\dagger(\mathbf{r} - \mathbf{r}_d) e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}_d} J_K(\mathbf{k}_1, \mathbf{k}_2) \vec{S}_{\mathbf{r}_d} \cdot \vec{S}_{\mathbf{k}_1, \mathbf{k}_2} T(\mathbf{r} - \mathbf{r}_d) &= \sum_{\mathbf{r}, \mathbf{k}_1, \mathbf{k}_2} e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}} J_K(\mathbf{k}_1, \mathbf{k}_2) \vec{S}_{\mathbf{r}} \cdot \vec{S}_{\mathbf{k}_1, \mathbf{k}_2} \\ &= \frac{J_K}{Z} \sum_{\mathbf{r}} \sum_{\mathbf{r}' \rightarrow \text{NN of } \mathbf{r}} \vec{S}_{\mathbf{r}} \cdot \vec{S}_{\mathbf{r}'}, \\ &= \frac{2J_K}{Z} \sum_{\langle \mathbf{r}, \mathbf{r}' \rangle} \vec{S}_{\mathbf{r}} \cdot \vec{S}_{\mathbf{r}'}, \end{aligned} \quad (3.14)$$

where we used the second equation from 3.10. Finally, the bath interaction term W maps onto another Hubbard term

$$\begin{aligned} \sum_{\mathbf{r}, \mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} T^\dagger(\mathbf{r} - \mathbf{r}_d) e^{i(\mathbf{k}_1 - \mathbf{k}_2 + \mathbf{k}_3 - \mathbf{k}_4) \cdot \mathbf{r}_d} W(\{\mathbf{k}_i\}) c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \sigma'}^\dagger c_{\mathbf{k}_4, \sigma'} T(\mathbf{r} - \mathbf{r}_d) \\ = \sum_{\mathbf{r}, \mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} e^{i(\mathbf{k}_1 - \mathbf{k}_2 + \mathbf{k}_3 - \mathbf{k}_4) \cdot \mathbf{r}} W(\{\mathbf{k}_i\}) c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \sigma'}^\dagger c_{\mathbf{k}_4, \sigma'} \\ = \frac{W}{2Z} \sum_{\mathbf{r}} \sum_{\mathbf{r}' \rightarrow \text{NN of } \mathbf{r}} n_{\mathbf{r}', \sigma} n_{\mathbf{r}', \sigma'} \\ = \frac{W}{2} \sum_{\mathbf{r}} n_{\mathbf{r}, \sigma} n_{\mathbf{r}, \sigma'}, \end{aligned} \quad (3.15)$$

where we used the fact that each point $\mathbf{r}'$ (which we relabel as $\mathbf{r}$) appears Z times in the summation.

The complete lattice Hamiltonian, after applying the translation operators, is

$$H_{\text{latt}} = \varepsilon_d \sum_{\mathbf{r}, \sigma} n_{\mathbf{r}, \sigma} + U_d \sum_{\mathbf{r}} n_{\mathbf{r}, \uparrow} n_{\mathbf{r}, \downarrow} - \frac{2V}{Z} \sum_{\langle \mathbf{r}, \mathbf{r}' \rangle} c_{\mathbf{r}, \sigma}^\dagger c_{\mathbf{r}', \sigma} + \frac{2J_K}{Z} \sum_{\langle \mathbf{r}, \mathbf{r}' \rangle} \vec{S}_{\mathbf{r}} \cdot \vec{S}_{\mathbf{r}'} - \frac{W}{2} \sum_{\mathbf{r}} (n_{\mathbf{r}, \uparrow} - n_{\mathbf{r}, \downarrow})^2. \quad (3.16)$$

This is clearly an extended Hubbard model, with both 1-particle and spin-exchange nearest-neighbour hybridisation. The general form of the model is

$$H_{\text{ex-Hub}} = -t \sum_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger c_{j,\sigma} + \text{h.c.}) + J \sum_{\langle i,j \rangle} \mathbf{S}_i \cdot \mathbf{S}_j + U \sum_i n_{i,\uparrow} n_{i,\downarrow} - \mu \sum_{i,\sigma} n_{i,\sigma} , \quad (3.17)$$

such that the effective couplings resulting from the tiling operation is

$$t = \frac{2V}{Z}, \quad J = \frac{2J_K}{Z}, \quad U = U_d - W, \quad \mu = -\varepsilon_d + \frac{W}{2} . \quad (3.18)$$

4.3.3 Form Of The Eigenstates: Bloch's Theorem

The tiled Hamiltonian defined in eq. 3.11 is manifestly symmetric under discrete translation, given that it's expressed in terms of the translation operators. The fact that the Hamiltonian H_{latt} commutes with the manybody translation operator implies that the total crystal momentum is a conserved quantity. We will now use this to write down a “tiling” relation between the eigenstates of the auxiliary model and those of the lattice model.

In the tight-binding approach to lattice problems, the full Hamiltonian is obtained by translating the dynamics of the localised orbitals at each site over the entire lattice. The full eigenstate $|\Psi\rangle$ is obtained by constructing linear combinations of the eigenstates $|\psi(i)\rangle$ of the local Hamiltonians. According to Bloch's theorem, $|\Psi\rangle$ is of the form $|\Psi_{n,\mathbf{K}}\rangle = \sum_i e^{i\mathbf{K}\cdot\mathbf{r}_i} |\psi_n(i)\rangle$, where $\mathbf{K}$ is the single-particle crystal momentum, n is the band-index that labels various states within the local orbital, and $\mathbf{r}_i$ sums over the positions of the local Hamiltonians. Bloch's theorem ensures that eigenstates satisfy the following relation under a translation operation by an arbitrary number of lattice spacings $n\mathbf{a}$:

$$T^\dagger(l\mathbf{a}) |\Psi_{n,\mathbf{K}}\rangle = \sum_i e^{i\mathbf{K}\cdot\mathbf{r}_i} |\psi_n(i+l)\rangle = e^{-il\mathbf{K}\cdot\mathbf{a}} |\Psi_{n,\mathbf{K}}\rangle . \quad (3.19)$$

As discussed earlier, the eigenstates $|\Psi_{n,\mathbf{K}}\rangle$ of the lattice Hamiltonian obtained using eq. 3.11 enjoy the discrete translation symmetry of the lattice, and therefore enjoy a *many-body* Bloch's theorem [Stoyanova(2006)]. This means that the *local* eigenstates $|\psi_n(\mathbf{r}_d)\rangle$ (with the impurity located at an arbitrary position $\mathbf{r}_d$) of the unit cell auxiliary model Hamiltonian $H_{\text{clust}}(\mathbf{r}_d)$ defined in eq. 3.9 can be used to construct eigenstates of the lattice Hamiltonian. The index $n(= 0, 1, \dots)$ in the subscript indicates that it is the n^{th} eigenstate of the auxiliary model. The following combination of the auxiliary model eigenstates satisfies a many-particle equivalent of Bloch's theorem [Stoyanova(2006)]:

$$|\Psi_s\rangle \equiv |\Psi_{\mathbf{K},n}\rangle = \frac{1}{\sqrt{N}} \sum_{\mathbf{r}_d} e^{i\mathbf{K}\cdot\mathbf{r}_d} |\psi_n(\mathbf{r}_d)\rangle , \quad (3.20)$$

where N is the total number of lattice sites and $\mathbf{r}_d$ is summed over all lattice spacings. The set of $n = 0$ states form the lowest band in the spectrum of the lattice, while higher values of n produce the more energetic bands.

4.4 Properties of the Bloch states

4.4.1 Translation invariance

It is easy to verify that $\Psi_{\vec{k}}(\{\mathbf{r}_k\})$ transforms like Bloch functions under translation by a displacement $\mathbf{r}$:

$$\begin{aligned}\Psi_{\vec{k}}(\{\mathbf{r}_k + \mathbf{r}\}) &= \frac{1}{\lambda_{|\vec{k}\rangle} \mathcal{Z} N} \sum_{\mathbf{r}_i, \mathbf{a}} e^{i\vec{k} \cdot \vec{R}_i} \psi_{\text{aux}}(\mathbf{r}_i - \mathbf{r}, \mathbf{a}) = \frac{1}{\lambda_{|\vec{k}\rangle} \mathcal{Z} N} \sum_{\mathbf{r}_j, \mathbf{a}} e^{i\vec{k} \cdot (\vec{R}_j + \mathbf{r})} \psi_{\text{aux}}(\mathbf{r}_j, \mathbf{a}) \\ &= e^{i\vec{k} \cdot \vec{R}} \Psi_{\vec{k}}(\{\mathbf{r}_k\})\end{aligned}\quad (4.1)$$

In the last equation, we transformed $\mathbf{r}_i \rightarrow \mathbf{r}_j = \mathbf{r}_i - \mathbf{r}$. Note that the argument $\mathbf{a}$ does not change under a translation of the system, because that vector always represents the difference between the impurity lattice position and its nearest neighbours, irrespective of the absolute position of the impurity.

The wavefunction can be even brought into the familiar Bloch function form:

$$\begin{aligned}\Psi_{\vec{k}}^n(\{\mathbf{r}_k\}) &= \sum_{\mathbf{r}_i, \mathbf{a}} \frac{e^{i\vec{k} \cdot \vec{R}_i}}{\lambda_{|\vec{k}\rangle} \mathcal{Z} N} \psi_{\text{aux}}(\{\mathbf{r}_d - \mathbf{r}_i, \mathbf{r}_0 - \mathbf{r}_i - \mathbf{a}\}) \\ &= \frac{e^{i\vec{k} \cdot \frac{1}{N} \sum_k \vec{r}_k}}{\lambda_{|\vec{k}\rangle} \mathcal{Z} N} \sum_{\mathbf{r}_i, \mathbf{a}} e^{-i\vec{k} \cdot (\frac{1}{N} \sum_k \mathbf{r}_k - \vec{R}_i)} \psi_{\text{aux}}(\{\mathbf{r}_d - \mathbf{r}_i, \mathbf{r}_0 - \mathbf{r}_i - \mathbf{a}\}) \\ &= e^{i\vec{k} \cdot \vec{r}_{\text{COM}}} \eta_{\vec{k}}^n(\{\mathbf{r}_k\})\end{aligned}\quad (4.2)$$

where $\mathbf{r}_{\text{COM}} = \frac{1}{N} \sum_k \mathbf{r}_k$ is the center-of-mass coordinate and

$$\eta_{\vec{k}}^n(\{\mathbf{r}_k\}) = \frac{1}{\lambda_{|\vec{k}\rangle} \mathcal{Z} N} \sum_{\mathbf{r}_i} e^{-i\vec{k} \cdot (\frac{1}{N} \sum_k \mathbf{r}_k - \vec{R}_i)} \psi_{\text{aux}}^n(\{\mathbf{r}_k - \mathbf{r}_i\}), \quad (4.3)$$

is the translation symmetric function. This form of the eigenstate allows the interpretation that tuning the Bloch momentum $\vec{k}$ corresponds to a translation of the center of mass of the system (or in this case, of the auxiliary models that comprise the system).

4.4.2 Orthonormality

It is straightforward to show that these states form an orthonormal basis. We start by writing down the inner product of two distinct such states:

$$\begin{aligned}\langle \Psi_{\vec{k}'} | \Psi_{\vec{k}} \rangle &= \frac{1}{\lambda_{|\vec{k}'\rangle}^* \lambda_{|\vec{k}\rangle} \mathcal{Z}^2 N^2} \sum_{\mathbf{r}_i, \mathbf{r}_j, \mathbf{a}, \mathbf{a}'} e^{i(\vec{k} \cdot \vec{R}_i - \vec{k}' \cdot \vec{R}_j)} \langle \psi_{\text{aux}}(\mathbf{r}_j, \mathbf{a}') | \psi_{\text{aux}}(\mathbf{r}_i, \mathbf{a}) \rangle \\ &= \frac{1}{\lambda_{|\vec{k}'\rangle}^* \lambda_{|\vec{k}\rangle} \mathcal{Z}^2 N^2} \sum_{\mathbf{r}_i, \mathbf{r}_j, \mathbf{a}, \mathbf{a}'} e^{i(\vec{k} \cdot \vec{R}_i - \vec{k}' \cdot \vec{R}_j)} \langle \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}') T^\dagger(\mathbf{r}_0 - \mathbf{r}_j) T(\mathbf{r}_0 - \mathbf{r}_i) | \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}) \rangle\end{aligned}\quad (4.4)$$

At this point, we insert a complete basis of momentum eigenkets $1 = \sum_{\vec{q}} |\vec{q}\rangle \langle \vec{q}|$ to resolve the translation operators:

$$\begin{aligned}
 \langle \Psi_{\vec{k}'} | \Psi_{\vec{k}} \rangle &= \frac{1}{\lambda_{|\vec{k}'}^* \lambda_{|\vec{k}} \mathcal{Z}^2 N^2} \sum_{\mathbf{r}_i, \mathbf{r}_j, \mathbf{a}, \mathbf{a}', \vec{q}} e^{i(\vec{k} \cdot \vec{R}_i - \vec{k}' \cdot \vec{R}_j)} \langle \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}') T(\mathbf{r}_j - \mathbf{r}_i) |\vec{q}\rangle \langle \vec{q}| \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}) \rangle \\
 &= \frac{1}{\lambda_{|\vec{k}'}^* \lambda_{|\vec{k}} \mathcal{Z}^2 N^2} \sum_{\mathbf{r}_i, \mathbf{r}_j, \mathbf{a}, \mathbf{a}', \vec{q}} e^{i(\vec{k} \cdot \vec{R}_i - \vec{k}' \cdot \vec{R}_j)} \langle \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}') e^{i\vec{q} \cdot (\mathbf{r}_j - \mathbf{r}_i)} |\vec{q}\rangle \langle \vec{q}| \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}) \rangle \\
 &= \frac{1}{|\lambda_{|\vec{k}}|^2 \mathcal{Z}^2} \delta_{\vec{k}, \vec{k}'} \sum_{\mathbf{a}, \mathbf{a}'} \langle \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}') | \vec{k} \rangle \langle \vec{k} | \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}) \rangle
 \end{aligned} \tag{4.5}$$

At the last step, we used the summation form of the Kronecker delta function: $\sum_{\mathbf{r}_i} e^{i\mathbf{r}_i \cdot (\vec{k} - \vec{q})} \sum_{\mathbf{r}_j} e^{i\mathbf{r}_j \cdot (\vec{q} - \vec{k}')} = N \delta_{\vec{k}, \vec{q}} N \delta_{\vec{k}', \vec{q}}$. The final remaining step is to identify that the inner products inside the summation are actually independent of the direction $\mathbf{a}, \mathbf{a}'$ of the connecting vector, and both are equal to the normalisation factor $\lambda_{|\vec{k}}|$:

$$\langle \Psi_{\vec{k}'} | \Psi_{\vec{k}} \rangle = \frac{1}{|\lambda_{|\vec{k}}|^2 \mathcal{Z}^2} \delta_{\vec{k}, \vec{k}'} \sum_{\mathbf{a}, \mathbf{a}'} |\lambda_{|\vec{k}}|^2 = \frac{1}{|\lambda_{|\vec{k}}|^2 \mathcal{Z}^2} \delta_{\vec{k}, \vec{k}'} w^2 |\lambda_{|\vec{k}}|^2 = \delta_{\vec{k}, \vec{k}'} \tag{4.6}$$

This concludes the proof of orthonormality.

4.4.3 Eigenstates

To demonstrate that these are indeed eigenstates of the bulk Hamiltonian, we will calculate the matrix elements of the full Hamiltonian between these states:

$$\begin{aligned}
 \langle \Psi_{\vec{k}'} | H_{\text{H-H}} | \Psi_{\vec{k}} \rangle &= \frac{1}{\lambda_{|\vec{k}'}^* \lambda_{|\vec{k}} \mathcal{Z}^2 N^2} \sum_{\mathbf{r}_i, \mathbf{r}_j, \mathbf{r}_k, \mathbf{a}, \mathbf{a}'} e^{i(\vec{k} \cdot \vec{R}_i - \vec{k}' \cdot \vec{R}_j)} \langle \psi_{\text{aux}}(\mathbf{r}_j, \mathbf{a}') | H_{\text{aux}}(\mathbf{r}_k) | \psi_{\text{aux}}(\mathbf{r}_i, \mathbf{a}) \rangle \\
 &= \frac{1}{\lambda_{|\vec{k}'}^* \lambda_{|\vec{k}} \mathcal{Z}^2 N^2} \sum_{\mathbf{r}_i, \mathbf{r}_j, \mathbf{r}_k, \mathbf{a}, \mathbf{a}'} e^{i(\vec{k} \cdot \vec{R}_i - \vec{k}' \cdot \vec{R}_j)} \langle \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}') | T_{\mathbf{r}_0 - \mathbf{r}_j}^\dagger T_{\mathbf{r}_0 - \mathbf{r}_k} H_{\text{aux}}(\mathbf{r}_0) T_{\mathbf{r}_0 - \mathbf{r}_k}^\dagger T_{\mathbf{r}_0 - \mathbf{r}_i} | \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}) \rangle \\
 &= \frac{1}{\lambda_{|\vec{k}'}^* \lambda_{|\vec{k}} \mathcal{Z}^2 N^2} \sum_{\mathbf{r}_i, \mathbf{r}_j, \mathbf{r}_k, \mathbf{a}, \mathbf{a}'} e^{i(\vec{k} \cdot \vec{R}_i - \vec{k}' \cdot \vec{R}_j)} \langle \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}') | T_{\mathbf{r}_k - \mathbf{r}_j}^\dagger H_{\text{aux}}(\mathbf{r}_0) T_{\mathbf{r}_k - \mathbf{r}_i} | \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}) \rangle
 \end{aligned} \tag{4.7}$$

To resolve the translation operators, we will now insert the momentum eigenstates $|\vec{k}\rangle$ of the full model: $1 = \sum_{\vec{q}} |\vec{q}\rangle \langle \vec{q}|$ in between the operators.

$$\begin{aligned}
 \langle \Psi_{\vec{k}'} | H_{\text{H-H}} | \Psi_{\vec{k}} \rangle &= \sum_{\substack{\mathbf{r}_i, \mathbf{r}_j, \mathbf{r}_k, \\ \mathbf{a}, \mathbf{a}', \vec{q}, \vec{q}'}} \frac{e^{i(\vec{k} \cdot \vec{R}_i - \vec{k}' \cdot \vec{R}_j)}}{\lambda_{|\vec{k}'\rangle}^* \lambda_{|\vec{k}\rangle} \mathcal{Z}^2 N^2} \langle \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}') | \vec{q}' \rangle \langle \vec{q}' | T_{\mathbf{r}_k - \mathbf{r}_j}^\dagger H_{\text{aux}}(\mathbf{r}_0) T_{\mathbf{r}_k - \mathbf{r}_i} | \vec{q} \rangle \langle \vec{q} | \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}) \rangle \\
 &= \sum_{\substack{\mathbf{r}_i, \mathbf{r}_j, \mathbf{r}_k, \\ \mathbf{a}, \mathbf{a}', \vec{q}, \vec{q}'}} \frac{e^{i(\vec{k} \cdot \vec{R}_i - \vec{k}' \cdot \vec{R}_j)}}{\lambda_{|\vec{k}'\rangle}^* \lambda_{|\vec{k}\rangle} \mathcal{Z}^2 N^2} \langle \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}') | \vec{q}' \rangle e^{-i\vec{q}' \cdot (\mathbf{r}_k - \mathbf{r}_j)} e^{i\vec{q}' \cdot (\mathbf{r}_k - \mathbf{r}_i)} \langle \vec{q}' | H_{\text{aux}}(\mathbf{r}_0) | \vec{q} \rangle \langle \vec{q} | \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}) \rangle \\
 &= \frac{N}{\lambda_{|\vec{k}'\rangle}^* \lambda_{|\vec{k}\rangle} \mathcal{Z}^2} \sum_{\mathbf{a}, \mathbf{a}', \vec{q}, \vec{q}'} \delta_{\vec{k}, \vec{q}} \delta_{\vec{q}, \vec{q}'} \delta_{\vec{q}', \vec{k}} \langle \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}') | \vec{q}' \rangle \langle \vec{q}' | H_{\text{aux}}(\mathbf{r}_0) | \vec{q} \rangle \langle \vec{q} | \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}) \rangle \\
 &= \frac{N}{|\lambda_{|\vec{k}\rangle}|^2 \mathcal{Z}^2} \delta_{kk'} \sum_{\mathbf{a}, \mathbf{a}'} \langle \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}') | \vec{k} \rangle \langle \vec{k} | H_{\text{aux}}(\mathbf{r}_0) | \vec{k} \rangle \langle \vec{k} | \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}) \rangle \\
 &= \frac{N}{|\lambda_{|\vec{k}\rangle}|^2 \mathcal{Z}^2} \delta_{kk'} \langle \vec{k} | H_{\text{aux}}(\mathbf{r}_0) | \vec{k} \rangle \underbrace{\sum_{\mathbf{a}, \mathbf{a}'} \langle \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}') | \vec{k} \rangle \langle \vec{k} | \psi_{\text{aux}}(\mathbf{r}_0, \mathbf{a}) \rangle}_{w^2 |\lambda_{|\vec{k}\rangle}|^2} \\
 &= N \delta_{kk'} \langle \vec{k} | H_{\text{aux}}(\mathbf{r}_0) | \vec{k} \rangle
 \end{aligned} \tag{4.8}$$

The momentum eigenket $|\vec{k}\rangle$ can be expressed as a sum over position eigenkets at varying distances from $\mathbf{r}_0$. This form allows a systematic method of improving the energy eigenvalue estimate, because the interacting in the impurity model is extremely localised, so the overlap between two auxiliary models decreases rapidly with distance. The majority of the contribution will come from the state at $\mathbf{r}_0$, and improvements are then made by considering auxiliary models at further distances.

4.5 Mapping Greens Functions from Auxiliary Model to Lattice Model

4.5.1 One-particle k -space Greens functions

In the previous part, we proposed a form for the eigenstates $|\Psi_s\rangle$ of the bulk Hamiltonian in terms of the ground-states $|\psi_n\rangle$ of the auxiliary models. In this section, we will relate one-particle Greens functions of the lattice model to those of the auxiliary model. We define the retarded time-domain lattice k -space Greens function at zero temperature as

$$G(\mathbf{k}\sigma; t) = -i\theta(t) \langle \Psi_{\text{gs}} | \{c_{\mathbf{k}\sigma}(t), c_{\mathbf{k}\sigma}^\dagger\} | \Psi_{\text{gs}} \rangle. \tag{5.1}$$

where $|\Psi_{\text{gs}}\rangle$ is the lattice groundstate. Time evolution of the operators is governed by the lattice Hamiltonian H_{latt} :

$$c_{\mathbf{k}\sigma}(t) = e^{itH_{\text{latt}}} c_{\mathbf{k}\sigma} e^{-itH_{\text{latt}}}. \tag{5.2}$$

By using eq. 3.8 and the fact that the translation operators commute with H_{latt} , we have the relation

$$T(\vec{a}) c(\mathbf{k})(t) T^\dagger(\vec{a}) = e^{-i\mathbf{k} \cdot \vec{a}} c(\mathbf{k})(t), \quad (5.3)$$

which combined with eq. 3.8 gives

$$T(\mathbf{a}) c(\mathbf{k})(t) c^\dagger(\mathbf{k}) T^\dagger(\mathbf{b}) = e^{i\mathbf{k} \cdot (\mathbf{b} - \mathbf{a})} c(\mathbf{k})(t) T^\dagger(\mathbf{b} - \mathbf{a}) c^\dagger(\mathbf{k}). \quad (5.4)$$

In order to make use of the mapping between auxiliary model and lattice model Hamiltonians, we would like to express this lattice Greens function in terms of auxiliary model Greens functions – objects that can be computed purely from within a single auxiliary model. This will be a major simplification in our computations, because it is easier to work with a quantum impurity model, as compared to an interacting lattice model.

To make progress towards this, we consider the equivalent object

$$G(\mathbf{k}\sigma; t) = -i\theta(t) \frac{1}{2} \left[\langle \Psi_{\text{gs}} | \{c_{\mathbf{k}\sigma}(t), c_{\mathbf{k}\sigma}^\dagger\} | \Psi_{\text{gs}} \rangle + \langle \Psi_{\text{gs}} | \{c_{\mathbf{k}\sigma}, c_{\mathbf{k}\sigma}^\dagger(-t)\} | \Psi_{\text{gs}} \rangle \right]. \quad (5.5)$$

We consider only the first term for now. We express the lattice groundstate in terms of the auxiliary model groundstates (using eq. 3.20):

$$\langle \Psi_{\text{gs}} | \{c_{\mathbf{k}\sigma}(t), c_{\mathbf{k}\sigma}^\dagger\} | \Psi_{\text{gs}} \rangle = \frac{1}{N} \sum_{\mathbf{r}_1, \mathbf{r}_2} \langle \psi_{\text{gs}}(\mathbf{r}_2) | \{c_{\mathbf{k}\sigma}(t), c_{\mathbf{k}\sigma}^\dagger\} | \psi_{\text{gs}}(\mathbf{r}_1) \rangle. \quad (5.6)$$

We take one of the terms in the anticommutator and proceed to simplify it:

$$\begin{aligned} \sum_{\mathbf{r}_1, \mathbf{r}_2} \langle \psi_{\text{gs}}(\mathbf{r}_2) | c_{\mathbf{k}\sigma}(t) c_{\mathbf{k}\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_1) \rangle &= \sum_{\mathbf{r}_1, \mathbf{r}_2} \langle \psi_{\text{gs}}(\mathbf{r}_d) | T(\mathbf{r}_2 - \mathbf{r}_d) c_{\mathbf{k}\sigma}(t) c_{\mathbf{k}\sigma}^\dagger T^\dagger(\mathbf{r}_1 - \mathbf{r}_d) | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &= \sum_{\mathbf{r}_1, \mathbf{r}_2} e^{i\mathbf{k} \cdot (\mathbf{r}_1 - \mathbf{r}_2)} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{k}\sigma}(t) T^\dagger(\mathbf{r}_1 - \mathbf{r}_2) c_{\mathbf{k}\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &= N \sum_{\mathbf{r}} e^{i\mathbf{k} \cdot \mathbf{r}} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{k}\sigma}(t) T^\dagger(\mathbf{r}) c_{\mathbf{k}\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle. \end{aligned} \quad (5.7)$$

where $\mathbf{r}_d$ is a reference lattice site, and we used eq. 5.4 in the second step. The other term in the anticommutator gives

$$\begin{aligned} \sum_{\mathbf{r}_1, \mathbf{r}_2} \langle \psi_{\text{gs}}(\mathbf{r}_2) | c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma}(t) | \psi_{\text{gs}}(\mathbf{r}_1) \rangle &= \sum_{\mathbf{r}_1, \mathbf{r}_2} \langle \psi_{\text{gs}}(\mathbf{r}_d) | T(\mathbf{r}_2 - \mathbf{r}_d) c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma}(t) T^\dagger(\mathbf{r}_1 - \mathbf{r}_d) | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &= \sum_{\mathbf{r}_1, \mathbf{r}_2} e^{-i\mathbf{k} \cdot (\mathbf{r}_1 - \mathbf{r}_2)} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{k}\sigma}^\dagger T^\dagger(\mathbf{r}_1 - \mathbf{r}_2) c_{\mathbf{k}\sigma}(t) | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &= N \sum_{\mathbf{r}} e^{-i\mathbf{k} \cdot \mathbf{r}} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{k}\sigma}^\dagger T^\dagger(\mathbf{r}) c_{\mathbf{k}\sigma}(t) | \psi_{\text{gs}}(\mathbf{r}_d) \rangle. \end{aligned} \quad (5.8)$$

To resolve the time evolution of the operators, we insert $1 = \sum_m |\psi_m(\mathbf{r}_d)\rangle \langle \psi_m(\mathbf{r}_d)|$ into the equation. We again work with just the first term of the anticommutator for convenience:

$$\begin{aligned}
 & \sum_{\mathbf{r}_1, \mathbf{r}_2} \langle \psi_{\text{gs}}(\mathbf{r}_2) | c_{\mathbf{k}\sigma}(t) c_{\mathbf{k}\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_1) \rangle \\
 &= N \sum_{\mathbf{r}, n_1, n_2} e^{i\mathbf{k} \cdot \mathbf{r}} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{k}\sigma}(t) | \psi_{n_1}(\mathbf{r}_d) \rangle \langle \psi_{n_1}(\mathbf{r}_d) | T^\dagger(\mathbf{r}) | \psi_{n_2}(\mathbf{r}_d) \rangle \langle \psi_{n_2}(\mathbf{r}_d) | c_{\mathbf{k}\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\
 &= N \sum_{\mathbf{r}, n_1, n_2} e^{i\mathbf{k} \cdot \mathbf{r}} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{k}\sigma}(t) | \psi_{n_1}(\mathbf{r}_d) \rangle \langle \psi_{n_1}(\mathbf{r}_d) | \psi_{n_2}(\mathbf{r}_d + \mathbf{r}) \rangle \langle \psi_{n_2}(\mathbf{r}_d) | c_{\mathbf{k}\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle .
 \end{aligned} \tag{5.9}$$

The overlap $\langle \psi_{n_1}(\mathbf{r}_d) | \psi_{n_2}(\mathbf{r}_d + \mathbf{r}) \rangle$ between the eigenstates of the auxiliary models at $\mathbf{r}_d$ and $\mathbf{r}_d + \mathbf{r}$ are appreciable only for $n_1 = n_2$. This is because, for small $\mathbf{r}$, the auxiliary models are very similar and the states are therefore almost orthogonal. For large $\mathbf{r}$, the overlap is again suppressed because the auxiliary model eigenstates are localised around the impurity. We therefore make the simplifying assumption: $\langle \psi_{n_1}(\mathbf{r}_d) | \psi_{n_2}(\mathbf{r}_d + \mathbf{r}) \rangle = \delta_{n_1, n_2} \langle \psi_{n_1}(\mathbf{r}_d) | \psi_{n_1}(\mathbf{r}_d + \mathbf{r}) \rangle = \delta_{n_1, n_2} F_{n_1}(\mathbf{r})$. Substituting this, we get

$$\sum_{\mathbf{r}_1, \mathbf{r}_2} \langle \psi_{\text{gs}}(\mathbf{r}_2) | c_{\mathbf{k}\sigma}(t) c_{\mathbf{k}\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_1) \rangle = N \sum_n F_n(\mathbf{k}) \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{k}\sigma}(t) | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | c_{\mathbf{k}\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle , \tag{5.10}$$

where we have defined the Fourier transform $F_n(\mathbf{k}) = \sum_{\mathbf{r}} e^{i\mathbf{k} \cdot \mathbf{r}} F_n(\mathbf{r})$ of the overlap function.

We now consider more carefully the transition operator $c_{\mathbf{k}\sigma}$ for the 1-particle excitation giving rise to the above Greens function. Within our auxiliary model approach, gapless excitations within the lattice model are represented by gapless excitations of the impurity site, specifically those that try to screen the impurity site. As a result, the uncoordinated scattering process $c_{\mathbf{k}\sigma}$ for the lattice model must be replaced by a coherent excitation $\mathcal{T}$ within the impurity model that captures those gapless excitations that occur in connection with the impurity, and projects out the uncorrelated excitations that take place even when the impurity site is decoupled from the bath.

In order to construct this auxiliary model $\mathcal{T}$ -matrix, we allow for all excitations of the impurity site in conjunction with those of the k -state:

$$\mathcal{T}_{\mathbf{k}, \sigma} = [c_{\mathbf{r}_d \sigma}^\dagger + S_{\mathbf{r}_d}^\sigma + C_{\mathbf{r}_d}^+] c_{\mathbf{k}\sigma} , \tag{5.11}$$

where $S_{\mathbf{r}_d}^+$ and $C_{\mathbf{r}_d}^+$ are spin and charge flip operators on the impurity site that cause transition from $|\downarrow\rangle \rightarrow |\uparrow\rangle$ and $|0\rangle \rightarrow |\uparrow\downarrow\rangle$ respectively. The first term encodes the local component of the k -space Greens function by creating an excitation on the impurity site (which is then translated across the lattice), and the rest of the terms encode the non-local components by tracking impurity-bath excitations.

Apart from modifying the excitation operator, we will now simplify the time-evolution as well. The first overlap on the right-hand side of eq. 5.10 expands to

$$\langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k}, \sigma}(t) | \psi_n(\mathbf{r}_d) \rangle = \langle \psi_{\text{gs}}(\mathbf{r}_d) | e^{i(H_{\text{aux}}(\mathbf{r}_d) + H_{\text{cav}}(\mathbf{r}_d))t} \mathcal{T}_{\mathbf{k}, \sigma} e^{-i(H_{\text{aux}}(\mathbf{r}_d) + H_{\text{cav}}(\mathbf{r}_d))t} | \psi_n(\mathbf{r}_d) \rangle , \tag{5.12}$$

where $H_{\text{aux}}(\mathbf{r}_d)$ (eq. 2.3) is the auxiliary model Hamiltonian consisting of the cluster Hamiltonian $H_{\text{clust}}(\mathbf{r}_d)$ along with all one-particle terms H_{1p} in the Hamiltonian (in a typical lattice model, this

will be the tight-binding hopping terms), and $H_{\text{cav}}(\mathbf{r}_d)$ is the “cavity” Hamiltonian consisting of all interaction (2-particle and beyond) terms in the Hamiltonian. Together, they create the full lattice Hamiltonian: $H_{\text{latt}} = H_{\text{aux}}(\mathbf{r}_d) + H_{\text{cav}}(\mathbf{r}_d)$. Using the Zassenhaus form of the Baker-Campbell-Hausdorff formula, we have

$$e^{i(H_{\text{aux}}(\mathbf{r}_d) + H_{\text{cav}}(\mathbf{r}_d))t} \approx e^{iH_{\text{aux}}(\mathbf{r}_d)t} e^{iH_{\text{cav}}(\mathbf{r}_d)t} e^{[H_{\text{aux}}(\mathbf{r}_d), H_{\text{cav}}(\mathbf{r}_d)]t^2/2}, \quad (5.13)$$

where we have dropped the higher-order terms because the commutators involve only the boundary degrees of freedom and become subsequently smaller as the order increases. One can think of the operator $e^{iH_{\text{cav}}(\mathbf{r}_d)t} e^{[H_{\text{aux}}(\mathbf{r}_d), H_{\text{cav}}(\mathbf{r}_d)]t^2/2}$ as renormalising the auxiliary model $\mathcal{T}$ -matrix through hybridisation with the rest of the system and formally leading to a new $\mathcal{T}$ -matrix:

$$\mathcal{T}_{\mathbf{k},\sigma}^* = e^{iH_{\text{cav}}(\mathbf{r}_d)t} e^{[H_{\text{aux}}(\mathbf{r}_d), H_{\text{cav}}(\mathbf{r}_d)]t^2/2} \mathcal{T}_{\mathbf{k},\sigma} e^{-[H_{\text{aux}}(\mathbf{r}_d), H_{\text{cav}}(\mathbf{r}_d)]t^2/2} e^{-iH_{\text{cav}}(\mathbf{r}_d)t}. \quad (5.14)$$

One can also absorb the modulating phase factor $F_n(\mathbf{k})$ (defined above and appearing in eq. 5.10) into this renormalisation:

$$\mathcal{T}_{\mathbf{k},\sigma}^* = e^{iH_{\text{cav}}(\mathbf{r}_d)t} e^{[H_{\text{aux}}(\mathbf{r}_d), H_{\text{cav}}(\mathbf{r}_d)]t^2/2} \mathcal{T}_{\mathbf{k},\sigma} F_n(\mathbf{k}) \mathcal{P}_n(\mathbf{r}_d) e^{-[H_{\text{aux}}(\mathbf{r}_d), H_{\text{cav}}(\mathbf{r}_d)]t^2/2} e^{-iH_{\text{cav}}(\mathbf{r}_d)t}, \quad (5.15)$$

where $\mathcal{P}_n(\mathbf{r}_d)$ projects onto the n^{th} eigenstate $|\psi_n(\mathbf{r}_d)\rangle$.

Since the overlap is calculated in the eigenstates of the auxiliary model at $\mathbf{r}_d$, we adopt the zeroth order approximation of ignoring this renormalisation. Later, we will specify a prescription of systematically improving our computation. The overlap expression becomes

$$\langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k},\sigma}(t) | \psi_n(\mathbf{r}_d) \rangle = \langle \psi_{\text{gs}}(\mathbf{r}_d) | e^{i(H_{\text{clust}}(\mathbf{r}_d) + H_{1p})t} \mathcal{T}_{\mathbf{k},\sigma} e^{-i(H_{\text{clust}}(\mathbf{r}_d) + H_{1p})t} | \psi_n(\mathbf{r}_d) \rangle. \quad (5.16)$$

To make the problem tractable, we treat the effect of H_{1p} at the level of an expectation value and note that the effect of the transition operator is to create a hole of momentum $\mathbf{k}$ (eq. 5.11). Let the Hamiltonian H_{1p} be diagonal in certain modes $c_{\nu,\sigma}$ with energies ϵ_ν , and the operator $c_{\mathbf{k},\sigma}$ can be expressed in terms of these modes: $c_{\mathbf{k},\sigma} = \sum_\nu C_{\nu,\mathbf{k}} c_{\nu,\sigma}$. Expanding the $\mathcal{T}$ -matrix in this fashion gives

$$\langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k},\sigma}(t) | \psi_n(\mathbf{r}_d) \rangle = \sum_\nu C_{\nu,\mathbf{k}} \langle \psi_{\text{gs}}(\mathbf{r}_d) | e^{i(H_{\text{clust}}(\mathbf{r}_d) + H_{1p})t} \mathcal{T}_{\nu,\sigma} e^{-i(H_{\text{clust}}(\mathbf{r}_d) + H_{1p})t} | \psi_n(\mathbf{r}_d) \rangle. \quad (5.17)$$

Each excitation $\mathcal{T}_{\nu,\sigma}$ therefore leads to an energy cost $-\epsilon_\nu$. Moreover, $|\psi_n(\mathbf{r}_d)\rangle$ and $|\psi_{\text{gs}}(\mathbf{r}_d)\rangle$ are eigenstates of $H_{\text{clust}}(\mathbf{r}_d)$ with energies E_n and E_0 . This leads to the following simplified expression:

$$\langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k},\sigma}(t) | \psi_n(\mathbf{r}_d) \rangle = \sum_\nu C_{\nu,\mathbf{k}} e^{-i(E_n - E_0 + \epsilon_\nu)t} \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\nu,\sigma} | \psi_n(\mathbf{r}_d) \rangle. \quad (5.18)$$

Substituting the expression back into 5.10 and then into 5.6, we get an expression for one half of the first part of the retarded Greens function (the full form is in eq. 5.5):

$$\begin{aligned} & -i\theta(t) \langle \Psi_{\text{gs}} | c_{\mathbf{k}\sigma}(t) c_{\mathbf{k}\sigma}^\dagger | \Psi_{\text{gs}} \rangle \\ & = -i\theta(t) \sum_{\nu,n} C_{\nu,\mathbf{k}} e^{-i(E_n - E_0 + \epsilon_\nu)t} \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\nu,\sigma} | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k},\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle. \end{aligned} \quad (5.19)$$

Working through the other term of eq. 5.6 with the same strategy, we get the full expression:

$$\begin{aligned}
 & -i\theta(t) \langle \Psi_{\text{gs}} | \{c_{\mathbf{k}\sigma}(t), c_{\mathbf{k}\sigma}^\dagger\} | \Psi_{\text{gs}} \rangle \\
 & = -i\theta(t) \sum_{\nu, n} C_{\nu, \mathbf{k}} e^{-i(E_n - E_0 + \epsilon_\nu)t} \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\nu, \sigma} | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k}, \sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\
 & \quad - i\theta(t) \sum_{\nu, n} C_{\nu, \mathbf{k}} e^{i(E_n - E_0 - \epsilon_\nu)t} \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k}, \sigma}^\dagger | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\nu, \sigma} | \psi_{\text{gs}}(\mathbf{r}_d) \rangle .
 \end{aligned} \tag{5.20}$$

Finally, we consider the other term in eq. 5.5, $-i\theta(t) \langle \Psi_{\text{gs}} | \{c_{\mathbf{k}\sigma}, c_{\mathbf{k}\sigma}^\dagger(-t)\} | \Psi_{\text{gs}} \rangle$. Proceeding in the same way, eq. 5.18 is replaced by

$$\begin{aligned}
 \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k}, \sigma}^\dagger(-t) | \psi_{\text{gs}}(\mathbf{r}_d) \rangle & = \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k}, \sigma}(-t) | \psi_n(\mathbf{r}_d) \rangle^* \\
 & = \sum_{\nu} C_{\nu, \mathbf{k}}^* e^{-i(E_n - E_0 + \epsilon_\nu)t} \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\nu, \sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle .
 \end{aligned} \tag{5.21}$$

Evaluating this term exactly as before and changing from time to frequency domain using $g(\omega) = \int dt e^{i\omega T} f(t)$ gives

$$\begin{aligned}
 G(\mathbf{k}, \sigma; \omega) & = \sum_{\nu, n} C_{\nu, \mathbf{k}} \left[\frac{\langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\nu, \sigma} | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k}, \sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle}{\omega - (E_n - E_0 + \epsilon_\nu)} \right. \\
 & \quad \left. + \frac{\langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k}, \sigma}^\dagger | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\nu, \sigma} | \psi_{\text{gs}}(\mathbf{r}_d) \rangle}{\omega + (E_n - E_0 - \epsilon_\nu)} \right] + \text{h.c.}
 \end{aligned} \tag{5.22}$$

We recognise the two parts of the lattice Greens function as a retarded off-diagonal Greens functions calculated within the auxiliary model at $\mathbf{r}_d$, but at shifted frequencies $\omega - \epsilon_\nu$:

$$G(\mathbf{k}, \sigma; \omega) = \sum_{\nu} \left[C_{\nu, \mathbf{k}} \mathcal{G}(\mathcal{T}_{\nu, \sigma}, \mathcal{T}_{\mathbf{k}, \sigma}^\dagger; \omega - \epsilon_\nu) + C_{\nu, \mathbf{k}}^* \mathcal{G}(\mathcal{T}_{\mathbf{k}, \sigma}, \mathcal{T}_{\nu, \sigma}^\dagger; \omega - \epsilon_\nu) \right] \tag{5.23}$$

We now discuss how to systematically improve this estimate. One way is to include the effect of other auxiliary models in the overlap function $F_n(\mathbf{r})$. This will of course require us to calculate an off-diagonal Greens function. As a concrete example, upon including the effects of the auxiliary models nearest to the reference point $\mathbf{r}_d$, the expression for the Greens function becomes

$$\begin{aligned}
 G(\mathbf{k}, \sigma; \omega) & = \sum_{\nu, n} C_{\nu, \mathbf{k}} \left[\frac{\langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\nu, \sigma}^* | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k}, \sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle}{\omega - \epsilon_\nu - (E_n - E_0)} \right. \\
 & \quad \left. + \frac{\langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k}, \sigma}^\dagger | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\nu, \sigma}^* | \psi_{\text{gs}}(\mathbf{r}_d) \rangle}{\omega - \epsilon_\nu + (E_n - E_0)} \right] ,
 \end{aligned} \tag{5.24}$$

with $\tau_{\nu, \sigma}^* = \mathcal{T}_{\nu, \sigma}(F_n(0) + \sum_{\mathbf{a}} F_n(\mathbf{a}) e^{i\mathbf{k} \cdot \mathbf{a}})$ where $\mathbf{a}$ sums over the primitive lattice vectors. Another way to include more non-local contributions is by considering the effects of the cavity Hamiltonian in eq. 5.15:

$$\mathcal{T}_{\mathbf{k}, \sigma}^*(t) = e^{iH_{\text{cav}}(\mathbf{r}_d)t} (\mathcal{T}_{\mathbf{k}, \sigma} + t^2/2 [[H_{\text{aux}}(\mathbf{r}_d), H_{\text{cav}}(\mathbf{r}_d)], \mathcal{T}_{\mathbf{k}, \sigma}]) e^{-iH_{\text{cav}}(\mathbf{r}_d)t} . \tag{5.25}$$

The time-domain Greens function associated with this operator will then need to be calculated, which can then be used to calculate frequency-domain functions.

4.5.2 One-particle real-space Greens functions

We now work with Greens functions for real-space excitations. The retarded Greens function in this case is

$$G(\mathbf{r}\sigma; t) = -i\theta(t) \langle \Psi_{\text{gs}} | \{c_{\mathbf{r}_d+\mathbf{r},\sigma}(t), c_{\mathbf{r}_d,\sigma}^\dagger\} | \Psi_{\text{gs}} \rangle . \quad (5.26)$$

Since our lattice model has discrete translation symmetry, we instead calculate the equivalent quantity

$$G(\mathbf{r}\sigma; t) = -i\theta(t) \frac{1}{2} \sum_{\mathbf{r}_d} [\langle \Psi_{\text{gs}} | \{c_{\mathbf{r}_d+\mathbf{r},\sigma}(t), c_{\mathbf{r}_d,\sigma}^\dagger\} | \Psi_{\text{gs}} \rangle + \langle \Psi_{\text{gs}} | \{c_{\mathbf{r}_d+\mathbf{r},\sigma}, c_{\mathbf{r}_d,\sigma}^\dagger(-t)\} | \Psi_{\text{gs}} \rangle] . \quad (5.27)$$

Similar to the k -space Greens functions, we consider only the first term for now, and express the groundstates as a superposition of auxiliary model Greens functions:

$$\begin{aligned} \sum_{\mathbf{r}_d} \langle \Psi_{\text{gs}} | c_{\mathbf{r}_d+\mathbf{r},\sigma}(t) c_{\mathbf{r}_d,\sigma}^\dagger | \Psi_{\text{gs}} \rangle &= \frac{1}{N} \sum_{\mathbf{R}, \Delta, \mathbf{r}_d} \langle \psi_{\text{gs}}(\mathbf{R} + \Delta) | c_{\mathbf{r}_d+\mathbf{r},\sigma}(t) c_{\mathbf{r}_d,\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{R}) \rangle \\ &= \frac{1}{N} \sum_{\mathbf{R}, \Delta} \langle \psi_{\text{gs}}(-\mathbf{r}_d) | c_{\mathbf{r}-\mathbf{R}-\Delta,\sigma}(t) T(\Delta) c_{-\mathbf{R},\sigma}^\dagger | \psi_{\text{gs}}(-\mathbf{r}_d) \rangle . \end{aligned} \quad (5.28)$$

Local Greens function: $\mathbf{r} = 0$

We first consider the case of local Greens function, by setting $\mathbf{r} = 0$. We also change the dummy variable $\mathbf{r}_d$, $\mathbf{R}$ and Δ to $-\mathbf{r}_d$, $\mathbf{R}$ and $-\Delta$ to make the expressions tidier:

$$\sum_{\mathbf{r}_d} \langle \Psi_{\text{gs}} | c_{\mathbf{r}_d,\sigma}(t) c_{\mathbf{r}_d,\sigma}^\dagger | \Psi_{\text{gs}} \rangle = \frac{1}{N} \sum_{\mathbf{R}, \Delta, \mathbf{r}_d} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{R}+\Delta,\sigma}(t) T^\dagger(\Delta) c_{\mathbf{R},\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle . \quad (5.29)$$

We split the full summation into four parts, based on whether $\mathbf{R} = \mathbf{r}_d$ and whether $\Delta + \mathbf{R} = \mathbf{r}_d$:

$$\begin{aligned} \sum_{\mathbf{r}_d} \langle \Psi_{\text{gs}} | c_{\mathbf{r}_d,\sigma}(t) c_{\mathbf{r}_d,\sigma}^\dagger | \Psi_{\text{gs}} \rangle &= \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{r}_d,\sigma}(t) c_{\mathbf{r}_d,\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\Delta \neq 0} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{r}_d+\Delta,\sigma}(t) T^\dagger(\Delta) c_{\mathbf{r}_d,\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{R} \neq \mathbf{r}_d} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{r}_d,\sigma}(t) T^\dagger(\mathbf{r}_d - \mathbf{R}) c_{\mathbf{R},\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{R} \neq \mathbf{r}_d, \Delta \neq \mathbf{r}_d} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\Delta,\sigma}(t) T^\dagger(\Delta - \mathbf{R}) c_{\mathbf{R},\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle , \end{aligned} \quad (5.30)$$

where we have used the fact that the location of the auxiliary model does not affect the correlation calculated within the auxiliary model in order to cancel out the summation over $\mathbf{r}_d$ with the factor of $1/N$. The reason for splitting the expression into these parts is to explicitly write out which operators act on the impurity site $\mathbf{r}_d$ and which act on other real-space sites away from the impurity

site. We will now replace the real-space excitation operators away from the impurity site with the associated T -matrices, following the discussion above eq. 5.11:

$$\mathcal{T}_{r,\sigma} = \left[\left(\sum_{\sigma'} c_{r_d\sigma}^\dagger + \text{h.c.} \right) + (S_{r_d}^+ + \text{h.c.}) + (C_{r_d}^+ + \text{h.c.}) \right] c_{r\sigma} . \quad (5.31)$$

We also insert a complete basis of states of the auxiliary model at $\mathbf{r}_d$, $1 = \sum_n |\psi_n(\mathbf{r}_d)\rangle \langle \psi_n(\mathbf{r}_d)|$, generating an overlap factor F_n as before:

$$\begin{aligned} \sum_{r_d} \langle \Psi_{\text{gs}} | c_{r_d,\sigma}(t) c_{r_d,\sigma}^\dagger | \Psi_{\text{gs}} \rangle &= \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{r_d,\sigma}(t) c_{r_d,\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{r}_1, \mathbf{r}_2 \neq \mathbf{r}_d, n} F_n(\mathbf{r}_1 - \mathbf{r}_2)^* \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathbf{r}_1,\sigma}(t) | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\mathbf{r}_2,\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle . \end{aligned} \quad (5.32)$$

Several terms have been dropped because those contributions vanish for systems where the eigenstates conserve total spin S_z .

We next treat the time-evolution of the operators exactly as in the k -space Greens function. We replace the one-particle Hamiltonian H_{1p} with its local component H_{loc} (such as any on-site term of inter-layer hopping, etc). As done previously, we will make use of the eigenstates $\nu(\mathbf{r})$ of H_{loc} , with energy $\epsilon_\nu^{\text{loc}}$. Without repeating the details, we state the final result here, for the frequency-domain real-space local Greens function:

$$\begin{aligned} G_{\text{loc}}(\omega) &= \frac{1}{2} \sum_{\nu} [C_{\nu, r_d} \mathcal{G}(\nu(r_d)\sigma, c_{r_d,\sigma}^\dagger; \tilde{\omega}) + C_{\nu, r_d}^* \mathcal{G}(c_{r_d,\sigma}, \nu^\dagger(r_d)\sigma; \tilde{\omega})] \\ &+ \frac{1}{2} \sum_{\mathbf{r}_1, \mathbf{r}_2 \neq \mathbf{r}_d, \nu} F_n(\mathbf{r}_1 - \mathbf{r}_2)^* [C_{\mathbf{r}_1, \nu} \mathcal{G}(\mathcal{T}_{\nu(\mathbf{r}_1)\sigma}, \mathcal{T}_{\mathbf{r}_2\sigma}^\dagger; \tilde{\omega}) + C_{\mathbf{r}_1, \nu}^* \mathcal{G}(\mathcal{T}_{\mathbf{r}_1\sigma}, \mathcal{T}_{\nu(\mathbf{r}_2)\sigma}^\dagger; \tilde{\omega})] , \end{aligned} \quad (5.33)$$

where the hermitian conjugate in each part comes from considering the second term in eq. 5.27. The shifted frequency $\tilde{\omega} = \omega - \epsilon_\nu^{\text{loc}}$ captures the effect of local terms in H_{1p} ; the shift is defined as $e^{i\epsilon_\nu^{\text{loc}}t} c_\nu^\dagger = e^{iH_{\text{loc}}t} c_\nu^\dagger e^{-iH_{\text{loc}}t}$. The operator $\nu^\dagger(\mathbf{r})$ represents the eigenstates of H_{loc} at any site $\mathbf{r}$. The coefficients $C_{\nu, \mathbf{r}}$ express the eigenstates in terms of local states: $c_{\mathbf{r}} = \sum_{\nu} C_{\nu, \mathbf{r}} c_\nu(\mathbf{r})$.

Of course, the states ν will differ from the local states $c_{\mathbf{r}}$ only if there are multiple local orbitals (say α) that can hybridise among each other. In such a case, one can also think of an inter-orbital Greens function, defined as

$$G_{\text{loc}}(\alpha, \alpha'; t) = -i\theta(t) \langle \Psi_{\text{gs}} | \{c_{r_d,\sigma,\alpha}(t), c_{r_d,\sigma,\alpha'}^\dagger\} | \Psi_{\text{gs}} \rangle . \quad (5.34)$$

Such a Greens function can also be calculated from our approach:

$$\begin{aligned} G_{\text{loc}}(\alpha, \alpha'; \omega) &= \frac{1}{2} \sum_{\nu} [C_{\nu, \alpha} \mathcal{G}(\nu(r_d, \alpha)\sigma, c_{r_d,\sigma,\alpha'}^\dagger; \tilde{\omega}) + C_{\nu, \alpha}^* \mathcal{G}(c_{r_d,\sigma,\alpha}, \nu^\dagger(r_d, \alpha')\sigma; \tilde{\omega})] \\ &+ \frac{1}{2} \sum_{\mathbf{r}_1, \mathbf{r}_2 \neq \mathbf{r}_d, \nu} F_n(\mathbf{r}_1 - \mathbf{r}_2)^* [C_{\alpha, \nu} \mathcal{G}(\mathcal{T}_{\nu(\mathbf{r}_1), \sigma, \alpha}, \mathcal{T}_{\mathbf{r}_2, \sigma, \alpha'}^\dagger; \tilde{\omega}) + C_{\alpha', \nu}^* \mathcal{G}(\mathcal{T}_{\mathbf{r}_1, \sigma, \alpha}, \mathcal{T}_{\nu(\mathbf{r}_2), \sigma, \alpha'}^\dagger; \tilde{\omega})] , \end{aligned} \quad (5.35)$$

Non-local Greens function: $\mathbf{r} \neq 0$

We next consider the case of non-local Greens function, where the excitations propagate a non-zero distance $\mathbf{r}$:

$$\sum_{\mathbf{r}_d} \langle \Psi_{\text{gs}} | c_{\mathbf{r}_d+\mathbf{r},\sigma}(t) c_{\mathbf{r}_d,\sigma}^\dagger | \Psi_{\text{gs}} \rangle = \frac{1}{N} \sum_{\mathbf{R}, \Delta, \mathbf{r}_d} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{R}+\Delta+\mathbf{r},\sigma}(t) T^\dagger(\Delta) c_{\mathbf{R},\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle . \quad (5.36)$$

To bring this expression closer to the one for the local Greens function, we redefine $\Delta + \mathbf{r} \rightarrow \Delta$:

$$\sum_{\mathbf{r}_d} \langle \Psi_{\text{gs}} | c_{\mathbf{r}_d+\mathbf{r},\sigma}(t) c_{\mathbf{r}_d,\sigma}^\dagger | \Psi_{\text{gs}} \rangle = \frac{1}{N} \sum_{\mathbf{R}, \Delta, \mathbf{r}_d} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{R}+\Delta,\sigma}(t) T^\dagger(\Delta) T(\mathbf{r}) c_{\mathbf{R},\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle . \quad (5.37)$$

This allows to again split the full summation based on whether $\mathbf{R} = \mathbf{r}_d$ and whether $\Delta + \mathbf{R} = \mathbf{r}_d$:

$$\begin{aligned} \sum_{\mathbf{r}_d} \langle \Psi_{\text{gs}} | c_{\mathbf{r}_d+\mathbf{r},\sigma}(t) c_{\mathbf{r}_d,\sigma}^\dagger | \Psi_{\text{gs}} \rangle &= \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{r}_d,\sigma}(t) T(\mathbf{r}) c_{\mathbf{r}_d,\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\Delta \neq 0} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{r}_d+\Delta,\sigma}(t) T^\dagger(\Delta - \mathbf{r}) c_{\mathbf{r}_d,\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{R} \neq \mathbf{r}_d} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{r}_d,\sigma}(t) T^\dagger(\mathbf{r}_d - \mathbf{R} - \mathbf{r}) c_{\mathbf{R},\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{R} \neq \mathbf{r}_d, \Delta \neq \mathbf{r}_d} \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\Delta,\sigma}(t) T^\dagger(\Delta - \mathbf{R} - \mathbf{r}) c_{\mathbf{R},\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle . \end{aligned} \quad (5.38)$$

Introducing Fourier transforms of the operators and inserting eigenstates of the auxiliary models leads to

$$\begin{aligned} &\sum_{\mathbf{r}_d} \langle \Psi_{\text{gs}} | c_{\mathbf{r}_d+\mathbf{r},\sigma}(t) c_{\mathbf{r}_d,\sigma}^\dagger | \Psi_{\text{gs}} \rangle \\ &= \sum_n F_n^*(\mathbf{r}) \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{r}_d,\sigma}(t) | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | c_{\mathbf{r}_d,\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{k}, n} F'_n(\mathbf{k}) F_n^*(\mathbf{r}) \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k},\sigma}(t) | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | c_{\mathbf{r}_d,\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{k}, n} F'_n(\mathbf{k})^* F_n^*(\mathbf{r}) \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{r}_d,\sigma}(t) | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k},\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{k}_1, \mathbf{k}_2} F'_n(\mathbf{k}_1) F'_n(\mathbf{k}_2)^* F_n^*(\mathbf{r}) \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k}_1,\sigma}(t) | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\mathbf{k}_2,\sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle , \end{aligned} \quad (5.39)$$

The net difference from the diagonal Greens function is the presence of the overlap factor $F_n^*(\mathbf{r})$.

4.5.3 Equal-Time Ground State Correlators

Apart from dynamical correlations such as Greens functions, we would also like to use our approach to calculate static correlation functions in terms of auxiliary model quantities. We first consider a

real-space correlation that tracks the propagation of a general excitation described by the operator $\mathcal{O}(\mathbf{r})$ over a distance Δ :

$$C_{\mathcal{O}}(\mathbf{r}) = \langle \Psi_{\text{gs}} | \mathcal{O}(\mathbf{r}_d + \mathbf{r}) \mathcal{O}^\dagger(\mathbf{r}_d) | \Psi_{\text{gs}} \rangle . \quad (5.40)$$

We first consider a local correlation: $\mathbf{r} = 0$. To obtain a tractable expression for this, we note that the quantity being computed here is almost identical to the one computed in eq. 5.28, the only difference being the time-evolution of the operator. For the local correlation, we therefore assume the result of eq. 5.31, dropping the time evolution because the correlators in this section are calculated at equal time. We also transform from k -space back to real-space because there are no time-evolution operators which need extended states in order to be resolved. Putting all this together, we get

$$\begin{aligned} C_{\mathcal{O}}^{\text{loc}} &= \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{O}(\mathbf{r}_d) \mathcal{O}^\dagger(\mathbf{r}_d) | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{r} \neq 0, n} F_n(\mathbf{r}) \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathcal{O}(\mathbf{r}+\mathbf{r}_d)} | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{O}^\dagger(\mathbf{r}_d) | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{r} \neq 0, n} F_n(\mathbf{r}) \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{O}(\mathbf{r} + \mathbf{r}_d) | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\mathcal{O}(\mathbf{r}_d)}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{r}_1 \neq 0, \mathbf{r}_2 \neq 0, n} F_n(\mathbf{r}) \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathcal{O}(\mathbf{r}_1+\mathbf{r}_d)} | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\mathcal{O}(\mathbf{r}_2+\mathbf{r}_d)}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle , \end{aligned} \quad (5.41)$$

where the transition operator $\mathcal{T}_{\mathcal{O}(\mathbf{r}+\mathbf{r}_d)}$ represents impurity-bath correlation:

$$\mathcal{T}_{\mathcal{O}(\mathbf{r}+\mathbf{r}_d), \sigma} = [c_{\mathbf{r}_d \sigma}^\dagger + S_{\mathbf{r}_d}^\sigma + C_{\mathbf{r}_d}^+] \mathcal{O}(\mathbf{r} + \mathbf{r}_d) , \quad (5.42)$$

We next consider non-local correlations over a distance Δ . Adapting eq. 5.39 for the non-local real-space Greens function in the same manner, we get

$$\begin{aligned} C_{\mathcal{O}}(\Delta) &= \sum_n F_n^*(\Delta) \langle \psi_{\text{gs}}(\mathbf{r}_d) | c_{\mathbf{r}_d, \sigma}(t) | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | c_{\mathbf{r}_d, \sigma}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{r} \neq 0, n} F_n(\mathbf{r} + \Delta) \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathcal{O}(\mathbf{r}+\Delta+\mathbf{r}_d)} | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{O}^\dagger(\mathbf{r}_d) | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{r} \neq 0, n} F_n(\mathbf{r} + \Delta) \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{O}(\mathbf{r} + \Delta + \mathbf{r}_d) | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\mathcal{O}(\mathbf{r}_d)}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle \\ &+ \sum_{\mathbf{r}_1 \neq 0, \mathbf{r}_2 \neq 0, n} F_n(\mathbf{r} + \Delta) \langle \psi_{\text{gs}}(\mathbf{r}_d) | \mathcal{T}_{\mathcal{O}(\mathbf{r}_1+\Delta+\mathbf{r}_d)} | \psi_n(\mathbf{r}_d) \rangle \langle \psi_n(\mathbf{r}_d) | \mathcal{T}_{\mathcal{O}(\mathbf{r}_2+\mathbf{r}_d)}^\dagger | \psi_{\text{gs}}(\mathbf{r}_d) \rangle , \end{aligned} \quad (5.43)$$

4.6 Entanglement Measures

We will now describe the prescription of calculating entanglement measures of the lattice model from within our auxiliary model treatment. In this section, we are interested mainly in two such measures, the entanglement entropy and the mutual information. Given a pure state $|\Psi\rangle$ describing

the complete system, the entanglement entropy $S_{\text{EE}}(\nu)$ of a subsystem ν quantifies the entanglement of ν with the rest of the subsystem, and is defined as

$$S_{\text{EE}}(\nu) = -\text{Tr} [\rho(\nu) \log \rho(\nu)], \quad \rho(\nu) = \text{Tr}_\nu [|\Psi\rangle \langle \Psi|] \quad (6.1)$$

where $\text{Tr}[\cdot]$ is the trace operator, and $\rho(\nu)$ is the reduced density matrix (RDM) for the subsystem ν obtained by taking the partial trace Tr_ν (over the states of ν) of the full density matrix $\rho = |\Psi\rangle \langle \Psi|$. If the subsystem ν describes local regions in real space (or states in k -space), we might be interested in the entanglement between two such subsystems ν_1 and ν_2 . The correct measure to quantify such entanglement is the mutual information:

$$I_2(\nu_1, \nu_2) = S_{\text{EE}}(\nu_1) + S_{\text{EE}}(\nu_2) - S_{\text{EE}}(\nu_1 \cup \nu_2), \quad (6.2)$$

where $\nu_1 \cup \nu_2$ is a larger subsystem formed by combining ν_1 and ν_2 .

We start by recalling the decomposition of entanglement entropy in terms of multipartite information as laid out in Subsection 2.8.2. Given a partitioning of the system into mutually exclusive subsystems $\{\nu_i\}$, the average entanglement entropy $S_{\text{EE}}^{(1)}$ over all subsystems can be expressed as

$$S_{\text{EE}}^{(1)} = \frac{1}{N} \sum_{n=2}^N (-1)^n \binom{N}{n} \bar{I}_n, \quad (6.3)$$

$$\bar{I}_n = \frac{(N-n)!n!}{N!} \sum_{S_n \in C_n(S_N)} I_n(S_n),$$

where $\bar{I}_n$ is the average multipartite information I_n over all distinct combinations of n subsystems.

We first take a look at real-space entanglement. Real-space entanglement measures can be used to probe delocalisation-localisation transitions. The simplest such measure is the entanglement of a lattice site in real-space. Since the local entanglement entropy will be uniform at each lattice site for a system with translation invariance, it suffices to calculate the real-space averaged entanglement entropy $S_{\text{EE}}^{(1)}$. The corresponding multipartite measures $\bar{I}_n$ involve collections of lattice sites. We write such a measure as

$$\bar{I}_n = \frac{(N-n)!n!}{N!} \frac{1}{N} \sum_{I(\mathbf{r}_d)} \sum_{B^{n-1}(\mathbf{r}_d)} I_n(I(\mathbf{r}_d); B^{n-1}(\mathbf{r}_d)), \quad (6.4)$$

where $I(\mathbf{r}_d)$ sums over all lattice sites in the form of an impurity site and $B^{n-1}(\mathbf{r}_d)$ considers all possible subsets of size $n-1$ of the remaining bath sites. Together, this choice corresponds to an auxiliary model with the impurity placed at $\mathbf{r}_d$. The additional factor of $1/N$ takes care of the overcounting arising from each choice appearing in N auxiliary models. We can therefore write the multipartite information as an average over auxiliary models, each auxiliary model contributing an average I_n over its bath sites:

$$\bar{I}_n = \frac{n}{N^2} \sum_{\mathbf{r}_d} I_n^{(\text{tot})}(\mathbf{r}_d). \quad (6.5)$$

The quantity $I_n^{(\text{tot})}(\mathbf{r}_d)$ is defined as $I_n^{(\text{tot})}(\mathbf{r}_d) = \frac{(N-n)!(n-1)!}{(N-1)!} \sum_{B^{n-1}(\mathbf{r}_d)} I_n(I(\mathbf{r}_d); B^{n-1}(\mathbf{r}_d))$. By translation invariance, such a measure will be uniform across auxiliary models and it suffices to

calculate the measure for a single auxiliary model located at a reference point $\mathbf{r}_d$; the whole expression therefore simplifies to

$$S_{\text{EE}}^{(1)} = \frac{1}{N} \sum_{n=2}^N (-1)^n \binom{N-1}{n-1} I_n^{(\text{tot})}(\mathbf{r}_d) = \frac{1}{N} \sum_{n=2}^N (-1)^n \sum_{B^{n-1}(\mathbf{r}_d)} I_n(I(\mathbf{r}_d); B^{n-1}(\mathbf{r}_d)) . \quad (6.6)$$

For example, truncating the summation to $n = 3$, we have the explicit expression

$$S_{\text{EE}}^{(1)} = \frac{N-1}{N} I_2^{(\text{tot})}(\mathbf{r}_d) - \frac{(N-1)(N-2)}{2N} I_3^{(\text{tot})}(\mathbf{r}_d) \quad (6.7)$$

Chapter 5

Mott Criticality as the Confinement Transition of of a Pseudogap-Mott Metal: Insights on Mott Transition and Pseudogap in Two Dimensions

5.1 Introduction

The Landau quasiparticle paradigm [Landau(1957)] underpins our understanding of metallic behaviour, where the screening of long-ranged repulsive interactions between delocalised electrons leads to well-defined excitations that carry the same quantum numbers as free electrons. This remarkable phenomenon - known as adiabatic continuity - is captured by excitations described by poles of the single-particle Greens function of the interacting system. But what defines metallicity when quasiparticles are no longer well-defined [Phillips et al.(2013)Phillips, Langley, and Hutasoit]? Can a surface of zeros of the Greens function [Dzyaloshinskii(2003)], rather than poles forming a Fermi surface, define an entirely new kind of metal? Could such an exotic state serve as the precursor to Mott localisation driven purely by repulsion, and without invoking magnetic order [Mott(1949)]? These questions necessitate a radical departure in understanding conduction in correlated electron systems beyond the quasiparticle paradigm [Phillips et al.(2022)Phillips, Hussey, and Abbamonte].

The anomalous pseudogap (PG) phase, observed in the cuprates, exhibits nodal-antinodal dichotomy with spectral weight concentrated on Fermi arcs around the nodal regions [Loeser et al.(1996)Loeser, Shen, Dessau, Marshall, Park, Fournier, and Kapitulnik, Norman et al.(1998)Norman, Ding, Randeria, Campuzano, Yokoya, Takeuchi, Takahashi, Mochiku, Kadowaki, Guptasarma et al., Hashimoto et al.(2014)Hashimoto, Vishik, He, Devereaux, and Shen]. In the PG phase of such Mott systems [Taillefer(2010), Fradkin et al.(2015)Fradkin, Kivelson, and Tranquada, Sato et al.(2017)Sato, Kasahara, Murayama, Kasahara, Moon, Nishizaki, Loew, Porras, Keimer, Shibauchi et al., Doiron-Leyraud et al.(2017)Doiron-Leyraud, Cyr-Choinière, Badoux, Ataei, Collignon, Gourgout, Dufour-

Beauséjour, Tafti, Laliberté, Boulanger et al., Mukhopadhyay et al.(2019)Mukhopadhyay, Sharma, Kim, Edkins, Hamidian, Eisaki, ichi Uchida, Kim, Lawler, Mackenzie et al.,Gupta et al.(2021)Gupta, McMahon, Sutarto, Shi, Gong, Wei, Shen, He, Ma, Dragomir et al.,Horio et al.(2025)Horio, Sakai, Suzuki, Nonaka, Hashimoto, Lu, Shen, Ohgi, Konno, Adachi et al.,Kyung et al.(2006)Kyung, Kancharla, Sénéchal, Tremblay, Civelli, and Kotliar, Macridin et al.(2006)Macridin, Jarrell, Maier, Kent, and D’Azevedo, Sakai et al.(2009b)Sakai, Motome, and Imada, Werner et al.(2009)Werner, Gull, Parcollet, and Millis, Sakai et al.(2010)Sakai, Motome, and Imada, Lin et al.(2010)Lin, Gull, and Millis, Gull et al.(2010)Gull, Ferrero, Parcollet, Georges, and Millis, Gull and Millis(2012), Mirzaei et al.(2013)Mirzaei, Stricker, Hancock, Berthod, Georges, van Heumen, Chan, Zhao, Li, Greven et al.,Gull et al.(2013)Gull, Parcollet, and Millis,Mukherjee and Lal(2020d),Hille et al.(2020b)Hille, Rohe, Honerkamp, and Andergassen, Jiang et al.(2022)Jiang, Liu, Klein, Wang, Sun, Chubukov, and Meng, Scheurer et al.(2018)Scheurer, Chatterjee, Wu, Ferrero, Georges, and Sachdev, Krien et al.(2022)Krien, Worm, Chalupa-Gantner, Toschi, and Held, Kitatani et al.(2023)Kitatani, Si, Worm, Tomczak, Arita, and Held, Sorella(2023), Schäfer et al.(2021)Schäfer, Wentzell, Šimkovic, He, Hille, Klett, Eckhardt, Arzhang, Harkov, Le Régent et al., White and Scalapino(1998), Ido et al.(2018)Ido, Ohgoe, and Imada,Proust and Taillefer(2019),Ponsioen et al.(2019)Ponsioen, Chung, and Corboz, Xu et al.(2022)Xu, Shi, Vitali, Qin, and Zhang], the single-particle Green’s function develops surfaces of zeros, also known as Luttinger surfaces [Dave et al.(2013)Dave, Phillips, and Kane,Dzyaloshinskii(2003)], that signals a fundamental obstruction to quasiparticle formation. As interaction strength increases, these zero surfaces proliferate, fragmenting the Fermi surface into nodal arcs [Sakai et al.(2009b)Sakai, Motome, and Imada, Dave et al.(2013)Dave, Phillips, and Kane], violating the Luttinger sum rule by reconfiguring Fermi volume [Seki and Yunoki(2017)], and breaking an emergent $\mathbb{Z}_2$ symmetry associated with the conservation of spin currents in Fermi liquids [Anderson and Haldane(2001), Huang et al.(2022)Huang, Nave, and Phillips]. Unlike conventional instabilities driven by poles, this phenomenon marks a topological breakdown of Fermi liquid theory [Coleman et al.(2001b)Coleman, Pépin, Si, and Ramazashvili,Phillips et al.(2013)Phillips, Langley, and Hutasoit]. We show that this breakdown arises from a strongly entangled, scale-invariant state of quantum matter, one that serves as the parent metal to the Mott insulator in a half-filled lattice model of strongly correlated electrons. This offers a concrete, controlled framework to understand both the non-quasiparticle character of the pseudogap phase and its continuous evolution into a repulsion-driven insulator, in full alignment with Mott’s original vision [Mott(1949)].

Our approach involves developing a renormalization group (RG)-based impurity-to-lattice framework that reconstructs the low-energy physics of a strongly correlated system from the fixed-point structure of a quantum impurity model (Fig. 5.1(a)). Starting from a dynamical Anderson impurity embedded in a correlated bath, we solve the system using a unitary RG (URG) approach [Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)]. We then construct a translation-invariant lattice Hamiltonian through a many-body tiling procedure that preserves non-local correlations and inherits momentum-resolved self-energies and renormalized couplings from the impurity solution. The resulting model is an *extended*-Hubbard Hamiltonian on a two-dimensional square lattice at half-filling, with both on-site repulsion and nearest-neighbour spin exchange interactions. These correlations, derived directly from the impurity dynamics, enable tracking the evolution from the FL to the Mott insulator (MI) via an intermediate PG phase.




Figure 5.1: Schematic representation of (a) theoretical framework adopted by us (see main text for explanation of symbols), (b) phases obtained for the extended Hubbard model in two dimensions at $T = 0$, and passage between them.

A central insight of our work is that the PG phase hosts a NFL state governed by Kondo frustration and doublon–holon deconfinement [Keimer et al.(2015)Keimer, Kivelson, Norman, Uchida, and Zaanen] (Fig. 5.1(b)). A correlation scale (W) in the conduction bath suppresses Kondo screening (with coupling J) beyond a critical ratio of W/J , triggering the emergence of Luttinger surfaces in the lattice model. This breakdown of quasiparticles begins at the antinodes and progressively engulfs the nodal regions, driving a continuous transformation into the MI. The resulting NFL exhibits a pseudogapped density of states that vanishes as ω^2 as $\omega \rightarrow 0$, multipartite long-range entanglement, and a universal scattering rate of the form $\sim (a + b\omega^2)^{-1}$ that grows as $\omega \rightarrow 0$. At the critical endpoint of this phase, the URG flow identifies a singular NFL described by the Hatsugai–Kohmoto (HK) model [Hatsugai and Kohmoto(1992), Baskaran(1991), Huang et al.(2022)Huang, Nave, and Phillips], with fully deconfined holon–doublon excitations.

The emergence of Luttinger surfaces alters the anomaly structure associated with the generalized symmetry of the FS [Altshuler et al.(1998)Altshuler, Chubukov, Dashevskii, Finkel’stein, and Morr, Heath and Bedell(2020a), Nave et al.(2025)Nave, Zhao, and Phillips]. The corresponding topological charge is defined via the Luttinger–Ward functional. This charge is modified by Green’s function zeros, signalling a shift in the underlying anomaly. Within our framework, this reorganization grants topological protection to the RG flows terminating in the PG regime, ensuring the stability of its NFL excitations. These are adiabatically connected to the Hatsugai–Kohmoto model [Hatsugai and Kohmoto(1992), Baskaran(1991)], reinforcing the interpretation of the PG as a distinct quantum phase. We thus identify the NFL state as a new gapless, long-range entangled

phase of correlated matter - the *Mott metal* - whose deconfined holon-doublon excitations are eventually confined across the Mott transition [Mott(1949)].

While we demonstrate our approach for an extended Hubbard model, the underlying framework is broadly applicable to correlated quantum systems. The impurity-plus-tiling construction accommodates multi-orbital degrees of freedom, frustrated geometries, and nontrivial band topology. By deriving lattice Hamiltonians directly from renormalized local dynamics - without relying on mean-field approximations - this method preserves nonlocal correlations and enhances momentum-space resolution. Coupled with the unitary renormalisation group method as the impurity solver, this approach provides access to one-particle as well as two-particle correlations, entanglement measures and several dynamical quantities, making the method very versatile. It thus provides a versatile and controlled route for engineering strongly correlated models, particularly in regimes where PG formation, NFL scaling, and Mott physics intersect.

We now describe the layout of the work. In section 5.2, we formulate the lattice-embedded auxiliary quantum impurity model and sketch the unitary renormalisation group (URG) method employed to analyse the impurity model. In Chapter 4, we have already formulated the tiling procedure by which to map the impurity Hamiltonian, eigenstates and observables into a lattice model of strongly correlated electrons via the usage of many-body translation operators. Section 5.5 displays the destruction of the Fermi liquid from the breakdown of Kondo screening, leading to a pseudogapped non-Fermi liquid phase of matter. The novel properties of this phase are rigorously established in section 5.6. In section 5.7, we demonstrate that the continuous transition between the pseudogap and Mott insulating phases is described by the exactly solvable Hatsugai-Kohmoto model. We demonstrate in section 5.10 the universal organising principle that is responsible for the emergence of the pseudogap non-Fermi liquid phase. This also unveils the pseudogap as the strongly coupled long-range entangled parent metal of the Mott insulator. We conclude in section 7.10. Finally, we present appendices containing the derivation of the URG analysis of the impurity model, and the exactly solvable model at the critical point.

5.2 Auxiliary Model and URG analysis

To capture the interplay between Kondo physics and Mott localization, our approach relies on a two-dimensional auxiliary model consisting of an Anderson impurity embedded in a minimally-correlated bath, which we solve the system using a unitary RG (URG) approach [Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)]. The impurity model and URG analysis are laid out in this section. In order to make contact with a lattice model, we construct a translation-invariant Hamiltonian from the impurity model through a many-body tiling procedure: this preserves non-local correlations, as well as allows for momentum-resolved self-energies and renormalized couplings generated from the impurity solution. For the impurity model chosen in this work, tiling results in an *extended*-Hubbard Hamiltonian on a two-dimensional square lattice at particle-hole symmetry (half-filling). The interaction terms within the lattice model are a result of analogous correlations present in the auxiliary model. This is laid out next in Section ??.

5.2.1 Lattice-embedded Impurity Model

The auxiliary model consists of a correlated impurity site (with a double occupancy cost U) embedded within a correlated conduction bath defined on a square lattice (see Fig. 5.1(a)). The impurity site hybridises with the conduction bath sites adjacent to it through one-particle hybridisation V as well as a spin-exchange interaction J arising from fluctuations in local spin densities of the electrons. The conduction bath has minimal correlations in the form of a local correlation term W on the sites that are adjacent to the impurity.

The Hamiltonian of the auxiliary model studied here is

$$\mathcal{H}_{\text{aux}} = H_{\text{imp}} + H_{\text{coup}} + H_{\text{cbath}}, \quad (2.1)$$

where the particle-hole symmetric impurity term is

$$H_{\text{imp}} = -\frac{U}{2} (\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2, \quad (2.2)$$

describing on-site Coulomb interactions at the impurity. Here, $c_{d\sigma}^\dagger$ creates an electron with spin σ at the impurity site, and $\hat{n}_{d\sigma} = c_{d\sigma}^\dagger c_{d\sigma}$ denotes the corresponding number operator.

The impurity couples to four nearest-neighbour bath sites $c_{Z\sigma}$ through both hybridization and Kondo exchange:

$$H_{\text{coup}} = \frac{J}{2} \sum_{\sigma_1, \sigma_2, Z} \mathbf{S}_d \cdot c_{Z\sigma_1}^\dagger \boldsymbol{\tau}_{\sigma_1 \sigma_2} c_{Z\sigma_2} - V \sum_{\sigma, Z} \left(c_{Z\sigma}^\dagger c_{d\sigma} + \text{h.c.} \right), \quad (2.3)$$

where $\boldsymbol{\tau}$ are the Pauli matrices, and Z indexes the four neighbouring bath sites. The half-filled bath includes nearest neighbour hopping and on-site correlations on the site neighbouring the impurity:

$$H_{\text{cbath}} = \sum_{\mathbf{k}} \epsilon_{\mathbf{k}} c_{\mathbf{k},\sigma}^\dagger c_{\mathbf{k},\sigma} - \frac{W}{2} \sum_Z (\hat{n}_{Z\uparrow} - \hat{n}_{Z\downarrow})^2, \quad (2.4)$$

where $\epsilon_{\mathbf{k}} = -2t(\cos k_x + \cos k_y)$, and W parametrizes local repulsion in the bath that frustrates Kondo screening and drives the system towards local-moment formation. A crucial feature of this construction is that the Kondo coupling acquires a momentum structure upon Fourier transforming:

$$J_{\mathbf{k},\mathbf{k}'} = \frac{J}{2} [\cos(k_x - k'_x) + \cos(k_y - k'_y)], \quad (2.5)$$

which respects the C_4 symmetry of the square lattice.

5.2.2 The unitary renormalisation group method

In order to obtain the various low-energy phases of our impurity model, we perform a scaling analysis of the associated Hamiltonian using the recently developed unitary renormalisation group (URG) method [19,39,40]. The method has been applied successfully on a wide variety of problems of correlated fermions [Pal et al.(2019)Pal, Mukherjee, and Lal, 2, Mukherjee and Lal(2020c), Patra and

Lal(2021), Mukherjee et al.(2022)Mukherjee, Mukherjee, Vidhyadhiraja, Taraphder, and Lal, Patra et al.(2023b)Patra, Mukherjee, and Lal]. The method proceeds by resolving quantum fluctuations in high-energy degrees of freedom, leading to a low-energy Hamiltonian with renormalised couplings and new emergent degrees of freedom. Typically, for a system with Fermi energy ϵ_F and bandwidth E_N , the sequence of isoenergetic shells $\{E_{(j)}\}$, $E_j \in [E_0, E_N]$ define the states whose quantum fluctuations we sequentially resolve. The momentum states lying on shells E_N that are far away from the Fermi surface comprise the UV states, while those on shells near the Fermi surface comprise the IR states.

As a result of the URG transformations, the Hamiltonian $H_{(j)}$ at a given RG step j involves scattering processes between the k -states that have energies lower than $D_{(j+1)}$. The unitary transformation $U_{(j)}$ is then defined so as to remove the number fluctuations of the currently most energetic set of states $D_{(j)}$ [Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)]:

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger, \text{ such that } [H_{(j-1)}, \hat{n}_j] = 0. \quad (2.6)$$

The eigenvalue of $\hat{n}_j$ has, thus, been rendered an integral of motion (IOM) under the RG transformation.

The unitary transformations can be expressed in terms of a generator $\eta_{(j)}$ that has fermionic algebra [Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)]:

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 + \eta_{(j)} - \eta_{(j)}^\dagger \right), \quad \{ \eta_{(j)}, \eta_{(j)}^\dagger \} = 1, \quad (2.7)$$

where $\{\cdot\}$ is the anticommutator. The unitary operator $U_{(j)}$ that appears in Eq. (2.7) can be cast into the well-known general form $U = e^{\mathcal{S}}$, $\mathcal{S} = \frac{\pi}{4} \left(\eta_{(j)}^\dagger - \eta_{(j)} \right)$ that a unitary operator can take, defined by an anti-Hermitian operator $\mathcal{S}$. The generator $\eta_{(j)}$ is given by the expression [Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)]

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - \text{Tr}(H_{(j)} \hat{n}_j)} c_j^\dagger \text{Tr}(H_{(j)} c_j). \quad (2.8)$$

The operators $\eta_{(j)}, \eta_{(j)}^\dagger$ behave as the many-particle analogues of the single-particle field operators $c_j, c_j^\dagger$ - they change the occupation number of the single-particle Fock space $|n_j\rangle$. The important operator $\hat{\omega}_{(j)}$ originates from the quantum fluctuations that exist in the problem because of the non-commutation of the kinetic energy terms and the interaction terms in the Hamiltonian:

$$\hat{\omega}_{(j)} = H_{(j-1)} - H_{(j)}^i. \quad (2.9)$$

$H_{(j)}^i$ is the part of $H_{(j)}$ that commutes with $\hat{n}_j$ but does *not* commute with at least one $\hat{n}_l$ for $l < j$. The RG flow continues up to energy D^* , where a fixed point is reached from the vanishing of the RG function. Detailed comparisons of the URG with other methods (e.g., the functional RG, spectrum bifurcation RG etc.) can be found in Refs. [Mukherjee and Lal(2020c), Mukherjee and Lal(2020a)], and implementation on the present problem in Appendix A.

5.3 Derivation of unitary RG equations for the lattice-embedded impurity model

5.3.1 RG scheme

At any given step j of the RG procedure, we decouple the states $\{\mathbf{q}\}$ on the isoenergetic surface of energy ε_j . The diagonal Hamiltonian H_D for this step consists of all terms that do not change the occupancy of the states $\{\mathbf{q}\}$:

$$H_D^{(j)} = \varepsilon_j \sum_{q,\sigma} \tau_{q,\sigma} + \frac{1}{2} \sum_{\mathbf{q}} J_{\mathbf{q},\mathbf{q}} S_d^z (\hat{n}_{\mathbf{q},\uparrow} - \hat{n}_{\mathbf{q},\downarrow}) - \frac{1}{2} \sum_{\mathbf{q}} W_{\mathbf{q}} (\hat{n}_{\mathbf{q},\uparrow} - \hat{n}_{\mathbf{q},\downarrow})^2, \quad (3.1)$$

where $\tau = \hat{n} - 1/2$ and $W_{\mathbf{q}}$ is a shorthand for $W_{\mathbf{q},\mathbf{q},\mathbf{q}}$. The three terms, respectively, are the kinetic energy of the momentum states on the isoenergetic shell that we are decoupling, the spin-correlation energy between the impurity spin and the spins formed by these momentum states and, finally, the local correlation energy associated with these states arising from the W term. The off-diagonal part of the Hamiltonian on the other hand leads to scattering in the states $\{\mathbf{q}\}$. We now list these terms, classified by the coupling they originate from.

Arising from the Kondo spin-exchange term

$$\begin{aligned} T_{KZ1}^\dagger + T_{KZ1} &= \frac{1}{2} \sum_{\mathbf{k},\mathbf{q},\sigma} \sigma J_{\mathbf{k},\mathbf{q}} S_d^z [c_{\mathbf{q}\sigma}^\dagger c_{\mathbf{k},\sigma} + \text{h.c.}], \\ T_{KZ2}^\dagger + T_{KZ2} &= \frac{1}{2} \sum_{\mathbf{q},\sigma} \sigma J_{\mathbf{q},\mathbf{q}} S_d^z [c_{\mathbf{q}\sigma}^\dagger c_{\mathbf{q},\sigma} + \text{h.c.}], \\ T_{KT1}^\dagger + T_{KT1} &= \frac{1}{2} \sum_{\mathbf{k},\mathbf{q}} J_{\mathbf{k},\mathbf{q}} \left[S_d^+ \left(c_{\mathbf{q}\downarrow}^\dagger c_{\mathbf{k}\uparrow} + c_{\mathbf{k}\downarrow}^\dagger c_{\mathbf{q}\uparrow} \right) + \text{h.c.} \right], \\ T_{KT2}^\dagger + T_{KT2} &= \frac{1}{2} \sum_{\mathbf{q}} J_{\mathbf{q},\mathbf{q}} \left[S_d^+ \left(c_{\mathbf{q}\downarrow}^\dagger c_{\mathbf{q}\uparrow} + c_{\mathbf{q}\downarrow}^\dagger c_{\mathbf{q}\uparrow} \right) + \text{h.c.} \right], \end{aligned} \quad (3.2)$$

Arising from spin-preserving scattering within conduction bath

$$\begin{aligned}
 T_{P1}^\dagger + T_{P1} &= - \sum_{\mathbf{q} \in \varepsilon_j} \sum_{\mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4 < \varepsilon_j} \sum_{\sigma} \left[W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \sigma}^\dagger c_{\mathbf{k}_4, \sigma} + \text{h.c.} \right] \\
 T_{P2}^\dagger + T_{P3} &= - \sum_{\mathbf{q} \in \varepsilon_j} \sum_{\mathbf{k}_2 < \varepsilon_j} \sum_{\sigma} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{q}, \mathbf{q}} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} n_{\mathbf{q}, \sigma} - \sum_{\mathbf{q} \in \varepsilon_j} \sum_{\mathbf{k}_1 < \varepsilon_j} \sum_{\sigma} W_{\mathbf{k}_1, \mathbf{q}, \mathbf{q}, \mathbf{q}} c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{q}, \sigma} n_{\mathbf{q}, \sigma} \\
 T_{P4} &= - \sum_{\mathbf{q} \in \varepsilon_j} \sum_{\mathbf{k}_2, \mathbf{k}_3 < \varepsilon_j} \sum_{\sigma} W_{\mathbf{q}, \mathbf{q}, \mathbf{k}_2, \mathbf{k}_3} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{q}, \sigma} c_{\mathbf{k}_2, \sigma}^\dagger c_{\mathbf{k}_3, \sigma} \\
 T_{P5} &= - \sum_{\mathbf{q} \in \varepsilon_j} \sum_{\mathbf{k}_2, \mathbf{k}_3 < \varepsilon_j} \sum_{\sigma} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{q}} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \sigma}^\dagger c_{\mathbf{q}, \sigma} \\
 &= + \sum_{\mathbf{q} \in \varepsilon_j} \sum_{\mathbf{k}_2, \mathbf{k}_3 < \varepsilon_j} \sum_{\sigma} W_{\mathbf{q}, \mathbf{k}_3, \mathbf{k}_2, \mathbf{q}} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{q}, \sigma} c_{\mathbf{k}_2, \sigma}^\dagger c_{\mathbf{k}_3, \sigma} \\
 &= -T_{P4}
 \end{aligned} \tag{3.3}$$

Arising from spin-flip scattering within conduction bath

$$\begin{aligned}
 T_{F1}^\dagger + T_{F1} &= \sum_{\mathbf{q} \in \varepsilon_j} \sum_{\mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4 < \varepsilon_j} \sum_{\sigma} \left[W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \bar{\sigma}}^\dagger c_{\mathbf{k}_4, \bar{\sigma}} + \text{h.c.} \right] \\
 T_{F2} &= \sum_{\mathbf{q}, \mathbf{q}' \in \varepsilon_j} \sum_{\mathbf{k}_2, \mathbf{k}_3 < \varepsilon_j} \sum_{\sigma} W_{\mathbf{q}, \mathbf{q}', \mathbf{k}_2, \mathbf{k}_3} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{q}', \sigma} c_{\mathbf{k}_2, \bar{\sigma}}^\dagger c_{\mathbf{k}_3, \bar{\sigma}} \\
 T_{F3} &= \sum_{\mathbf{q}, \mathbf{q}' \in \varepsilon_j} \sum_{\mathbf{k}_2, \mathbf{k}_3 < \varepsilon_j} \sum_{\sigma} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{q}'} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \bar{\sigma}}^\dagger c_{\mathbf{q}', \bar{\sigma}} \\
 T_{F4}^\dagger + T_{F4} &= \sum_{\mathbf{q}, \mathbf{q}' \in \varepsilon_j} \sum_{\mathbf{k}_1 < \varepsilon_j} \sum_{\sigma} \left[W_{\mathbf{q}, \mathbf{q}, \mathbf{q}', \mathbf{k}_1} n_{\mathbf{q}, \sigma} c_{\mathbf{q}', \bar{\sigma}}^\dagger c_{\mathbf{k}_1, \bar{\sigma}} + \text{h.c.} \right]
 \end{aligned} \tag{3.4}$$

In all of the terms $T_{P[i]}$ and $T_{F[i]}$, the factor of $1/2$ in front has been cancelled out by a factor of 2 coming from the multiple possibilities of arranging the momentum labels. We will henceforth ignore T_{P4} and T_{P5} because they cancel each other out.

The renormalisation of the Hamiltonian is constructed from the general expression

$$\Delta H^{(j)} = H_X \frac{1}{\omega - H_D} H_X. \tag{3.5}$$

The states on the isoenergetic shell $\pm|\varepsilon_j|$ come in particle-hole pairs $(\mathbf{q}, \mathbf{q})$ with energies of opposite signs (relative to the Fermi energy). If $\mathbf{q}$ is defined as the hole state (unoccupied in the absence of quantum fluctuations), it will have positive energy, while the particle state $\mathbf{q}$ will be of negative energy and hence below the Fermi surface. To be more specific, given a state $\mathbf{q}$ with energy $\pm|\varepsilon_j|$, we define its particle-hole transformed counterpart as the state $\mathbf{q} = \boldsymbol{\pi} + \mathbf{q}$, having energy $\mp|\varepsilon_j|$ and residing in the opposite quadrant of the Brillouin zone. Given this definition, we have the important property that

$$\begin{aligned}
 J_{\mathbf{k}, \mathbf{q}} &= -J_{\mathbf{k}, \mathbf{q}}, \\
 W_{\{\mathbf{k}\}, \mathbf{q}} &= -W_{\{\mathbf{k}\}, \mathbf{q}}.
 \end{aligned} \tag{3.6}$$




Figure 5.2: Particle and hole states.

5.3.2 Renormalisation of the bath correlation term W

The bath correlation term W can undergo renormalisation only via scattering processes arising from itself. Irrespective of whether the state $\mathbf{q}$ being decoupled is in a particle or hole configuration in the initial many-body state, the propagator $G = 1/(\omega - H_D)$ of the intermediate excited state is uniform, and equal to

$$G_W = 1/(\omega - |\varepsilon_j|/2 + W_{\mathbf{q}}/2) \quad , \quad (3.7)$$

where $W_{\mathbf{q}}$ is the same whether $\mathbf{q}$ is above or below the Fermi surface. The $|\varepsilon_j|/2$ in H_D arises from the excited nature of the state after the initial scattering process.

Scattering arising purely from spin-preserving processes

In this subsection, we calculate the renormalisation to W arising from the terms T_{P1} , T_{P2} and T_{P3} . The first term is

$$\begin{aligned} T_{P1}^\dagger G_W T_{P3} &= \sum_{\sigma} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} \sum_{\mathbf{q}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \sigma}^\dagger c_{\mathbf{k}_4, \sigma} G_W W_{\mathbf{k}_1, \mathbf{q}, \mathbf{q}, \mathbf{q}} c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{q}, \sigma} n_{\mathbf{q}, \sigma} \\ &= - \sum_{\sigma} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \sigma}^\dagger c_{\mathbf{k}_4, \sigma} \sum_{\mathbf{q} \in \text{PS}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} G_W W_{\mathbf{k}_1, \mathbf{q}, \mathbf{q}, \mathbf{q}} . \end{aligned} \quad (3.8)$$

The operators acting on the states being decoupled contract to form a number operator $n_{\mathbf{q}, \sigma}$ which projects the sum over $\mathbf{q}$ into the states that are initially occupied (particle sector, PS).

The second such contribution is obtained by flipping the sequence of scattering processes:

$$\begin{aligned} T_{P3} G_W T_{P1}^\dagger &= \sum_{\sigma} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} \sum_{\mathbf{q}} W_{\mathbf{k}_1, \mathbf{q}, \mathbf{q}, \mathbf{q}} c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{q}, \sigma} n_{\mathbf{q}, \sigma} G_W W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \sigma}^\dagger c_{\mathbf{k}_4, \sigma} \\ &= \sum_{\sigma} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \sigma}^\dagger c_{\mathbf{k}_4, \sigma} \sum_{\mathbf{q} \in \text{HS}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} G_W W_{\mathbf{k}_1, \mathbf{q}, \mathbf{q}, \mathbf{q}} . \end{aligned} \quad (3.9)$$

By virtue of eq. 3.6, the product of couplings $W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} G_W W_{\mathbf{k}_1, \mathbf{q}, \mathbf{q}, \mathbf{q}}$ is the same irrespective of whether $\mathbf{q}$ belongs to the particle or hole sector. The two contributions therefore cancel each other.

Moreover, the remaining contributions $T_{P3}^\dagger G_W T_{P1}$ and $T_{P1} G_W T_{P2}^\dagger$ are effectively hermitian conjugates of the two contributions considered above, and therefore also cancel each other.

Scattering arising from spin-flip processes

We now come to the processes that involve spin-flips. Considering T_{F1} and T_{F4} first, we get

$$\begin{aligned}
 T_{F1}^\dagger G_W T_{F4} &= \sum_{\sigma} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} \sum_{\mathbf{q}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \bar{\sigma}}^\dagger c_{\mathbf{k}_4, \bar{\sigma}} G_W W_{\mathbf{k}_1, \mathbf{q}, \mathbf{q}, \mathbf{q}} c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{q}, \sigma} n_{\mathbf{q}, \bar{\sigma}} \\
 &= - \sum_{1,2,3,4} \sum_{\sigma} c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \bar{\sigma}}^\dagger c_{\mathbf{k}_4, \bar{\sigma}} \sum_{\mathbf{q} \in \text{PS}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_4, \mathbf{k}_4} G_W W_{\mathbf{k}_1, \mathbf{q}, \mathbf{q}, \mathbf{q}} , \\
 T_{F4} G_W T_{F1}^\dagger &= \sum_{\sigma} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} \sum_{\mathbf{q}} W_{\mathbf{k}_1, \mathbf{q}, \mathbf{q}, \mathbf{q}} c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{q}, \sigma} n_{\mathbf{q}, \bar{\sigma}} G_W W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \bar{\sigma}}^\dagger c_{\mathbf{k}_4, \bar{\sigma}} \\
 &= \sum_{1,2,3,4} \sum_{\sigma} c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \bar{\sigma}}^\dagger c_{\mathbf{k}_4, \bar{\sigma}} \sum_{\mathbf{q} \in \text{HS}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_4, \mathbf{k}_4} G_W W_{\mathbf{k}_1, \mathbf{q}, \mathbf{q}, \mathbf{q}} .
 \end{aligned} \tag{3.10}$$

By the same arguments as in the previous subsection, these terms cancel each other out. Their hermitian conjugate contributions $T_{F1} G_W T_{F4}^\dagger$ and $T_{F4}^\dagger G_W T_{F1}$ also cancel out. The other two terms are T_{F2} and T_{F3} , and their contributions also cancel out for the same reason:

$$\begin{aligned}
 T_{F2} G_W T_{F2} &= \sum_{\sigma} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} \sum_{\mathbf{q}} W_{\mathbf{q}, \mathbf{q}, \mathbf{k}_3, \mathbf{k}_4} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{q}, \sigma} c_{\mathbf{k}_3, \bar{\sigma}}^\dagger c_{\mathbf{k}_4, \bar{\sigma}} G_W W_{\mathbf{q}, \mathbf{q}, \mathbf{k}_1, \mathbf{k}_2} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{q}, \sigma} c_{\mathbf{k}_1, \bar{\sigma}}^\dagger c_{\mathbf{k}_2, \bar{\sigma}} \\
 &= \sum_{1,2,3,4} \sum_{\sigma} c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \bar{\sigma}}^\dagger c_{\mathbf{k}_4, \bar{\sigma}} \sum_{\mathbf{q} \in \text{PS}} W_{\mathbf{q}, \mathbf{q}, \mathbf{k}_3, \mathbf{k}_4} G_W W_{\mathbf{q}, \mathbf{q}, \mathbf{k}_1, \mathbf{k}_2} , \\
 T_{F3} G_W T_{F3} &= \sum_{\sigma} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4} \sum_{\mathbf{q}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{q}} c_{\mathbf{q}, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \bar{\sigma}}^\dagger c_{\mathbf{q}, \bar{\sigma}} G_W W_{\mathbf{q}, \mathbf{k}_4, \mathbf{k}_1, \mathbf{q}} c_{\mathbf{q}, \bar{\sigma}}^\dagger c_{\mathbf{k}_4, \bar{\sigma}} c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{q}, \sigma} \\
 &= - \sum_{1,2,3,4} \sum_{\sigma} c_{\mathbf{k}_1, \sigma}^\dagger c_{\mathbf{k}_2, \sigma} c_{\mathbf{k}_3, \bar{\sigma}}^\dagger c_{\mathbf{k}_4, \bar{\sigma}} \sum_{\mathbf{q} \in \text{PS}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_3, \mathbf{q}} G_W W_{\mathbf{q}, \mathbf{k}_4, \mathbf{k}_1, \mathbf{q}} ,
 \end{aligned} \tag{3.11}$$

Scattering involving both spin-flip and spin-preserving processes

These processes involve the combination of terms like T_{P1} with T_{F4} , and T_{P2} with T_{F1} . These again cancel each other out for the same reasons as outline above.

Net renormalisation for the bath correlation term

Since all the contributions cancel out in pairs, the bath correlation term W is *marginal*.

5.3.3 Renormalisation of the Kondo scattering term J

We focus on the renormalisation of the spin-flip part of the Kondo interaction. For these processes, the intermediate many-body state always involves the impurity spin being anti-correlated with the conduction electron spin, such that the propagator for that state is $G_J = 1/(\omega - |\varepsilon_j|/2 + J_{\mathbf{q}}/4 + W_{\mathbf{q}}/2)$.

Impurity-mediated spin-flip scattering purely through Kondo-like processes

The following processes arising from the Kondo term renormalise the spin-flip interaction:

$$\begin{aligned}
 T_{KT1}^\dagger G_J (T_{KZ1} + T_{KZ1}^\dagger) &= \frac{1}{4} \sum_{\mathbf{k}_1, \mathbf{k}_1, \mathbf{q}} J_{\mathbf{q}, \mathbf{k}_2} S_d^+ \left[-c_{\mathbf{q}\downarrow}^\dagger c_{\mathbf{k}_2\uparrow} G_J c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{q}\downarrow} + c_{\mathbf{k}_2\downarrow}^\dagger c_{\mathbf{q}\uparrow} G_J c_{\mathbf{q}\uparrow}^\dagger c_{\mathbf{k}_1\uparrow} \right] J_{\mathbf{k}_1, \mathbf{q}} S_d^z \\
 &= -\frac{1}{8} \sum_{\mathbf{k}_1, \mathbf{k}_1, \mathbf{q}} J_{\mathbf{q}, \mathbf{k}_2} S_d^+ \left[c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{k}_2\uparrow} G_J n_{\mathbf{q}\downarrow} + c_{\mathbf{k}_2\downarrow}^\dagger c_{\mathbf{k}_1\uparrow} (1 - n_{\mathbf{q}\uparrow}) G_J \right] J_{\mathbf{k}_1, \mathbf{q}} \\
 &= -\frac{1}{8} \sum_{\mathbf{k}_1, \mathbf{k}_1} S_d^+ c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{k}_2\uparrow} \sum_{\mathbf{q} \in \text{PS}} [J_{\mathbf{q}, \mathbf{k}_2} J_{\mathbf{k}_1, \mathbf{q}} + J_{\mathbf{q}, \mathbf{k}_1} J_{\mathbf{k}_2, \mathbf{q}}] G_J .
 \end{aligned} \tag{3.12}$$

In getting the final expression, we used the sigma matrix relation $S_d^+ S_d^z = -\frac{1}{2} S_d^+$, and absorbed the projector $1 - n_{\mathbf{q}\uparrow}$ into the sum over the particle sector by replacing q with its particle-hole transformed counterpart $\bar{q}$. An identical contribution is obtained by switching the sequence of processes:

$$\begin{aligned}
 (T_{KZ1} + T_{KZ1}^\dagger) G_J T_{KT1}^\dagger &= \frac{1}{4} \sum_{\mathbf{k}_1, \mathbf{k}_1, \mathbf{q}} J_{\mathbf{k}_1, \mathbf{q}} S_d^z \left[-c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{q}\downarrow} G_J c_{\mathbf{q}\downarrow}^\dagger c_{\mathbf{k}_2\uparrow} + c_{\mathbf{q}\uparrow}^\dagger c_{\mathbf{k}_1\uparrow} G_J c_{\mathbf{k}_2\downarrow}^\dagger c_{\mathbf{q}\uparrow} \right] J_{\mathbf{q}, \mathbf{k}_2} S_d^+ \\
 &= -\frac{1}{8} \sum_{\mathbf{k}_1, \mathbf{k}_1} S_d^+ c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{k}_2\uparrow} \sum_{\mathbf{q} \in \text{PS}} [J_{\mathbf{q}, \mathbf{k}_2} J_{\mathbf{k}_1, \mathbf{q}} + J_{\mathbf{q}, \mathbf{k}_1} J_{\mathbf{k}_2, \mathbf{q}}] G_J .
 \end{aligned} \tag{3.13}$$

Scattering processes involving interplay between the Kondo interaction and conduction bath interaction

Looking at $T_{KT1}^\dagger$ first, we have

$$T_{KT1}^\dagger G_J (T_{F4} + T_{F4}^\dagger) = \frac{1}{2} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{q}} J_{\mathbf{k}_2, \mathbf{q}} S_d^+ \left(c_{\mathbf{q}\downarrow}^\dagger c_{\mathbf{k}_2\uparrow} G_J W_{\mathbf{q}, \mathbf{q}, \mathbf{k}_1, \mathbf{q}} n_{\mathbf{q}\uparrow} c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{q}\downarrow} + c_{\mathbf{k}_2\downarrow}^\dagger c_{\mathbf{q}\uparrow} G_J W_{\mathbf{q}, \mathbf{q}, \mathbf{q}, \mathbf{k}_1} n_{\bar{q}\downarrow} c_{\mathbf{q}\uparrow}^\dagger c_{\mathbf{k}_1\uparrow} \right) . \tag{3.14}$$

For either of the two choices of the functional form of W , it is easy to show that $W_{\mathbf{q}, \mathbf{q}, \mathbf{k}_1, \mathbf{q}} = W_{\mathbf{q}, \mathbf{q}, \mathbf{q}, \mathbf{k}_1}$.

$$T_{KT1}^\dagger G_J (T_{F4} + T_{F4}^\dagger) = \frac{1}{2} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{q}} J_{\mathbf{k}_2, \mathbf{q}} W_{\mathbf{q}, \mathbf{q}, \mathbf{k}_1, \mathbf{q}} G_J S_d^+ \left[-c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{k}_2\uparrow} n_{\mathbf{q}\downarrow} n_{\mathbf{q}\uparrow} + c_{\mathbf{k}_2\downarrow}^\dagger c_{\mathbf{k}_1\uparrow} (1 - n_{\mathbf{q}\uparrow}) n_{\bar{q}\downarrow} \right] . \tag{3.15}$$

Another contribution is obtained by switching the sequence of the scattering processes:

$$\begin{aligned}
 (T_{F4} + T_{F4}^\dagger) G_J T_{KT1}^\dagger &= \frac{1}{2} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{q}} \left(W_{\mathbf{q}, \mathbf{q}, \mathbf{k}_1, \mathbf{q}} n_{\bar{q}\uparrow} c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{q}\downarrow} G_J c_{\mathbf{q}\downarrow}^\dagger c_{\mathbf{k}_2\uparrow} + W_{\mathbf{q}, \mathbf{q}, \mathbf{q}, \mathbf{k}_1} n_{\mathbf{q}\downarrow} c_{\mathbf{q}\uparrow}^\dagger c_{\mathbf{k}_1\uparrow} G_J c_{\mathbf{k}_2\downarrow}^\dagger c_{\mathbf{q}\uparrow} \right) J_{\mathbf{k}_2, \mathbf{q}} S_d^+ \\
 &= \frac{1}{2} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{q}} \left(c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{k}_2\uparrow} n_{\bar{q}\uparrow} (1 - n_{\mathbf{q}\downarrow}) - c_{\mathbf{k}_2\downarrow}^\dagger c_{\mathbf{k}_1\uparrow} n_{\mathbf{q}\downarrow} n_{\mathbf{q}\uparrow} \right) W_{\mathbf{q}, \mathbf{q}, \mathbf{k}_1, \mathbf{q}} G_J J_{\mathbf{k}_2, \mathbf{q}} S_d^+
 \end{aligned} \tag{3.16}$$

The two contributions (eqs. 3.15 and 3.16) arising from T_{KT1} cancel each other.

We now consider the other spin-exchange process $T_{KT2}^\dagger$. One such contribution is

$$\begin{aligned}
 T_{KT2}^\dagger G_J T_{F3} &= \frac{1}{2} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{q}} J_{\mathbf{q}, \mathbf{q}} S_d^+ \left(c_{\mathbf{q}\downarrow}^\dagger c_{\mathbf{q}\uparrow} G_J c_{\mathbf{q}\uparrow}^\dagger c_{\mathbf{k}_2\uparrow} c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{q}\downarrow} + c_{\mathbf{q}\downarrow}^\dagger c_{\mathbf{q}\uparrow} G_J c_{\mathbf{q}\uparrow}^\dagger c_{\mathbf{k}_2\uparrow} c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{q}\downarrow} \right) W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_1, \mathbf{q}} \\
 &= -\frac{1}{2} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{q}} S_d^+ c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{k}_2\uparrow} [n_{\mathbf{q}\downarrow}(1 - n_{\mathbf{q}\uparrow}) + n_{\mathbf{q}\downarrow}(1 - n_{\mathbf{q}\uparrow})] J_{\mathbf{q}, \mathbf{q}} G_J W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_1, \mathbf{q}} \\
 &= -\frac{1}{2} \sum_{\mathbf{k}_1, \mathbf{k}_2} S_d^+ c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{k}_2\uparrow} \sum_{\mathbf{q} \in \text{PS}} (J_{\mathbf{q}, \mathbf{q}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_1, \mathbf{q}} + J_{\mathbf{q}, \mathbf{q}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_1, \mathbf{q}}) G_J.
 \end{aligned} \tag{3.17}$$

An identical contribution is obtained from the reversed term:

$$\begin{aligned}
 T_{F3} G_J T_{KT2}^\dagger &= \frac{1}{2} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{q}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_1, \mathbf{q}} \left(c_{\mathbf{q}\uparrow}^\dagger c_{\mathbf{k}_2\uparrow} c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{q}\downarrow} G_J c_{\mathbf{q}\downarrow}^\dagger c_{\mathbf{q}\uparrow} + c_{\mathbf{q}\uparrow}^\dagger c_{\mathbf{k}_2\uparrow} c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{q}\downarrow} G_J c_{\mathbf{q}\downarrow}^\dagger c_{\mathbf{q}\uparrow} \right) J_{\mathbf{q}, \mathbf{q}} S_d^+ \\
 &= -\frac{1}{2} \sum_{\mathbf{k}_1, \mathbf{k}_2} S_d^+ c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{k}_2\uparrow} \sum_{\mathbf{q} \in \text{PS}} (J_{\mathbf{q}, \mathbf{q}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_1, \mathbf{q}} + J_{\mathbf{q}, \mathbf{q}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_1, \mathbf{q}}) G_J.
 \end{aligned} \tag{3.18}$$

Net renormalisation to the Kondo interaction

Combining the results from eqs. 3.12, 3.13, 3.17 and 3.18, as well as using the properties $J_{\mathbf{q}, \mathbf{k}_1} J_{\mathbf{k}_2, \mathbf{q}} = J_{\mathbf{q}, \mathbf{k}_2} J_{\mathbf{k}_1, \mathbf{q}} = J_{\mathbf{k}_2, \mathbf{q}} J_{\mathbf{q}, \mathbf{k}_1}$ and $J_{\mathbf{q}, \mathbf{q}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_1, \mathbf{q}} = J_{\mathbf{q}, \mathbf{q}} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_1, \mathbf{q}}$, the total renormalisation in the momentum-resolved Kondo coupling $J^{(j)}$ at the j^{th} step amounts to

$$\Delta J_{\mathbf{k}_1, \mathbf{k}_2}^{(j)} = - \sum_{\mathbf{q} \in \text{PS}} \frac{J_{\mathbf{k}_2, \mathbf{q}}^{(j)} J_{\mathbf{q}, \mathbf{k}_1}^{(j)} + 4 J_{\mathbf{q}, \mathbf{q}}^{(j)} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_1, \mathbf{q}}}{\omega - \frac{1}{2} |\varepsilon_j| + J_{\mathbf{q}}^{(j)}/4 + W_{\mathbf{q}}/2} \tag{3.19}$$

A representation of the RG phase diagram in the $-W/t$ vs. J/t plane of couplings has been provided in Fig.5.4(a) from detailed numerical computations of eq.(??). We also provide a schematic representation of RG flows to various fixed point theories in the J vs. W plane of couplings in Fig.5.3. This schematic figure shows the existence of three distinct phases (Fermi liquid metal, Mott Insulator, and Pseudogap-Mott metal) described by distinct fixed point theories of the RG flows. The Fermi liquid metal and Mott Insulator phases are distinguished from the Pseudogap-Mott metal by separatrices of the RG flows.

5.4 Reconstructing of lattice model from auxiliary model

To reconstruct a translationally invariant lattice Hamiltonian from the impurity model (with Hamiltonian $\mathcal{H}_{\text{aux}}$), we define a many-body tiling prescription that embeds the auxiliary impurity system uniformly across the two-dimensional square lattice. Thus, the tiling process systematically derives the full lattice dynamics from the renormalized fixed-point structure of the impurity. As described




Figure 5.3: Renormalisation group (RG) flow diagram of the lattice model as obtained from a unitary RG analysis of the lattice-embedded impurity model. For $|W/J| < |W/J|_{PG}$, the flows lead to a Fermi liquid phase, while for $|W/J| < |W/J|_c$, they lead to a Mott insulator. For values in between, the system flows to a Mott metal phase that has a pseudogap in the density of states.




Figure 5.4: (a) Phase diagram of impurity model at strong coupling in U in terms of competing dimensionless Kondo (J/t) and bath correlation (W/t) couplings. Colourbar represents the fraction of states on the Fermi surface replaced by zeros of the Greens function (Luttinger surface). A pseudogap phase (shaded region) is observed between the local Fermi liquid (red region) and local moment (blue region) phases (b) k -space-resolved spin-spin correlation $\chi_s(d, \mathbf{k}) = \langle \mathbf{S}_d \cdot \mathbf{S}_{\mathbf{k}} \rangle$ in the pseudogap phase of the impurity model. Antinodal regions are observed to decouple from Kondo screening of the impurity. (c) Upon tiling, this leads to a k -space-resolved antinodal gap in the electronic density of states of the lattice model, corresponding to Luttinger surfaces of zeros.




Figure 5.5: Left: Spin correlations χ_s for the tiled model, between the nodal point $k_N = (-\pi/2, -\pi/2)$ and an arbitrary k -point on the Fermi surface. In the PG, the spin-correlations near the antinode vanish, indicating that they have been removed from the metallic excitations, while the correlations near the nodal point become enhanced because of the increasingly correlated nature of the metal. Right: Mutual information I_2 between an arbitrary k -state and the nodal point. The nodal points remain strongly entangled with each other all the way through the PG.




Figure 5.6: Initiation of the decoupling of $J_{\mathbf{k},\mathbf{k}'}$ (positive, negative and zeros shown in red, blue and white respectively), with $\mathbf{k}$ (black circle) at low-energies for $\mathbf{k}$ corresponding to the node (a), antinode (b) and a point mid-way between them (c) on the top right arm of the FS respectively with tuning W/J . As dictated by the symmetry of $J_{\mathbf{k},\mathbf{k}'}$, the decoupling for a given $\mathbf{k}$ proceeds via the appearance of zeros (white patches) of $J_{\mathbf{k},\mathbf{k}'}$ for $\mathbf{k}'$ initially on the nodal regions of adjacent arms, and progresses gradually towards the antinodes. The decoupling ends with the onset of the pseudogap.

in Section 5.2, the auxiliary model consists of a single impurity embedded in a correlated bath at the location $\mathbf{r}_d$:

$$H(\mathbf{r}_d) = H_{\text{imp}} + H_{\text{cbath}} + H_{\text{imp-cbath}} . \quad (4.1)$$

The form of various terms in eq.(4.1) have been defined in eqs.(2.2), (2.3) and (2.4) in Section 5.2.

The tiled lattice Hamiltonian is generated by symmetrically translating this impurity model:

$$\mathcal{H}_{\text{tiled}} = \mathcal{T}[H(\mathbf{r}_d)] = \sum_{\mathbf{r}} T^\dagger(\mathbf{r} - \mathbf{r}_d) H(\mathbf{r}_d) T(\mathbf{r} - \mathbf{r}_d), \quad (4.2)$$

where $T(\mathbf{a})$ are many-body translation operators [Stoyanova(2006)]:

$$T^\dagger(\mathbf{a}) \mathcal{O}(\{\mathbf{r}_i\}) T(\mathbf{a}) = \mathcal{O}(\{\mathbf{r}_i + \mathbf{a}\}). \quad (4.3)$$

Applying this to each term yields:

$$\begin{aligned} \mathcal{T}[H_{\text{cbath}}] &= -\frac{(N-2)t}{\sqrt{\mathcal{Z}}} \sum_{\langle \mathbf{r}_i, \mathbf{r}_j \rangle; \sigma} (c_{\mathbf{r}_i \sigma}^\dagger c_{\mathbf{r}_j \sigma} + \text{h.c.}) \\ &\quad - \frac{1}{2} W \sum_{\mathbf{r}} (\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow})^2, \\ \mathcal{T}[H_{\text{imp}}] &= -\frac{U}{2} \sum_{\mathbf{r}} (\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow})^2, \\ \mathcal{T}[H_{\text{imp-cbath}}] &= \sum_{\langle \mathbf{r}_i, \mathbf{r}_j \rangle} \left[\frac{2J}{\mathcal{Z}} \mathbf{S}_{\mathbf{r}_i} \cdot \mathbf{S}_{\mathbf{r}_j} - \frac{2V}{\sqrt{\mathcal{Z}}} \sum_{\sigma} (c_{\mathbf{r}_i \sigma}^\dagger c_{\mathbf{r}_j \sigma} + \text{h.c.}) \right], \end{aligned} \quad (4.4)$$

with local spin operators defined as

$$\mathbf{S}_{\mathbf{r}} = \sum_{\sigma, \sigma'} c_{\mathbf{r}\sigma}^\dagger \boldsymbol{\tau}_{\sigma\sigma'} c_{\mathbf{r}\sigma'}. \quad (4.5)$$

To avoid overcounting of the non-interacting bath terms, we subtract extra copies of the kinetic energy term. The reconstructed extended-Hubbard model then reads:

$$\begin{aligned} \mathcal{H}_{\text{tiled}} &= -\frac{1}{\sqrt{\mathcal{Z}}} \tilde{t} \sum_{\langle \mathbf{r}_i, \mathbf{r}_j \rangle; \sigma} (c_{\mathbf{r}_i \sigma}^\dagger c_{\mathbf{r}_j \sigma} + \text{h.c.}) \\ &\quad + \frac{\tilde{J}}{\mathcal{Z}} \sum_{\langle \mathbf{r}_i, \mathbf{r}_j \rangle} \mathbf{S}_{\mathbf{r}_i} \cdot \mathbf{S}_{\mathbf{r}_j} - \frac{1}{2} \tilde{U} \sum_{\mathbf{r}} (\hat{n}_{\mathbf{r}\uparrow} - \hat{n}_{\mathbf{r}\downarrow})^2, \end{aligned} \quad (4.6)$$

where the effective lattice couplings are given by:

$$\tilde{t} = t + 2V, \quad \tilde{U} = U + W, \quad \tilde{J} = 2J. \quad (4.7)$$

A non-zero chemical potential in the bath and/or the impurity can be included to study the effects of doping. Crucially, the eigenstates of $\mathcal{H}_{\text{tiled}}$ obey a many-body generalization of Bloch's theorem [Stoyanova(2006)], which allows for the exact computation of momentum-resolved observables (presented here for a 77×77 $\mathbf{k}$ -space Brillouin zone grid). We shall demonstrate below that the lattice embedding and tiling procedures provide a controlled route to capture low-energy NFL behavior and Mott criticality directly from a quantum impurity model.

5.5 Low-energy theory for the lattice-embedded impurity model

5.5.1 RG Phase Diagram

A detailed picture of the PG in the impurity model is obtained from a momentum-resolved breakdown of Kondo screening in the strong-coupling regime $U \gg t$ phase of H_{aux} . A unitary renormalisation group (URG) scaling analysis [Mukherjee and Lal(2020a)] obtains a flow equation of the Kondo coupling $J_{\mathbf{k}_1, \mathbf{k}_2}^{(j)}$ (see Appendix A for details)

$$\Delta J_{\mathbf{k}_1, \mathbf{k}_2}^{(j)} = - \sum_{\mathbf{q} \in \text{PS}} \frac{J_{\mathbf{k}_2, \mathbf{q}}^{(j)} J_{\mathbf{q}, \mathbf{k}_1}^{(j)} + 4 J_{\mathbf{q}, \mathbf{q}}^{(j)} W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_1, \mathbf{q}}}{\omega - \frac{1}{2} |\varepsilon_j| + J_{\mathbf{q}}^{(j)} / 4 + W_{\mathbf{q}} / 2}, \quad (5.1)$$

where ε_j is the energy of the shell being decoupled at the j^{th} step, the sum is over all occupied momentum states $\mathbf{q}$ of the energy shell ε_j , and $\mathbf{q} = \mathbf{q} + \boldsymbol{\pi}$ is the particle-hole transformed state associated with $\mathbf{q}$. The bath interaction coupling $W_{\mathbf{q}, \mathbf{k}_2, \mathbf{k}_1, \mathbf{q}}$ is found to be marginal under these transformations. While we present a detailed numerical evaluation of the RG equation for $J_{\mathbf{k}_1, \mathbf{k}_2}$ (eq.(5.1)) below, it is clear that the frustration of Kondo screening due to charge fluctuations (for attractive bath interactions $W < 0$) leads to the Mott transition [Mukherjee et al.(2023)Mukherjee, Vidhyadhiraja, Taraphder, and Lal].

A detailed numerical computation of the URG equation in eq. (5.1) leads to the phase diagram shown in Fig. 5.4(a). Upon tuning the ratio of the bath and Kondo interactions (W/J) from zero to negative values, the following phases emerge in the impurity model from the competition between J and W in eq. (5.1): (i) for $W/J < (W/J)_{\text{PG}}$, an LFL phase (red region), where the entire FS participates in Kondo screening, (ii) for $\frac{W}{J} \in [(\frac{W}{J})_{\text{PG}}, (\frac{W}{J})_c]$ (shaded region), a local PG phase where disconnected parts of the FS around the node participate in Kondo screening, and (iii) a local moment phase for $\frac{W}{J} > (\frac{W}{J})_c$ (blue region), where the impurity remains unscreened at low-energies. These can be visualised from spin correlations, $\chi_s(d, \mathbf{k}) = \langle \mathbf{S}_d \cdot \mathbf{S}_{\mathbf{k}} \rangle$, as shown in Fig. 5.4(b) for the PG. The values $(W/J)_{\text{PG}}$ and $(W/J)_c$ are therefore the entry into and exit from the PG phase.

Mapping onto the lattice model via tiling, we observe that RG-induced nodal-antinodal dichotomy in $J_{\mathbf{k}, \mathbf{k}'}$ in the impurity model is the microscopic origin of the PG in the lattice model: the extinction of Kondo coherence translates into spectral zeros in the lattice Green's function (i.e., Luttinger surfaces) at antinodal momenta (Fig. 5.4(c)). This establishes that the $T = 0$ Mott transition of the 2D *extended*-Hubbard model proceeds from FL to MI through an intervening PG phase. In Fig. 5.5, similar insights obtain from the spin-correlation $\chi_s(\mathbf{k}, k_N) = \langle \mathbf{S}_{\mathbf{k}} \cdot \mathbf{S}_{k_N} \rangle$ in k -space (where k_N is the nodal momentum) and mutual information $I_2(\mathbf{k} : k_N)$ between two momentum states: the PG displays Fermi arcs in the nodal regions along with vanishing correlation and entanglement in the antinodal regions.

5.5.2 Unravelling of Kondo Screening

The $\mathbf{k}$ -space anisotropy of Kondo breakdown can be visualized in terms of zeros of $J_{\mathbf{k}_N, \mathbf{k}}$, involving spin-flip scattering between the node $\mathbf{k}_N = (\pi/2, \pi/2)$ and a general wavevector $\mathbf{k}$. For any W/J , the $\mathcal{C}_4$ lattice symmetry dictates that $J_{\mathbf{k}_N, \mathbf{k}}$ vanishes if $\mathbf{k}$ belongs to any of the antinodes or adjacent

nodes. Tuning W/J towards $(W/J)_{\text{PG}}$ leads to an unravelling of the Kondo screening: $J_{\mathbf{k}_N, \mathbf{k}}$ for $\mathbf{k}$ close to the adjacent nodes turns RG-irrelevant first, and a patch of zeros subsequently appears in $J_{\mathbf{k}_N, \mathbf{k}}$ around this point (Fig. 5.6(a)). Tuning W/J further extends the patch of zeros towards the antinodes (Fig. 5.6(b) and 5.6(c)). Kondo screening thus unravels by a systematic decoupling of all $J_{\mathbf{k}_1, \mathbf{k}_2}$ that connect adjacent quadrants of the Brillouin zone.

Precisely at $W/J = (W/J)_{\text{PG}}$, the antinode joins this connected region of zeros in $J_{\mathbf{k}_1, \mathbf{k}_2}$, marking the decoupling of the antinodes from all other points in the neighbourhood of the FS. This is an interaction-driven Lifshitz transition of the FS, and marks the entry into a PG phase possessing Fermi arcs [Wu et al.(2018)Wu, Scheurer, Chatterjee, Sachdev, Georges, and Ferrero]. Importantly, it coincides with an emergent two-channel Kondo (2CK) impurity model, where each channel corresponds to a pair of Fermi arcs on opposite faces of the conduction bath FS (see Subsection 5.5.3). The 2CK nature of the PG is guaranteed by the symmetry of $J_{\mathbf{k}, \mathbf{k}'}$:

$$J_{\mathbf{k}, \mathbf{k}'} = -J_{\mathbf{k}+\mathbf{Q}, \mathbf{k}'} = -J_{\mathbf{k}, \mathbf{k}'+\mathbf{Q}}, \quad \mathbf{Q} = (\pi, \pi). \quad (5.2)$$

The PG expands by shrinking these disconnected Fermi arcs towards the respective nodes, leading to nodal metals whose disappearance heralds the Mott transition.

5.5.3 Emergence of a two-channel Kondo Model

As the bath interaction W is tuned through the L-PG phase, the four nodal points in the Brillouin zone are the last to decouple from the impurity. This allows us to write down a simpler Kondo model near the transition, focusing on scattering processes involving the nodal region. Specifically, for the Kondo term $J_{\mathbf{k}_1, \mathbf{k}_2}$, we find that only the following classes of scattering processes survive close to the critical point:

- Both momenta in a neighbourhood around $\mathbf{k}_N$, where $\mathbf{k}_N$ can be any of the four nodal points,
- One momentum from the neighbourhood around $\mathbf{k}_N$, while the other from the neighbourhood around $\mathbf{k}_N + (\pi, \pi)$.

The second class is tied to the first through a symmetry of the Hamiltonian (eq. ??). The crucial feature of these processes is that each node only interacts with the other node directly opposite to it across the origin. That is, $(\pi/2, \pi/2)$ and $(-\pi/2, -\pi/2)$ only interact between themselves, while $(\pi/2, -\pi/2)$ and $(-\pi/2, \pi/2)$ form their own pair of connected regions.

Let $S^\pm$ be the set of momentum states in a small window around nodes along $k_x = \pm k_y$. As discussed immediately above, the Kondo spin-exchange term close to the transition does not lead to scattering of any k -state from S^+ to S^- . This is verified by calculating the ratio $\max \left\{ J_{k_1^+, k_2^-} \right\} / \max \left\{ J_{k_1^+, k_2^+} \right\}$, where the maximum in the numerator is calculated from all the low-energy (fixed point) Kondo scattering processes that connect the two sets $S^\pm$, while the denominator maximum is from within the set S^+ . This ratio vanishes (left panel of Fig. 5.9) within the local pseudogap regime, indicating that no direct Kondo scattering process persists between the two sectors $S^\pm$ at low energies, deep within the pseudogap regime. We call the value of W at which this ratio vanishes W^* .




Figure 5.7: Ratio between the maximum values of the fixed point values of the Kondo couplings for scattering processes between and within the regions S^+ and S^- . The vanishing of the fixed point ratio indicates that the pseudogap regime features a singular low-energy effective theory, where the Fermi surface neighbourhood splits into two disconnected regions with no explicit term in the Hamiltonian connecting them directly.

The decoupling of the k -space regions $S^\pm$ is captured by the following simplified low-energy Hamiltonian:

$$H_{\text{eff}} = H(S^+) + H(S^-); \quad H(S) = \sum_{\mathbf{q} \in S, \sigma} \varepsilon_{\mathbf{q}} c_{\mathbf{q}\sigma}^\dagger c_{\mathbf{q}\sigma} + \sum_{\mathbf{q}_1 \in S, \mathbf{q}_2 \in S} \sum_{\alpha, \beta} J^*(\mathbf{q}_1, \mathbf{q}_2) \mathbf{S}_d \cdot \sigma_{\alpha\beta} c_{\mathbf{q}_1\alpha}^\dagger c_{\mathbf{q}_2\beta}, \quad (5.3)$$

where $H(S^\pm)$ represents the dynamics of momentum states residing in regions $S^\pm$. The only source of information transfer between the two regions is the fact that they interact with the same impurity local moment. Eq. 5.3 tells us that the low-energy dynamics of the embedded eSIAM close to the Kondo-breakdown transition is governed by an emergent **two-channel Kondo model**, with the conduction bath states in the sets $S^\pm$ making up the two channels.

5.5.4 Momentum-resolved Dynamical Spectral Weight Transfer

Passage through the PG phase is accompanied by a highly structured transfer of spectral weight across the FS. Strong charge fluctuations develop between the nodal and antinodal regions of the FS in the PG regime of the impurity model (Fig. 5.8(a)), as captured by the correlator:

$$\chi_c(\mathbf{k}_1, \mathbf{k}_2) = \left\langle c_{\mathbf{k}_1\uparrow}^\dagger c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{k}_2\downarrow} c_{\mathbf{k}_2\uparrow} + \text{h.c.} \right\rangle. \quad (5.4)$$

These fluctuations dynamically redistribute low-energy spectral weight from the antinodes to higher energies, leading to selective gap formation. Accordingly, the Luttinger surfaces of the PG [Dzyaloshinskii(2003)] coincides with the appearance of poles of the lattice model self-energy $\Sigma(\mathbf{k}, \omega = 0)$ near the antinodes; these poles approach the nodes on tuning towards the Mott transition (Fig. 5.8(b)). This mirrors the coalescing of finite-frequency poles of the self-energy poles towards zero frequency in the underlying impurity model (Fig. 5.8(c)).




Figure 5.8: (a) Enhanced charge correlations $\chi_c(\mathbf{k}_1, \mathbf{k}_2) = \langle c_{\mathbf{k}_1\uparrow}^\dagger c_{\mathbf{k}_1\downarrow}^\dagger c_{\mathbf{k}_2\downarrow} c_{\mathbf{k}_2\uparrow} + \text{h.c.} \rangle$ between the nodal and antinodal regions, signalling Kondo breakdown in the pseudogap phase of the impurity model. (b) In turn, the breakdown leads to the gapping of the antinodal regions in the lattice model, seen from the appearance of poles in the imaginary part of the self-energy. (c) The imaginary part of the impurity self-energy $\Sigma''(\omega > 0)$ possesses a pole at non-zero ω in the PG that moves towards $\omega = 0$ as the Mott transition is approached.

5.6 Non-Fermi liquid excitations within the Pseudogap

5.6.1 Pseudogapped spectral function and singular self-energy

In the PG regime, the nature of gapless Fermi arcs changes dramatically. We have already argued that the low-energy dynamics of these gapless Fermi arcs are governed by an underlying two-channel Kondo (2CK) impurity model [Tsvetick and Wiegmann(1985), Emery and Kivelson(1992)]. This is consistent with the rapid fall of the impurity quasiparticle residue Z_{imp} (Fig. 5.9(a)) from finite values in the FL phase to vanishingly small values just before the onset of the PG. Inset shows the simultaneous emergence of increasingly uncompensated local magnetic moments upon traversing the PG phase. The accompanying impurity spectral function of the gapless arcs show a pseudogapped behaviour for $\omega \simeq 0$, with a rapid fall in the spectral weight at $\omega = 0$ upon traversing the PG (Fig. 5.9(c)). The collapse of the Kondo resonance into a pseudogapped spectral function is accompanied by the redistribution of spectral weight [Dzyaloshinskii(2003)] in the impurity spectral function (Fig. 5.9(b)) from $\omega \sim 0$ to the Hubbard sidebands at finite frequencies $\omega \simeq \pm 3$ (in units of the bandwidth).

Concomitant with this is the emergence of a zero-frequency peak in the imaginary part of the self-energy of the NFL in the PG phase(Fig. 5.10(a)),

$$-1/\Sigma''(\omega) \sim 1/\Sigma''(0) + \omega^\beta. \quad (6.1)$$

Remarkably, we find that the exponent $\beta = 2$ characterises the NFL for the entire PG phase (see Fig. 5.10(b)), including the critical end-point. This is in stark contrast with the $\Sigma''(\omega) \sim \omega^2$ for the FL (Fig. 5.10(b)). A similar analysis reveals the power-law scaling for the low-frequency behaviour of the impurity spectral function in Fig.5.11. We observe that $A_d(\omega) \sim A_d(0) + \omega^2$ for the pseudogap phase, while $1/A_d(\omega) \sim 1/A_d(0) + \omega^2$ for the Fermi liquid (inset of Fig. 5.11).

5.6.2 Finite-Frequency Scattering Rate and Optical Response Through the Pseudogap

All $\Sigma''(\omega)$ for the NFL are observed to lie above the Mott-Ioffe-Regel (MIR) bound [Gunnarsson et al.(2003)Gunnarsson, Calandra, and Han, Hussey et al.(2004)Hussey, Takenaka, and Takagi] (dashed blue line in Fig. 5.10(a)), while those for the FL lie below. The MIR bound is the maximum expected scattering rate in metals when the mean free path approaches the lattice spacing:

$$-2\Sigma''_{\text{MIR}} = 1/\tau_{\text{MIR}}, \quad \tau_{\text{MIR}} = l_{\text{min}}/v_F, \quad (6.2)$$

where l_{min} is the minimum mean free path in metals (equal to one lattice spacing) and τ_{MIR} is the associated lifetime of quasiparticles close to the FS (with Fermi velocity v_F). The height of the zero-frequency peak rises by almost 4 orders of magnitude from the start of the PG till its end at the Mott transition point (Fig. 5.10(c)); the dramatic growth of the peak height very near the Mott critical point coincides with the coalescing of the finite-frequency poles of the self-energy into a single pole at zero-frequency, signalling the singular nodal NFL present at the Mott quantum critical point.

To further probe the breakdown of coherent transport in the pseudogap (PG) regime, we examine the product of the impurity spectral function $A(\omega)$ and the scattering lifetime $\tau(\omega)$, which approximates the optical conductivity $\sigma(\omega)$ via a Drude-type relation. As shown in the main panel of Fig. ??, this quantity exhibits a sharp Drude peak at $\omega = 0$ in the Fermi liquid (FL) regime (blue), consistent with long-lived quasiparticles. In stark contrast, the PG regime (green, red, orange) shows a pronounced suppression of low-frequency conductivity and the emergence of a finite-frequency peak centered near $\omega/D \sim 0.3-0.4$ (where D is the kinetic energy bandwidth).

Strikingly, this peak shifts very little across the PG regime (see inset), despite a deepening spectral gap and diverging self-energy. Its position lies well below the Mott gap ($\sim 2.5D$), suggesting that it represents an emergent mid-infrared (MIR) excitation intrinsic to the Mott metal characterising the PG. This feature closely mirrors the experimentally observed MIR peak in high- T_c cuprates [Herr et al.(1987)Herr, Kamarás, Porter, Doss, Tanner, Bonn, Greedan, Stager, and Timusk, Orenstein et al.(1987)Orenstein, Thomas, Rapkine, Bethea, Levine, Cava, Rietman, and Johnson, Uchida et al.(1991)Uchida, Ido, Takagi, Arima, Tokura, and Tajima], which has been variously attributed to excitations of electrons bound to vacancies [Thomas et al.(1992)Thomas, Rapkine, Cooper, Cheong, Cooper, Schneemeyer, and Waszczak], or the dressing of holes by spin excitations [Dagotto(1994)]. Here, its robust energy scale and insensitivity to increasing incoherence imply a universality persisting through the PG, likely rooted in short-range entanglement between deconfined doublons and holons excitations of the Mott metal. This is further evidence that the PG metal is not simply a degraded Fermi liquid, but a distinct phase whose optical response is driven by emergent degrees of freedom.

5.6.3 Non-local correlations in the pseudogap

In Fig. 5.12(a) and 5.12(b), the spin-flip correlations and mutual information between the impurity spin and conduction bath sites respectively are observed to undergo a crossover within the PG, from a short-ranged behaviour at its onset, to a long-ranged behaviour as the Mott transition approaches.




Figure 5.9: (a) Suppression of quasiparticle residue (Z_{imp}) as the impurity model is tuned towards the Mott transition. An initial drastic fall in Z_{imp} is observed for $W/J \lesssim (W/J)_{\text{PG}}$ from 0.3 to around 0.05, signalling the destruction of the FL with unravelling of Kondo screening. A steady decrease in Z_{imp} is observed in passage through the PG, and is vanishingly small close to the Mott transition due to a divergent self-energy. Inset shows the growth of unscreened impurity magnetic moment in the pseudogap phase, signalling the breakdown of Kondo screening. (b) Impurity Spectral function in the FL and PG phases. The evolution of the central peak from Kondo resonance to pseudogap is accompanied by dynamical spectral weight transfer to the Hubbard side bands at finite frequencies $\omega \simeq \pm 3$ (in units of the bandwidth). (c) Zoom-in of the central (Kondo) resonance of the impurity spectral function. While undisturbed in the Fermi liquid phase, it splits into a pseudogap in the PG phase, with its height diminishing rapidly as the Mott transition is approached.

The entanglement is also observed to be multipartite in nature: in Fig. 5.12(c), the quantum Fisher information (QFI) [Hauke et al.(2016)Hauke, Heyl, Tagliacozzo, and Zoller] computed for the ground state wavefunction using an operator corresponding to the sum of local spin-flip exchange processes shows a jump at the onset of the PG. Further, the FL is observed to possess bi-partite entanglement while the NFL of the PG phase displays pentapartite entanglement [?, ?].

These striking results imply that the Mott transition observed by us lies beyond the local quantum criticality scenario [Si et al.(2001)Si, Rabello, Ingersent, and Smith]. Instead, we observe the PG phase to be a novel state of strongly interacting quantum matter emergent from the breakdown of local Kondo screening. This state is described by a quantum critical Fermi surface [Senthil(2008)] with NFL Fermi arcs that display increasingly critical behaviour, i.e., dynamics described by non-local quantum fluctuations, and excitations that become truly long-ranged close to the transition. It is tempting to conjecture that the long-range entangled critical state captured in Fig. 5.12(b) reflects the existence of a highly efficient scrambler [Sekino and Susskind(2008), Hayden and Preskill(2007), Bentsen et al.(2019)Bentsen, Hashizume, Buyskikh, Davis, Daley, Gubser, and Schleier-Smith], i.e., a state in which entanglement spreads among its constituents at the fastest rate permissible by quantum mechanics.

5.7 Exactly solvable nodal Non-Fermi liquid at Mott Criticality

Very close to the transition, the excitations of the nodal NFL correspond to those of a Hatsugai-Kohmoto model [Baskaran(1991), Hatsugai and Kohmoto(1992)]. This insight is obtained from a perturbation-theoretic treatment of the RG fixed point Hamiltonian of the impurity model for $W/J \lesssim (W/J)_{\text{PG}}$, by considering the effects of a small fixed point Kondo scattering probability J^* in the backdrop of a larger bath interaction parameter $|W|$. This yields the HK model [Baskaran(1991), Hatsugai and Kohmoto(1992)] as the singular part of the effective Hamiltonian arising from forward scattering processes.

As the bath interaction W is tuned through the L-PG phase, the nodal region is the last to decouple from the impurity. This allows us to write down a simpler Kondo model near the transition, where only the nodal region is hybridising with the impurity spin through Kondo interactions. This is done by retaining only those scattering processes $\mathbf{k}_1 \rightarrow \mathbf{k}_2$ that originate from and end at k -points within a small neighborhood of width $|\mathbf{q}|$ around the four nodal points: $\mathbf{k}_1, \mathbf{k}_2 \in \mathbf{N} + \mathbf{q}$, $|\mathbf{q}| \ll \pi$, where $\mathbf{N}$ can be any one of the four nodal points $\mathbf{N}_1 = (\pi/2, \pi/2)$, $\mathbf{N}_2 = (-\pi/2, \pi/2)$ and $\mathbf{N}_1 + \mathbf{Q}_1$ and $\mathbf{N}_2 + \mathbf{Q}_2$, where $\mathbf{Q}_1 = (-\pi, -\pi)$ and $\mathbf{Q}_2 = (\pi, -\pi)$ are the two nesting vectors. We assume that the window of $\mathbf{q}$ is small enough so that the fixed point Kondo coupling values $J^*(\mathbf{q}_1, \mathbf{q}_2)$ for the scattering processes involving $\mathbf{q}_1$ and $\mathbf{q}_2$ can be replaced by an average value J^* .

With these considerations, the simplified low-energy model near the transition describing the Kondo scattering processes can be written as

$$\tilde{H}_{\text{imp-cbath}} = J^* \frac{1}{2} \sum_{l=1,2} \sum_{\mathbf{q}_1, \mathbf{q}_2} \sum_{\alpha, \beta} \mathbf{S}_d \cdot \boldsymbol{\sigma}_{\alpha\beta} \left(c_{\mathbf{N}_l + \mathbf{q}_1, \alpha}^\dagger c_{\mathbf{N}_l + \mathbf{q}_2, \beta} + c_{\mathbf{N}_l + \mathbf{Q}_l + \mathbf{q}_1, \alpha}^\dagger c_{\mathbf{N}_l + \mathbf{Q}_l + \mathbf{q}_2, \beta} - c_{\mathbf{N}_l + \mathbf{Q}_l + \mathbf{q}_1, \alpha}^\dagger c_{\mathbf{N}_l + \mathbf{q}_2, \beta} - c_{\mathbf{N}_l + \mathbf{q}_1, \alpha}^\dagger c_{\mathbf{N}_l + \mathbf{Q}_l + \mathbf{q}_2, \beta} \right). \quad (7.1)$$

The label l can take values 1 or 2, allowing us to consider both the decoupled channels in the PG (associated with $\mathbf{N}_1$ and $\mathbf{N}_2$). It also labels the nesting vectors $\mathbf{Q}_l$ associated with the two sets. α and β indicate spin indices, and $\mathbf{q}_1$ and $\mathbf{q}_2$ represent incoming and outgoing momenta in the scattering processes.

Because of the decoupling of the channel $l = 1$ and $l = 2$, we consider only the $l = 1$ channel for the rest of the calculations in this section. In order to simplify the Hamiltonian, we define new fermionic operators

$$\psi_{\mathbf{q}, \sigma} = \frac{1}{\sqrt{2}} (c_{\mathbf{N}_1 + \mathbf{q}, \sigma} + c_{\mathbf{N}_1 + \mathbf{Q}_1 - \mathbf{q}, \sigma}), \quad \phi_{\mathbf{q}, \sigma} = \frac{1}{\sqrt{2}} (c_{\mathbf{N}_1 + \mathbf{q}, \sigma} - c_{\mathbf{N}_1 + \mathbf{Q}_1 - \mathbf{q}, \sigma}), \quad (7.2)$$

The operator satisfy fermionic anticommutation relations. For convenience, we define new number operators for the sum and relative degrees of freedom:

$$s_{\mathbf{q}, \sigma} = \psi_{\mathbf{q}}^\dagger \psi_{\mathbf{q}}, \quad r_{\mathbf{q}, \sigma} = \phi_{\mathbf{q}}^\dagger \phi_{\mathbf{q}}. \quad (7.3)$$

We then have the following useful relation between the corresponding number operators $n = c^\dagger c$:

$$n_{\mathbf{N}_1 + \mathbf{q}, \sigma} + n_{\mathbf{N}_1 + \mathbf{Q}_1 - \mathbf{q}, \sigma} = s_{\mathbf{q}, \sigma} + r_{\mathbf{q}, \sigma}. \quad (7.4)$$

In terms of these new degrees of freedom, the auxiliary model Hamiltonian takes the form

$$\tilde{H} = -\frac{1}{2}W \sum_{\mathbf{q},\sigma} r_{\mathbf{q},\sigma} + W \sum_{\mathbf{q}_1,\mathbf{q}_2} r_{\mathbf{q}_1,\uparrow} r_{\mathbf{q}_2,\downarrow} + \sum_{\mathbf{q}_1,\mathbf{q}_2,\alpha,\beta} J^* \mathbf{S}_d \cdot \boldsymbol{\sigma}_{\alpha\beta} \phi_{\mathbf{q}_1\alpha}^\dagger \phi_{\mathbf{q}_2\beta} + \sum_{\mathbf{q},\sigma} \varepsilon_{\mathbf{N}_1+\mathbf{q}} (\psi_{\mathbf{q},\sigma}^\dagger \phi_{\mathbf{q},\sigma} + \phi_{\mathbf{q},\sigma}^\dagger \psi_{\mathbf{q},\sigma}), \quad (7.5)$$

where we have considered a simplified form of the bath interaction (for the channel $l = 1$), taking into account the density-density correlations in k -space.

For bath interaction strength close to the critical value ($W \lesssim W_c$), the fixed point coupling value J^* is much smaller than W . In order to obtain the gapless excitations of the system arising from the presence of the impurity site, we integrate out the impurity dynamics via a Schrieffer-Wolff transformation. The perturbation term $\mathcal{V}$ then consists of Hamiltonian terms that modify the impurity configuration,

$$\mathcal{V} = \sum_{\mathbf{q},\sigma} \varepsilon_{\mathbf{N}_1+\mathbf{q}} (\psi_{\mathbf{q},\sigma}^\dagger \phi_{\mathbf{q},\sigma} + \phi_{\mathbf{q},\sigma}^\dagger \psi_{\mathbf{q},\sigma}) + \sum_{\mathbf{q}_1,\mathbf{q}_2} J^* S_d^+ \phi_{\mathbf{q}_1\downarrow}^\dagger \phi_{\mathbf{q}_2\uparrow} + \text{h.c.}, \quad (7.6)$$

while the “non-interacting” Hamiltonian is

$$H_D = -\frac{1}{2}W \sum_{\mathbf{q},\sigma} r_{\mathbf{q},\sigma} + W \sum_{\mathbf{q}_1,\mathbf{q}_2} r_{\mathbf{q}_1,\uparrow} r_{\mathbf{q}_2,\downarrow}. \quad (7.7)$$

For the present Hamiltonian, the low-energy state is the one that minimises the bath interaction term $\tilde{H}_{\text{bath-int}}$. High-energy states are obtained by applying, on the state $|L\rangle$, the excitation operator $\phi_{\mathbf{q}_1,\sigma}^\dagger \phi_{\mathbf{q}_2,\bar{\sigma}}$ or its hermitian conjugate.

The complete second-order renormalised Hamiltonian is

$$\Delta\tilde{H} = \sum_{\mathbf{q},\sigma} \epsilon_{\mathbf{q}} r_{\mathbf{q},\sigma} + \mathcal{U} \sum_{\mathbf{q},\sigma} r_{\mathbf{q}\sigma} r_{\mathbf{q}\bar{\sigma}} + \mathcal{U} \sum_{\mathbf{q}_1 \neq \mathbf{q}_2, \sigma} \left[r_{\mathbf{q}_1\sigma} r_{\mathbf{q}_2\bar{\sigma}} + \phi_{\mathbf{q}_1,\bar{\sigma}}^\dagger \phi_{\mathbf{q}_1,\sigma}^\dagger \phi_{\mathbf{q}_2,\sigma} \phi_{\mathbf{q}_2,\bar{\sigma}} \right]. \quad (7.8)$$

where $\epsilon_{\mathbf{q}} = \text{sign}(\varepsilon_{\mathbf{N}_1+\mathbf{q}}) \frac{\varepsilon_{\mathbf{N}_1+\mathbf{q}}^2}{-W}$ and $\mathcal{U} = \frac{J^{*2}}{4W}$.

The $\mathbf{q}_1 = \mathbf{q}_2$ component of the Hamiltonian shows the emergence of the exactly solvable Hatsugai-Kohmoto model [Baskaran(1991), Hatsugai and Kohmoto(1992)] at the critical point. The correlation term $\mathcal{U} \sum_{\mathbf{q},\sigma} r_{\mathbf{q}\sigma} r_{\mathbf{q}\bar{\sigma}}$ leads to a transfer of spectral weight across the Fermi surface and separates the available states into three classes:

$$\langle n_{\mathbf{q}} \rangle = \begin{cases} 2, & \epsilon_{\mathbf{q}} < -|\mathcal{U}/2|, \\ 1, & |\mathcal{U}/2| > \epsilon_{\mathbf{q}} > -|\mathcal{U}/2|, \\ 0, & |\mathcal{U}/2| < \epsilon_{\mathbf{q}}. \end{cases} \quad (7.9)$$

In contrast to a Fermi liquid, there is a highly degenerate single-occupied region in the middle, and this gives rise to non-Fermi liquid excitations.

We now discuss the effects of the $\mathbf{q}_1 \neq \mathbf{q}_2$ component. The first term partially lifts the degeneracy of the central singly-occupied region and allows only zero magnetisation configurations. The second term creates gapped excitations involving the regions of zero and double occupancy; these




Figure 5.10: (a) Imaginary part of impurity self-energy $\Sigma''(\omega)$ as a function of frequency ω for Fermi liquid (FL, $|W/J| < 1.79$) and pseudogapped phases (NFL, $|W/J| \geq 1.79$). $\Sigma''(\omega)$ falls to zero as $\omega \rightarrow 0$ for the FL, while it attains a peak in the pseudogap. All $\Sigma''(\omega)$ for the NFL are observed to lie above the Mott-Ioffe-Regel (MIR) bound (dashed blue line), while those for the FL lie below. (b) Scaling of $\Sigma''(\omega)$ with frequency for Fermi liquid and pseudogapped phases. The FL self-energy fits to $\Sigma'' \sim \omega^\alpha$ with $\alpha \approx 2$, vanishing as $\omega \rightarrow 0$, while the NFL self-energy grows as $\Sigma'' \approx a - \omega^\beta$ for small ω , with $\beta \approx 2$. Remarkably, the NFL exponent remains mostly unchanged through the entirety of the PG phase. (c) Variation of the zero-frequency imaginary self-energy $-\Sigma''(\omega = 0)$ with ω in the FL and pseudogap phases (entry into the PG is marked by the vertical dashed line). The inset shows the same but in linear scale, in order to display the dramatic rise (by almost 30 times) on entering the PG.

represent subdominant pairing fluctuations of the nodal non-Fermi liquid. We recall that similar fluctuations across the Fermi surface were observed in our calculations of charge correlations in Fig. 5.8 (a), and are consistent with pairing fluctuations observed from quantum Monte Carlo simulations and found to give rise to quantum-critical behavior and PG physics [Jiang et al.(2022)Jiang, Liu, Klein, Wang, Sun, Chubukov, and Meng].

Consequently, the resulting NFL metal of the lattice model involves long-lived excitations of multiple $\mathbf{k}$ -states, and manifest in the form of a divergent one-particle self-energy at the non-interacting FS [Phillips et al.(2020)Phillips, Yeo, and Huang]: $\Sigma_{\mathbf{q}}(\omega) = -\mathcal{U}^2/4(\omega - \epsilon_{\mathbf{q}})$, such that $\Sigma_{\epsilon_{\mathbf{q}}=0}(\omega \rightarrow 0) \rightarrow \infty$. This zero-frequency self-energy pole presages the transition into a Mott insulating phase, where it marks a hard gap in the spectral function for charge excitations. This is consistent with our findings for the lattice (Fig. 5.8(c)) and impurity self-energies (Fig. 5.10(c)). The nodal Mott metal thus comprises of a Greens function zero at the Fermi energy together with an anisotropic massless Dirac dispersion, leading to a non-zero Chern number [Vafeek and Vishwanath(2014b), Morimoto and Nagaosa(2016), Flores-Calderón and Hooley(2025)]. This topological index survives into the Mott insulator.

For small but non-zero values of $\omega - \epsilon_{\mathbf{k}}$, we obtain a quasiparticle residue that vanishes with ω , $Z_{\text{imp}} \sim \omega^2/\mathcal{U}^2$. The scattering rate of this singular NFL possesses a sharp peak at the FS ($\omega = \epsilon_{\mathbf{k}_F}$):

$$\Gamma \sim \mathcal{U}^2 \delta(\omega - \epsilon_{\mathbf{k}}), \quad (7.10)$$

consistent with the sharply peaked Lorentzian $\Sigma''_{\mathbf{k}}(\omega) \sim \omega^{-2}$ captured in Fig. 5.10(a). This quantum critical NFL metal is an example of a strongly coupled scale-invariant form of quantum matter. The exact solution for eq.(??) reveals the presence of low-energy excitations comprised of




Figure 5.11: Left: Impurity spectral function $A_d(\omega)$ in the local Fermi liquid (inset) and pseudogap (main). The PG spectral function is fit to a power law (with the $\omega = 0$ contribution subtracted, see Fig. 5.10(c) for the full spectral function). The exponent of the power law comes out to be $n \approx 2$ throughout the PG. The local FL spectral function (inset) is fit to a lorentzian (by fitting the inverse to a power law). Right: Optical conductivity $\sigma(\omega)$ for excitations of the impurity model, approximated through the product of the carrier density (taken from the density of states $A(\omega)$) and the lifetime $\tau(\omega)$ (obtained from the inverse scattering rate $-1/\Sigma''(\omega)$, where Σ'' is the imaginary part of the self-energy). The Fermi liquid (blue) shows a sharp Drude peak at $\omega = 0$ due to the presence of long-lived quasiparticles. The PG (green, red, orange) shows a shifted Drude peak at a non-zero frequency which is reminiscent of the mid-infrared (MIR) peak seen experimentally in the cuprates [Herr et al.(1987)Herr, Kamarás, Porter, Doss, Tanner, Bonn, Greedan, Stager, and Timusk, Orenstein et al.(1987)Orenstein, Thomas, Rapkine, Bethea, Levine, Cava, Rietman, and Johnson, Uchida et al.(1991)Uchida, Ido, Takagi, Arima, Tokura, and Tajima]. Inset shows the robustness of the position of the MIR peak through the PG, indicating it as a universal feature of the Mott metal.

holons and doublons [Hatsugai and Kohmoto(1992)]. These features point to the nodal NFL as a long-ranged and multipartite entangled, strongly interacting scale-invariant state of quantum matter [Georgi(2007a), Georgi(2007b), Phillips et al.(2013)Phillips, Langley, and Hutasoit, Phillips(2014)] that are completely disconnected from the quasiparticles of the FL. Following the arguments laid out in [Phillips et al.(2022)Phillips, Hussey, and Abbamonte], at finite temperatures, such a scale-invariant highly entangled non-Fermi liquid is likely associated with Planckian dissipation (i.e., $\Sigma''(T) \sim k_B T$) and a resistivity that varies linearly with temperature ($\rho \sim T$) [?, Legros et al.(2019)Legros, Benhabib, Tabis, Laliberté, Dion, Lizaire, Vignolle, Vignolles, Raffy, Li et al., Hartnoll and Mackenzie(2022)]. This sits well with the possibility that this state is also a highly efficient scrambler [Sekino and Susskind(2008), Hayden and Preskill(2007), Bentsen et al.(2019)Bentsen, Hashizume, Buyskikh, Davis, Daley, Gubser, and Schleier-Smith]. Additionally, as mentioned above, we observe that the nodal metal possesses pairing fluctuations that can become dominant upon doping [Phillips et al.(2020)Phillips, Yeo, and Huang].




Figure 5.12: (a) Spin correlation $\langle \mathbf{S}_d \cdot \mathbf{S}_r \rangle$ and (b) mutual information $I_2(d, r)$ between the impurity spin and conduction bath local spin density as a function of the distance $\mathbf{r}$ between them, normalised against the value at $r = 1$. Both decay very quickly in the FL phase (blue), but show long-ranged behaviour in the NFL phase (green and red), extending to the edges of the system at the critical point (red). (c) Evolution of the Quantum Fisher Information F_Q for a nearest-neighbour spin-flip operator $\mathcal{O} = \sum_{i \in \text{odd}} (S_i^+ S_{i+1}^- + \text{h.c.})$ through the first Lifshitz transition and the pseudogap. The vertical dashed line marks the onset of the PG. To the left of it, the QFI in the Fermi liquid phase shows at most bipartite entanglement ($F_Q < 2$ (below blue dashed line)), while the PG shows the presence of multipartite entanglement upto 5 parties ($F_Q > 4$ (red dashed line)).

5.8 Heisenberg Model as a low-energy description of the tiled Mott insulator

For simplicity, we consider two impurity sites labelled 1 and 2 associated with two nearest-neighbour auxiliary models. The ground state subspace is four-fold degenerate:

$$|\Psi_L\rangle = \{|\sigma_1, \sigma_2\rangle\}, \quad \sigma_i = \pm 1, \quad (8.1)$$

where σ_i is the spin state of site i . This ground state is derived from the following "zeroth order" Hamiltonian that emerges in the local moment phase of the auxiliary models when all scattering processes between the impurity and conduction bath are RG-irrelevant:

$$H_0 = -\frac{U}{2} \sum_{i=1,2} (n_{i\uparrow} - n_{i\downarrow})^2; \quad (8.2)$$

the local correlation on the impurity site becomes the largest scale in the problem in this phase and pushes the $|n_i = 2\rangle$ and $|n_i = 0\rangle$ states to high energies. This then defines the high-energy subspace for our calculation:

$$|\Psi_H\rangle = |C_1, C_2\rangle, \quad (8.3)$$

where C_i can take values 0 or 2, indicating that the state i is either empty or full, respectively. Both the double and hole states exist at a charge gap of the order of $U/2$ above the low-energy singly-occupied subspace defined by the states $|\Psi_L\rangle$.

In order to allow virtual fluctuations that can lift the large ground state degeneracy and lower the energy, we consider (perturbatively) the effects of an irrelevant single-particle hybridisation that

connects the nearest-neighbour sites. This perturbation Hamiltonian is therefore of the form

$$H_t = \sum_{\omega} V(\omega) \mathcal{P}(\omega) \sum_{\sigma} \left(c_{1\sigma}^{\dagger} c_{2\sigma} + \text{h.c.} \right), \quad (8.4)$$

where $V(\omega)$ only acts on states at the energy scale ω ; the renormalisation of V is encoded in the fact that $V(\omega)$ is largest for the excited states and vanishes at low-energies: $V(\omega \rightarrow 0) = 0$.

In order to obtain a low-energy effective Hamiltonian for the impurity sites arising from this hybridisation, we integrate out H_t via a Schrieffer-Wolff transformation. This leads to the following second-order Hamiltonian:

$$H_{\text{eff}} = \mathcal{P}_L H_t G \mathcal{H}_t \mathcal{P}_L. \quad (8.5)$$

The operator $\mathcal{P}_L$ projects onto the low-energy subspace $|\Psi_L\rangle$ - this ensures that we remain in the low-energy subspace at the beginning and at the end of the total process. The Greens function $G = (E_L - H_0)^{-1}$ incorporates the excitation energy to go from the low-energy subspace $|\Psi\rangle_L$ (of energy E_L) to the excited subspace $|\Psi\rangle_H$ of energy $E_L + U/2$. Substituting the form of the perturbation Hamiltonian and the excitation energy into the above expression gives

$$H_{\text{eff}} = \frac{V_H^2}{-U/2} \sum_{\sigma, \sigma'} \left[c_{1\sigma}^{\dagger} c_{2\sigma} c_{2\sigma'}^{\dagger} c_{1\sigma'} + c_{2\sigma}^{\dagger} c_{1\sigma} c_{1\sigma'}^{\dagger} c_{2\sigma'} \right]. \quad (8.6)$$

where $V_H \equiv V(\omega \rightarrow U/2)$ is the impurity-bath hybridisation at energy scales of the order of the Mott gap, in the sense of an RG flow. Terms with consecutive creation or annihilation operators on the same site are prohibited because each site is singly-occupied in the ground state. It is now easy to cast this Hamiltonian into a more recognizable form. For $\sigma' = \sigma$, we get

$$\sum_{\sigma} \delta_{\sigma, \sigma'} c_{1\sigma}^{\dagger} c_{2\sigma} c_{2\sigma'}^{\dagger} c_{1\sigma'} = \sum_{\sigma} (n_{1\sigma} - n_{1\sigma} n_{2\sigma}), \quad (8.7)$$

while $\sigma = -\sigma' = \pm 1$ gives

$$\sum_{\sigma} \delta_{\sigma, -\sigma'} c_{1\sigma}^{\dagger} c_{2\sigma} c_{2\sigma'}^{\dagger} c_{1\sigma'} = - (S_1^+ S_2^- + \text{h.c.}). \quad (8.8)$$

For the latter expression, we introduced the local spin-flip operators $S_i^{\pm}$. The expression above it can also be cast into spin variables, using the equations

$$\begin{aligned} \frac{1}{2} \sum_{\sigma} n_{i\sigma} &= \frac{1}{2}, \\ \frac{1}{2} \sum_{\sigma} \sigma n_{i\sigma} &= S_i^z, \end{aligned} \quad (8.9)$$

where the first equation is simply the condition of half-filling at each site, and the second equation is the definition of the local spin operator in z -direction. Adding and subtracting the equations gives $n_{i\sigma} = \frac{1}{2} + \sigma S_i^z$.

Substituting everything back into eq. 8.6 and dropping constant terms gives

$$H_{\text{eff}} = 2 \frac{V_H^2}{U/2} (2S_1^z S_2^z + S_1^+ S_2^- + S_1^- S_2^+) = J_{\text{eff}} \mathbf{S}_1 \cdot \mathbf{S}_2, \quad (8.10)$$

where the effective antiferromagnetic Heisenberg coupling is $J_{\text{eff}} = \frac{8V_H^2}{U}$.

5.9 Effective Hamiltonian for excitons in the Mott insulator

The Mott insulating phase is characterised by a large onsite correlation U that forbids long-lived coherent charge propagation. This suggests a natural separation of the total Hamiltonian into a large zeroth order part arising from the atomic limit of the Hubbard model, followed by a “small” perturbative part due to irrelevant hopping processes:

$$H_0 = -\frac{U}{2} \sum_i (n_{i\uparrow} - n_{i\downarrow})^2, \quad H_1 = -t \sum_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger c_{j,\sigma} + \text{h.c.}) . \quad (9.1)$$

The zeroth order ground state is of course half-filled, each site being singly-occupied:

$$|\Psi_0\rangle = \otimes_i |\sigma_i\rangle , \quad (9.2)$$

where σ_i can be ± 1 and indicates that each site hosts a local moment that can be either up or down, but doubles or holes are prohibited. This also reflects the large degeneracy ($\sim 2^N$) associated with such a state.

We first note that a non-zero hopping probability t indicates that the ground state cannot be just $|\Psi_0\rangle$, simply because $|\Psi_0\rangle$ is not an eigenstate of H_1 . In order to consider hopping processes that start from the half-filled ground state, we define a projector $P_H = \otimes_i P_H(i) = \otimes_i (n_{i\uparrow} + n_{i\downarrow} - 2n_{i\uparrow}n_{i\downarrow})$ that annihilates the hole or double state at any site. The only allowed processes are therefore

$$H_{\text{eff}} = H_1 P_H = -t \sum_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger P_H(i) c_{j,\sigma} P_H(j) + \text{h.c.}) . \quad (9.3)$$

The action of the projector on any local operator is easy to calculate:

$$\begin{aligned} c_{j,\sigma} P_H(j) &= c_{j\sigma} (n_{j\sigma} + n_{j\bar{\sigma}} - 2n_{j\sigma}n_{j\bar{\sigma}}) = c_{j\sigma} (1 - n_{j\bar{\sigma}}) , \\ c_{i,\sigma}^\dagger P_H(i) &= c_{i,\sigma}^\dagger (n_{j\sigma} + n_{j\bar{\sigma}} - 2n_{j\sigma}n_{j\bar{\sigma}}) = c_{i,\sigma}^\dagger n_{i\bar{\sigma}} , \end{aligned} \quad (9.4)$$

The effective Hamiltonian for such hopping processes from the correlated ground state is therefore

$$H_{\text{eff}} = -t \sum_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger c_{j,\sigma} n_{i\bar{\sigma}} (1 - n_{j\bar{\sigma}}) + \text{h.c.}) . \quad (9.5)$$

In order to identify the propagating degrees of freedom, we define an exciton creation operator $b^\dagger$:

$$b_{i,\delta}^\dagger = B \sum_\sigma c_{i,\sigma}^\dagger c_{i+\delta,\sigma} n_{i\bar{\sigma}} (1 - n_{i+\delta,\bar{\sigma}}) , \quad (9.6)$$

where δ is a primitive lattice vector. [CHECK THE ALGEBRA.] This operator creates an exciton: a bound holon-doublon pair between the sites i and $i + \delta$. In terms of these operators, the Hamiltonian becomes

$$H_{\text{eff}} = -tB \sum_{i,\delta} (b_{i,\delta}^\dagger + \text{h.c.}) . \quad (9.7)$$

5.10 Pseudogap as a strongly coupled phase of quantum matter

We now unveil an organising principle that leads to the remarkable properties observed above for the strongly interacting NFL of the PG phase. The existence of a sharp connected FS at $T = 0$ can be understood as the existence of a topologically protected manifold of gapless chiral excitations in $\mathbf{k}$ -space at the FS [Heath and Bedell(2020a)]. The FS is characterised by a topological index corresponding to an anomaly in the quantum many-body theory for electrons, and can be understood as a generalised symmetry of such a system [Nave et al.(2025)Nave, Zhao, and Phillips, McGreevy(2023)]. A theorem by Luttinger and Ward [Luttinger and Ward(1960)] shows that a count of the physical charge (known as Luttinger’s volume) is identical to the topological index (a so-called homotopy charge known as Luttinger’s count) even in the presence of electronic interactions that do not disturb the FS. We will now argue that the emergence of antinodal Luttinger surfaces involve a disconnection of the FS (into Fermi arcs) and that, by following La Nave et al. [Nave et al.(2025)Nave, Zhao, and Phillips], the accompanying change in its topological properties leads to the existence of gapless NFL excitations that are non-local in nature.

The antinodal Luttinger surfaces arise from the splitting of double poles of the single-particle Greens function on the FS into poles lying on opposite complex half-planes, together with zeros that are pinned at the FS. These changes in the analytic structure of the single-particle Greens function have important consequences. First, the emergent zeros break a $\mathbb{Z}_2$ symmetry of the FS [Anderson and Haldane(2001), Huang et al.(2022)Huang, Nave, and Phillips, Nave et al.(2025)Nave, Zhao, and Phillips]. This symmetry is guaranteed within FL theory because the presence of quasiparticles vanishingly close to the FS allows the interchange of spin and charge degrees of freedom, promoting the separate $SU(2)$ symmetries of spin and charge to the larger symmetry of $O(4) = SU(2) \times SU(2) \times \mathbb{Z}_2$ [Anderson and Haldane(2001)]. The insertion of zeros of the Greens function at the Fermi surface in the PG, and the associated splitting of the Greens function, breaks this $\mathbb{Z}_2$ symmetry by placing the pole for the spin excitation in one half of the complex plane while placing that for the charge excitation in the other half [Su and Martin(2025), Huang et al.(2022)Huang, Nave, and Phillips].

Second, they signal a divergent electronic self-energy as a function of the wavevector $\mathbf{k}$, render ill-behaved the Luttinger-Ward functional of the interacting electronic problem, and violate the generalised symmetry encoded within it. The changes in the pole structure change the Luttinger count topological invariant, while the zeros give rise to an additional topological contribution (linked to the Adler-Bell-Jackiw-type chiral anomaly) [Adler(1969), Bell and Jackiw(1969), Altshuler et al.(1998)Altshuler, Chubukov, Dashevskii, Finkel’stein, and Morr, Flores-Calderón and Hooley(2025)]. As a consequence of the half-filled particle-hole symmetric nature of the system at hand, Luttinger’s volume is preserved upon taking into account topological contributions from both the Luttinger count *and* the zeros [Seki and Yunoki(2017)]. Importantly, La Nave et al. [Nave et al.(2025)Nave, Zhao, and Phillips] argue, following recent developments in understanding generalised symmetries [Casini and Magán(2021)], that the additional anomaly arising from the Luttinger surfaces guarantees the existence of gapless NFL excitations that are non-local in nature.

Thus, the drastic change in nature of the real-space excitations - from locally well-defined Landau quasiparticles of the FL to the increasingly nonlocal excitations of the NFL Fermi arcs in the PG phase - appears to be dictated by a topological principle and, therefore, robust under renormalisation.

This signals the NFL Fermi arcs of the PG as an emergent phase of strongly interacting quantum matter - which we dub the *Mott metal* - that is the parent metal of the MI. Such an identification of the pseudogap as a distinct phase is backed up by angle-resolved photoemission spectroscopy (ARPES) experiments [Hashimoto et al.(2014)Hashimoto, Vishik, He, Devereaux, and Shen]; the anomalous momentum, temperature and doping dependence of the gap function observed in the cuprates suggests that the pseudogap maybe described by an order parameter distinct from that of the superconductor [Hashimoto et al.(2012)Hashimoto, He, Vishik, Schmitt, Moore, Lu, Yoshida, Eisaki, Hussain, Devereaux et al., Vishik et al.(2009)Vishik, Nowadnick, Lee, Shen, Moritz, Devereaux, Tanaka, Sasagawa, and Fujii]. The same topological principle connects the nonlocal unparticle-like gapless excitations [Georgi(2007a), Georgi(2007b), Phillips et al.(2013)Phillips, Langley, and Hutasoit, Phillips(2014)] of the scale-invariant nodal NFL of the HK model observed precisely at the Mott critical point to those of the rest of the PG phase, e.g., a universal scaling of $\Sigma''_{\mathbf{k}}(\omega) \sim (a + \omega^2)^{-1}$ (Fig. 5.10(b)) and local density of states $A_d(\omega) \sim \omega^2$ (Fig. 5.11) of the NFL throughout the PG phase, such that the value of $\Sigma''_{\mathbf{k}}(\omega = 0)$ value continues to grow upon violating the MIR bound (Fig. 5.10(c)).

5.11 Conclusions

Our analysis reveals that the Mott transition proceeds via a continuous evolution through a PG regime characterized by a singular NFL metal - the Mott metal - with deconfined holon–doublon excitations confined to nodal Fermi arcs, and a scaling of it’s spectral features with universal power-law exponents. As the system approaches criticality, this metallic phase exhibits increasingly non-local correlations and a divergent self-energy, signalling the breakdown of Landau quasiparticles, formation of local moments and the onset of long-range quantum entanglement. Anchored in two-channel Kondo dynamics at intermediate scales and governed by Hatsugai–Kohmoto physics near the critical endpoint, the Mott metal provides a unified framework for understanding anomalous metallicity in strongly correlated systems. It’s fate under finite doping presents a compelling direction for future investigation. More complicated models can be studied by suitably extending the tiling approach and choosing an appropriate auxiliary model. Electron-phonon coupling can be taken into account by studying an Anderson-Holstein model that couples the impurity site to local lattice modes [Hewson and Meyer(2001)]. The effects of disorder can be studied by averaging quantities over various disorder configurations of the conduction bath. The versatility of the unitary RG solver and the impurity model-based construction makes it convenient to explore a large variety of lattice models.

5.12 Appendix: Effective action for the Hubbard-Heisenberg model and the eSIAM

In order to better understand the similarities and differences between the tiling approach outlined here and the approach adopted in dynamical mean-field approximations, we compute the effective action for the local impurity as well as the impurity-zeroth site system, in both the auxiliary and bulk

models. This will be done in the limit of infinite coordination number, in order to create a tractable effective theory.

5.12.1 Local theory for the Hubbard-Heisenberg model

We will first work with the Hubbard-Heisenberg model. Let us recall that within the eSIAM, most of the dynamics is governed by the physics of the impurity site and the bath site coupled to the impurity. Accordingly, we will first obtain an effective action within the Hubbard-Heisenberg model for the small system consisting of a local site and all its nearest-neighbours.

We choose a certain local site that we call 0, whose nearest-neighbours will be denoted as $\{\bar{0}\}$. The action for the full Hubbard-Heisenberg model can be formally separated into three parts:

$$S_{\text{H-H}} = S_{0,\bar{0}} + S^{(0,\bar{0})} + S_{\text{int}}, \quad (12.1)$$

where $S_{0,\bar{0}}$ represents the part of the action that involves only the sites 0 and $\{\bar{0}\}$, $S^{(0,\bar{0})}$ represents the part that has all the sites apart from 0 and $\{\bar{0}\}$, while S_{int} contains all terms that connect these two parts. These three parts have the forms

$$\begin{aligned} S_{0,\bar{0}} = & \sum_{i=0,\bar{0}} \sum_{\sigma} \int_0^{\beta} d\tau c_{i\sigma}^{\dagger}(\tau) \partial_{\tau} c_{i\sigma}(\tau) - t \sum_{\bar{0},\sigma} \int_0^{\beta} d\tau \left[c_{0\sigma}^{\dagger}(\tau) c_{\bar{0}\sigma}(\tau) + \text{h.c.} \right] \\ & - \frac{U}{2} \sum_{i=0,\bar{0}} \int_0^{\beta} d\tau (n_{i\uparrow}(\tau) - n_{i\downarrow}(\tau))^2 + J \sum_{\bar{0}} \int_0^{\beta} d\tau \vec{S}_0(\tau) \cdot \vec{S}_{\bar{0}}(\tau), \end{aligned} \quad (12.2)$$

$$\begin{aligned} S^{(0,\bar{0})} = & \sum_{j \neq (0,\bar{0})} \sum_{\sigma} \int_0^{\beta} d\tau c_{j\sigma}^{\dagger}(\tau) \partial_{\tau} c_{j\sigma}(\tau) - t \sum_{\langle j,l \rangle \neq (0,\bar{0}),\sigma} \int_0^{\beta} d\tau \left[c_{j\sigma}^{\dagger}(\tau) c_{l\sigma}(\tau) + \text{h.c.} \right] \\ & - \frac{U}{2} \sum_{j \neq (0,\bar{0})} \int_0^{\beta} d\tau (n_{j\uparrow}(\tau) - n_{j\downarrow}(\tau))^2 + J \sum_{\langle j,l \rangle \neq (0,\bar{0})} \int_0^{\beta} d\tau \vec{S}_l(\tau) \cdot \vec{S}_j(\tau), \end{aligned} \quad (12.3)$$

$$\begin{aligned} S_{\text{int}} = & -t \sum_{i \in \bar{0}} \sum_{j \in \text{NN of } i} \sum_{\sigma} \int_0^{\beta} d\tau \left(c_{i\sigma}^{\dagger}(\tau) c_{j\sigma}(\tau) + \text{h.c.} \right) + J \sum_{i \in \bar{0}} \sum_{j \in \text{NN of } i} \int_0^{\beta} d\tau \vec{S}_i(\tau) \cdot \vec{S}_j(\tau) \\ = & S_{\text{int}}^{\text{hop}} + S_{\text{int}}^{\text{spin}} \end{aligned} \quad (12.4)$$

In order to obtain an effective theory $S_{\text{eff}}^{0,\bar{0}}$ purely for the local system $(0, \bar{0})$, we need to trace over all the other degrees of freedom. This partial trace will be carried out over the states of the so-called ‘‘cavity’’ system $S^{(0,\bar{0})}$. The effective action can be constructed by tracing over scattering processes of all orders that leave the final action diagonal in the system $(0, \bar{0})$. The formal expression can be written as

$$S_{\text{eff}}^{0,\bar{0}} = S_{0,\bar{0}} + \sum_{n=1}^{\infty} \langle (S_{\text{int}}^{\text{hop}})^{2n} \rangle_{(0,\bar{0})} + \sum_{n=1}^{\infty} \langle (S_{\text{int}}^{\text{spin}})^{2n} \rangle_{(0,\bar{0})} \quad (12.5)$$

where, as mentioned before, the average is carried out in the cavity system $S^{(0,\bar{0})}$. Each hopping term in the expansion is of the form

$$\begin{aligned}
 & \left(S_{\text{int}}^{\text{hop}} \right)^{2n} \\
 &= t^{2n} \sum_{\sigma} \sum_{(i_1, i'_1), \dots, (i_n, i'_n)} \sum_{(j_1, j'_1), \dots, (j_n, j'_n)} \int_0^{\beta} d\tau_1 \dots d\tau_n d\tau'_1 \dots d\tau'_{j_n} \\
 & c_{i_1\sigma}^{\dagger}(\tau_{i_1}) c_{i_2\sigma}^{\dagger}(\tau_2) \dots c_{i_n\sigma}^{\dagger}(\tau_n) c_{i'_1\sigma}(\tau'_1) c_{i'_2\sigma}(\tau'_2) \dots c_{i'_n\sigma}(\tau'_n) \left\langle c_{j_1\sigma}(\tau_1) c_{j_2\sigma}(\tau_2) \dots c_{j_n\sigma} c_{j'_1\sigma}^{\dagger}(\tau'_1) \dots c_{j'_n\sigma}^{\dagger}(\tau'_n) \right\rangle_{(0,\bar{0})} \\
 &= t^{2n} \sum_{\sigma} \sum_{(i_1, i'_1), \dots, (i_n, i'_n)} \sum_{(j_1, j'_1), \dots, (j_n, j'_n)} \int_0^{\beta} d\tau_1 \dots d\tau'_{j_n} c_{i_1\sigma}^{\dagger}(\tau_{i_1}) \dots c_{i_n\sigma}^{\dagger}(\tau_n) c_{i'_1\sigma}(\tau'_1) \dots c_{i'_n\sigma}(\tau'_n) \\
 & G_n^{(0,\bar{0})}(\tau_1 \dots \tau_n; \tau'_1 \dots \tau'_n) ,
 \end{aligned} \tag{12.6}$$

where the indices i_1 through i_n and their primed counterparts run through $\bar{0}$, while j_l ($\in \{j_1, \dots, j_n\}$) runs through the nearest-neighbours of the corresponding i_l index. The n -particle Greens function $G_n^{(0,\bar{0})}(\tau_1 \dots \tau_n; \tau'_1 \dots \tau'_n)$ for the cavity system is defined in the usual fashion

$$G_n^{(0,\bar{0})}(\tau_1 \dots \tau_n; \tau'_1 \dots \tau'_n) = \left\langle c_{j_1\sigma}(\tau_1) c_{j_2\sigma}(\tau_2) \dots c_{j_n\sigma} c_{j'_1\sigma}^{\dagger}(\tau'_1) \dots c_{j'_n\sigma}^{\dagger}(\tau'_n) \right\rangle_{(0,\bar{0})} \tag{12.7}$$

In the limit of infinite coordination number, only the $n = 1$ term survives:

$$\begin{aligned}
 \lim_{Z \rightarrow \infty} \sum_{n=1}^{\infty} \left\langle \left(S_{\text{int}}^{\text{hop}} \right)^{2n} \right\rangle_{(0,\bar{0})} &= t^2 \sum_{\sigma} \sum_{i, i'} \int_0^{\beta} d\tau d\tau' c_{i\sigma}^{\dagger}(\tau) c_{i'\sigma}(\tau') \sum_{j, j'} G_1^{(0,\bar{0})}(j\tau; j'\tau') \\
 &= \sum_{\sigma} \sum_{i, i'} \int_0^{\beta} d\tau d\tau' \Delta_{ii',\sigma}(\tau - \tau') c_{i\sigma}^{\dagger}(\tau) c_{i'\sigma}(\tau') ,
 \end{aligned} \tag{12.8}$$

where we have defined the bath hybridisation function $\Delta_{ii',\sigma}(\tau - \tau')$ as

$$\Delta_{ii',\sigma}(\tau - \tau') = t^2 \sum_{j \in \text{NN of } i} \sum_{j' \in \text{NN of } i'} G_1^{(0,\bar{0})}(j\sigma\tau; j'\sigma\tau') . \tag{12.9}$$

Under similar approximations, the spin term in the action reduces to

$$\lim_{Z \rightarrow \infty} \sum_{n=1}^{\infty} \left\langle \left(S_{\text{int}}^{\text{spin}} \right)^{2n} \right\rangle_{(0,\bar{0})} = \sum_{i, i'} \int_0^{\beta} d\tau d\tau' \chi(\tau - \tau') \vec{S}_i(\tau) \cdot \vec{S}_{i'}(\tau') , \tag{12.10}$$

where the susceptibility $\chi_{ii'}(\tau - \tau')$ is defined as

$$\chi_{ii'}(\tau - \tau') = J^2 \sum_{j \in \text{NN of } i} \sum_{j' \in \text{NN of } i'} \vec{S}_j(\tau) \cdot \vec{S}_{j'}(\tau') . \tag{12.11}$$

The effective action for the $0, \bar{0}$ system, in the limit of large coordination number, simplifies to

$$\begin{aligned}
 S_{\text{eff}}^{0, \bar{0}} = & \sum_{i=0, \bar{0}} \sum_{\sigma} \int_0^{\beta} d\tau \, c_{i\sigma}^{\dagger}(\tau) \partial_{\tau} c_{i\sigma}(\tau) - t \sum_{\bar{0}, \sigma} \int_0^{\beta} d\tau \, \left[c_{0\sigma}^{\dagger}(\tau) c_{\bar{0}\sigma}(\tau) + \text{h.c.} \right] \\
 & - \frac{U}{2} \sum_{i=0, \bar{0}} \int_0^{\beta} d\tau \, (n_{i\uparrow}(\tau) - n_{i\downarrow}(\tau))^2 + J \sum_{\bar{0}} \int_0^{\beta} d\tau \, \vec{S}_0(\tau) \cdot \vec{S}_{\bar{0}}(\tau) \\
 & + \sum_{i, i'} \int_0^{\beta} d\tau d\tau' \left[\sum_{\sigma} \Delta_{ii', \sigma}(\tau - \tau') c_{i\sigma}^{\dagger}(\tau) c_{i'\sigma}(\tau') + \chi_{ii'}(\tau - \tau') \vec{S}_i(\tau) \cdot \vec{S}_{i'}(\tau') \right]
 \end{aligned} \tag{12.12}$$

One can also write down an effective action purely for the local site 0. The sites $\bar{0}$ will now be a part of the environment action $S^{(0)}$ instead of the system action, but since the system is thermodynamically large, the cavity action can be assumed to remain the same, such the expectation values are calculated in the same system. As a result, the hybridisation and susceptibility functions remain unchanged. S_{int} is now made of terms that connect 0 and $\bar{0}$, so the transformation from $S^{(0, \bar{0})}$ can be made by replacing the set $\{j\}$ with the set $\bar{0}$, and the set $\bar{0}$ itself will be replaced by the local site 0. We quote the final form of the local effective action:

$$\begin{aligned}
 S_{\text{eff}}^0 = & \sum_{\sigma} \int_0^{\beta} d\tau \, c_{0\sigma}^{\dagger}(\tau) \partial_{\tau} c_{0\sigma}(\tau) - \frac{U}{2} \int_0^{\beta} d\tau \, (n_{0\uparrow}(\tau) - n_{0\downarrow}(\tau))^2 \\
 & + \int_0^{\beta} d\tau d\tau' \left[\sum_{\sigma} \Delta_{00, \sigma}(\tau - \tau') c_{0\sigma}^{\dagger}(\tau) c_{0\sigma}(\tau') + \chi_{00}(\tau - \tau') \vec{S}_0(\tau) \cdot \vec{S}_0(\tau') \right]
 \end{aligned} \tag{12.13}$$

5.12.2 Local theory for the extended SIAM

The Hamiltonian for the extended SIAM model is shown in eq. ???. We will denote the impurity with the label d and the bath sites with the label c . Among the bath sites, we will represent the site coupled to the impurity with z . The full action has the form

$$\begin{aligned}
 S_{\text{ES}} = & \sum_{d, c} \sum_{\sigma} \int_0^{\beta} d\tau \, c_{i\sigma}^{\dagger}(\tau) \partial_{\tau} c_{i\sigma}(\tau) - t \sum_{\langle i, j \rangle \in c, \sigma} \int_0^{\beta} d\tau \, \left[c_{i\sigma}^{\dagger}(\tau) c_{j\sigma}(\tau) + \text{h.c.} \right] \\
 & - V \sum_{\sigma} \int_0^{\beta} d\tau \, \left[c_{d\sigma}^{\dagger}(\tau) c_{z\sigma}(\tau) + \text{h.c.} \right] + J \int_0^{\beta} d\tau \, \vec{S}_d(\tau) \cdot \vec{S}_z(\tau) \\
 & - \frac{U}{2} \int_0^{\beta} d\tau \, (n_{d\uparrow}(\tau) - n_{d\downarrow}(\tau))^2 - \frac{W}{2} \int_0^{\beta} d\tau \, (n_{z\uparrow}(\tau) - n_{z\downarrow}(\tau))^2,
 \end{aligned} \tag{12.14}$$

As in the Hubbard-Heisenberg model, we first obtain an effective action for the pair of sites (d, z) . As in the previous calculation, this will again generate a hybridisation function $\Delta_{zz, \sigma}$ for the bath site nearest to the zeroth site. No susceptibility will be generated however, because there is no

spin-exchange coupling within the bath. The net result is

$$\begin{aligned}
 S_{\text{eff}}^{d,z} = & \sum_{d,c} \sum_{\sigma} \int_0^{\beta} d\tau \, c_{i\sigma}^{\dagger}(\tau) \partial_{\tau} c_{i\sigma}(\tau) - V \sum_{\sigma} \int_0^{\beta} d\tau \, \left[c_{d\sigma}^{\dagger}(\tau) c_{z\sigma}(\tau) + \text{h.c.} \right] + J \int_0^{\beta} d\tau \, \vec{S}_d(\tau) \cdot \vec{S}_z(\tau) \\
 & - \frac{U}{2} \int_0^{\beta} d\tau \, (n_{d\uparrow}(\tau) - n_{d\downarrow}(\tau))^2 - \frac{W}{2} \int_0^{\beta} d\tau \, (n_{z\uparrow}(\tau) - n_{z\downarrow}(\tau))^2 \\
 & + \int_0^{\beta} d\tau d\tau' \sum_{\sigma} \Delta_{zz,\sigma}(\tau - \tau') c_{z\sigma}^{\dagger}(\tau) c_{z\sigma}(\tau')
 \end{aligned} \tag{12.15}$$

This hybridisation $\Delta_{zz,\sigma}$ is calculated in the cavity model of the eSIAM, obtained by removing the impurity and the zeroth site from the full model; it is a completely non-interacting system.

One can go one step further and also remove the zeroth site in order to obtain a theory for the impurity site. With this choice, the cavity model now also includes the zeroth site, and hence contains the correlation term associated with W . The one-particle connection V will lead to a modified hybridisation $\mathcal{F}$ for this effective theory; the Kondo terms will generate a susceptibility. The resultant action is

$$\begin{aligned}
 S_{\text{eff}}^d = & \sum_{\sigma} \int_0^{\beta} d\tau \, c_{d\sigma}^{\dagger}(\tau) \partial_{\tau} c_{d\sigma}(\tau) - \frac{U}{2} \int_0^{\beta} d\tau \, (n_{d\uparrow}(\tau) - n_{d\downarrow}(\tau))^2 \\
 & + \int_0^{\beta} d\tau d\tau' \left[\sum_{\sigma} \mathcal{F}_{\sigma}(\tau - \tau') c_{i\sigma}^{\dagger}(\tau) c_{i'\sigma}(\tau') + \chi_d(\tau - \tau') \vec{S}_i(\tau) \cdot \vec{S}_{i'}(\tau') \right]
 \end{aligned} \tag{12.16}$$

where $\mathcal{F}$ is defined in terms of the local correlator of the zeroth site

$$\mathcal{F}_{\sigma}(\tau - \tau') = V^2 G_1^{(d)}(z\sigma\tau; z\sigma\tau') = V^2 \langle c_{z\sigma}(\tau) c_{z\sigma}^{\dagger}(\tau') \rangle_{(d)} \tag{12.17}$$

with the average computed in the interacting cavity model that has the impurity site removed. The interaction arises from the W -term. The susceptibility is also calculated in this interacting cavity model, and follows the same definition as in the bulk model:

$$\chi_d(\tau - \tau') = J^2 \vec{S}_z(\tau) \cdot \vec{S}_z(\tau') . \tag{12.18}$$

Chapter 6

Competing tendencies In Heavy Fermion Systems: Kondo screening vs. Mott localisation

6.1 Bilayer extended Hubbard model: Impurity model

[Mukherjee et al.(2023)] Mukherjee, Vidhyadhiraja, Taraphder, and Lal] We approach the heavy-fermion problem by starting from a bilayer extended Hubbard model, consisting of two layers (f and c). Towards studying this lattice model, we adopt a two-layer impurity problem that hosts a correlated impurity site in each layer (S_f and S_d):

$$H_{\text{aux}} = H_{1\text{p}} + H_f + H_d + H_{fd} , \quad (1.1)$$

where $H_{1\text{p}}$ is the Hamiltonian for the non-interacting itinerant electrons of either layer,

$$H_{1\text{p}} = -t \sum_{\sigma, \alpha} \sum_{\langle i, j \rangle} \left(c_{i, \sigma, \alpha}^\dagger c_{j, \sigma, \alpha} + \text{h.c.} \right) - \sum_{i, \sigma, \alpha} \mu_\alpha n_{i, \sigma, \alpha} , \quad (1.2)$$

such that α sums over the two layers f and d . H_f and H_d describe the dynamics of the correlated impurity sites (and their local neighbourhood) in each layer:

$$\begin{aligned} H_f = & -\frac{U_f}{2} (n_{f, \uparrow} - n_{f, \downarrow})^2 + \sum_{Z \in \text{NN}} \left[-V \sum_{\sigma} \left(c_{f, \sigma}^\dagger c_{Z, \sigma, f} + \text{h.c.} \right) + \frac{J}{2} \sum_{\alpha, \beta} \mathbf{S}_f \cdot \boldsymbol{\sigma}_{\alpha\beta} c_{Z, \alpha, f}^\dagger c_{Z, \beta, f} \right. \\ & \left. - \frac{W_f}{2} (n_{Z, \uparrow, f} - n_{Z, \downarrow, f})^2 \right] , \\ H_d = & -\eta_d n_d - \frac{U_d}{2} (n_{d, \uparrow} - n_{d, \downarrow})^2 + \sum_{Z \in \text{NN}} \left[-V \sum_{\sigma} \left(c_{d, \sigma}^\dagger c_{Z, \sigma, d} + \text{h.c.} \right) + \frac{J}{2} \sum_{\alpha, \beta} \mathbf{S}_d \cdot \boldsymbol{\sigma}_{\alpha\beta} c_{Z, \alpha, d}^\dagger c_{Z, \beta, d} \right] , \end{aligned} \quad (1.3)$$

where $c_{\nu,\sigma,\alpha}^\dagger$ can refer to creation operator for either the f -layer. Finally, H_{fd} represents the inter-layer hybridisation:

$$H_{fc} = -V_\perp \sum_{\sigma} \left(c_{d,\sigma}^\dagger c_{f,\sigma} + \text{h.c.} \right). \quad (1.4)$$

6.2 Tiling Into Bilayer Extended Hubbard Model

Tiling the impurity model leads to a bilayer extended Hubbard model on the 2D square lattice. In order to tile our auxiliary model to restore translation invariance, we identify the cluster which acts as the “unit” cell of translation. As discussed in the text surrounding eq. 3.9, a good choice for the unit cell is the impurity site and its surrounding correlated environment. Placing the impurity sites at $\mathbf{r}_d$ in each layer, we get

$$H_{\text{clust}}(\mathbf{r}_d) = H_f(\mathbf{r}_d) + H_d(\mathbf{r}_d) + H_{fc}(\mathbf{r}_d). \quad (2.1)$$

The details of tiling a cluster Hamiltonian were already shown in Sec. 4.3.2; the general idea is to use manybody translation operators (details in Sec. 4.3.1) to center the cluster Hamiltonian at various sites of the lattice, such that combining all such terms leads to a periodised Hamiltonian. we write down the final result here:

$$\begin{aligned} H_{\text{latt}} &= \sum_{\mathbf{r}} T^\dagger(\mathbf{r}) H_{\text{clust}}(\mathbf{r}_d) T(\mathbf{r}) \\ &= -\frac{2V}{Z} \sum_{\langle i,j \rangle, \sigma} \left(c_{i,\sigma}^\dagger c_{j,\sigma} + \text{h.c.} \right) - \sum_i \left[\eta_d n_{i,d} + \frac{U_d}{2} (n_{i,\uparrow,d} - n_{i,\downarrow,d})^2 \right] + \frac{2J}{Z} \sum_{\langle i,j \rangle} \mathbf{S}_{i,d} \cdot \mathbf{S}_{j,d} \\ &\quad - \frac{2V}{Z} \sum_{\langle i,j \rangle, \sigma} \left(f_{i,\sigma}^\dagger f_{j,\sigma} + \text{h.c.} \right) - \frac{U_f - W_f}{2} \sum_i (n_{i,\uparrow,f} - n_{i,\downarrow,f})^2 + \frac{2J}{Z} \sum_{\langle i,j \rangle} \mathbf{S}_{i,f} \cdot \mathbf{S}_{j,f} \\ &\quad - V_\perp \sum_{i,\sigma} (f_{i,\sigma}^\dagger c_{i,\sigma} + \text{h.c.}) \end{aligned} \quad (2.2)$$

Tiling the impurity model therefore leads to a bilayer extended Hubbard model, with effective couplings that depend on the auxiliary model couplings. The value of η_d defines the effective chemical potential of the d -layer; a non-zero value moves the layer away from half-filling. The f -layer is pinned at half-filling.

Following Sec. 4.5.1, the retarded k -space Greens function in either layer can be expressed in terms of auxiliary model Greens functions:

$$G(\mathbf{k}, \sigma, \alpha; \omega) = \sum_{\nu} C_{\nu, \mathbf{k} \alpha} \mathcal{G}(\mathcal{T}_{\nu, \sigma, \alpha}; \omega - \mathcal{E}_{\nu}(\mathbf{k})) \quad (2.3)$$

where $\alpha = f, d$ denotes the layer in which the Greens function is being calculated, the modes ν indicate the operators which diagonalise the Hamiltonian H_{1p} , with energies $\mathcal{E}_{\nu}(\mathbf{k})$:

$$H_{1p} = -\frac{2V}{Z} \sum_{\langle i,j \rangle, \sigma} \left(c_{i,\sigma}^\dagger c_{j,\sigma} + \text{h.c.} + f_{i,\sigma}^\dagger f_{j,\sigma} + \text{h.c.} \right) - V_\perp \sum_{i,\sigma} (f_{i,\sigma}^\dagger c_{i,\sigma} + \text{h.c.}) - \sum_i \eta_d n_{i,d}, \quad (2.4)$$

and the $\mathcal{T}$ –matrices are defined in eq. 5.11:

$$\mathcal{T}_{\nu,\sigma,\alpha} = \left[\left(\sum_{\sigma'} c_{\alpha\sigma}^\dagger + \text{h.c.} \right) + (S_\alpha^+ + \text{h.c.}) + (C_\alpha^+ + \text{h.c.}) \right] c_{\nu,\sigma}, \quad (2.5)$$

In order to calculate $\mathcal{E}_\nu(\mathbf{k})$, we introduce the k –space modes via a Fourier transform:

$$c_{i,\sigma}^\dagger = \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} e^{-i\mathbf{k} \cdot \mathbf{r}_i} c_{\mathbf{k},\sigma}^\dagger, \quad f_{i,\sigma}^\dagger = \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} e^{-i\mathbf{k} \cdot \mathbf{r}_i} f_{\mathbf{k},\sigma}^\dagger. \quad (2.6)$$

In terms of these modes, the 1–particle Hamiltonian decouples in k –space:

$$H_{1p} = \sum_{\mathbf{k},\sigma} \left[(\epsilon_{\mathbf{k}} - \eta_d)(n_{\mathbf{k},\sigma,d} + n_{\mathbf{k},\sigma,f}) - V_\perp (c_{\mathbf{k},\sigma}^\dagger f_{\mathbf{k},\sigma} + \text{h.c.}) \right]. \quad (2.7)$$

The diagonal modes are given by

$$\nu_{\mathbf{k},\sigma,\pm}^\dagger = C_{\pm,\mathbf{k}} c_{\mathbf{k},\sigma}^\dagger + \sqrt{1 - C_{\pm,\mathbf{k}}^2} f_{\mathbf{k},\sigma}^\dagger, \quad (2.8)$$

the coefficients obtained from diagonalising the matrix

$$H_{1p}(\mathbf{k}) = \begin{pmatrix} \epsilon_{\mathbf{k}} & -V_\perp \\ -V_\perp & \epsilon_{\mathbf{k}} - \eta_d \end{pmatrix}. \quad (2.9)$$

The energies are given by $\mathcal{E}_\pm(\mathbf{k}) = \epsilon_{\mathbf{k}} - \eta_d/2 \pm \Delta$, with $\epsilon_{\mathbf{k}}$ being $-\frac{4V}{Z}(\cos k_x + \cos k_y)$ for a 2D square lattice and $\Delta = \sqrt{V_\perp^2 + \eta_d^2/4}$. Substituting into the expression for the lattice Greens function, we get

$$\begin{aligned} G(\mathbf{k}, d; \omega) = & C_{+,\mathbf{k}} \mathcal{G}(\mathcal{T}_{\mathbf{k}+,d}, \mathcal{T}_{\mathbf{k},d}^\dagger; \omega - \mathcal{E}_+(\mathbf{k})) + C_{-,\mathbf{k}} \mathcal{G}(\mathcal{T}_{\mathbf{k}-,d}, \mathcal{T}_{\mathbf{k},d}^\dagger; \omega - \mathcal{E}_-(\mathbf{k})) \\ & + C_{+,\mathbf{k}}^* \mathcal{G}(\mathcal{T}_{\mathbf{k},d}, \mathcal{T}_{\mathbf{k}+,d}^\dagger; \omega - \mathcal{E}_+(\mathbf{k})) + C_{-,\mathbf{k}}^* \mathcal{G}(\mathcal{T}_{\mathbf{k},d}, \mathcal{T}_{\mathbf{k}-,d}^\dagger; \omega - \mathcal{E}_-(\mathbf{k})). \end{aligned} \quad (2.10)$$

Following eq. 5.33, we can similarly write down an expression the local lattice Greens function for the f – and d –electrons. For that, we define the local diagonal modes $\nu_\pm$ obtained from the local Hamiltonian

$$H_{\text{loc}} = \begin{pmatrix} 0 & -V_\perp \\ -V_\perp & 0 - \eta_d \end{pmatrix}, \quad (2.11)$$

and their associated energies $\varepsilon_\pm$. Within the impurity model, these local bonding/antibonding modes can be constructed on the impurity sites ($\nu_\pm$) or on the neighbouring bath sites, ($\nu_\pm^Z$). The local Greens function can be expressed in terms of these modes:

$$\begin{aligned} G_{dd}(\omega) = & \frac{1}{2} \sum_{\nu} C_{\nu,d} [\mathcal{G}(\nu\sigma, d\sigma; \omega - \varepsilon_\nu) + \mathcal{G}(d\sigma, \nu\sigma; \omega - \varepsilon_\nu)] \\ & + \frac{1}{2} \sum_{\nu^Z} C_{\nu^Z,d} [\mathcal{G}(\mathcal{T}_{Zd,\sigma}, \mathcal{T}_{\nu^Z,\sigma}^\dagger; \omega - \varepsilon_{\nu^Z}) + \mathcal{G}(\mathcal{T}_{\nu^Z,\sigma}, \mathcal{T}_{Zd,\sigma}^\dagger; \omega - \varepsilon_{\nu^Z})]. \end{aligned} \quad (2.12)$$

The presence of local hybridisation between the d - and f - orbitals makes the inter-orbital Greens function important as well. Following eq. 5.35, we have

$$G_{fd}(\omega) = \frac{1}{2} \sum_{\nu} [C_{\nu,f} \mathcal{G}(\nu\sigma, d\sigma; \omega - \varepsilon_{\nu}) + C_{\nu,d} \mathcal{G}(f\sigma, \nu\sigma; \omega - \varepsilon_{\nu})] + \frac{1}{2} \sum_{\nu Z} [C_{\nu Z, Z_f} \mathcal{G}(\mathcal{T}_{\nu Z, \sigma}, \mathcal{T}_{Z_d, \sigma}; \omega - \varepsilon_{\nu Z}) + C_{\nu Z, Z_d} \mathcal{G}(\mathcal{T}_{Z_f, \sigma}, \mathcal{T}_{\nu Z, \sigma}; \omega - \varepsilon_{\nu Z})] . \quad (2.13)$$

Using these, we can construct the hybridised Greens function associated with the states ν :

$$G_{\nu+}(\omega) = C_{+,d}^2 G_{dd}(\omega) + C_{+,f}^2 G_{ff}(\omega) + C_{+,d} C_{+,f} [G_{df}(\omega) + G_{fd}(\omega)] , \quad (2.14)$$

6.3 Unitary RG Analysis of The Bilayer Lattice-Embedded ESIAM

In the limit of large U_{α} , we carry out a Schrieffer-Wolff transformation and work with the following low-energy Hamiltonian:

$$H_{\text{aux}} = \sum_{\mathbf{k}, \sigma, \alpha} \epsilon_{\mathbf{k}, \alpha} n_{\mathbf{k}, \sigma, \alpha} + \sum_{\alpha} \sum_{Z \in \text{NN}} \left[\frac{1}{2} J_{\alpha} \sum_{\sigma, \sigma'} \mathbf{S}_{\alpha} \cdot \boldsymbol{\sigma}_{\alpha\beta} c_{Z, \sigma, \alpha}^{\dagger} c_{Z, \sigma', \alpha} - \frac{W_{\alpha}}{2} (n_{Z, \uparrow, \alpha} - n_{Z, \downarrow, \alpha})^2 \right] + J \mathbf{S}_f \cdot \mathbf{S}_d . \quad (3.1)$$

The Kondo coupling and bath interaction acquire k -dependence:

$$J(\mathbf{k}, \mathbf{q}) = \frac{J}{2} [\cos(\mathbf{k}_x - \mathbf{q}_x) + \cos(\mathbf{k}_y - \mathbf{q}_y)] , \quad (3.2)$$

$$W(\mathbf{k}, \mathbf{q}, \mathbf{k}', \mathbf{q}') = \frac{W}{2} [\cos(\mathbf{k}_x - \mathbf{q}_x + \mathbf{k}'_x - \mathbf{q}'_x) + \cos(\mathbf{k}_y - \mathbf{q}_y + \mathbf{k}'_y - \mathbf{q}'_y)] .$$

The particle-hole transformed state $\bar{\mathbf{q}}$ associated with $\mathbf{q}$ is defined as $\bar{\mathbf{q}} = \mathbf{q} + (\pi, \pi)$ if $\mathbf{q}$ is in the first or third quadrant and $\mathbf{q} + (\pi, -\pi)$ is in the second or fourth quadrant.

In order to study the low-energy physics of the impurity model, we iteratively integrate out high-energy degrees of freedom using the unitary RG method. Let the momentum states being decoupled from the two layers be $|\mathbf{q}, \sigma, f\rangle$ and $|\mathbf{q}, \sigma, d\rangle$ if they are above the Fermi energy (unoccupied in the low-energy configuration) and $|\bar{\mathbf{q}}, \sigma, f\rangle$ and $|\bar{\mathbf{q}}, \sigma, d\rangle$ if they are below the Fermi energy (occupied in the low-energy configuration) For every shell of conduction bath states of width ΔD integrated out, the Hamiltonian couplings renormalise by folding in the effects of virtual excitations to these states. We already have the renormalisation group equations for the couplings J_{α} in the case of $J = 0$:

$$\frac{\Delta J_f(\mathbf{k}_1, \mathbf{k}_2)}{\Delta D} = - \int_{UV} d\mathbf{q} \rho(\mathbf{q}) [J_f(\mathbf{k}_1, \mathbf{q}) J_f(\mathbf{q}, \mathbf{k}_2) + 4 J_f(\mathbf{q}, \bar{\mathbf{q}}) W_f(\mathbf{k}_1, \mathbf{q}, \bar{\mathbf{q}}, \mathbf{k}_2)] G_f^{(0)}(\mathbf{q}) , \quad (3.3)$$

where $\mathbf{q}$ sums over the UV states being integrated out, $\rho(\mathbf{q})$ is the density of states for the momentum states being integrated out and $G_f^{(0)}$ is the propagator for the excited intermediate state (for $J = 0$):

$$G_f^{(0)}(\mathbf{q}) = \frac{1}{\omega - |\epsilon_f(\mathbf{q})|/2 + J_f(\mathbf{q})/4 + W_f(\mathbf{q})/2} . \quad (3.4)$$

An identical equations holds for ΔJ_d , with all f -quantities replaced with d -quantities. Switching on J modifies the excitation energy in the propagator:

$$G_f(\mathbf{q}) = \frac{1}{1/G_f^{(0)} - J\vec{S}_f \cdot \vec{S}_d}. \quad (3.5)$$

This expression can be simplified by noting that the operator in the denominator has only two eigenvalues, $-3/4$ in the singlet sector and $1/4$ in the triplet sector. Defining the projectors $\mathcal{P}_0$ and $\mathcal{P}_1$ for the two sectors, we have

$$\vec{S}_f \cdot \vec{S}_d = \frac{-3}{4}\mathcal{P}_0 + \frac{1}{4}\mathcal{P}_1, \quad \mathcal{P}_0 + \mathcal{P}_1 = \mathbb{I}. \quad (3.6)$$

Using these relations, we can write

$$\begin{aligned} G_f(\mathbf{q}) &= \frac{1}{1/G_f^{(0)} + 3J/4}\mathcal{P}_0 + \frac{1}{1/G_f^{(0)} - J/4}\mathcal{P}_1 \\ &= \left(\frac{1/4}{1/G_f^{(0)} + 3J/4} + \frac{3/4}{1/G_f^{(0)} - J/4} \right) \mathbb{I} + \left(\frac{1}{1/G_f^{(0)} + 3J/4} - \frac{1}{1/G_f^{(0)} - J/4} \right) \vec{S}_f \cdot \vec{S}_d. \end{aligned} \quad (3.7)$$

Noting that only the first operator renormalises the Kondo interaction, we get the following modified RG equation for the Kondo coupling J_α :

$$\begin{aligned} \frac{\Delta J_\alpha}{\Delta D} &= \int_{UV} d\mathbf{q} \rho(\mathbf{q}) [J_\alpha(\mathbf{k}_1, \mathbf{q})J_\alpha(\mathbf{q}, \mathbf{k}_2) + J_\alpha(\mathbf{q}, \bar{\mathbf{q}})W_\alpha(\mathbf{k}_1, \mathbf{q}, \bar{\mathbf{q}}, \mathbf{k}_2)] \left(\frac{1}{4} \frac{1}{1/G_\alpha^{(0)}(\mathbf{q}) + 3J/4} \right. \\ &\quad \left. + \frac{3}{4} \frac{1}{1/G_\alpha^{(0)}(\mathbf{q}) - J/4} \right), \end{aligned} \quad (3.8)$$

We now turn to the renormalisation of J , arising from hybridisation of f - and d -layers. In order to capture coherent scattering processes between the two layers, we project onto the following rotated basis of excited states:

$$\begin{aligned} |H\rangle_\pm &= \frac{1}{\sqrt{2}} \left(|H(\mathbf{q})\rangle_f \pm |H(\mathbf{q})\rangle_d \right), \\ \mathcal{P}(\mathbf{q})_H &= |H\rangle_+ \langle H|_+ + |H\rangle_- \langle H|_- , \end{aligned} \quad (3.9)$$

where $|H(\mathbf{q})\rangle_\alpha$ is the state obtained upon exciting both states in the layer α : $|H(\mathbf{q})\rangle_\alpha = c_{\mathbf{q},\alpha}^\dagger c_{\bar{\mathbf{q}},\alpha} |L\rangle_f |L\rangle_d$, and $\mathcal{P}(\mathbf{q})_H$ projects onto this rotated excited basis. We focus on the spin-flip component $J S_f^+ S_d^-$ of the Hamiltonian. There are two kinds of processes that renormalise this coupling. We first consider one that involves a spin-flip of the d -layer followed by a spin-flip of the f -layer:

$$\Delta H = \frac{1}{N^2} \sum_{\mathbf{q} \in UV} \langle L | \frac{1}{2} J_f(\bar{\mathbf{q}}, \mathbf{q}) S_f^+ c_{\mathbf{q}-\downarrow, f}^\dagger c_{\mathbf{q}+\uparrow, f} \mathcal{P}(\mathbf{q})_H G \mathcal{P}(\mathbf{q})_H \frac{1}{2} J_d(\mathbf{q}, \bar{\mathbf{q}}) S_d^- c_{\mathbf{q}\uparrow, d}^\dagger c_{\bar{\mathbf{q}}\downarrow, d} | L \rangle, \quad (3.10)$$

where $|L\rangle = |L\rangle_f |L\rangle_d$ is the initial configuration for both layers. G is the propagator for the excited state:

$$G = \frac{1}{\omega - H_D} , \quad (3.11)$$

where H_D is the diagonal part of the Hamiltonian corresponding to the excited(intermediate) state. The diagonal part consists of the kinetic energy and the Ising part of the Kondo interaction:

$$H_D = \sum_{\mathbf{q}, \sigma \in \mathbf{q}_{\pm}, \alpha} \epsilon_{\alpha}(\mathbf{q}) \tau_{\mathbf{q}, \sigma} + \frac{1}{2} \sum_{\alpha} J_{\alpha} S_{\alpha}^z \sum_{\mathbf{q}, \sigma} \sigma n_{\mathbf{q}, \sigma, \alpha} + J \vec{S}_f \cdot \vec{S}_d . \quad (3.12)$$

Calculating the matrix elements of the propagator in the rotated basis gives

$$\mathcal{P}(\mathbf{q})_H G \mathcal{P}(\mathbf{q})_H = \frac{1}{2} (G_f + G_d) (|H\rangle_+ \langle H|_+ + |H\rangle_- \langle H|_-) + \frac{1}{2} (G_f - G_d) (|H\rangle_+ \langle H|_- + |H\rangle_- \langle H|_+) , \quad (3.13)$$

where

$$G_{\alpha}(\mathbf{q}) = \frac{1}{\omega - \frac{1}{2} [\epsilon_{\alpha}(\mathbf{q}) - \epsilon_{\alpha}(\bar{\mathbf{q}})] + \frac{1}{2} J_{\alpha}(\mathbf{q}) + W_{\alpha}(\mathbf{q}) - \frac{1}{4} J} . \quad (3.14)$$

In eq. 3.10, since the first process is an excitation into a d -state and the second process is a de-excitation from an f -state, the expression can be simplified into

$$\begin{aligned} \langle L | S_f^+ c_{\mathbf{q}-\downarrow, f}^{\dagger} c_{\mathbf{q}+\uparrow, f} \mathcal{P}(\mathbf{q})_H G \mathcal{P}(\mathbf{q})_H S_d^- c_{\mathbf{q}+\uparrow, d}^{\dagger} c_{\mathbf{q}-\downarrow, d} | L \rangle &= S_f^+ S_d^- \langle H_f | \mathcal{P}(\mathbf{q})_H G \mathcal{P}(\mathbf{q})_H | H_d \rangle \\ &= \frac{1}{2} (G_f + G_d) S_f^+ S_d^- . \end{aligned} \quad (3.15)$$

The net Hamiltonian renormalisation is finally a sum over the contribution from each mode:

$$\Delta H = \frac{1}{8N^2} \sum_{\mathbf{q} \in UV} J_f(\bar{\mathbf{q}}, \mathbf{q}) J_d(\mathbf{q}, \bar{\mathbf{q}}) (G_f(\mathbf{q}) + G_d(\mathbf{q})) S_f^+ S_d^- . \quad (3.16)$$

Converting the sums to integrals (assuming a density of states $\rho(\mathbf{q})$) gives the final expression

$$\frac{\Delta H}{\Delta D} = \frac{1}{8N} S_f^+ S_d^- \int_{UV} d\mathbf{q} \rho(\mathbf{q}) J_f(\bar{\mathbf{q}}, \mathbf{q}) J_d(\mathbf{q}, \bar{\mathbf{q}}) (G_f(\mathbf{q}) + G_d(\mathbf{q})) . \quad (3.17)$$

The time-reversed of this process can be obtained simply by exchanging the d -interaction with the f -interaction. Since the renormalisation itself is symmetric, the contribution is equal to what we obtained here. The total renormalisation in the coupling for the term $S_f^+ S_d^-$ is therefore

$$\begin{aligned} \frac{\Delta J}{\Delta D} &= \frac{1}{2} \int_{UV} d\mathbf{q} \rho(\mathbf{q}) J_f(\bar{\mathbf{q}}, \mathbf{q}) J_d(\mathbf{q}, \bar{\mathbf{q}}) (G_f(\mathbf{q}) + G_d(\mathbf{q})) \\ &= \frac{1}{2} \int_{UV} d\mathbf{q} \rho(\mathbf{q}) \left[\frac{J_f(\bar{\mathbf{q}}, \mathbf{q}) J_d(\mathbf{q}, \bar{\mathbf{q}})}{\omega - \frac{1}{2} \bar{\epsilon}_f(\mathbf{q}) + \frac{1}{2} J_f(\mathbf{q}) + W_f(\mathbf{q}) - \frac{1}{4} J} + \frac{J_f(\bar{\mathbf{q}}, \mathbf{q}) J_d(\mathbf{q}, \bar{\mathbf{q}})}{\omega - \frac{1}{2} \bar{\epsilon}_d(\mathbf{q}) + \frac{1}{2} J_d(\mathbf{q}) + W_d(\mathbf{q}) - \frac{1}{4} J} \right] , \end{aligned} \quad (3.18)$$

where $\bar{\epsilon}_{\alpha}(\mathbf{q}) = \epsilon_f(\mathbf{q}) - \epsilon_f(\bar{\mathbf{q}})$ is the total excitation energy for the pair of states $\mathbf{q}, \bar{\mathbf{q}}$.




Figure 6.1: RG phase diagrams for various Kondo couplings: f -layer (left), d -layer (middle) and inter-layer (right). For each plot, the background color (quantified by the right colorbar) tracks the fixed point value of the coupling as a function of the inter-layer couplings $J_{\perp}$ and the f -layer bath interaction $|W_f|$. The marker color complements this by tracking whether the coupling is relevant, intermediate-coupling or irrelevant in the RG sense, by taking the values 1, 1/2 or 0 respectively.

Since the bath interaction doesn't renormalise, its functional form remains unchanged and can be substituted into the RG equations. In fact, we can write the expressions in a compact matrix form for efficient computation:

$$\frac{\Delta J_{\alpha}}{\Delta D} = -J_{\alpha} \mathcal{G}_{\alpha} J_{\alpha}^{\dagger} - 4\mathcal{W}_{\alpha} \text{Tr} [\mathcal{G}'_{\alpha}] , \quad (3.19)$$

where all objects are matrices. J_{α} is the Kondo coupling matrix. $\mathcal{G}_{\alpha}$ is a diagonal matrix with matrix elements

$$\mathcal{G}_{\alpha}[q, q] = \begin{cases} \rho(q) \left(\frac{1}{8} \frac{1}{1/G_{\alpha}^{(0)}(\mathbf{q}) + \mu/2 + 3J/4} + \frac{3}{8} \frac{1}{1/G_{\alpha}^{(0)}(\mathbf{q}) + \mu/2 - J/4} \right), & \mathbf{q} \in \text{PS} , \\ \rho(q) \left(\frac{1}{8} \frac{1}{1/G_{\alpha}^{(0)}(\mathbf{q}) - \mu/2 + 3J/4} + \frac{3}{8} \frac{1}{1/G_{\alpha}^{(0)}(\mathbf{q}) - \mu/2 - J/4} \right), & \mathbf{q} \in \text{HS} , \end{cases} \quad (3.20)$$

if q is in the UV subspace and zero otherwise. $\mathcal{W}_{\alpha}$ is the bath interaction matrix whose matrix elements are given by $\mathcal{W}_{\alpha}[k, k'] = \frac{W}{2} [\cos(k_x - k'_x) + \cos(k_y - k'_y)]$. Finally, $\mathcal{G}'_{\alpha}$ is another diagonal matrix whose matrix elements are defined as $\mathcal{G}'_{\alpha}[q, q] = \mathcal{G}[q, q] J_{\alpha}[q, \bar{q}]$.

The RG equation for the inter-layer coupling J can be written as

$$\frac{\Delta J}{\Delta D} = \frac{1}{2} \mathbf{J} \cdot \mathbf{G} , \quad (3.21)$$

where $\mathbf{J}$ and $\mathbf{G}$ are vectors defined by the elements $\mathbf{J}[q] = J_f(\bar{\mathbf{q}}, \mathbf{q}) J_d(\mathbf{q}, \bar{\mathbf{q}})$ and

$$\mathbf{G}[q] = \rho(\mathbf{q}) \left[\frac{1}{\omega - \frac{1}{2}\bar{\epsilon}_f(\mathbf{q}) + \frac{1}{2}J_f(\mathbf{q}) + W_f(\mathbf{q}) - \frac{1}{4}J} + \frac{1}{\omega - \frac{1}{2}\bar{\epsilon}_d(\mathbf{q}) + \frac{1}{2}J_d(\mathbf{q}) + W_d(\mathbf{q}) - \frac{1}{4}J} \right] . \quad (3.22)$$

6.4 Renormalisation Group Flows of Kondo Couplings: RG Phase Diagram

We first analyse the model by obtaining unitary RG flows of three kinds of Kondo couplings. Fig. 6.1 shows the fixed point values of the couplings under renormalisation to a low-energy theory. We view

these values as a function of the f -layer bath interaction W_f and inter-layer Kondo coupling $J_\perp$. In each panel, the right (left) colorbar quantifies the background (marker) color of the plot. The background displays the fixed point value of each coupling whereas the marker color tracks whether the couplings is relevant (1), intermediate coupling (0.5) or irrelevant (0).

The Kondo coupling J_d (middle panel) of the d -layer remains relevant for the entirety of the chosen parameter regime. The Kondo coupling J_f (left panel) of the f -layer becomes RG-irrelevant for larger values of $|W_f|$; this leads to an in-plane Mott transition of the kind observed in our earlier work, where the local Fermi liquid excitations of the f -impurity give way to a pseudogapped non-Fermi liquid Mott metal phase which ultimately destabilises into a local moment phase. The RG flow of J_f does not appear to be significantly affected by the variation of $J_\perp$. Concomitant with the irrelevance of J_f under increasing $|W_f|$, the inter-plane coupling $J_\perp$ (right panel) becomes RG-relevant, as seen from the darkening markers as we go to higher $|W_f|$.

These simulations motivate an RG-based phase diagram with the following regimes: (i) weak $J_\perp$, weak $|W_f|$: decoupled metallic layers, each layer participating in transport independently, (ii) strong $J_\perp$, strong $|W_f|$: Kondo insulator regime, where the local moments of the f -layer hybridise with the itinerant electrons of the d -layer to form a band insulator with the chemical potential situated in the hybridisation gap, and (iii) weak $J_\perp$, strong $|W_f|$: the so-called RKKY regime, where the f -layer forms local moments which decouple from the d -layer and are then free to order magnetically, and (iv) strong $J_\perp$, weak $|W_f|$: this is similar to the Kondo insulator regime, except here the f -electrons remain itinerant and it is they who hybridise with the d -electrons to again form two bands separated by a hybridisation gap.

6.5 Results at half-filling




Figure 6.2: Left: Real-space spin correlations for the auxiliary model. Background colour (right colourbar) represents that between the f -impurity and its nearest-neighbour sites, while the marker colour (left colourbar) represents that between the f -impurity and the d -impurity. While the former shrinks as $J_\perp$ or $|W_f|$ is increased, the latter grows. This marks a transfer of entanglement from intra-plane channel into inter-plane channel. Right:




Figure 6.3: Spectral functions of the tiled model, for excitations in the bonding (two on the left) and antibonding (two on the right) basis, defined by the Greens function $G_{k,\pm} = G(k_d \pm k_f; \omega)$. There is no hybridisation for $J_\perp = 0$ (first and third plots), and the only effect of W_f is to shrink the height of the central resonance. Introducing a non-zero $J_\perp$ splits the central resonance into an upper band (second from left) and a lower band (first from right). This leads to a Kondo insulator, where $\omega = 0$ resides inside the hybridisation gap.




Figure 6.4: Spectral functions A_d and A_f of the auxiliary model, for excitations in the d - and f -layers respectively. Both show a central Kondo resonance for $J_\perp = 0$ which splits into two bands upon adding a non-zero $J_\perp$. Increasing $|W_f|$ to large values gaps out the local Fermi liquid excitations of A_f .

Chapter 7

Holography of Entanglement in 2D Free Fermions: Emergent Geometry and Fermionic Criticality

7.1 Introduction

The manuscript is structured in the following manner. Sec. 7.2 describes the system we work with and defines certain physical ideas and measures of entanglement that are employed in later sections. In Sec. 7.3, we define the scaling transformations mentioned earlier, and discuss its implications on the scaling of entanglement in the system under study. In Sections 7.4, 7.5 and 7.6, we probe the holographic nature of the entanglement, unveiling an emergent spatial dimension in the process and obtain several characteristics of this emergent geometry. Section 7.7 discusses the consequences of a critical Fermi surface on the holographic space, and how the emergent curvature can be linked to a convergence parameter associated with the RG flows. In Section 7.9, we probe the topological aspects of the entanglement, and discuss its implications for the system at hand. We conclude with a discussion of our results in Sec. 7.10 and point out some open questions.

Summary of our main results

For the convenience of the reader, we present below a summary of the main results obtained by us.

1. Quantum information measures (such as entanglement entropy and multipartite information) reveal a hierarchical structure in the nature of entanglement within the system. We relate geometric parameters like distance and curvature of the holographic dimension to an RG beta function of the mass gap, providing thereby an explicit analytic manifestation of the holographic principle. We show that a change in the boundedness of the space corresponds to a change in certain topological winding numbers.

2. We go on to argue that this topological transition and the underlying critical Fermi surface coincides with the formation of a quantum wormhole geometry that connects the UV and the IR of the emergent dimension. The additional (conformal) symmetry at the transition links the emergent metric and the stress-energy tensor.




Figure 7.1: The model on the left consists of a system of non-interacting electrons placed on a torus with a flux threading through it. The model on the right consists of a sum of $1 + 1$ -dimensional systems, each consisting of massive electrons placed on a ring; the $1 + 1$ -D systems are decoupled from each other. The mass arises from the flux. These two models can be shown to be equivalent to each other [Chung and Peschel(2000), Arias et al.(2015)Arias, Blanco, and Casini, Chen et al.(2017)Chen, Witczak-Krempa, Faulkner, and Fradkin, Murciano et al.(2020)Murciano, Ruggiero, and Calabrese].

3. We show that the geometry-independent part of the entanglement is invariant under appropriate scaling transformations for a system of fixed number density, and is related to the Luttinger volume of the gapless fermionic system. In the presence of a strong transverse magnetic field, the topological Luttinger volume in the metallic state transforms into the Chern number $\mathcal{C}$ of the insulating integer quantum Hall state.

7.2 Preliminaries and definitions

7.2.1 The system

In this work, we focus on the case of Dirac fermions in two spatial dimensions described by the Lagrangian (in natural units $\hbar = c = 1$) $\mathcal{L} = \int dx dy \bar{\psi}(x, y, t) (i\gamma^\mu \partial_\mu - m) \psi(x, y, t)$. The system is placed on a 2-torus of dimensions $L_x \times L_y$ with periodic boundary conditions, such that $\psi(x, y, t) = \psi(x + L_x, y, t) = \psi(x, y + L_y, t)$. Entanglement measures will be calculated in the presence of a gauge field A that transforms the Lagrangian by shifting the momenta: $i\partial_\mu \rightarrow i\partial_\mu + eA$. We choose the vector potential A such that, in any particular setup, it couples to just one component of the momentum (say, the x -component). This results in $\phi = eAL_x/2\pi$ units of flux quantum threading the torus in the y -direction. One of our goals is to express the Luttinger volume V_L in terms of multi-partite measures of entanglement $\left(I_{\{\mathcal{A}_i\}}^g(\phi)\right)$ between multiple subsystems $\{\mathcal{A}_i, i = 1, 2, \dots, g\}$ [Patra et al.(2022)Patra, Basu, and Lal], in the presence of the flux ϕ . The precise definition of I^g and the method of choosing the subsystems will be given later. In order to calculate $I_{\{\mathcal{A}_i\}}^g$, however, we find it convenient to take the thin torus limit: $(L_y/L_x) \rightarrow \infty$, keeping

l/L_y fixed. Importantly, this operation does not affect quantities such as V_L as long as we keep the number density unchanged.

In order to calculate entanglement measures, we will express the $2 + 1$ -dimensional system as a sum of an infinite number of massive $1 + 1$ -dimensional systems [Chung and Peschel(2000), Arias et al.(2015)Arias, Blanco, and Casini, Chen et al.(2017)Chen, Witczak-Krempa, Faulkner, and Fradkin, Murciano et al.(2020)Murciano, Ruggiero, and Calabrese]. In the presence of a gauge field $\vec{A} = A\hat{x}$, the complete Lagrangian (in natural units) is

$$\mathcal{L} = \int dx dy \bar{\psi}(x, y) [\gamma^\mu (i\partial_\mu - eA_\mu) - m] \psi(x, y), \quad (2.1)$$

where $A_\mu = A\delta_{\mu,x}$ and $\phi = eAL_x/2\pi$. The integral ranges over the surface area of the torus, and the time dependence of the fields has been suppressed. Due to the periodic boundary conditions (PBCs) in the x -direction, the momenta k_x are quantised: $k_x^n = \frac{2\pi n}{L_x}, n \in \mathbb{Z}$. We then expand the fields $\psi(x, y)$ in these momenta:

$$\psi(x, y) = \sum_{n=-\infty}^{\infty} e^{ik_x^n x} \psi(k_x^n, y). \quad (2.2)$$

In the absence of the gauge field, the Lagrangian for $2 + 1$ -dimensional massive non-interacting relativistic electrons (in natural units) is

$$\mathcal{L} = \int dx dy \bar{\psi}(x, y) [\gamma^\mu (i\partial_\mu - eA_\mu) - m] \psi(x, y). \quad (2.3)$$

The integral ranges over the area of the torus on which the electrons reside, and the time dependence of the fields has been suppressed. Each component $\psi^a, a = 0, 1, 2, 3$, of the Dirac spinor $\psi = (\psi^0 \ \psi^1 \ \psi^2 \ \psi^3)$ satisfies the Dirac equation

$$[\gamma^\mu (i\partial_\mu - eA_\mu) - m] \psi^a(x, y) = 0, \quad (2.4)$$

which is equivalent to the form

$$\int dx \bar{\psi}^a(x, y) [\gamma^\nu (i\partial_\nu + eA_\nu) + m] [\gamma^\mu (i\partial_\mu - eA_\mu) - m] \psi^a(x, y) = 0. \quad (2.5)$$

Due to the periodic boundary conditions (PBCs) in the x -direction, the momenta k_x are quantised: $k_x^n = \frac{2\pi n}{L_x}, n \in \mathbb{Z}$. We expand the fields $\psi^a(x, y)$ in these momenta:

$$\psi^a(x, y) = \sum_{n=-\infty}^{\infty} e^{ik_x^n x} \psi^a(k_x^n, y). \quad (2.6)$$

Writing the Dirac equation in terms of the dual fields $\psi^a(k_x^n, y)$ gives

$$\sum_{m,n} (\bar{\psi}^a(k_x^m, y) e^{-ik_x^m x}) [\gamma^\nu (i\partial_\nu + eA_\nu) + m] [\gamma^\mu (i\partial_\mu - eA_\mu) - m] (e^{ik_x^n x} \psi^a(k_x^n, y)) = 0. \quad (2.7)$$

Evaluating the right operation gives

$$[\gamma^\mu(i\partial_\mu - eA_\mu) - m] (e^{ik_x^n} \psi^a(k_x^n, y)) = e^{ik_x^n} [\gamma^\mu(i\partial_\mu - eA_\mu) - (\gamma^x(k_x^n + eA_x) + m)] \psi^a. \quad (2.8)$$

Similarly, the left operation gives

$$\begin{aligned} (\bar{\psi}^a(k_x^m, y) e^{-ik_x^m x}) [\gamma^\nu(i\partial_\nu + eA_\nu) + m] &= [[\gamma^\nu(i\partial_\nu - eA_\nu) + m] (e^{ik_x^m x} \gamma_0 \psi^a)]^\dagger \\ &= \bar{\psi}^a e^{-ik_x^m x} [\gamma_\nu(i\partial^\nu + eA_\nu) - (\gamma^x(k_x^m + eA_x) - m)]. \end{aligned} \quad (2.9)$$

Substituting these into the equation eq.(2.7) gives

$$\sum_n \int dy \bar{\psi}^a(k_x^n, y) [-(i\partial_\mu + eA_\mu)(i\partial_\mu - eA_\mu) + (k_x^n + eA_x)^2 + m^2] \psi^a(k_x^n, y) = 0, \quad (2.10)$$

where we have used $\int dx e^{ix(k_x^m - k_x^n)} = \delta_{m,n}$. In terms of these dual fields, the Lagrangian can be shown to be equivalent to a tower of massive $1 + 1$ -dimensional fermions (see Fig. 7.1):

$$\mathcal{L} = \sum_{k_x^n} \int dy \bar{\psi}(k_x^n, y) (\gamma^\mu(i\partial_\mu - eA_\mu) - M(n, \phi)) \psi(k_x^n, y), \quad (2.11)$$

with an effective mass $M_{n,\phi}$ given by

$$M_{n,\phi} = \sqrt{m^2 + (k_x^n + eA)^2} = \sqrt{m^2 + \frac{4\pi^2}{L_x^2} (n + \phi)^2}. \quad (2.12)$$

This decoupling ensures that the various modes $n \in \mathbb{Z}$ are disentangled from one another, such that the total density matrix ρ_{2D} of the $2 + 1$ -dimensional system can be written as a product of the density matrices ρ_{1D}^n for each of the modes:

$$\rho_{2D} = \prod_{n=-\infty}^{\infty} \rho_{1D}^n. \quad (2.13)$$

We will focus primarily on the case of massless $2 + 1$ -dimensional electrons by setting $m = 0$, such that the effective mass is given by $M_{n,\phi} = \frac{2\pi}{L_x} |n + \phi|$. Certain results will, however, be generalised to the case of massive fermions ($m \neq 0$). While the massless case corresponds to the gapless excitations of a Fermi liquid or those at the surface of a topological insulator, the massive case corresponds to gapped or insulating states of matter, e.g., the Bogoliubov-de Gennes quasiparticles above an s -wave superconducting gap. Massive Dirac fermions have also been observed experimentally in several materials like ultrathin films [Lu et al.(2010)Lu, Shan, Yao, Niu, and Shen], topological insulating crystals [Zhao et al.(2021)Zhao, Trang, Li, Chen, Yue, Bake, Tan, Wang, Nancarrow, Edmonds et al.] and van der Waals heterostructures [Hunt et al.(2013)Hunt, Sanchez-Yamagishi, Young, Yankowitz, LeRoy, Watanabe, Taniguchi, Moon, Koshino, Jarillo-Herrero et al.]. They are also emergent at low-energies, for instance, in fermionic nonlinear σ -models [Abanov and Wiegmann(2000)], and can also be realised near the IR fixed point of the kagome Heisenberg quantum antiferromagnets placed in an external magnetic field along the z -direction [Pal et al.(2019)Pal, Mukherjee, and Lal].




Figure 7.2: Visual depiction of the subsets $\mathcal{A}_z(j)$ employed by us in this work, for $z = 1$ and $z = -1$. As j is changed by 1, the spacing between the modes in the subset increases by a factor of 2^{j^z} . Top: $z = 1$. The initial subset $\mathcal{A}_z(0)$ has k -states at the modes $n = 0, 1, 2, 3, 4$. Upon changing j by 1, the new subset obtained has double the spacing between the modes ($2^{j^z} = 1/2^j$) of the previous number of modes. That is, if $j = 0$ has $0, 1, 2, 3, 4$, then $j = 1$ has $0, 2, 4$, $j = 2$ has $0, 4$ and $j = 3$ has 0 . Bottom: $z = -1$. In this case, the number of modes get double at each state ($2^{-j^z} = 2^j$). We start with k -modes present only at $n = 0$, but the scaling transformations lead to $j = 1$ having the modes $0, 4$, $j = 2$ having $0, 2, 4$, and so on.

7.2.2 Entanglement measures

We will work with two measures of entanglement discussed below. Given the density matrix $\rho = |\Psi\rangle\langle\Psi|$ for the ground state $|\Psi\rangle$ of the complete system, the entanglement entropy (EE) of a set of subsystems $\{\mathcal{A}_i, i = 1, 2, \dots, g\}$ with the rest of the system is given by:

$$S_{\{\mathcal{A}_i\}} = -\text{Tr} [\rho_{\{\mathcal{A}_i\}} \ln \rho_{\{\mathcal{A}_i\}}]; \quad \rho_{\{\mathcal{A}_i\}} = \text{Tr}_{\{\mathcal{A}_i\}} [\rho]. \quad (2.14)$$

$\text{Tr}[\cdot]$ is the trace operation over all degrees of freedom, while $\text{Tr}_{\{\mathcal{A}_i\}}[\cdot]$ is the partial trace over only the states corresponding to the set of subsystems $\{\mathcal{A}_i\}$. $\rho_{\{\mathcal{A}_i\}}$ is referred to as the reduced density matrix for the rest of the system (complement of the set $\{\mathcal{A}_i\}$). Using EE, one can define the mutual information $I(\{\mathcal{A}_i\} : \{\mathcal{B}_i\})$ between two subsystems A and B :

$$I_2(\{\mathcal{A}_i\} : \{\mathcal{B}_i\}) = S(\{\mathcal{A}_i\}) + S(\{\mathcal{B}_i\}) - S(\{\mathcal{A}_i\} \cup \{\mathcal{B}_i\}), \quad (2.15)$$

where $S(\{\mathcal{A}_i\} \cup \{\mathcal{B}_i\})$ is the EE of $\{\mathcal{A}_i\} \cup \{\mathcal{B}_i\}$ with the rest. Higher order measures of information among N subsystems can also be defined similarly (see, e.g., [Patra et al.(2022)Patra, Basu, and Lal]). Such multi-partite measures of quantum information convey the degree of correlations present within the system - the presence of N -particle correlations ensures all information measures up to the N -partite information will be non-zero.

One final object of interest in this context is the entanglement Hamiltonian $\mathcal{H}(\{\mathcal{A}_i\})$ for a subsystem $\{\mathcal{A}_i\}$, defined through the relation [Li and Haldane(2008), Calabrese and Lefevre(2008)]

$$\rho_{\{\mathcal{A}_i\}} = \frac{e^{-\mathcal{H}(\{\mathcal{A}_i\})}}{\text{Tr}(e^{-\mathcal{H}(\{\mathcal{A}_i\})})}. \quad (2.16)$$

The entanglement Hamiltonian is essentially the spectrum of the reduced density matrix, and contains additional information on the nature of correlations beyond the entanglement entropy. For the

system we are considering (described in the previous subsection), the total reduced density matrix can be decomposed into decoupled 1D modes (eq. (2.13)). This implies that the total entanglement Hamiltonian is simply a sum of the individual Hamiltonians for the 1D modes:

$$\rho_{2D} = \prod_{n=-\infty}^{\infty} \rho_{1D}^n \sim \prod_{n=-\infty}^{\infty} e^{-\mathcal{H}_{1D}^n(\{\mathcal{A}_i\})} = e^{-\mathcal{H}_{2D}(\{\mathcal{A}_i\})}, \quad (2.17)$$

where $\mathcal{H}_{2D} = \sum_n \mathcal{H}_{1D}^n$.

7.2.3 Luttinger volume

The Luttinger volume V_L is defined as the number of $\vec{k}$ -space points inside the Fermi surface [Luttinger(1960), Luttinger and Ward(1960)] :

$$V_L = \int_{G_{\vec{k}}(\omega=0) > 0} \frac{1}{(2\pi)^d} d\vec{k} = \int_0^{k_F} \frac{1}{(2\pi)^d} d\vec{k}. \quad (2.18)$$

The integral (referred to as Luttinger's integral) runs over all $\vec{k}$ -space points where the $\omega = 0$ Greens function is positive, and the Fermi momentum k_F is defined as the set of points in $\vec{k}$ -space where $G_{\vec{k}}(\omega = 0)$ changes sign. Luttinger's integral can be shown to be topological in nature [Seki and Yunoki(2017), Oshikawa(2000)], promoting Luttinger's volume to the status of a topological invariant that characterises the system. For systems with long-lived excitations close to the Fermi surface, V_L is equal to the average number density of electrons in the system, leading to Luttinger's theorem [Luttinger(1960), Luttinger and Ward(1960), Heath and Bedell(2020a)]. For systems with gapped excitations in the momentum window $k \leq k^*$ above the ground state, the violation ΔV_L of Luttinger's theorem can be quantified through a count of the bound states [Mukherjee and Lal(2020a)]:

$$\Delta V_L = \int_0^{k_F} \frac{1}{(2\pi)^d} d\vec{k} - \int_{k^*}^{k_F} \frac{1}{(2\pi)^d} d\vec{k} = \int_0^{k^*} \frac{1}{(2\pi)^d} d\vec{k} \quad (2.19)$$

7.3 Entanglement hierarchy in mixed momentum and real space

Before beginning our calculations of entanglement measures, we define the subsystem(s) whose entanglement we want to obtain. The subsystems we will use in this work are of the following kind:

$$\mathcal{A}_z(i) = \tilde{A}_z(i) \times A_z(i), \quad (3.1)$$

where i takes integral values starting from zero, z can take all integral values apart from 0 and $\times$ represents the Cartesian product between the set of real space points $\tilde{A}_z(i) = \{0 \leq y \leq l\}$ and the set of momentum space points

$$A_z(i) = \left\{ k_x^n \right\}; \quad k_x^n = \frac{2\pi}{L_x} n; \quad \max(\{k_x^n\}) = \frac{2\pi N}{L_x};$$

$$n \in \{-N, -N + t_z(i), \dots, -t_z(i), 0, t_z(i), N\}; \quad t_z(i) = 2^{i^z}, \quad (3.2)$$

and where $t_z(i) = 2^{iz}$ is the difference between consecutive values of n . The subset $\mathcal{A}_z(i)$ is therefore a region in mixed momentum and real spaces, spanned by the variables k_x and y respectively. In y -space, the region is of length l and bounded by the points $y = 0$ and $y = l$. In k_x -space, $\mathcal{A}_z(i)$ comprises the points $\left\{0, \pm 2^{iz} \frac{2\pi}{L_x}, \pm 2^{iz} \frac{4\pi}{L_x}, \dots\right\}$ such that $|k_x| \leq \frac{2\pi N}{L_x}$. For example, for the case of $z = 1$ and N being a multiple of 4, we have

$$\begin{aligned} A_1(0) &= \left[-\frac{2\pi}{L_x}N, -\frac{2\pi}{L_x}(N-1), \dots, -\frac{2\pi}{L_x}2, -\frac{2\pi}{L_x}1, 0, \frac{2\pi}{L_x}1, \frac{2\pi}{L_x}2, \dots, \frac{2\pi}{L_x}(N-1), \frac{2\pi}{L_x}N \right], \\ A_1(1) &= \left[-\frac{2\pi}{L_x}N, -\frac{2\pi}{L_x}(N-2), \dots, -\frac{2\pi}{L_x}4, -\frac{2\pi}{L_x}2, 0, \frac{2\pi}{L_x}2, \frac{2\pi}{L_x}4, \dots, \frac{2\pi}{L_x}(N-2), \frac{2\pi}{L_x}N \right], \\ A_1(2) &= \left[-\frac{2\pi}{L_x}N, -\frac{2\pi}{L_x}(N-4), \dots, -\frac{2\pi}{L_x}8, -\frac{2\pi}{L_x}4, 0, \frac{2\pi}{L_x}4, \frac{2\pi}{L_x}8, \dots, \frac{2\pi}{L_x}(N-4), \frac{2\pi}{L_x}N \right]. \end{aligned} \quad (3.3)$$

Note that N represents the edge of the Brillouin zone, and will be sent to infinity when we calculate quantities in the continuum limit. In Fig. 7.2, we have shown two such families of subsets $\mathcal{A}_z(0), \mathcal{A}_z(1), \mathcal{A}_z(2), \mathcal{A}_z(3), \dots$, for $z = 1$ and $z = -1$.

We note here that the transformations for $z < 0$ correspond to increasing the density (fine-graining) of k_x -states by adding more momentum states while keeping the bounds fixed. This involves reducing the interval in k_x -space, and hence equivalent to increasing the system size L_x . This is similar to methods such as numerical renormalisation group treatment of the Kondo problem [Wilson(1975b)] where an increasing number of states are added to the system at increasingly larger distances from the impurity site, such that the system asymptotically goes to the thermodynamic limit. The other case of $z > 0$ is, on the contrary, a decimation process in k_x -space, and equivalent to reducing the system size. Indeed, it will be shown later that the sequence of Hamiltonians generated in the process (for both positive and negative z) are related to one other by renormalisation group transformations.

Within a given set $\mathcal{A}_z(i)$, the distance between adjacent k -space points is $\frac{2\pi}{L_x}t_z(i)$ where $t_z(i) = 2^{iz}$. Choosing $i = 0$ keeps the set of k_x -space points unchanged, such that we recover the entire Lagrangian of eq. (2.1) in the x -direction for system size L_x , while the y -direction is constrained to length l . This means that $\mathcal{A}_z(0)$ is just an annulus that wraps around the x -direction of the torus and extends to a length l longitudinally in the y -direction (shown in Fig. 7.3). Along with the period $t_z(i)$, one can also define the fraction of maximum points present in the set $\mathcal{A}_z(i)$ as

$$f_z(i) = 1/t_z(i) = 2^{-iz}. \quad (3.4)$$

For example, for $z > 0$, $f_z(i) = 2^{-iz}$ expresses the fact that in going from the set at $j = 0$ to the set at $j = i$, the number of elements in the set is scaled down by a factor of 2^{iz} (see Fig. 7.4).

We now define the shorthand notation $S_z(i)(\phi) \equiv S_{\mathcal{A}_z(i)}(\phi)$ to denote the EE of the subsystem $\mathcal{A}_z(i)$ in the presence of the Aharonov-Bohm flux ϕ . In order to calculate $S_z(i)$, we start with $i = 0$. As mentioned earlier, this is essentially the EE between an annulus of length l and the rest of the 2-torus. Since the density matrices are in product, the total EE is just a sum of the EE arising from each mode:

$$S_{\mathcal{A}_z(0)}(\phi) \equiv S_z(0)(\phi) = \lim_{N \rightarrow \infty} \sum_{n=-N}^N S_{\mathcal{A}_z(0)}^n, \quad (3.5)$$




Figure 7.3: Top: Depiction of the subset $\mathcal{A}_z(i)$ for the choice $i = 0$. Setting $i = 0$ leads to unchanged length L_x along x and the constraint $0 \leq y \leq l$ in the y -direction. This creates an annulus (red region) of height l and circumference L_x .

Bottom: Typical k -modes present in the subset $\mathcal{A}_z(i)$ for $z = 1$ and $i = 2$. This is because, if $i = 0$ has the modes $n = 0, 1, 2, 3, 4, \dots$, increasing i by 2 increases the spacing between the modes to $2^{2z} = 4$, so that the mode just after $n = 0$ is $n = 4$, and the one following that is $n = 8$, and so on.

where $\tilde{A}_z(0)$ acts as a projection of the annulus on the $k_x - y$ plane, and S^n indicates that the EE is calculated for the n^{th} mode. Each massive fermionic mode has a correlation length-scale $\xi_n = 1/M_{n,\phi}$. We choose l to be much larger than the largest correlation length: $l \gg \max(\{\xi_n, \forall n\})$. Using the expression for ϕ and the quantisation of the momenta k_x^n , this condition can be expressed as

$$l \gg \frac{L_x}{2\pi} \times \max \left(\left\{ \frac{1}{|n + \phi|} \right\} \right) = \frac{L_x}{2\pi \{\phi\}}, \quad (3.6)$$

where $\{\phi\} = \phi - \lfloor \phi \rfloor$ is the fractional part of the flux ϕ in units of the flux quantum. Since we are in the thin torus limit, L_x must remain finite, and the maximum correlation can diverge only if $\{\phi\} = 0$. We ensure that l is larger than the correlation lengths by choosing ϕ to be necessarily non-integral, as this guarantees that the correlation lengths remain finite. Then, the EE of the annulus is given by [Arias et al.(2015)Arias, Blanco, and Casini, Chen et al.(2017)Chen, Witczak-Krempa, Faulkner, and Fradkin]

$$S_z(0)(\phi) = c \left[\alpha \frac{L_x}{\epsilon} - \ln |2 \sin(\pi \phi)| \right], \quad (3.7)$$

where c is the conformal central charge and ϵ is a UV cutoff; c is $1/3$ for a Dirac fermion. The computation of $S_z(0)(\phi)$ uses the EE $S_{\tilde{A}_z(0)}^n$ of a 1-D chain of fermions with a mass $M_{n,\phi}$ [Calabrese and Cardy(2004), Casini and Huerta(2009b)]: $S_{\tilde{A}_z(0)}^n = c \ln \frac{L_x}{2\pi\epsilon} - c \ln |n + \phi|$. On summing over n and regularising the UV cutoff, the first term of eq.(3.7) produces the leading area-law term of $S_0(\phi)$ [Chen et al.(2017)Chen, Witczak-Krempa, Faulkner, and Fradkin]:

$$\sum_{n=-\infty}^{\infty} c \ln \frac{L_x}{2\pi\epsilon} \rightarrow c \alpha \frac{L_x}{\epsilon}, \quad (3.8)$$




Figure 7.4: A plot of the fraction of states remaining upon performing i number of transformations on the starting Hamiltonian. Obtained by plotting the expression $f = 2^{-iz}$, for a range of i , and for two values of z (red and blue). This fraction decreases for $z > 0$ (blue curve), and increases for $z < 0$ (red curve).

where α is a cutoff-dependent non-universal constant. The flux-dependent term in $S_{\tilde{A}_z(0)}^n$ leads to the sub-leading term of $S_z(0)(\phi)$, and requires a regularisation using the Hurwitz zeta function [Chen et al.(2017)Chen, Witczak-Krempa, Faulkner, and Fradkin]:

$$\sum_{n=-\infty}^{\infty} \ln |n + \phi| = \sum_{n=-\infty}^{\infty} [\ln |n + \phi| + \ln |n + (1 - \phi)|] = -\frac{\partial \zeta(\phi)}{\partial \phi} - \frac{\partial \zeta(1 - \phi)}{\partial \phi}, \quad (3.9)$$

where $\zeta(\phi) = \sum_{n=-\infty}^{\infty} (n + \phi)^{-1}$ is the Hurwitz zeta function. The derivative converges to $\frac{\partial \zeta(\phi)}{\partial \phi} = \ln \Gamma(\phi) - \frac{1}{2} \ln 2\pi$, as long as the zero mode $n + \phi = 0$ is excluded from the summation. The latter condition is guaranteed as we have chosen ϕ to be a non-integer. Γ is the Gamma function, satisfying $\Gamma(\phi)\Gamma(1 - \phi) = \pi / \sin(\pi\phi)$. Substituting the derivative gives [Arias et al.(2015)Arias, Blanco, and Casini, Chen et al.(2017)Chen, Witczak-Krempa, Faulkner, and Fradkin, Crépel et al.(2021)Crépel, Hackenbroich, Regnault, and Estienne]

$$\sum_{n=-\infty}^{\infty} \ln |n + \phi| = \ln 2\pi - \ln [\Gamma(\phi)\Gamma(1 - \phi)] = \ln |2 \sin(\pi\phi)|. \quad (3.10)$$

For a general $\mathcal{A}_z(i)$, there are certain modifications to the expression in eq. (3.7). For the first term in eq.(3.7), we have

$$\lim_{N \rightarrow \infty} \sum_{\substack{n=-N, \\ -N+t_z(i), \dots}}^N c \ln \frac{L_x}{2\pi\epsilon} = \lim_{N \rightarrow \infty} \sum_{\substack{n'=-\frac{N}{t_z(i)}, \\ -\frac{N}{t_z(i)}+1, \dots}}^{\frac{N}{t_z(i)}} c \ln \frac{L_x}{2\pi\epsilon} = t_z^{-1}(i) \lim_{N \rightarrow \infty} \sum_{n=-N}^N c \ln \frac{L_x}{2\pi\epsilon} \rightarrow c f_z(i) \frac{\alpha L_x}{\epsilon}. \quad (3.11)$$

Similarly, for the second term, we have

$$\begin{aligned}
 \lim_{N \rightarrow \infty} \sum_{n=-N, \dots, -N+t_z(i), \dots}^N \ln |n + \phi| &= \lim_{N \rightarrow \infty} \sum_{n'=-\frac{N}{t_z(i)}, \dots, -\frac{N}{t_z(i)}+1, \dots}^{\frac{N}{t_z(i)}} \ln |n' t_z(i) + \phi| = \lim_{N \rightarrow \infty} \sum_{n'=-\frac{N}{t_z(i)}, \dots, -\frac{N}{t_z(i)}+1, \dots}^{\frac{N}{t_z(i)}} \ln (f_z(i) |n' + f_z(i) \phi|) \\
 &= \sum_{n'=-\infty}^{\infty} \ln |n' + f_z(i) \phi| ,
 \end{aligned} \tag{3.12}$$

where we have assumed that $t_z(i)$ is finite, and dropped the constant part $\ln f_z(i)$ (which can be absorbed into the UV cutoff ϵ in eq. (3.11)). Following eq. (3.10), this gives

$$\lim_{N \rightarrow \infty} \sum_{n=-N, \dots, -N+t_z(i), \dots}^N \ln |n + \phi| = \ln |2 \sin (\pi f_z(i) \phi)| . \tag{3.13}$$

Combining both parts, we obtain the expression for the EE that will be employed in what lies ahead:

$$S_z(i)(\phi) = c f_z(i) \frac{\alpha L_x}{\epsilon} - c \ln |2 \sin (\pi f_z(i) \phi)| . \tag{3.14}$$

It is evident from the expression in eq. (3.14) that, as modes are added into the subsystem, the L_x –dependent geometric part of the EE increases, leading to a *hierarchical structure* in the entanglement. A similar result was previously shown to hold for geometric measures of entanglement of quantum-mechanical systems with eigenstates such as the W and GHZ states and their symmetric superpositions [Blasone et al.(2008)Blasone, Dell’Anno, De Siena, and Illuminati]. The variation of the geometric part with the subsystem index j is shown in Fig. 7.4.

The above-mentioned hierarchy in entanglement measures leads to an interesting property for the entanglement: the combined entanglement entropy of any number of systems $\{\mathcal{A}_z(i)\}$ is equal to the entanglement entropy of the most dense system among them. For $z > 0$, the sets become smaller as the RG index increases, while for $z < 0$, the sets become larger as the RG index increases. Thus, we may write

$$S_{\{\mathcal{A}_z(i)\}} = \theta(z) S_{\min\{i\}, z} + \theta(-z) S_{\max\{i\}, z} , \tag{3.15}$$

The mutual information $I^2(i, j)$ between the systems at two RG steps i and j can then be computed:

$$I^2(i, j) \equiv S_z(i) + S_{j,z} - S_{i \cup j, z} = \theta(-z) S_{\min(i, j), z} + \theta(z) S_{\max(i, j), z} . \tag{3.16}$$

In light of this subsystem-based hierarchical structure, we can also consider the g –partite information $I_{\{\mathcal{A}_z(i)\}}^g$ among g sets $\{\mathcal{A}_z(i)\}$. I^g is defined as [Patra et al.(2022)Patra, Basu, and Lal]

$$I_{\{\mathcal{A}_z(i)\}}^g = \sum_{r=1}^g (-1)^{r+1} \sum_{\substack{\beta \in \mathcal{P}(\{\mathcal{A}_z(i)\}) \\ |\beta|=r}} S_{\beta} , \tag{3.17}$$

where $\mathcal{P}(\{\mathcal{A}_z(i)\})$ is the power set of $\{\mathcal{A}_z(i)\}$, and $|\beta| = r$ indicates that the sum is carried out only over those sets β that have r number of elements in them. For example, for $g = 3$, we have

$$I^3 = S_{\mathcal{A}_z(0)} + S_{\mathcal{A}_z(1)} + S_{\mathcal{A}_z(2)} - S_{\mathcal{A}_z(0) \cup \mathcal{A}_z(1)} - S_{\mathcal{A}_z(1) \cup \mathcal{A}_z(2)} - S_{\mathcal{A}_z(2) \cup \mathcal{A}_z(0)} + S_{\mathcal{A}_z(0) \cup \mathcal{A}_z(1) \cup \mathcal{A}_z(2)} . \quad (3.18)$$

For our choice of subsystems, the g -partite information can be shown to be equal to the EE of the *intersection set* of the concerned g -sets (details in [Appendix A](#)). For the case of $z > 0$, this takes the form

$$I_{\{\mathcal{A}_z(i)\}}^g(\phi) = S_g(\phi) = cf_{g,z} \alpha \frac{L_x}{\epsilon} - c \ln |2 \sin(\pi f_{g,z} \phi)| . \quad (3.19)$$

Since the g -partite information is simply the EE of the g^{th} subsystem, the hierarchy in the EE can thus be seen to lead to multiple levels of correlation in the system, corresponding to an increasing number of parties entering the set.

Finally, we note that the geometric (L_x -dependent) part of the EE in eq. (3.14), as well as any of the information measures I^g (eq. (3.19)), has the same form even if we start with a non-zero mass $m > 0$ [Chen et al.(2017)Chen, Witczak-Krempa, Faulkner, and Fradkin]. This ensures that the above observations regarding the hierarchical structure of the entanglement hold even for gapped electronic systems.

7.4 EHM, the Ryu-Takayanagi bound and holographic inequalities

As we will now show, the approach laid out above to the creation of subsystems amounts essentially to performing a sequence of renormalisation group (RG) transformations on the momentum modes of the Hamiltonian in the x -direction. We consider here the general case of massive Dirac electrons $M_{n,\phi} = \sqrt{m^2 + \frac{\pi^2}{L_x^2} (n + \phi)^2}$. Further, we define the superset $\mathcal{A}_z^{(0)}$ of k_x -states: $\mathcal{A}_z^{(0)} = \bigcup_{i=0}^{\infty} \mathcal{A}_z(i)$, such that all k_x -states that can appear at any point of the sequence $\{\mathcal{A}_z(i); i = 0, 1, 2, \dots\}$ are present within it. This superset can now be used to define a parent Hamiltonian $H(0)$ that consists of all the modes in the x -direction:

$$H(0) = \sum_{k_x \in \mathcal{A}_z^{(0)}} H(k_x) , \quad (4.1)$$

where $H(k_x)$ is the Hamiltonian for a massive Dirac fermion that extends over a length L_y in the single spatial dimension, and having mass $M_{n,\phi}$. The Hamiltonian at any given step of the RG is then obtained by projecting onto appropriate sets of momenta. The set of momentum at step j is given by $\mathcal{A}_{j,z} = \frac{2\pi}{L_x} \times \{0, \pm t_z(j), \pm, 2t_z(j), \dots\}$ with $t_z(j) = 2^{jz}$. The projector onto this set is given by

$$P_{j,z} = \sum_{k_x^1=0,1} \sum_{k_x^2=0,1} \dots \sum_{k_x^N=0,1} |\hat{n}_{k_x^1} \hat{n}_{k_x^2} \dots\rangle \langle \hat{n}_{k_x^1} \hat{n}_{k_x^2} \dots| , \quad k_x^j \in \mathcal{A}_{j,z} , \quad (4.2)$$

and the Hamiltonian at the corresponding step is then obtained as

$$H_z(i) = P_z(i) H(0) P_z(i) . \quad (4.3)$$




Figure 7.5: Top: Modes forming the subsystem $\mathcal{A}_z(i)$ (red circles) form the boundary between $\mathcal{A}_z(i)$ and its complement (blue circles). Bottom: Scaling of the subsystem entanglement entropy (red), and its upper bound (blue), as the RG transformations are performed (denoted by increasing index i). The bound is saturated at large i .

In this formulation of the RG, the index i represents the RG step, while the parameter z indicates the *strength of the scaling* mechanism: for $z > 0$, a higher value of z means that the states of the Hamiltonian become rarer as the index i increases, while for $z < 0$, the states become denser with increasing i . Importantly, we will see subsequently that the parameter z corresponds also to the anomalous dimension of the coupling that leads to a spectral gap in the electronic dispersion, dictating its growth or decay under RG evolution (eq.(5.6)).

Since this is a free theory, the k -modes are decoupled, and the projection operation preserves the spectrum of the remaining degrees of freedom. The renormalised Hamiltonian (eq. (4.3)) can then be said to constitute an exact holographic mapping (EHM) [Qi(2013), Lee and Qi(2016)] in the direction of the RG transformations. It is also pertinent to note that the sequence of transformations in eq. (4.3) can be thought of as a simple and specific example of a unitary renormalisation group (URG) procedure [Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)] that first disentangles and then projects out a certain set of degrees of freedom. Because the operation is unitary, the URG preserves the spectrum in the process. In the present case, the disentanglement operation is unity. The URG has already been shown to lead to a holographic renormalisation of the Hilbert space geometry which can be seen through the evolution of the Fubini-Study metric under the URG transformations [Mukherjee and Lal(2020a)]. Our approach differs from MERA implementations like those in refs. [Evenbly and Vidal(2009), Evenbly and White(2016)] in the fact that while those are involve a real space renormalisation scheme, we perform our scaling transformations in momentum space.

More evidence for the holographic nature of the construction comes from the fact that the entanglement entropy for both the massless and massive cases satisfy the bound provided by the Ryu-Takayanagi proposal that relates the entanglement of a conformal field theory (CFT) with the geometry of its gravity dual [Ryu and Takayanagi(2006a), Ryu and Takayanagi(2006b)]:

$$S_{\mathcal{A}_z(i)} \leq \mathcal{N}_z(i) \max(\{S^n\}) . \quad (4.4)$$

In the above constraint, $\max(\{S^n\})$ is the maximum single entropy of all the modes n residing in $\mathcal{A}_z(i)$, and $\mathcal{N}$ is the area of the minimal surface that separates the subsystem $\mathcal{A}_z(i)$ from the rest. In order to see how this is satisfied in the presence case, we note that since the subsystem $\mathcal{A}_z(i)$ is interspersed among the rest of the system (see left panel of Fig. 7.5), the minimal surface is as large as the subsystem itself. The area of this surface is therefore given by the number of modes in $\mathcal{A}_z(i)$. Using this information, the right hand side of eq. (4.4) can be written as

$$\mathcal{N}_z(i) \max(\{S^n\}) = \max(\{S^n\}) N f_z(i) = \max(\{S^n\}) \sum_{n \in \mathcal{A}_z(i)} . \quad (4.5)$$

Following eq. (2.13), the left hand side of eq. (4.4) can, on the other hand, be written as

$$S_{\mathcal{A}_z(i)} = \sum_{n' \in \mathcal{A}_z(i)} S^{n'} \leq \sum_{n' \in \mathcal{A}_z(i)} \max(\{S^n\}) . \quad (4.6)$$

Comparing eqs. (4.5) and (4.6), we recover the Ryu-Takayanagi condition of eq. (4.4). The scaling of both the left and right hand sides of eq. (4.4) has been shown in the right panel of Fig. 7.5.

Entanglement measures like entanglement entropy and mutual information (associated with disjoint intervals) have been shown to satisfy a number of inequalities in quantum field theories with gravity duals. We now consider some of them here, and show that the theory considered by us here also satisfies these inequalities. One such constraint is the monogamy of mutual information, which says that for three disjoint regions A_1, A_2 and A_3 , we must have $I_2(A_1 : A_2) + I_2(A_1 : A_3) \leq I_2(A_1 : A_2 \cup A_3)$, where $I_2(A_1 : A_2) = S(A_1) + S(A_2) - S(A_1 \cup A_2)$ [Hayden et al.(2013)Hayden, Headrick, and Maloney]. For this, we consider three disjoint regions, all of the kind depicted in the left panel of fig. 7.3, but at different positions $y_i \in [l_i, l_i + L], i = 1, 2, 3$. We assume that the distance $l_i - l_{i+1}$ between adjacent regions is much larger than the correlation length. Calabrese and Cardy have shown that the total entanglement entropy of the union of such disjoint regions is obtained by multiplying the entanglement entropy of any one region with the number of disjoint intervals. Putting all this together, we get

$$\begin{aligned} S(A_1) &= S(A_2) = S(A_3), \\ S(A_1 \cup A_2) &= S(A_1 \cup A_3) = 2S(A_1), \\ S(A_1 \cup A_2 \cup A_3) &= 3S(A_1) , \end{aligned} \quad (4.7)$$

leading to the above inequality being satisfied:

$$I_2(A_1 : A_2) + I_2(A_1 : A_3) - I_2(A_1 : A_2 \cup A_3) = 0 . \quad (4.8)$$

We next consider a cyclic family of inequalities in terms of conditional entropies, introduced and proved in Ref. [Bao et al.(2015)Bao, Nezami, Ooguri, Stoica, Sully, and Walter]: $\sum_{i=1}^{2k+1} S(A_i | A_{i+1} \dots A_{i+k}) \geq S(A_1 \dots A_{2k+1})$, where $S(A_i | A_j) = S(A_i \cup A_j) - S(A_j)$. Using the results of eq. (4.7), we get $\sum_{i=1}^{2k+1} S(A_i | A_{i+1} \dots A_{i+k}) = \sum_{i=1}^{2k+1} S(A_i) = (2k+1)S(A_1)$ and $S(A_1 \dots A_{2k+1}) = (2k+1)S(A_1)$, leading to $\sum_{i=1}^{2k+1} S(A_i | A_{i+1} \dots A_{i+k}) = S(A_1 \dots A_{2k+1})$, which also satisfies the inequality. In both cases, we find that the inequality is satisfied by saturating the bound; this is a consequence of the fact that the intervals are separated by distances larger than the correlation length and the total entanglement entropies are simply a sum of the individual entropies.

7.5 Holographic geometry of the RG flow

7.5.1 RG evolution as the emergent dimension

To obtain a more specific description of the RG flow as the emergence of a geometric space, we introduce a measure of distance. For this, we will employ another bipartite measure of entanglement - the mutual information I^2 , as defined in eq. (2.15), due to its non-negative nature. We focus on the massless case $m = 0$ for the time-being. Inspired by Refs. [Van Raamsdonk(2010), Lee and Qi(2016), Hyatt et al.(2017)Hyatt, Garrison, and Bauer, Mukherjee and Lal(2022b)], we adopt the following definition for distance:

$$d_z(i, j) \equiv \ln \frac{I_{\max}^2}{I_z^2(i : j)}, \quad (5.1)$$

where $I_z^2(i : j) \equiv I^2(\mathcal{A}_z(i), \mathcal{A}_{j,z})$ is the mutual information between the two mentioned sets, and $I_{\max}^2$ is the mutual information between two maximally entangled systems. Accordingly, $d_z(j)$ is zero (i.e., the mutual information is maximum) at $j = 0$ if $z > 0$ and $j = \infty$ if $z < 0$, and this distance increases (decreases) with the decrease (increase) in the mutual information as the scaling transformations are performed.

We compute the distance between the points i and j to obtain the scaling of the distance $d_z(j)$. We assume $i < j$ without loss of generality. At each step, we tune the flux ϕ so as to remove the flux-dependent part from the EE. Following eq. (3.16), the mutual information is of the form $\theta(-z)S_i + \theta(z)S_j$, such that the distance is given by

$$d_z(i, j) = \theta(-z) \ln \frac{S_{\max}}{S_z(i)} + \theta(z) \ln \frac{S_{\max}}{S_{j,z}} = \theta(z)j^z \ln 2 + \theta(-z)i^z \ln 2. \quad (5.2)$$

For $z > 0$, the distance increases as j increases (keeping i fixed), as the k -space points are made more coarse-grained under the RG transformations. On the other hand, for $z < 0$, the distance decreases as i is increased (keeping j fixed). The rate of increase or decrease depends on the value of z . This scaling of the distance is shown in the left panel of Fig. 7.6.

At any given RG step j , we define two subsystems $\mathcal{B}_j^1$ and $\mathcal{B}_j^2$. The first subsystem $\mathcal{B}_j^1$ is the entire set of states $\mathcal{A}_{j,z}$ available at the j^{th} RG step, while the second subsystem $\mathcal{B}_j^2$ is the set comprising those modes that remain to be considered at the next RG step if $z > 0$, and the set comprising only the modes that were present in the previous step for $z < 0$. This scheme is shown in the right panel of Fig. 7.6. The mutual information between these two sets defines a measure of distance y_j at the j^{th} RG step:

$$y_z(j) \equiv \ln \frac{I_{\max}^2}{I_z^2(\mathcal{B}_j^1 : \mathcal{B}_j^2)} = \begin{cases} (j+1)^z \ln 2, & z > 0, \\ (j-1)^z \ln 2, & z < 0. \end{cases} \quad (5.3)$$

The distance along the RG direction ($x_z(j)$) is, on the other hand, defined as the distance between the current RG step and the one with the maximum entanglement entropy:

$$x_z(j) \equiv \theta(z)d(0, j) + \theta(-z)d(j, \infty) = j^z \ln 2, \quad (5.4)$$

where $d(i, j)$ was defined in eq. (5.1).




Figure 7.6: Top: Variation of the information distance $d_z(0 : j)$ (defined along the RG direction) under the scaling transformations. Obtained by plotting the expression for the distance, as a function of the number of transformations j , for two values of z . For $z = 1$ (blue curve), the distance between two extreme points increases, showing a decrease in entanglement. For $z = -1$ (red curve), the distance decreases, showing an increase in entanglement. Bottom: Geometry of subsystems used for calculating the emergent curvature. The red blobs represent the subsystems generated along the emergent direction, through the scaling transformations: $\mathcal{A}_{\in j} \rightarrow \mathcal{B}_{j+1}^1$. The blue blobs represent subsystems generated within the $1+1$ -dimensional field theory, transverse to the emergent direction.

The connection of the x - and y -distances to the RG can be made manifest by relating them to the RG beta function. As we have seen above, even though we are working with massless Dirac fermions ($m = 0$), the Aharonov-Bohm flux and the dimensional reduction into $1+1$ -dimensional modes generate an effective mass $M_{n,\phi} = \frac{2\pi}{L_x}(n + \phi)$ for $n > 0$. At a specific RG step j , the n^{th} mode is given by $t_z(j) \times n$ on account of the RG transformations in k_x -space. With these points in mind, we define a coupling $g_z(j)$ that acts as a measure of the gap in the single-particle spectrum

$$g_z(j) \equiv \ln \frac{M_{n+1,\phi}(j) - M_{n,\phi}(j)}{2\pi/L_x} = \ln t_z(j) = j^z \ln 2. \quad (5.5)$$

The RG beta-function for the coupling $g_z(j)$ can then be written as

$$\beta_z(j) \equiv \frac{\Delta \log g_z}{\Delta j} = \log \frac{g_z(j+1)}{g_z(j)} = z \ln \left(1 + \frac{1}{j} \right), \quad (5.6)$$

where we have substituted $\Delta j = 1$. The beta function is positive for $z > 0$, indicating that the coupling $g_z(j)$ is RG-relevant in this regime (left panel of Fig. 7.7). On the other hand, the beta function is negative for $z < 0$, leading to irrelevant RG flows in this regime. The RG beta-function $\beta_z(j)$ itself decreases monotonically as j increases under the RG transformations (see Fig. 7.7): for a finite z , a fixed point is reached at $j \rightarrow \infty$ where the beta function goes to zero.

As we will now see, the flow of the RG beta functions can be related to the extremisation of a potential defined in the space of Hamiltonians. This potential is a function of the mutual information




Figure 7.7: Top: Depiction of the RG flow of the coupling $g_z(j)$. For $z < 0$ (blue and red curves), the coupling is RG-irrelevant and shows a decrease as j is increased. For $z > 0$ (gray and violet curves), the coupling is RG-relevant and shows an increase as j is increased. Bottom: Variation of RG beta function $\beta_z(j)$ for the coupling $g_z(j)$, as a function of step index j . For all values of z , the beta function converges to 0 as $j \rightarrow \infty$, indicating a fixed point of the coupling g .

$I_z^2(0 : j)$:

$$\beta_z(j) = -\Delta \left[\ln \left(\ln \frac{I_z^2(0 : j)}{I_{\max}^2} \right) \right]. \quad (5.7)$$

The potential given within the square brackets above corresponds to an RG monotone whose fixed point corresponds with the fixed point of the RG. For relevant flows, the above equation ensures that the function $\ln \left(\ln \frac{I_z^2(0:j)}{I_{\max}^2} \right)$ is always minimised under the RG flow, with the consequence that the trajectories ultimately lead to minimisation of the mutual information. This indicates that the relevance of the mass coupling leads in turn to the decrease of the correlation length and hence a lowering of entanglement between points that are separated in real space. On the other hand, for irrelevant flows, the above-mentioned function is maximised, implying that the RG trajectories now maximise the entanglement. This reflects the fact that the decreasing mass leads to increased correlation lengths and an approach towards criticality.

Given eqs.(5.3), (5.4) and (5.7), we can relate the distances x and y to the beta function β_z :

$$x_z = \left(e^{\frac{\beta_z}{z}} - 1 \right)^{-z} \ln 2, \quad (5.8)$$

$$y_z = \begin{cases} x_z e^{\beta_z} & z > 0, \\ x_z \left(2 - e^{\frac{\beta_z}{z}} \right)^z & z < 0. \end{cases} \quad (5.9)$$

The contrasting behaviour of the beta function depending on the sign of z indicates that it tracks the effective number of degrees of freedom that contribute to the entanglement (eq. (3.14)) at any given RG step. Indeed, we find that the effective number of degrees of freedom is the effective c -function (or central charge) $\tilde{c}_z(j)$ of the theory, taking the form $\tilde{c}_z(j) = f_z(j)c$. The beta function is thus related to the c -function as

$$\beta_z = \Delta \left[\ln \left(\ln \frac{c}{\tilde{c}_z(j)} \right) \right]. \quad (5.10)$$




Figure 7.8: Top: Variation of the effective central charge defined in eq. (5.10), under the RG transformations for relevant ($z = 1$, blue curve) and irrelevant ($z = -1$, red curve) flows. In the former case, the central charge starts from the fermionic gapless value of $1/3$ and vanishes towards the fixed point, indicating the flow to a gapped massive phase. In the latter case, it starts from the gapped value of 0, and increases to $1/3$ at the fixed point, indicating the flow to a gapless phase. Bottom: Variation of the curvature $\kappa_z(j)$ along the RG for multiple values of z . $z = 1$ leads to a flat space with zero curvature, while $z > 1$ and $z < 1$ (inset) lead to negative and positive curvature respectively.

For RG relevant flows, the c -function is found to diminish from the free fermion central charge $c = 1/3$, reaching the value of zero at the fixed point (where only the central $n = 0$ mode remains and the rest have been removed because of the large mass gap). The non-negative, non-increasing nature of the c -function, and its flow towards its minimum at the fixed point, leads to speculate that eq.(5.10) can be thought of as a holographic counterpart of Zamalodchikov’s c -theorem [Álvarez and Gómez(1999), Myers and Sinha(2010), Myers and Sinha(2011), Chu and Giataganas(2020)]. For RG irrelevant flows, the behaviour is reversed. The flow of the c -function is shown in the left panel of Fig. 7.8.

7.5.2 Embedding the emergent dimension on a metric space

We have shown above that the renormalisation group flow of the Hamiltonian leads to the emergence of an additional direction, quantified by the parameter x_j . This will now be used to construct a metric, in the following way. We already know that the scaling transformations lead to a graph of theories that are connected via edges that represent the mutual information between any two theories. As a result, this “information graph” can be mapped on to a new “distance graph” where the edges now represent the distance between the theories at the nodes. Finally, these distances can be used to define a metric for the distance graph: the metric between any two nodes will be the distance along the shortest path between these two nodes. This approach as well as some of the terminology are inherited from reference [Cao et al.(2017)Cao, Carroll, and Michalakis].

The RG flow can be represented as an “information graph” $G(V, E)$, where the systems $\mathcal{B}_j^1$ (defined above eq. (5.3)) constitute the set of vertices $V = \{V_j\}$ and the mutual information $I_z^2(j : j')$ between the systems $\mathcal{B}_j^1$ and $\mathcal{B}_{j'}^1$ constitute the edges E_{ij} between the vertices. This information

graph is characterised by an adjacency matrix whose elements are defined by the mutual information edge weights $I_z^2(j : j')$ [Singha Roy et al.(2020)Singha Roy, Santalla, Rodríguez-Laguna, and Sierra]. This graph structure is shown in Fig. (7.6), with the red spheres indicating the vertices. In order to define a metric space, we will map the graph G to another graph $\mathcal{G}$ that has the same vertices as G , but whose edges are weighted according to a well-defined notion of distance.

A path $\mathcal{P}$ in $\mathcal{G}$ is defined as a sequence of m vertices: $\mathcal{P}(V_1, V_2, \dots V_m)$. The distance along the path is defined as the sum of distances along every pair of consecutive vertices: $l(\mathcal{P}) = \sum_{i=1}^{m-1} d(V_i, V_{i+1})$, where $d(V_i, V_{i+1}) = d(i, i+1)$ (see eq. (5.1)). The *geodesic* between two points V_i and V_j is the path $\mathcal{P}$ which minimises the distance $l(\mathcal{P})$, and the distance along the geodesic then defines the *metric* $\tilde{d}(V_i, V_j)$ between two vertices V_i and V_j [Singha Roy et al.(2020)Singha Roy, Santalla, Rodríguez-Laguna, and Sierra, Cao et al.(2017)Cao, Carroll, and Michalakis]:

$$\tilde{d}(V_i, V_j) = \min_{\mathcal{P}} \{l(\mathcal{P})\} = \min_{\mathcal{P}} \left\{ \sum_{i=1}^{m-1} d(V_i, V_{i+1}) \right\}, \quad (5.11)$$

where the minimisation is carried out over all possible connected paths $\{\mathcal{P}\}$ between V_i and V_j , and m is the total number of vertices in a particular connected path $\mathcal{P}_m$. Following the expression for the distance $d(i, j)$ obtained in eq. (5.2), the metric takes the form

$$\tilde{d}(V_i, V_j) = \theta(z)j^z \ln 2 + \theta(-z)i^z \ln 2, \quad (i < j), \quad (5.12)$$

where we have assumed $i < j$, without loss of generality.

Finally, the metric dimension of the emergent space can be determined by finding the set with the lowest possible cardinality that can resolve the metric space [Cao et al.(2017)Cao, Carroll, and Michalakis, Bau and Beardon(2013)]. In order for a set to resolve the metric space, we must have $d(i, a) = d(j, a) \implies i = j$ for all points a in the set and for any pair of points i and j in the metric space. In our case, the singleton set $\mathcal{R} = \{0\}$ can resolve the metric space for $z > 0$, as $d(i, 0) = d(j, 0) \implies i^z = j^z \implies i = j$. Similarly, the singleton set $\mathcal{R} = \{\infty\}$ resolves the metric space for $z < 0$. As a result, we conclude that our emergent metric space is one-dimensional in nature.

7.6 Curvature of the emergent space

Further insight on the spatial geometry of the RG flows can be obtained by looking at the curvature generated by the RG transformations. In the plane of the directions x and y generated by the RG flow, the curvature can be written using the ideas of analytic geometry as

$$\kappa_z(j) = \frac{v'_z(j)}{[1 + v_z(j)^2]^{\frac{3}{2}}}, \quad (6.1)$$

where

$$\begin{aligned} v_z(j) &\equiv \frac{\Delta y_z(j)}{\Delta x_z(j)} = \frac{y_z(j+1) - y_z(j)}{x_z(j+1) - x_z(j)} \\ &= \begin{cases} \frac{(j+2)^z - (j+1)^z}{(j+1)^z - j^z} & z > 0, \\ \frac{(j)^z - (j-1)^z}{(j+1)^z - j^z} & z < 0, \end{cases} \end{aligned} \quad (6.2)$$

and

$$v'_z(j) \equiv \frac{v_z(j+1) - v_z(j)}{x_z(j+1) - x_z(j)}. \quad (6.3)$$

The curvature $\kappa_z(j)$ has been plotted against the RG step index j in right panel of Fig. 7.8. For $z = 1$, the first derivative $v_z(j)$ becomes unity, so the second derivative $v'_z(j)$ (and hence the curvature) vanishes, leading to a flat (Minkowski) space. For $z > 1$, the curvature becomes negative, while it is positive for $z < 0$. All the flows lead to an asymptotically flat space (i.e., the curvature vanishes) as $j \rightarrow \infty$.

The different signatures of the emergent curvatures can be attributed to the different ways in which the mass of the theory scales under the decimation procedure. For $z > 0$, the curvature defined in eq. (6.1) can be written in terms of the beta function:

$$\kappa_z(j) = \frac{[e^{\beta(j)} - 1]^{\frac{1}{2}} j^{-z}}{[e^{\beta(j)+\beta(j+1)} - 1]^{\frac{3}{2}}} \left[\frac{e^{\beta(j+2)} - 1}{1 - e^{-\beta(j+1)}} - \frac{e^{\beta(j+1)} - 1}{1 - e^{-\beta(j)}} \right]. \quad (6.4)$$

This equation relates directly a geometric quantity (i.e., the curvature) of the bulk emergent space with the RG beta function for the field theory on the boundary. From Fig. 7.7, as the beta functions flow towards zero, we see from eq. (6.4) that the first fraction inside the square brackets is the first to vanish, followed by the second fraction. The sequential vanishing of these two fractions prior to the vanishing of the denominator $e^{\beta(j)} - 1$ in eq.(6.4) leads to the asymptotic vanishing of the curvature towards the fixed point. Different choices of $z > 0$ can be thought to represent different relevant perturbations that drive the system towards *gapped fixed points*, leading to distinct curvatures of the bulk. By employing the relation $v_z(j) = \Delta S_{j+1,z} / \Delta S_{j,z}$, eq. (6.1) can also be cast in terms of the entanglement entropy $S_{j,z}$, $S_{j+1,z}$ and $S_{j+2,z}$. The resulting non-linear relationship between $\kappa_z(j)$ and $S_{j,z}$ obtained is, however, in contrast to the linear dependence obtained by Cao et al. [Cao et al.(2017)Cao, Carroll, and Michalakis] between entanglement perturbations on the Hilbert space and the curvature of the resultant holographic space.

7.6.1 Topological nature of the sign of curvature

We will now argue that the signature of the curvature κ corresponds to a topological quantum number, and that a change in the signature as z is tuned through 1 amounts to a topological translation. For this, we first point out (following eq. (6.1)) that the sign of the curvature is determined purely by the sign of $\gamma_z(j) \equiv 1 - v_z(j+1)/v_z(j)$ and $\alpha_z(j) = y_z(j+1) - y_z(j)$:

$$\begin{aligned} \kappa_z(j) &= \frac{v'_z(j)}{[1 + v_z(j)^2]^{\frac{3}{2}}} = - \frac{\alpha_z(j) \gamma_z(j)}{(\Delta x_z(j))^2 [1 + v_z(j)^2]^{\frac{3}{2}}} \\ \implies \text{sign} [\kappa_z(j)] &= -\text{sign} [\alpha_z(j)] \text{sign} [\gamma_z(j)]. \end{aligned} \quad (6.5)$$




Figure 7.9: Top: Variation of γ_z along the RG for multiple values of $z > 0$. All trajectories for $z > 1$ intersect the line $y = \gamma^*$. Bottom: Variation of α_z along the RG for multiple values of $z < 0$. All trajectories intersect the line $y = \alpha^*$

We now promote j to a continuous complex variable with the real part being greater than or equal to 1: $\text{Re}[j] \geq 1$. The quantities $\alpha_z(j)$ and $\gamma_z(j)$ can be written as

$$\alpha_z(j) = \begin{cases} [(j+2)^z - (j+1)^z] \log 2, & z > 0 \\ [j^z - (j-1)^z] \log 2, & z < 0 \end{cases}, \quad (6.6)$$

$$\gamma_z(j) = \begin{cases} 1 - \frac{[(j+3)^z - (j+2)^z][(j+1)^z - j^z]}{[(j+2)^z - (j+1)^z]^2}, & z > 0 \\ 1 - \frac{[(j+1)^z - j^z]^2}{[j^z - (j-1)^z][(j+2)^z - (j+1)^z]}, & z < 0. \end{cases} \quad (6.7)$$

As shown in Fig. 7.9(left panel), the function γ_z is positive and monotonically decreasing for $z > 1$ or $z \leq -1$, and vanishes identically for $z = 1$. On the other hand, $\alpha_z(j)$ is positive for $z > 1$, zero at $z = 1$ and negative for $z \leq -1$ (see Fig. 7.9(right panel)). Combining this, we find that the sign of the curvature can be written as:

$$\text{sign}[\kappa_z] = \begin{cases} -1, & z \geq 1 \\ 1, & z \leq -1 \end{cases} = \begin{cases} -\text{sign}[\gamma_z(j)], & z \geq 1 \\ -\text{sign}[\alpha_z(j)], & z \leq -1. \end{cases} \quad (6.8)$$

We start by considering the region $z \geq 1$. Since $z = 0$ is excluded by definition, the denominator is always non-zero. This indicates that $\gamma_z(j)$ is continuous and monotonically decreasing in this range: the maximum value of $\{\gamma_z(j), j \in [1, \infty]\}$ occurs at $j = 1$. Further, the maximum value $\gamma_z(1)$ increases with the value of z , ranging between $\gamma_2(2)$ at $z = 2$ and unity at $z \rightarrow \infty$. These two features are demonstrated in the left panel of Fig. 7.9, and imply that all curves $\{\gamma_z(j) : j\}$ will take values both greater than and less than the value $\gamma^* \equiv \gamma_2(2)$ (dotted line in left panel of Fig. 7.9). Thus, the plot shows that there always exists a real value j_z^* for any given integer value $z > 1$ such that $\gamma_z(j_z^*) = \gamma^* = \gamma_2(2)$. This is precisely what ceases to happen for $z = 1$, as that is the only curve in the left panel of Fig. 7.9 that does not pass through γ^* .

Indeed, the difference between the curvature for the $z = 1$ and $z > 1$ classes can be captured by expressing the presence of the solution j_z^* in the form of a winding number. For this, we use the




Figure 7.10: Top: Contours $\gamma_z(\mathcal{C})$ for multiple values of $z > 0$. As z increases, the contours extend in the positive $\text{Re}[\gamma_z]$ direction. Bottom: Contours $\alpha_z(\mathcal{C})$ for multiple values of $z > 0$. As z increases, the contours extend in the negative $\text{Re}[\gamma_z]$ direction.

fact that the sign of $\gamma_j(z)$ is given by the following integral:

$$\text{sign}[\gamma_z(j)] \Big|_{z \geq 1} = \frac{1}{2\pi i} \oint_{\mathcal{C}} dj \frac{\partial}{\partial j} \ln[\gamma_z(j) - \gamma^*], \quad (6.9)$$

where the contour $\mathcal{C}$ extends from $j = 1$ to $j = \infty$, and is shown in the left panel of Fig. 7.12. As the value of z is tuned from 2 to 1, the pole at j_z^* disappears and the integral reduces to zero. As shown in Fig. 7.10, the integral can thus be expressed as the winding number of the curve $\gamma_z(\mathcal{C}(j))$ that counts the number of times the contour $\gamma_z(\mathcal{C})$ winds around the singularity at γ^* :

$$\text{sign}[\gamma_z(j)] = \frac{1}{2\pi i} \oint_{\mathcal{C}} dj \frac{\partial(\gamma_z(j) - \gamma^*)}{\gamma_z(j) - \gamma^*} = \frac{1}{2\pi i} \oint_{\gamma_z(\mathcal{C})} \frac{d\gamma_z}{\gamma_z - \gamma^*}. \quad (6.10)$$

This is easily seen from the following argument. If we write a general complex variable y in the form $y = re^{i\theta}$ and consider a general contour C that starts from $r = r_0, \theta = 0$ and ends at $r = r_0, \theta = 2w\pi$ (such that the winding number of the curve is w), integrals of the form given above simplify and return the winding number itself:

$$\frac{1}{2\pi i} \oint_C \frac{dy}{y} = \frac{1}{2\pi i} \int_{r_0}^{r_0} \frac{dr}{r} + \frac{1}{2\pi} \int_0^{2w\pi} d\theta = w, \quad (6.11)$$

where we used $dy = y \left(\frac{dr}{r} + d\theta \right)$.

The winding numbers take integer values, and are topological in nature: they are robust against geometric deformations, and change only when the curve crosses the singularity at $\gamma = \gamma^*$. Thus, for $z > 1$, the contour $\gamma_z(\mathcal{C})$ encloses the singularity at γ^* exactly once, leading to a winding number of unity. As z is lowered towards $z = 2$, the contours becomes smaller such that, in order to go from $z = 2$ to $z = 1$, the contour has to be moved across the singularity. This constituting a change in the topology of the contour: for $z = 1$, the contour no longer encloses the pole and the winding number is zero. In fact, at $z = 1$, the contour itself (depicted in blue) is of vanishing radius because $\gamma_1(j)$ has a constant value of zero. To complete the connection, we use eq. (6.5) to relate this topological winding number to the sign of the curvature:

$$\text{sign}[\kappa_z] \Big|_{z \geq 1} = -\mathcal{W}_z(\gamma^*), \quad (6.12)$$




Figure 7.11: Schematic depiction of the emergent geometries in the cases of gapless (top) and gapped (bottom) RG flows. While the gapless RG flow involves a growth in the entanglement between the UV and IR spaces, the latter leads to the vanishing of UV-IR entanglement. This results in the emergence of connected and disconnected geometries, respectively.

where $\mathcal{W}_z(\gamma^*)$ is the winding number for $\gamma_z(\mathcal{C})$ around γ^* .

One can make a very similar argument for the case of $z \leq -1$. Here, as shown in the right panel of Fig. 3.10, the relevant quantity is $\alpha_z(j)$. For each curve α_z , the minimum value is at $j = 1$. For all z , this minimum starting value is largest for $z = -1$, and all subsequent curves $\alpha_z(j)$ start from even lower values. If we define a value $\alpha^* \equiv \alpha_z(j = 1)/2$, it is then clear that all curves $\alpha_z(j)$ will pass through the value α^* (dotted line in right panel of Fig. 3.10) at some value of j . This is very similar to the situation discussed just above for the case of $z \geq 1$ (with γ^* playing the role of α^*); we, therefore, employ the same technique here. Taking the same contour $\mathcal{C}$ as defined above, we write the sign of α_z as the winding number of the curve $\alpha_z(\mathcal{C})$ around the point $\alpha_z = \alpha^*$

$$\text{sign} [\alpha_z(j)] \Big|_{z \leq -1} = -\mathcal{W}'_z(\alpha^*) = \frac{1}{2\pi i} \oint_{\alpha_z(\mathcal{C})} \frac{d\alpha_z}{\alpha_z - \alpha^*} . \quad (6.13)$$

For $z > 0$, α_z is positive, and the curve $\alpha_z(\mathcal{C})$ encloses points in the positive half of the real axis such that the winding number in the right panel of Fig. 7.10 is then zero. This indicates that the function $1 - 2\mathcal{W}'_z(\alpha^*)$ is unity for $z > 0$, leaving any other function in product with it unchanged. On the other hand, it is -1 for $z < 0$, returning the correct sign of $\alpha_z(j)$ in that range. Combining with eq. (6.12), the sign in the entire range can be expressed as

$$\text{sign} [\kappa_z] = \mathcal{W}_z(\gamma^*) \times [2\mathcal{W}'_z(\alpha^*) - 1] . \quad (6.14)$$

We can now summarise the topological nature of the sign of the curvature. For $z > 0$, the second winding number $\mathcal{W}'$ is 0. As z varies from 2 to 1, the other winding number $\mathcal{W}$ changes from 1 to 0, constituting the first topological change. In contrast, upon varying z from 1 to -1 , both winding numbers change: $\mathcal{W}$ from 0 to 1, and $\mathcal{W}'$ from 0 to 1. This constitutes the second topological change. Together, the two changes show that going from an open geometry (negative curvature) to a closed geometry (positive curvature) involves two topological transformations.

7.7 Entanglement Holography and Fermionic Criticality

7.7.1 Equivalence between a critical Fermi surface and a holographic worm-hole

As mentioned previously under eq. (4.3), the quantity z corresponds to the anomalous dimension of the spectral gap g_z in the effective field theory: a change in the sign of z corresponds to a change in the nature of the RG flow from relevance to irrelevance and vice-versa. The holographic nature of the RG flow dictates that such a change arises by a variation in some quantity in the parent interacting theory from which the effective field theory of non-interacting Dirac fermions can be thought to have originated. Indeed, such a change in the RG relevance has drastic consequences: a relevant spectral distance g_z implies a gap in the excitations at the fixed point, while an irrelevant g_z indicates the presence of gapless excitations. The presence or absence of a gap in the low-energy excitations determines the topology of the electronic Fermi surface through the change of a topological invariant, the Luttinger volume of the system [Luttinger(1960), Oshikawa(2000), Seki and Yunoki(2017), Mukherjee and Lal(2020a), Mukherjee and Lal(2020b)] (also see eq. (2.19)).

At the transition between the gapless (metallic) and gapped (insulating) states, the Fermi surface becomes critical, signalled by the vanishing lifetime of the gapless excitations close to it. Our formulation shows that such a gap-inducing transition and its associated quantum critical Fermi surface has consequences for the space generated along the RG flow. For the gapless case ($z < 0$), the degrees of freedom in the UV and the IR have finite entanglement, indicating that they are part of the same connected holographic space (see Figs. 7.6 and 7.7), while for the gapped case ($z > 0$), the entanglement vanishes into the IR, leading to a decoupling of these two regions and their separation into disconnected spaces [Van Raamsdonk(2010)]. These two scenarios are shown schematically in Fig. 7.11. It is therefore tempting to conclude that at the transition between the gapped and gapless states, a *wormhole geometry* connects the UV and the IR via some minimal entanglement [Van Raamsdonk(2010), Cao et al.(2017) Cao, Carroll, and Michalakis]. Our results also suggest that the presence of such a wormhole would then coincide with two topological transitions, the first being the change in the sign of the curvature (as shown in the previous subsection), and the second being the change in the topological Luttinger volume of the underlying quantum field theory (as discussed in the previous paragraph). We end by noting that we have provided, in a later section of this work, additional support for the topological nature of Luttinger's volume and its connection to the holographically generated space by relating it to multipartite entanglement (see eq. (9.17)).

7.7.2 Connecting modular energy with the metric

The notion of an entanglement Hamiltonian was introduced in eq. (2.16). At the transition between the gapless (metallic) and gapped (insulating) states, the system becomes critical, and the Dirac electrons are described by a CFT. It turns out that for such a CFT, the entanglement Hamiltonian takes a particularly simple form. In this subsection, we will relate the emergent metric to the entanglement Hamiltonian obtained at the IR fixed point, focusing on the theory at the transition.

For a 1D CFT, the entanglement Hamiltonian corresponding to a partition of length l is given

by [Bisognano and Wichmann(2008a), Bisognano and Wichmann(2008b)]

$$\mathcal{H}_{1D} = \frac{2\pi L}{c} \int_0^l dx \frac{L^2 - x^2}{2L^2} T_{00}(x), \quad (7.1)$$

where $T_{\mu\nu}$ is the stress-energy tensor of the CFT and c is the speed of light. We have already shown in eq. (2.17) that the entanglement Hamiltonian for the 2D theory is simply a sum of the 1D counterparts, owing to the decoupling of 1D modes in the Lagrangian. Working in a region where T_{00} is approximately uniform, the 2D entanglement Hamiltonian can be expressed as

$$\mathcal{H}_{2D} \sim \sum_n \mathcal{H}_{1D} = \frac{2\pi L^2}{3c} \sum_n T_{00}^n, \quad (7.2)$$

where T_{00}^n is the average value of T_{00} within the partition for the CFT associated with the n^{th} mode.

In order to make the connection with the emergent geometry, we use the entanglement first law [Blanco et al.(2013)Blanco, Casini, Hung, and Myers], which states that at first order, small variations ΔS of the entanglement entropy can be equated to the associated change in the expectation value of the entanglement Hamiltonian:

$$\Delta S(A) = \Delta I_2(A_0 : A) = \langle \Delta \mathcal{H} \rangle = \frac{2\pi L^2}{3c} \sum_n \langle T_{00}^n \rangle, \quad (7.3)$$

where $\langle \mathcal{H} \rangle = \text{Tr}(\rho \mathcal{H})$ is often referred to as the modular energy, and we have equated $\Delta S(A)$ and $\Delta I_2(A_0 : A)$ following eq. (3.16). Further, since the entanglement at the critical point is expected to be large, we can approximate the expression for the metric between points (i, j) obtained in subsection 7.5.2 as $\tilde{d}(i, j) = -\ln I^2(i : j)/I_{\text{max}}^2 \simeq |I^2(i : j)/I_{\text{max}}^2 - 1|$. Combining this equation with eq. (7.3) leads to the following interesting relation:

$$\Delta \tilde{d}(i, j) \simeq \frac{2\pi L^2}{3c} \sum_n \langle T_{00}^n \rangle. \quad (7.4)$$

The equation relates the change in a geometric measure (the emergent metric) with a mass/energy term. An analogous relation was obtained for states for which the entropy obeys an area law by Cao et al [Cao et al.(2017)Cao, Carroll, and Michalakis], connecting the spatial curvature and the stress tensor, ultimately leading to a semi-classical Einsteins field equation for gravity. Our formulation does not however lead to a relation involving the curvature, and we speculate that this is due to the violation of the area law in entanglement of fermionic critical systems. Instead, we show in the next section that the curvature in our formulation is related to the rate of convergence of the RG flows.

7.7.3 Kinematics of the RG flows

Further statements on the RG flow and the emergent holographic dimension can be made by studying quantities that can serve as candidates for the convergence parameter θ for the RG flows of




Figure 7.12: Top: The contour $\mathcal{C}$ used in eq. (6.9). It extends across all values of j , and the integral counts the number of poles of the function $1/(\gamma - \gamma^*)$ encircled by the contour. Bottom: Evolution of expansion parameter θ_z with distance x_z , leading to the fixed point at $\theta_z^* = 1/\sqrt{2}$.

g_z [Kar(2001), Kar and Sengupta(2007)]. We consider the following function as a candidate for the convergence parameter for the case of $z > 0$:

$$\theta_z(j) = \frac{1}{\sqrt{v_z^{-2} + 1}} = \left[1 + \left\{ \frac{1 - e^{-\beta_z(j)}}{e^{\beta_z(j+1)-1}} \right\}^2 \right]^{-\frac{1}{2}}. \quad (7.5)$$

The parameter θ_z has been plotted in Fig. 7.12, and the behaviour is observed to be very similar to that of the beta function shown in Fig. 7.7. Indeed, this similarity describes the holographic nature of the RG flow: the emergent geometry (as seen through θ_z) is dictated by the RG flow described by β .

The quantity θ can be interpreted as an estimate of the rate of change of area enclosed by a fixed number of RG trajectories. To see that this is the case, we note that by using $v_j = \Delta y / \Delta x$, the expansion parameter can be written as $\theta = \Delta y(j) / \sqrt{\Delta x^2 + \Delta y^2} \sim \Delta g(j+1) / \sqrt{\Delta x^2 + \Delta y^2}$. Considering RG flows for $g_z(j)$ corresponding to multiple z , the area enclosed by the curves is given by the maximum g_z , and Δg_z is then the change in the area under the RG transformation. The denominator $\sqrt{\Delta x^2 + \Delta y^2}$ is the distance covered in the emergent spacetime, rendering θ as the rate of change of the area.

The equation of motion for this expansion parameter can then be obtained as:

$$\frac{d\theta_z}{dx} = \frac{v_z^{-3}}{(1 + v_z^{-2})^{\frac{3}{2}}} \frac{dv_z}{dx_j} = \kappa_z. \quad (7.6)$$

The simple form of this relation makes it clear that for relevant flows (corresponding to a negative curvature $\kappa_z < 0$), θ_z flows to a minimum. Following Fig. 7.8, this is accompanied by the minimisation of the c -function to zero. It is, therefore, tempting to conclude that eq. (7.6) for the spatial evolution of the convergence parameter θ is dual to eq. (5.10) that controls the flow of the c -function of the quantum theory along the RG flow.

7.8 Extension to the massive case

We will now generalise several of the main results obtained above to the case of two-dimensional electrons with a single-particle mass-gap $M_{n,\phi} = \sqrt{m^2 + \frac{4\pi^2}{L_x^2}(n+\phi)^2}$. Given that they depend only on the geometric part of the entanglement (which remains unchanged on introducing the mass $M_{n,\phi}$), the following set of key qualitative results and observations discussed earlier continue to hold for the case of a dispersion with a mass-gap:

- the hierarchical structure of the entanglement content (e.g., the area-law term in eqns.(3.14) and (3.17)) and the fact that the entanglement satisfies the Ryu-Takayanagi bound (right panel of Fig.7.5),
- the dependence of the emergent distances x and y on the RG flow (Fig.7.6, eqns.(5.2) and (5.3)), and the fact that the RG beta function tracks the central charge (Fig.7.8),
- various qualitative and quantitative properties of the curvature (e.g., eq.(6.1)), including the topological nature of its sign (eq.(6.14)), and
- the existence of a convergence parameter that is related to the RG flow, and the equation that governs the evolution of the convergence parameter (eq.(7.6)).

The functional forms of certain other results do, however, undergo changes. For instance, the coupling $g_z(j)$ is given by the (more general) expression

$$\begin{aligned} g_z(n, j) &= \ln \frac{M_{n+1,\phi}(j) - M_{n,\phi}(j)}{2\pi/L_x} \\ &= \ln \left[\sqrt{\tilde{m}^2 + (t_z(j)(n+1) + \phi)^2} - \sqrt{\tilde{m}^2 + (t_z(j)n + \phi)^2} \right], \end{aligned} \quad (8.1)$$

where $\tilde{m} = mL_x/2\pi$ is a dimensionless parameter. We will work with the coupling for the mode $n = 0$:

$$g_z(0, j) = \ln \left[\sqrt{\tilde{m}^2 + (t_z(j) + \phi)^2} - \sqrt{\tilde{m}^2 + \phi^2} \right]. \quad (8.2)$$

In order to simplify the analysis, we work in the large mass limit $\frac{t_z + \phi}{\tilde{m}} \rightarrow 0$. In this limit, upon expanding the square root and retaining only the lowest order term, we obtain

$$g_z(0, j) = \ln \frac{[t_z(j) + \phi]^2 - \phi^2}{2\tilde{m}}. \quad (8.3)$$

The RG beta function $\beta = \ln(g_z(j+1)/g_z(j))$ can, in the large mass limit, then be related to the emergent distances x and y in terms of the coupling $g_z(j)$:

$$x_z(j) = \ln t_z(j) = \ln \left[\sqrt{2\tilde{m} e^{g_z(0,j)} + \phi^2} - \phi \right], \quad (8.4)$$

$$y_z(j) = \begin{cases} \ln \left[\sqrt{2\tilde{m} e^{g_z(0,j+1)} + \phi^2} - \phi \right], & z > 0, \\ \ln \left[\sqrt{2\tilde{m} e^{g_z(0,j-1)} + \phi^2} - \phi \right], & z < 0. \end{cases} \quad (8.5)$$




Figure 7.13: Contour C_j used to integrate eq. (9.3). The contour is chosen so as to encircle the simple pole of $1/Q$ at $\phi = j - 1/4$.

The quantitative dependence of the c -function on the RG beta function can also be expressed more generally as

$$e^{\beta_z(j)} = \left[\ln \frac{\left(\frac{\tilde{c}(j+1)}{c} + \phi \right)^2 - \phi^2}{2\tilde{m}} \right] \left[\ln \frac{\left(\frac{\tilde{c}(j)}{c} + \phi \right)^2 - \phi^2}{2\tilde{m}} \right]^{-1}. \quad (8.6)$$

Quantities such as the curvature κ and the convergence parameter θ can similarly be expressed in terms of the coupling $g_z(j)$.

7.9 Topological content of the entanglement spectrum

7.9.1 Luttinger's volume as a winding number: the massless case

We will now show how to obtain Luttinger's volume (eq.(2.18)) from the g -partite information entanglement measure I^g of a system of massless 2D electrons. The flux-dependent part of the g -partite information I^g can be used to probe the k -space volume V_L within the Fermi surface. We define a quantity $Q_g(\phi)$ that is periodic in the flux ϕ and independent of the geometry:

$$Q_g(\phi) = f_{g,z} \left[\frac{1}{\sqrt{2}} - \frac{1}{2} e^{-\frac{1}{c} (I_{\{\mathcal{A}_z(i)\}}^g(\phi) - I_{\{\mathcal{A}_z(i)\}}^g(\phi^{(0)} f_{g,z}))} \right] = f_{g,z} \left[\frac{1}{\sqrt{2}} - |\sin(\pi \phi f_{g,z})| \right], \quad (9.1)$$

where $\phi^{(0)} = \frac{1}{6}$ is the value at which the flux-dependent part of $S_0(\phi)$ vanishes. Further, from eq.(3.19), we have

$$I_{\{\mathcal{A}_z(i)\}}^g(\phi_g) - I_{\{\mathcal{A}_z(i)\}}^g(\phi_g^{(0)} f_{g,z}) = -c \ln(2 |\sin(\pi \phi f_{g,z})|). \quad (9.2)$$

Note that the non-differentiability of $Q_g(\phi)$ at $\phi f_{g,z} = 0, 1, 2, \dots$ is not a concern as ϕ is chosen to be non-integral, ensuring that $\phi f_{g,z}$ is also non-integral.

In order to express the number N_L^x of positive k_x -states available inside the Fermi surface in the x -direction in terms of $Q_g(\phi)$, we need to establish a relation between the flux ϕ and the Fermi surface. The k_x -states of interest to us are given by $k_x^n = \frac{2\pi n}{L_x}$, $n \in \mathbb{Z}$, $0 \leq k_x^n \leq k_x^F$, where k_x^F

is the x -component of the Fermi momentum $\vec{k}_F$. Assuming the flux-free Hamiltonian has time-reversal symmetry, there should be an equal number of states with the opposite momenta. Let n^* be the integer for which $k_x^{n^*} \leq k_x^F$ and $k_x^{n^*+1} > k_x^F$. Since the flux couples to the k_x -states as $k_x^n \rightarrow k_x^n + eA = \frac{2\pi n + 2\pi\phi}{L_x}$, tuning ϕ by 1 leads to the entire spectrum of k_x -states being shifted by one quantum number: $k_x^n \rightarrow \frac{2\pi(n+1)}{L_x} = k_x^{n+1}$. This means that the state $k_x^{n^*}$ moves to $k_x^{n^*+1}$, and hence out of the Fermi volume by crossing k_x^F . This flow of the spectrum is shown in Fig. 7.15.

If we now connect the system to a particle reservoir with the chemical potential set to zero, any k_x -state that moves out of the Fermi volume becomes unoccupied. We continue tuning the flux to a value ϕ^* , where the system becomes entirely devoid of any particles. The number of k_x -states that are below k_x^F can then be said to be given by the number of times ϕ goes through an integer value while changing from 0^+ to $\phi^* + 0^+$. Since we do not set the flux to an integer value, we can equivalently count the number of times ϕ goes through, say, three-quarters of an integer - $j - \frac{1}{4} : j = 1, 2, \dots$ - in changing from 0^+ to $\phi^* - 0^+$. We will now show that this count can be expressed in terms of Q_g . We begin by considering $Q_0(\phi) = \frac{1}{\sqrt{2}} - |\sin \pi\phi|$. At every three-quarter value of the flux, the quantity $\phi_j = j - \frac{1}{4} : j = 1, 2, \dots$, $|\sin \pi\phi_j|$ attains the value of $\sin \frac{\pi}{4} = \frac{1}{\sqrt{2}}$, such that $Q_0^{-1}(\phi)$ goes through a simple pole. Each such pole ϕ_j can be picked up via a complex integral around a contour $\mathcal{C}_j$ that encloses the pole:

$$-f_j \frac{1}{2\sqrt{2}i} \oint_{\mathcal{C}_j} \frac{d\phi}{Q_0(\phi)} = 1, \quad (9.3)$$

where f_j is either +1 or -1 and is defined through the equation $|\sin(\pi(j - \frac{1}{4}))| = f_j \sin(\pi(j - \frac{1}{4}))$, and $\mathcal{C}_j$ is a circle of diameter 1/4 centered at $\phi = j - \frac{1}{4}$ (shown in Fig. 7.13). The evaluation of the integral eq.(9.3) is shown in Appendix B. In order to drop the sign prefactor f_j , we reorient those contours for which f_j is +1, and such that they wind around in the clockwise direction. The contours are shown in the left panel of Fig. 7.14, and defined as

$$\mathcal{C}_j(\phi) = \left\{ \phi : \phi = j - \frac{1}{4} + \frac{1}{8}e^{i\theta}, \theta \in [0, -2\pi f_j] \right\}, \quad (9.4)$$

such that they host a singularity at their center. With this redefinition of the contours, we have

$$\frac{1}{2\sqrt{2}i} \oint_{\mathcal{C}_j} \frac{d\phi}{Q_0(\phi)} = 1. \quad (9.5)$$

Counting the number of three-quarter values then reduces to counting the number of poles of Q_0^{-1} in the above-mentioned range $\phi \in (0, \phi^*)$. In this way, we can now express the count N_L^x in terms of Q_0 :

$$N_L^x = \frac{1}{2} \sum_{j=1}^{\phi^*} \frac{1}{2\sqrt{2}i} \oint_{\mathcal{C}_j(\phi)} \frac{d\phi}{Q_0(\phi)} = \frac{1}{2\sqrt{2}} \sum_{j=1}^{\phi^*} \frac{1}{2i} \oint_{\mathcal{C}_j(\phi)} \frac{d\phi}{Q_0(\phi)}. \quad (9.6)$$

The factor of half in eq.(9.6) is required as we want to count only the number of positive momentum states (and which is equal to the number of negative momentum states).

In going from $Q_0(\phi)$ to a general $Q_g(\phi)$, the only change required is to redefine $\phi \rightarrow \phi_g = f_{g,z}\phi$. This implies that while the states in k -space have become more coarse-grained, the flux ϕ has become more fine-tuned by the same scaling factor. This is a reflection of the overall conservation of particle number density: keeping the chemical potential fixed, the increased spacing between the k_x -states implies that a smaller number of particles are now occupying an equally smaller volume. As the number density is conserved, so is the Luttinger volume (eq.(2.18)), and this shows that our approach is consistent with the holographic Luttinger theorem [Sachdev(2011)]. Thus, given that $\phi_g^* = \phi^*$, the expression for N_L^x remains practically unchanged:

$$N_L^x = \frac{1}{2\sqrt{2}} \sum_{j=1}^{\phi^*} \frac{1}{2i} \oint_{C_j(\phi_g)} \frac{d\phi_g}{\frac{1}{\sqrt{2}} - |\sin \pi \phi_g|}. \quad (9.7)$$

In order to rewrite N_L^x in terms of the flux ϕ , we start by making the substitution $\phi_g \rightarrow f_{g,z}\phi$:

$$N_L^x = \frac{1}{2\sqrt{2}} \sum_{j=1}^{\phi^*} \frac{1}{2i} \oint_{C_j(f_{g,z}\phi)} \frac{f_{g,z} d\phi}{\frac{1}{\sqrt{2}} - |\sin(\pi f_{g,z}\phi)|}. \quad (9.8)$$

In the above equation, we can recognise $\frac{f_{g,z}}{\frac{1}{\sqrt{2}} - |\sin(\pi f_{g,z}\phi)|}$ as $Q_g^{-1}(\phi)$. Also note that the contour can be written as

$$C_j(f_{g,z}\phi) \equiv C_j^g(\phi) = \left\{ \phi : \phi = t_{g,z} \left(j - \frac{1}{4} + \frac{e^{i\theta}}{8} \right), \theta \in [0, -2\pi f_j] \right\}. \quad (9.9)$$

These curves are again circles of radius $1/8$, but centered at $t_{g,z} \times (1 - \frac{1}{4})$, $t_{g,z} \times (2 - \frac{1}{4})$, $t_{g,z} \times (3 - \frac{1}{4})$ and so on. These are simply a generalisation of the curves defined in eq. (9.4); the curves for $t_{g,z} = 2$ are shown in the right panel of Fig. 7.14. Substituting this in eq.(9.8) gives

$$N_L^x = \frac{1}{2\sqrt{2}} \sum_{j=1}^{\phi^*} \frac{1}{2i} \oint_{C_j^g(\phi)} \frac{d\phi}{Q_g}. \quad (9.10)$$

For a spatially isotropic system, the Fermi volume is given by

$$V_L = v_2 \times \pi (N_L^x)^2 = \frac{1}{2} \frac{\pi^3}{V} \left[\sum_{j=1}^{\phi^*} \frac{1}{2i} \oint_{C_j^g(\phi)} \frac{d\phi}{Q_g(\phi)} \right]^2, \quad (9.11)$$

where V is the real space volume of the system, and $v_d = (2\pi)^d/V$ is the volume of the smallest unit in k -space. For a generally anisotropic Fermi volume, each direction in the d -dimensional k -space can be completely specified by an angular coordinate θ with respect to some predefined coordinate system. For each direction $\hat{\theta}$, we can insert a gauge field $\vec{A} = A\hat{\theta}$ and take the thin torus limit in this direction, obtaining thereby the g -partite information $I^g(\phi_\theta)$ for length l in the direction perpendicular to $\hat{\theta}$. With the θ -dependent I^g , one can then define the quantity $Q_g(\phi_\theta)$




Figure 7.14: Top: Contours C_j used to integrate over Q_0^{-1} in eq. (9.6). Tall vertical lines represent integer values of $\text{Re}(\phi)$. All contour start and end within two adjacent integer values. The final contour C_{ϕ^*} ends just short of $\text{Re}(\phi) = \phi^*$. Note the alternating direction of the contours brought about by the alternation of f_j in going from $j = 1$ to $j = 2$, and so on. Bottom: Contours C_j^g for $t_{g,z} = 2$, centered at $2(1 - \frac{1}{4})$, $2(2 - \frac{1}{4})$ and $2(3 - \frac{1}{4})$. Note the alternating directions of the contours.

which counts the number of states N_L^θ in the direction $\hat{\theta}$ inside the Fermi volume. In this way, the total Fermi volume can be obtained by integrating over θ :

$$V_L = \frac{\pi^2}{4V} \int_0^{2\pi} d\theta \left[\sum_{j=1}^{\phi_\theta^*} \frac{1}{2\pi i} \oint_{C_j^g(\phi)} \frac{d\phi}{Q_g(\phi)} \right]^2. \quad (9.12)$$

It is important to note that the quantity V_L is independent of the RG index g , and is thus identical in value for all g members of the hierarchy at the g th step of the coarse-graining process.

The topological nature of Luttinger volume can be made further manifest by expressing it in terms of winding numbers, following Ref. [Seki and Yunoki(2017)] for the case of interacting electrons. To do so, we point out that the integrand $Q_g^{-1}(\phi)$ can be written as a derivative:

$$\frac{1}{Q_g(\phi)} = \frac{\sqrt{2}}{\pi} \frac{\partial}{\partial \phi} \ln Y_j^g = \frac{\sqrt{2}}{\pi} \frac{1}{Y_j^g} \frac{\partial Y_j^g}{\partial \phi}, \quad (9.13)$$

where

$$Y_j^g(\phi) \equiv \left[\frac{f_j \tan 3\pi/8 - \tan(\pi f_{g,z}\phi/2)}{\tan(\pi f_{g,z}\phi/2) - \tan(\pi/8)} \right]^{f_j} \\ = \begin{cases} \frac{\tan 3\pi/8 - \tan(\pi f_{g,z}\phi/2)}{\tan(\pi f_{g,z}\phi/2) - \tan(\pi/8)}, & \text{when } j = \text{odd, and} \\ \frac{\tan(\pi f_{g,z}\phi/2) + \tan(\pi/8)}{-\tan 3\pi/8 - \tan(\pi f_{g,z}\phi/2)}, & \text{when } j = \text{even.} \end{cases} \quad (9.14)$$

Substituting this into the expression for 2-dimensional isotropic Luttinger volume in eq. (9.11), we obtain

$$V_L = \frac{\pi^3}{V} \left[\sum_{j=1}^{\phi^*} \frac{1}{2\pi i} \oint_{Y_j^g(C_j^g)} \frac{dY_j^g}{Y_j^g} \right]^2. \quad (9.15)$$

As before, the specific integral in the above expression is a winding number that counts the number of times the contour $Y_j^g(C_j^g)$ winds around the singularity at $Y_j^g = 0$. Note that these are the same




Figure 7.15: Spectral flow of the states within the Fermi volume. The red curve represents the Fermi surface. Upon tuning the flux ϕ , the k -space states (indicated in the form of dotted lines along the k_x -axis) are shifted towards the Fermi surface. The blue circles represent electrons with momenta below k_F , and are part of the ground state. On the other hand, the grey circles represent particles with momenta above k_F : they have moved out of the system, and into the reservoir connected to it (see discussion in main text). Each time ϕ changes by 1, the state nearest to the Fermi surface moves out of the Fermi volume. Tuning ϕ by n^* empties the entire Fermi volume.

singularities that exist at the centers of the contours $\mathcal{C}_j^g$ and that are picked up in, for example, eq. (9.3). To demonstrate this, we point out that the centers of the contours are given by $\phi = t_{g,z} (j - \frac{1}{4})$, $j = 1, 2, \dots, \phi^*$. At these values, we have

$$\tan\left(\frac{\pi f_{g,z}\phi}{2}\right) = \tan\left(\frac{\pi j}{2} - \frac{\pi}{8}\right) = \begin{cases} \tan \frac{3\pi}{8}, j = 1, 3, \dots \\ -\tan \frac{\pi}{8}, j = 2, 4, \dots \end{cases} \quad (9.16)$$

Comparing with eq. (9.14), we can see that the numerators of Y_g^j are zero at these values of ϕ for both even and odd values of j . This leads to $Y_g^j = 0$, and hence a singularity in $\frac{dY_g^j}{Y_g^j}$. These values of the flux are therefore the common singularities that are picked up by the poles of $Q_g^{-1}(\phi)$ as well as the winding numbers.

We now define w_g^j as the winding number for the curve $Y_g^j(\mathcal{C}_j^g)$ around $Y_g^j = 0$, and $\mathcal{W}(\phi^*) = \sum_{j=1}^{\phi^*} w_g^j$ as the total winding number for all the curves $\{Y_g^j(\mathcal{C}_j^g), j = 1, 2, \dots\}$. Then, the isotropic Fermi volume takes the compact form

$$V_L = \frac{\pi^3}{V} \mathcal{W}(\phi^*)^2. \quad (9.17)$$

7.9.2 A relation between the quantum hall Chern numbers and Luttinger's volume

Interestingly, the above method employed in using entanglement measures to extract the Luttinger volume can also be applied (with suitable modifications) to an integer quantum hall system (generated upon placing the system in a strong transverse magnetic field) to obtain the number of filled Landau levels (a Chern number). To begin, we imagine a system of free non-relativistic electrons in a cylinder geometry (fig. 7.16), and placed in a magnetic field $\nabla \times \vec{A} = \vec{B}$ that points radially outwards. The field $\vec{B}$ generates Φ flux quanta through the surface of the cylinder. An Aharonov-Bohm flux $\vec{A}' = -2\pi\phi/L_x\hat{x}$ is also placed in the x -direction (i.e., along the circumference of

the cylinder). We divide the cylinder into two subsystems A and B (fig. 7.16), such that tuning the dimensionless flux ϕ by 1 transfers a charge equal to the first Chern number, from A to B . The translation invariance in the x -direction ensures that the momentum k_x are good quantum numbers; for each value of k_x , we can define the matrix elements of the single-particle correlation matrix $G(k_y)$ for the subsystem A as [Peschel(2003), Alexandradinata et al.(2011)Alexandradinata, Hughes, and Bernevig]

$$G_{ij}^A(k_x) = \langle \psi_{\text{gs}} | c_{k_x,i}^\dagger c_{k_x,j} | \psi_{\text{gs}} \rangle , \quad (9.18)$$

where $|\psi_{\text{gs}}\rangle$ is the ground state and i, j represent the degrees of freedom other than k_x . The total matrix G^A is constructed by joining the smaller blocks $G^A(k_x)$. As ϕ is varied, the variation of the trace of the flux-dependent matrix $G^A(\phi)$ is linear in the flux ϕ : $\text{Tr}[G^A(\phi)] - \text{Tr}[G^A(0)] = -\frac{1}{e}\sigma_H\phi$. This implies that as long as there are extended states at sufficiently small energies, we will have spectral flow that can be captured through the integral:

$$-\frac{\sigma_H}{2\pi i e} \int_{\mathcal{C}_H} \frac{d\phi}{\text{Tr}[G^A(\phi)] - \text{Tr}[G^A(0)]} , \quad (9.19)$$

where $\mathcal{C}_H$ is a circle of radius $1/2$ in the complex plane centered at $\phi = 0$. This integral is zero if there is no spectral flow, i.e., the effect of the gauge field can be removed by a gauge transformation and the trace is insensitive to the flux. In order to count the number of filled Landau levels (essentially, the filling factor ν), we can now vary the flux quanta Φ (due to $\vec{B}$) so as to change the filling from ν to 0:

$$\nu = -\frac{\sigma_H}{2\pi i e} \sum_{\Phi'=\Phi}^{\Phi^{\max}} \int_{\mathcal{C}_H} \frac{d\phi}{\text{Tr}[G_{\Phi'}^A(\phi)] - \text{Tr}[G_{\Phi'}^A(0)]} . \quad (9.20)$$

The subscript Φ' on G^A indicates that the correlations depend on the variable magnetic field $\vec{B}'$. The upper limit $\Phi^{\max}$ is given by the lowest value of the magnetic field $\vec{B}'$ for which only one Landau level is filled. The magnetic field is varied such that exactly an integer number of Landau levels are filled at any given value. Importantly, the filling factor ν is known to be a Chern number $\mathcal{C}(\Phi)$ [Thouless et al.(1982)Thouless, Kohmoto, Nightingale, and Den Nijs, Niu et al.(1985)Niu, Thouless, and Wu].

We can also relate eq. (9.20) to the entanglement of the system by introducing the entanglement spectrum [Li and Haldane(2008), Laflorencie(2016a)], which refers to the set of eigenvalues $\{\lambda_\chi^A(k_x)\}$ of the reduced density matrix $\rho^A(k_x)$ for subsystem A corresponding to the momentum mode k_x in the x -direction and χ is a particular configuration of the degrees of freedom $\{i\}$ in the y -direction. We have the relation [Alexandradinata et al.(2011)Alexandradinata, Hughes, and Bernevig]

$$\lambda_\chi^A(k_x) = \text{Det} [P_\chi G^A(k_x) (1 - P_\chi) G^A(k_x)] , \quad (9.21)$$

where P_χ is an operator that projects on to the subset of states $\{j\}$ in $\{i\}$ that are occupied in the configuration χ . This relation between G^A and the eigenvalues λ then leads to the same conclusion as in the case without the magnetic field: the pole structure of the entanglement spectrum encodes the topological information of the system revealed in eq.(9.20). In the case of zero magnetic field, the topological quantity revealed from the pole structure was a winding number ($\mathcal{W}$), while in the present case it is a Chern number.




Figure 7.16: Cylindrical geometry for the integer quantum Hall system, with a magnetic field $\vec{B}$ pointing radially outwards through the surface of the cylinder. An Aharonov-Bohm vector potential acts in the x -direction (i.e., along the circumference). The entanglement is calculated between the subsystem A and its complement B .

Finally, we reveal the connection between the winding numbers $\mathcal{W}$ in the metallic case and the Chern numbers in the integer quantum Hall system. The degeneracy of each Landau level is equal to the number of flux quanta Φ generated by the magnetic field B . Since the total number of particles in the system is fixed, the electronic states of the filled Fermi sea that form the Luttinger volume (in the absence of the external B-field) are redistributed into the Landau levels with the appropriate degeneracy as a strong transverse magnetic field is turned on. Thus, we have the constraint that the total number of occupied states $\mathcal{C} \times \Phi$ in the quantum hall system ($\mathcal{C}$ being the Chern number and equal to the number of filled Landau levels) must be equal to $\pi (N_L^x)^2$, the total number of occupied states in the metallic system. Combining eqs. 9.10 and 9.17 with this discussion then allows us to relate $\mathcal{C}$ to the winding number $\mathcal{W}(\phi^*)$:

$$\mathcal{C} \times \Phi = \left\lfloor \frac{\pi}{4} \mathcal{W}(\phi^*)^2 \right\rfloor = \frac{V_L}{v_2}, \quad (9.22)$$

where $v_2 = 4\pi^2/V$ is the volume of the k -space unit cell (defined below eq. (9.10)) and $\lfloor \cdot \rfloor$ is the floor function and returns the largest integer less than or equal to the argument.

7.10 Conclusions and Outlook

In summary, we have shown that considering the renormalisation group flows of the spectral gap coupling of a system of non-interacting electrons in $2 + 1$ -dimensional theory leads to hierarchical arrangement of the entanglement with scale, and that the geometrical part of the entanglement can be described as the emergence of an additional spatial dimension, leading to holographic expressions for geometric parameters like distance and curvature. We show that changing the boundedness of the holographic space involves a topological transition with the Fermi surface of the boundary theory turning critical, and involves the generation of a geometric wormhole between the UV and IR degrees of freedom. For the massless case, the presence of an Aharonov-Bohm flux generates a geometry-independent term in the entanglement entropy which can be linked to the Luttinger volume of the gapless fermionic system. By placing the system of $2 + 1$ dimensional electrons in a strong magnetic field, we argue that the Chern number topological invariant for the ground states of the integer quantum Hall ground system are constrained by the value of the Luttinger volume in the underlying metallic system (i.e., prior to the insertion of the field).

How robust are our conclusions towards the inclusion of electronic correlations? Given the adiabatic continuity that links Fermi liquids to the non-interacting Fermi gas, i.e., the preservation of the k -space quantum numbers under adiabatic inclusion of (screened) inter-particle interactions, we expect our results to be equally valid for the many-particle entanglement of the excitations described by the fixed point effective theory for the Fermi liquid. This also suggests that our results for the wormhole geometry and the stress-energy tensor relation at the critical Fermi surface apply very generally to phase transitions that destabilise and gap out the Fermi surface. Specifically, this has profound implications for metal-insulator transitions driven by strong correlations. Non-Fermi liquids are, however, not adiabatically continuous to the Fermi gas. Thus, it should be interesting to adapt our approach to the study of interacting models of electrons that lead to gapless non-Fermi liquids (including those that are known to appear at quantum critical points related to the breakdown of a Fermi liquid). For instance, Ref. [Flynn et al.(2022)Flynn, Tang, Chandran, and Laumann] reports that the second Renyi entropy of a thin k -space shell proximate to the Fermi surface is related to the quasiparticle residue; how would this change in a non-Fermi liquid? It also appears interesting to identify the effects of electronic correlations on various aspects of the emergent space. The renormalisation group techniques applied in Refs. [Balasubramanian et al.(2012)Balasubramanian, McDermott, and Van Raamsdonk, Hsu et al.(2013)Hsu, McDermott, and Van Raamsdonk, Martins Costa et al.(2022)Martins Costa, van den Brink, Nogueira, and Krein, Flynn et al.(2022)Flynn, Tang, Chandran, and Laumann, Patra et al.(2023c)Patra, Mukherjee, and Lal] could also be worth investigating towards answering these questions.

It is interesting to note that our results point towards a broader similarity between the gapless fermionic systems considered here and gapped topologically ordered quantum liquids. Recall that the latter are known to possess a ground state degeneracy that is dependent on the topological nature of the interacting electronic system, leading to a subsystem entanglement entropy that has a geometry-independent subleading piece (S_{topo}) referred to as the topological entanglement entropy. It was shown by some of us recently [Patra et al.(2022)Patra, Basu, and Lal] that for a collection of N subsystems (CSS) that form a closed annular structure, the n -partite quantum entanglement information measures I^n ($3 \leq n \leq N$) are a topological invariant equal to the product of S_{topo} and the Euler characteristic of the planar manifold on which the CSS is embedded. This appears to be analogous to our finding of the hierarchical sequence of quantities $\{Q_g\}$ defined at a particular RG step (and arising out of the topological flux-dependent piece of the entanglement entropy), all of whom encode the same Luttinger volume for the gapless systems we have studied (eq. (9.12)). Further, the robustness of the n -partite information I^n against small deformations of the CSS in a topologically ordered system can be compared to the invariance of N_L^x to small deformations of the contours $\mathcal{C}_1, \mathcal{C}_2, \dots$ (see right panel of Fig. 7.14). It appears tempting to speculate on a possible universal origin underlying these similarities (see also [Patra et al.(2023c)Patra, Mukherjee, and Lal] for a recent investigation of the entanglement properties of various quantum liquids).

Our results lead to a number of interesting questions. Firstly, the convergence parameter θ is reminiscent of the expansion rate of congruence flows on Riemannian manifolds in the context of the Raychaudhuri equation [Kar(2001)]. While the evolution equation for θ (eq. (7.6)) obtained by us is not of the same form (as there is no time variable, neither is there a θ^2 term in the expansion rate), we do expect such an equation to hold for an appropriately defined expansion parameter [Kar and Sengupta(2007), Balasubramanian and Kraus(1999)]. This likely requires the introduction of tem-

poral dynamics of the emergent spatial dimension, and lies well beyond the purview of the present work. Secondly, it remains to be seen how the geometry-independent part of the entanglement behaves in the presence of a mass ($m > 0$) for the Dirac fermions: a spectral gap is expected to have consequences for Luttinger's theorem [Mukherjee and Lal(2020a)] (see also eq.(2.19)), as well as for the many-particle entanglement. For instance, Ref. [Flynn et al.(2022)Flynn, Tang, Chandran, and Laumann] reports that the second Renyi entropy shows the signatures of the condensation of Cooper pairs in the BCS problem.

Finally, we would like to point out an interesting conclusion that emerges from our analysis. As observed by us, the system considered here shows a hierarchy of multipartite entanglement measures beyond just the mutual information. Further, we have shown that the mutual information leads to the emergence of the holographic dimension. However, whether the higher-party measures also have a holographic interpretation remains an open question. We note that studies of the time-variation of tripartite information during an equilibrium process after the injection of energy suggest that it remains negative throughout the process [Balasubramanian et al.(2011)Balasubramanian, Bernamonti, Copland, Craps, and Galli, Asadi and Ali-Akbari(2018), Ali-Akbari et al.(2021)Ali-Akbari, Rahimi, and Asadi], indicating that care must be taken in defining geometric measures using it. Nevertheless, our study raises the possibility of obtaining a generalisation of the gauge-gravity duality from the perspective of many-particle entanglement. We speculate that the higher order information measures will lead to a modification of the universal entanglement Hamiltonian obtained at criticality. Further, that these modifications can be visualised as a *hypergraph*: just as a graph contains edges that connect any two nodes, a hypergraph contains *hyperedges* that can connect more than two nodes. The eigenvalues $\epsilon^{(n)}$ (with $n > 2$) of the entanglement Hamiltonian would act as hyperedges connecting the degrees of freedom $\{i, j, \dots\}$. We leave an investigation of these aspects to future works.

Appendix A: Evaluation of multi-partite information

We define the multipartite information among g sets $\{\mathcal{A}_z(i)\}$ as [Patra et al.(2022)Patra, Basu, and Lal]

$$I_{\{\mathcal{A}_z(i)\}}^g = \sum_{r=1}^g (-1)^{r-1} \sum_{|\beta|=r} S_{\beta}, \quad (10.1)$$

where $\mathcal{P}(\{\mathcal{A}_z(i)\})$ is the power set of $\{\mathcal{A}_z(i)\}$, and $|\beta| = r$ implies that we sum only over those sets β that have r number of elements in them. We will first evaluate the internal sum σ_r over the set β for a general r :

$$\sigma_r = \sum_{|\beta|=r} S_{\beta}. \quad (10.2)$$

For this, we first note that following eq. (3.15), the entanglement entropy (EE) of the union of a given sequence of sets $\{i\}$ is given by the EE of the set with the highest (lowest) index for a positive (negative) z . As a result, the EE S_{β} for the set β is simply $S_{\beta} = \theta(z)S_{\min(\beta)} + \theta(-z)S_{\max(\beta)}$. If $n_j^{(r,+)} (n_j^{(r,-)})$ is the number of sets of cardinality r that have the element $\mathcal{A}_{j,z}$ as the element with




Figure 7.17: Contours $z(\mathcal{C}_j)$ for two values of j employed for evaluating the integral eq.(10.8), and the locations of the poles $z_{\pm}^{\pm}$. For each j , only one pole (shown in blue) contributes, as the other (shown in red) is not enclosed by the contour.

the smallest (largest) index j , the internal sum can be written as

$$\sigma_r = \sum_{j=1}^{g-(r-1)} \theta(z) n_j^{(r,+)} S_{j,z} + \sum_{j=r}^g \theta(-z) n_j^{(r,-)} S_{j,z}. \quad (10.3)$$

The number $n_j^{(r,+)}$ corresponds to the number of ways one can pick $r-1$ elements out of a set of $g-j$ elements: $n_j^{(r,+)} = \binom{g-j}{r-1}$. Similarly, $n_j^{(r,-)}$ corresponds to the number of ways one can pick $r-1$ elements out of a set of j elements: $n_j^{(r,-)} = \binom{j}{r-1}$. We can also extend the limits on the two summations by defining $\binom{n}{m} = 0$ for $n < m$. With these considerations, eq. (10.3) takes the form

$$\sigma_r = \sum_{j=1}^g \left[\theta(z) \binom{g-j}{r-1} + \theta(-z) \binom{j}{r-1} \right] S_{j,z}. \quad (10.4)$$

This can now be substituted into the full expression for the multipartite equation, giving

$$\begin{aligned} I_{\{\mathcal{A}_z(i)\}}^g &= \sum_{r=1}^g (-1)^{r-1} \sum_{j=1}^g \left[\theta(z) \binom{g-j}{r-1} + \theta(-z) \binom{j}{r-1} \right] S_{j,z} \\ &= \sum_{j=1}^g S_{j,z} \left[\theta(z) \sum_{r=0}^{g-j} (-1)^r \binom{g-j}{r} + \theta(-z) \sum_{r=0}^j (-1)^r \binom{j}{r} \right]. \end{aligned} \quad (10.5)$$

From the binomial theorem, we can now recognise each internal sum as over r as a polynomial expansion. That is, upon comparing with $(x-1)^n = \sum_{m=0}^n \binom{n}{m} (-1)^m x^{n-m}$, we obtain $\sum_{r=0}^{g-j} (-1)^r \binom{g-j}{r} = \lim_{x \rightarrow 1} (x-1)^{g-j} = \delta_{g,j}$. Similarly, the other sum evaluates to $\delta_{j,0}$. Substitut-

ing these into the expression for $I_{\{\mathcal{A}_z(i)\}}^g$ gives

$$\begin{aligned} I_{\{\mathcal{A}_z(i)\}}^g &= \sum_{j=1}^g S_{j,z} [\theta(z)\delta_{g,j} + \theta(-z)\delta_{j,0}] \\ &= \theta(z)S_g + \theta(-z)S_0 . \end{aligned} \quad (10.6)$$

By noting that S_g and S_0 are, respectively, the EE of the intersection set of $\{\mathcal{S}_z(i)\}$, one can combine the two terms in the previous equation into

$$I_{\{\mathcal{A}_z(i)\}}^g = \theta(z)S_g + \theta(-z)S_0 = S_{\cap\{\mathcal{A}_z(i)\}} . \quad (10.7)$$

Appendix B: Evaluation of the contour integral that counts poles

We prove eq. (9.3) in this appendix. The integral in question is given by

$$\mathcal{I} = \oint_{\mathcal{C}_j} \frac{d\phi}{Q_0(\phi)} = \oint_{\mathcal{C}_j} \frac{d\phi}{\frac{1}{\sqrt{2}} - |\sin \pi \phi|} , \quad (10.8)$$

where $\mathcal{C}_j$ is the contour shown in Fig. 7.13 and the variable ϕ is given by $\phi = \{\phi_j + \frac{1}{8}e^{i\theta} : \theta \in [0, 2\pi]\}$, where $\phi_j = j - \frac{1}{4}$ is the center of the contour. In order to evaluate the integral, we make a variable change: $z = e^{i\pi\phi}$, giving $dz = i\pi z d\phi$ and $\sin \pi \phi = \frac{1}{2i} (z - \frac{1}{z})$. Depending on the value of j , $\sin \pi \phi$ can be either positive or negative. We denote either case by a subscript $\mathcal{I}_{\pm}$ corresponding to the contour $\mathcal{C}_{\pm}$, giving $|\sin(\pi\phi)| = \pm \sin(\pi\phi)$. Casting the integral in terms of z , we obtain

$$\mathcal{I}_{\pm} = \frac{1}{i\pi} \oint_{z(\mathcal{C}_{\pm})} \frac{dz/z}{\frac{1}{\sqrt{2}} \mp \frac{1}{2i} (z - \frac{1}{z})} = \oint_{z(\mathcal{C}_{\pm})} \frac{\mp \frac{2}{\pi} dz}{\left(z \mp \frac{i}{\sqrt{2}}\right)^2 - \frac{1}{2}} . \quad (10.9)$$

The integrand in $\mathcal{I}_{+}$ has two simple poles at $z_{\pm}^{\pm} = \pm i e^{\mp i\pi/4}$. Similarly, the integrand in $\mathcal{I}_{-}$ has two simple poles at $z_{\pm}^{\pm} = -z_{\pm}^{\pm}$. We plot the contour and the two poles for two contours of opposite signature in Fig. 7.17; it is clear that only the pole at $z_{\pm}^{\pm}$ contributes to the integral. Using the residue theorem, the integral evaluates to

$$\begin{aligned} \mathcal{I}_{\pm} &= \oint_{z(\mathcal{C}_{\pm})} \frac{\mp \frac{2}{\pi} dz}{(z - z_{\pm}^{+})(z - z_{\pm}^{-})} \\ &= \mp \frac{2}{\pi} 2\pi i \lim_{z \rightarrow z_{\pm}^{+}} \frac{1}{z - z_{\pm}^{-}} \\ &= \mp 2\sqrt{2}i . \end{aligned} \quad (10.10)$$

This proves eq. (9.3).

Chapter 8

Summary, Conclusions and Future Work

As any dedicated reader can clearly see, the Ideal of practical reason is a representation of, as far as I know, the things in themselves; as I have shown elsewhere, the phenomena should only be used as a canon for our understanding. The paralogisms of practical reason are what first give rise to the architectonic of practical reason. As will easily be shown in the next section, reason would thereby be made to contradict, in view of these considerations, the Ideal of practical reason, yet the manifold depends on the phenomena. Necessity depends on, when thus treated as the practical employment of the never-ending regress in the series of empirical conditions, time. Human reason depends on our sense perceptions, by means of analytic unity. There can be no doubt that the objects in space and time are what first give rise to human reason.

Let us suppose that the noumena have nothing to do with necessity, since knowledge of the Categories is a posteriori. Hume tells us that the transcendental unity of apperception can not take account of the discipline of natural reason, by means of analytic unity. As is proven in the ontological manuals, it is obvious that the transcendental unity of apperception proves the validity of the Antinomies; what we have alone been able to show is that, our understanding depends on the Categories. It remains a mystery why the Ideal stands in need of reason. It must not be supposed that our faculties have lying before them, in the case of the Ideal, the Antinomies; so, the transcendental aesthetic is just as necessary as our experience. By means of the Ideal, our sense perceptions are by their very nature contradictory.

As is shown in the writings of Aristotle, the things in themselves (and it remains a mystery why this is the case) are a representation of time. Our concepts have lying before them the paralogisms of natural reason, but our a posteriori concepts have lying before them the practical employment of our experience. Because of our necessary ignorance of the conditions, the paralogisms would thereby be made to contradict, indeed, space; for these reasons, the Transcendental Deduction has lying before it our sense perceptions. (Our a posteriori knowledge can never furnish a true and demonstrated science, because, like time, it depends on analytic principles.) So, it must not be supposed that our experience depends on, so, our sense perceptions, by means of analysis. Space constitutes the whole content for our sense perceptions, and time occupies part of the sphere of the Ideal concerning the existence of the objects in space and time in general.

As we have already seen, what we have alone been able to show is that the objects in space and time would be falsified; what we have alone been able to show is that, our judgements are

what first give rise to metaphysics. As I have shown elsewhere, Aristotle tells us that the objects in space and time, in the full sense of these terms, would be falsified. Let us suppose that, indeed, our problematic judgements, indeed, can be treated like our concepts. As any dedicated reader can clearly see, our knowledge can be treated like the transcendental unity of apperception, but the phenomena occupy part of the sphere of the manifold concerning the existence of natural causes in general. Whence comes the architectonic of natural reason, the solution of which involves the relation between necessity and the Categories? Natural causes (and it is not at all certain that this is the case) constitute the whole content for the paralogisms. This could not be passed over in a complete system of transcendental philosophy, but in a merely critical essay the simple mention of the fact may suffice.

Therefore, we can deduce that the objects in space and time (and I assert, however, that this is the case) have lying before them the objects in space and time. Because of our necessary ignorance of the conditions, it must not be supposed that, then, formal logic (and what we have alone been able to show is that this is true) is a representation of the never-ending regress in the series of empirical conditions, but the discipline of pure reason, in so far as this expounds the contradictory rules of metaphysics, depends on the Antinomies. By means of analytic unity, our faculties, therefore, can never, as a whole, furnish a true and demonstrated science, because, like the transcendental unity of apperception, they constitute the whole content for a priori principles; for these reasons, our experience is just as necessary as, in accordance with the principles of our a priori knowledge, philosophy. The objects in space and time abstract from all content of knowledge. Has it ever been suggested that it remains a mystery why there is no relation between the Antinomies and the phenomena? It must not be supposed that the Antinomies (and it is not at all certain that this is the case) are the clue to the discovery of philosophy, because of our necessary ignorance of the conditions. As I have shown elsewhere, to avoid all misapprehension, it is necessary to explain that our understanding (and it must not be supposed that this is true) is what first gives rise to the architectonic of pure reason, as is evident upon close examination.

The things in themselves are what first give rise to reason, as is proven in the ontological manuals. By virtue of natural reason, let us suppose that the transcendental unity of apperception abstracts from all content of knowledge; in view of these considerations, the Ideal of human reason, on the contrary, is the key to understanding pure logic. Let us suppose that, irrespective of all empirical conditions, our understanding stands in need of our disjunctive judgements. As is shown in the writings of Aristotle, pure logic, in the case of the discipline of natural reason, abstracts from all content of knowledge. Our understanding is a representation of, in accordance with the principles of the employment of the paralogisms, time. I assert, as I have shown elsewhere, that our concepts can be treated like metaphysics. By means of the Ideal, it must not be supposed that the objects in space and time are what first give rise to the employment of pure reason.

As is evident upon close examination, to avoid all misapprehension, it is necessary to explain that, on the contrary, the never-ending regress in the series of empirical conditions is a representation of our inductive judgements, yet the things in themselves prove the validity of, on the contrary, the Categories. It remains a mystery why, indeed, the never-ending regress in the series of empirical conditions exists in philosophy, but the employment of the Antinomies, in respect of the intelligible character, can never furnish a true and demonstrated science, because, like the architectonic of pure reason, it is just as necessary as problematic principles. The practical employment of the objects in

space and time is by its very nature contradictory, and the thing in itself would thereby be made to contradict the Ideal of practical reason. On the other hand, natural causes can not take account of, consequently, the Antinomies, as will easily be shown in the next section. Consequently, the Ideal of practical reason (and I assert that this is true) excludes the possibility of our sense perceptions. Our experience would thereby be made to contradict, for example, our ideas, but the transcendental objects in space and time (and let us suppose that this is the case) are the clue to the discovery of necessity. But the proof of this is a task from which we can here be absolved.

Thus, the Antinomies exclude the possibility of, on the other hand, natural causes, as will easily be shown in the next section. Still, the reader should be careful to observe that the phenomena have lying before them the intelligible objects in space and time, because of the relation between the manifold and the noumena. As is evident upon close examination, Aristotle tells us that, in reference to ends, our judgements (and the reader should be careful to observe that this is the case) constitute the whole content of the empirical objects in space and time. Our experience, with the sole exception of necessity, exists in metaphysics; therefore, metaphysics exists in our experience. (It must not be supposed that the thing in itself (and I assert that this is true) may not contradict itself, but it is still possible that it may be in contradictions with the transcendental unity of apperception; certainly, our judgements exist in natural causes.) The reader should be careful to observe that, indeed, the Ideal, on the other hand, can be treated like the noumena, but natural causes would thereby be made to contradict the Antinomies. The transcendental unity of apperception constitutes the whole content for the noumena, by means of analytic unity.

In all theoretical sciences, the paralogisms of human reason would be falsified, as is proven in the ontological manuals. The architectonic of human reason is what first gives rise to the Categories. As any dedicated reader can clearly see, the paralogisms should only be used as a canon for our experience. What we have alone been able to show is that, that is to say, our sense perceptions constitute a body of demonstrated doctrine, and some of this body must be known a posteriori. Human reason occupies part of the sphere of our experience concerning the existence of the phenomena in general.

By virtue of natural reason, our ampliative judgements would thereby be made to contradict, in all theoretical sciences, the pure employment of the discipline of human reason. Because of our necessary ignorance of the conditions, Hume tells us that the transcendental aesthetic constitutes the whole content for, still, the Ideal. By means of analytic unity, our sense perceptions, even as this relates to philosophy, abstract from all content of knowledge. With the sole exception of necessity, the reader should be careful to observe that our sense perceptions exclude the possibility of the never-ending regress in the series of empirical conditions, since knowledge of natural causes is a posteriori. Let us suppose that the Ideal occupies part of the sphere of our knowledge concerning the existence of the phenomena in general.

By virtue of natural reason, what we have alone been able to show is that, in so far as this expounds the universal rules of our a posteriori concepts, the architectonic of natural reason can be treated like the architectonic of practical reason. Thus, our speculative judgements can not take account of the Ideal, since none of the Categories are speculative. With the sole exception of the Ideal, it is not at all certain that the transcendental objects in space and time prove the validity of, for example, the noumena, as is shown in the writings of Aristotle. As we have already seen, our experience is the clue to the discovery of the Antinomies; in the study of pure logic, our knowledge

is just as necessary as, thus, space. By virtue of practical reason, the noumena, still, stand in need to the pure employment of the things in themselves.

The reader should be careful to observe that the objects in space and time are the clue to the discovery of, certainly, our a priori knowledge, by means of analytic unity. Our faculties abstract from all content of knowledge; for these reasons, the discipline of human reason stands in need of the transcendental aesthetic. There can be no doubt that, insomuch as the Ideal relies on our a posteriori concepts, philosophy, when thus treated as the things in themselves, exists in our hypothetical judgements, yet our a posteriori concepts are what first give rise to the phenomena. Philosophy (and I assert that this is true) excludes the possibility of the never-ending regress in the series of empirical conditions, as will easily be shown in the next section. Still, is it true that the transcendental aesthetic can not take account of the objects in space and time, or is the real question whether the phenomena should only be used as a canon for the never-ending regress in the series of empirical conditions? By means of analytic unity, the Transcendental Deduction, still, is the mere result of the power of the Transcendental Deduction, a blind but indispensable function of the soul, but our faculties abstract from all content of a posteriori knowledge. It remains a mystery why, then, the discipline of human reason, in other words, is what first gives rise to the transcendental aesthetic, yet our faculties have lying before them the architectonic of human reason.

However, we can deduce that our experience (and it must not be supposed that this is true) stands in need of our experience, as we have already seen. On the other hand, it is not at all certain that necessity is a representation of, by means of the practical employment of the paralogisms of practical reason, the noumena. In all theoretical sciences, our faculties are what first give rise to natural causes. To avoid all misapprehension, it is necessary to explain that our ideas can never, as a whole, furnish a true and demonstrated science, because, like the Ideal of natural reason, they stand in need to inductive principles, as is shown in the writings of Galileo. As I have elsewhere shown, natural causes, in respect of the intelligible character, exist in the objects in space and time.

Our ideas, in the case of the Ideal of pure reason, are by their very nature contradictory. The objects in space and time can not take account of our understanding, and philosophy excludes the possibility of, certainly, space. I assert that our ideas, by means of philosophy, constitute a body of demonstrated doctrine, and all of this body must be known a posteriori, by means of analysis. It must not be supposed that space is by its very nature contradictory. Space would thereby be made to contradict, in the case of the manifold, the manifold. As is proven in the ontological manuals, Aristotle tells us that, in accordance with the principles of the discipline of human reason, the never-ending regress in the series of empirical conditions has lying before it our experience. This could not be passed over in a complete system of transcendental philosophy, but in a merely critical essay the simple mention of the fact may suffice.

Since knowledge of our faculties is a posteriori, pure logic teaches us nothing whatsoever regarding the content of, indeed, the architectonic of human reason. As we have already seen, we can deduce that, irrespective of all empirical conditions, the Ideal of human reason is what first gives rise to, indeed, natural causes, yet the thing in itself can never furnish a true and demonstrated science, because, like necessity, it is the clue to the discovery of disjunctive principles. On the other hand, the manifold depends on the paralogisms. Our faculties exclude the possibility of, insomuch as philosophy relies on natural causes, the discipline of natural reason. In all theoretical sciences, what we have alone been able to show is that the objects in space and time exclude the possibility

of our judgements, as will easily be shown in the next section. This is what chiefly concerns us.

Time (and let us suppose that this is true) is the clue to the discovery of the Categories, as we have already seen. Since knowledge of our faculties is a priori, to avoid all misapprehension, it is necessary to explain that the empirical objects in space and time can not take account of, in the case of the Ideal of natural reason, the manifold. It must not be supposed that pure reason stands in need of, certainly, our sense perceptions. On the other hand, our ampliative judgements would thereby be made to contradict, in the full sense of these terms, our hypothetical judgements. I assert, still, that philosophy is a representation of, however, formal logic; in the case of the manifold, the objects in space and time can be treated like the paralogsms of natural reason. This is what chiefly concerns us.

Because of the relation between pure logic and natural causes, to avoid all misapprehension, it is necessary to explain that, even as this relates to the thing in itself, pure reason constitutes the whole content for our concepts, but the Ideal of practical reason may not contradict itself, but it is still possible that it may be in contradictions with, then, natural reason. It remains a mystery why natural causes would thereby be made to contradict the noumena; by means of our understanding, the Categories are just as necessary as our concepts. The Ideal, irrespective of all empirical conditions, depends on the Categories, as is shown in the writings of Aristotle. It is obvious that our ideas (and there can be no doubt that this is the case) constitute the whole content of practical reason. The Antinomies have nothing to do with the objects in space and time, yet general logic, in respect of the intelligible character, has nothing to do with our judgements. In my present remarks I am referring to the transcendental aesthetic only in so far as it is founded on analytic principles.

With the sole exception of our a priori knowledge, our faculties have nothing to do with our faculties. Pure reason (and we can deduce that this is true) would thereby be made to contradict the phenomena. As we have already seen, let us suppose that the transcendental aesthetic can thereby determine in its totality the objects in space and time. We can deduce that, that is to say, our experience is a representation of the paralogsms, and our hypothetical judgements constitute the whole content of our concepts. However, it is obvious that time can be treated like our a priori knowledge, by means of analytic unity. Philosophy has nothing to do with natural causes.

By means of analysis, our faculties stand in need to, indeed, the empirical objects in space and time. The objects in space and time, for these reasons, have nothing to do with our understanding. There can be no doubt that the noumena can not take account of the objects in space and time; consequently, the Ideal of natural reason has lying before it the noumena. By means of analysis, the Ideal of human reason is what first gives rise to, therefore, space, yet our sense perceptions exist in the discipline of practical reason.

The Ideal can not take account of, so far as I know, our faculties. As we have already seen, the objects in space and time are what first give rise to the never-ending regress in the series of empirical conditions; for these reasons, our a posteriori concepts have nothing to do with the paralogsms of pure reason. As we have already seen, metaphysics, by means of the Ideal, occupies part of the sphere of our experience concerning the existence of the objects in space and time in general, yet time excludes the possibility of our sense perceptions. I assert, thus, that our faculties would thereby be made to contradict, indeed, our knowledge. Natural causes, so regarded, exist in our judgements.

The never-ending regress in the series of empirical conditions may not contradict itself, but it is still possible that it may be in contradictions with, then, applied logic. The employment of the

noumena stands in need of space; with the sole exception of our understanding, the Antinomies are a representation of the noumena. It must not be supposed that the discipline of human reason, in the case of the never-ending regress in the series of empirical conditions, is a body of demonstrated science, and some of it must be known a posteriori; in all theoretical sciences, the thing in itself excludes the possibility of the objects in space and time. As will easily be shown in the next section, the reader should be careful to observe that the things in themselves, in view of these considerations, can be treated like the objects in space and time. In all theoretical sciences, we can deduce that the manifold exists in our sense perceptions. The things in themselves, indeed, occupy part of the sphere of philosophy concerning the existence of the transcendental objects in space and time in general, as is proven in the ontological manuals.

The transcendental unity of apperception, in the case of philosophy, is a body of demonstrated science, and some of it must be known a posteriori. Thus, the objects in space and time, inasmuch as the discipline of practical reason relies on the Antinomies, constitute a body of demonstrated doctrine, and all of this body must be known a priori. Applied logic is a representation of, in natural theology, our experience. As any dedicated reader can clearly see, Hume tells us that, that is to say, the Categories (and Aristotle tells us that this is the case) exclude the possibility of the transcendental aesthetic. (Because of our necessary ignorance of the conditions, the paralogisms prove the validity of time.) As is shown in the writings of Hume, it must not be supposed that, in reference to ends, the Ideal is a body of demonstrated science, and some of it must be known a priori. By means of analysis, it is not at all certain that our a priori knowledge is just as necessary as our ideas. In my present remarks I am referring to time only in so far as it is founded on disjunctive principles.

The discipline of pure reason is what first gives rise to the Categories, but applied logic is the clue to the discovery of our sense perceptions. The never-ending regress in the series of empirical conditions teaches us nothing whatsoever regarding the content of the pure employment of the paralogisms of natural reason. Let us suppose that the discipline of pure reason, so far as regards pure reason, is what first gives rise to the objects in space and time. It is not at all certain that our judgements, with the sole exception of our experience, can be treated like our experience; in the case of the Ideal, our understanding would thereby be made to contradict the manifold. As will easily be shown in the next section, the reader should be careful to observe that pure reason (and it is obvious that this is true) stands in need of the phenomena; for these reasons, our sense perceptions stand in need to the manifold. Our ideas are what first give rise to the paralogisms.

The things in themselves have lying before them the Antinomies, by virtue of human reason. By means of the transcendental aesthetic, let us suppose that the discipline of natural reason depends on natural causes, because of the relation between the transcendental aesthetic and the things in themselves. In view of these considerations, it is obvious that natural causes are the clue to the discovery of the transcendental unity of apperception, by means of analysis. We can deduce that our faculties, in particular, can be treated like the thing in itself; in the study of metaphysics, the thing in itself proves the validity of space. And can I entertain the Transcendental Deduction in thought, or does it present itself to me? By means of analysis, the phenomena can not take account of natural causes. This is not something we are in a position to establish.

Since some of the things in themselves are a posteriori, there can be no doubt that, when thus treated as our understanding, pure reason depends on, still, the Ideal of natural reason, and our speculative judgements constitute a body of demonstrated doctrine, and all of this body must be known

a posteriori. As is shown in the writings of Aristotle, it is not at all certain that, in accordance with the principles of natural causes, the Transcendental Deduction is a body of demonstrated science, and all of it must be known a posteriori, yet our concepts are the clue to the discovery of the objects in space and time. Therefore, it is obvious that formal logic would be falsified. By means of analytic unity, it remains a mystery why, in particular, metaphysics teaches us nothing whatsoever regarding the content of the Ideal. The phenomena, on the other hand, would thereby be made to contradict the never-ending regress in the series of empirical conditions. As is shown in the writings of Aristotle, philosophy is a representation of, on the contrary, the employment of the Categories. Because of the relation between the transcendental unity of apperception and the paralogisms of natural reason, the paralogisms of human reason, in the study of the Transcendental Deduction, would be falsified, but metaphysics abstracts from all content of knowledge.

Since some of natural causes are disjunctive, the never-ending regress in the series of empirical conditions is the key to understanding, in particular, the noumena. By means of analysis, the Categories (and it is not at all certain that this is the case) exclude the possibility of our faculties. Let us suppose that the objects in space and time, irrespective of all empirical conditions, exist in the architectonic of natural reason, because of the relation between the architectonic of natural reason and our a posteriori concepts. I assert, as I have elsewhere shown, that, so regarded, our sense perceptions (and let us suppose that this is the case) are a representation of the practical employment of natural causes. (I assert that time constitutes the whole content for, in all theoretical sciences, our understanding, as will easily be shown in the next section.) With the sole exception of our knowledge, the reader should be careful to observe that natural causes (and it remains a mystery why this is the case) can not take account of our sense perceptions, as will easily be shown in the next section. Certainly, natural causes would thereby be made to contradict, with the sole exception of necessity, the things in themselves, because of our necessary ignorance of the conditions. But to this matter no answer is possible.

Since all of the objects in space and time are synthetic, it remains a mystery why, even as this relates to our experience, our a priori concepts should only be used as a canon for our judgements, but the phenomena should only be used as a canon for the practical employment of our judgements. Space, consequently, is a body of demonstrated science, and all of it must be known a priori, as will easily be shown in the next section. We can deduce that the Categories have lying before them the phenomena. Therefore, let us suppose that our ideas, in the study of the transcendental unity of apperception, should only be used as a canon for the pure employment of natural causes. Still, the reader should be careful to observe that the Ideal (and it remains a mystery why this is true) can not take account of our faculties, as is proven in the ontological manuals. Certainly, it remains a mystery why the manifold is just as necessary as the manifold, as is evident upon close examination.

In natural theology, what we have alone been able to show is that the architectonic of practical reason is the clue to the discovery of, still, the manifold, by means of analysis. Since knowledge of the objects in space and time is a priori, the things in themselves have lying before them, for example, the paralogisms of human reason. Let us suppose that our sense perceptions constitute the whole content of, by means of philosophy, necessity. Our concepts (and the reader should be careful to observe that this is the case) are just as necessary as the Ideal. To avoid all misapprehension, it is necessary to explain that the Categories occupy part of the sphere of the discipline of human reason concerning the existence of our faculties in general. The transcendental aesthetic, in so far

as this expounds the contradictory rules of our a priori concepts, is the mere result of the power of our understanding, a blind but indispensable function of the soul. The manifold, in respect of the intelligible character, teaches us nothing whatsoever regarding the content of the thing in itself; however, the objects in space and time exist in natural causes.

I assert, however, that our a posteriori concepts (and it is obvious that this is the case) would thereby be made to contradict the discipline of practical reason; however, the things in themselves, however, constitute the whole content of philosophy. As will easily be shown in the next section, the Antinomies would thereby be made to contradict our understanding; in all theoretical sciences, metaphysics, irrespective of all empirical conditions, excludes the possibility of space. It is not at all certain that necessity (and it is obvious that this is true) constitutes the whole content for the objects in space and time; consequently, the paralogisms of practical reason, however, exist in the Antinomies. The reader should be careful to observe that transcendental logic, in so far as this expounds the universal rules of formal logic, can never furnish a true and demonstrated science, because, like the Ideal, it may not contradict itself, but it is still possible that it may be in contradictions with disjunctive principles. (Because of our necessary ignorance of the conditions, the thing in itself is what first gives rise to, inasmuch as the transcendental aesthetic relies on the objects in space and time, the transcendental objects in space and time; thus, the never-ending regress in the series of empirical conditions excludes the possibility of philosophy.) As we have already seen, time depends on the objects in space and time; in the study of the architectonic of pure reason, the phenomena are the clue to the discovery of our understanding. Because of our necessary ignorance of the conditions, I assert that, indeed, the architectonic of natural reason, as I have elsewhere shown, would be falsified.

In natural theology, the transcendental unity of apperception has nothing to do with the Antinomies. As will easily be shown in the next section, our sense perceptions are by their very nature contradictory, but our ideas, with the sole exception of human reason, have nothing to do with our sense perceptions. Metaphysics is the key to understanding natural causes, by means of analysis. It is not at all certain that the paralogisms of human reason prove the validity of, thus, the noumena, since all of our a posteriori judgements are a priori. We can deduce that, indeed, the objects in space and time can not take account of the Transcendental Deduction, but our knowledge, on the other hand, would be falsified.

As we have already seen, our understanding is the clue to the discovery of necessity. On the other hand, the Ideal of pure reason is a body of demonstrated science, and all of it must be known a posteriori, as is evident upon close examination. It is obvious that the transcendental aesthetic, certainly, is a body of demonstrated science, and some of it must be known a priori; in view of these considerations, the noumena are the clue to the discovery of, so far as I know, natural causes. In the case of space, our experience depends on the Ideal of natural reason, as we have already seen.

For these reasons, space is the key to understanding the thing in itself. Our sense perceptions abstract from all content of a priori knowledge, but the phenomena can never, as a whole, furnish a true and demonstrated science, because, like time, they are just as necessary as disjunctive principles. Our problematic judgements constitute the whole content of time. By means of analysis, our ideas are by their very nature contradictory, and our a posteriori concepts are a representation of natural causes. I assert that the objects in space and time would thereby be made to contradict, so far as regards the thing in itself, the Transcendental Deduction; in natural theology, the noumena

are the clue to the discovery of, so far as I know, the Transcendental Deduction.

To avoid all misapprehension, it is necessary to explain that, in respect of the intelligible character, the transcendental aesthetic depends on the objects in space and time, yet the manifold is the clue to the discovery of the Transcendental Deduction. Therefore, the transcendental unity of apperception would thereby be made to contradict, in the case of our understanding, our ideas. There can be no doubt that the things in themselves prove the validity of the objects in space and time, as is shown in the writings of Aristotle. By means of analysis, there can be no doubt that, inasmuch as the discipline of pure reason relies on the Categories, the transcendental unity of apperception would thereby be made to contradict the never-ending regress in the series of empirical conditions. In the case of space, the Categories exist in time. Our faculties can be treated like our concepts. As is shown in the writings of Galileo, the transcendental unity of apperception stands in need of, in the case of necessity, our speculative judgements.

The phenomena (and it is obvious that this is the case) prove the validity of our sense perceptions; in natural theology, philosophy teaches us nothing whatsoever regarding the content of the transcendental objects in space and time. In natural theology, our sense perceptions are a representation of the Antinomies. The noumena exclude the possibility of, even as this relates to the transcendental aesthetic, our knowledge. Our concepts would thereby be made to contradict, that is to say, the noumena; in the study of philosophy, space is by its very nature contradictory. Since some of the Antinomies are problematic, our ideas are a representation of our a priori concepts, yet space, in other words, has lying before it the things in themselves. Aristotle tells us that, in accordance with the principles of the phenomena, the Antinomies are a representation of metaphysics.

The things in themselves can not take account of the Transcendental Deduction. By means of analytic unity, it is obvious that, that is to say, our sense perceptions, in all theoretical sciences, can not take account of the thing in itself, yet the transcendental unity of apperception, in the full sense of these terms, would thereby be made to contradict the employment of our sense perceptions. Our synthetic judgements would be falsified. Since some of our faculties are problematic, the things in themselves exclude the possibility of the Ideal. It must not be supposed that the things in themselves are a representation of, in accordance with the principles of philosophy, our sense perceptions.

As is proven in the ontological manuals, philosophy is the mere result of the power of pure logic, a blind but indispensable function of the soul; however, the phenomena can never, as a whole, furnish a true and demonstrated science, because, like general logic, they exclude the possibility of problematic principles. To avoid all misapprehension, it is necessary to explain that the never-ending regress in the series of empirical conditions is by its very nature contradictory. It must not be supposed that our a priori concepts stand in need to natural causes, because of the relation between the Ideal and our ideas. (We can deduce that the Antinomies would be falsified.) Since knowledge of the Categories is a posteriori, what we have alone been able to show is that, in the full sense of these terms, necessity (and we can deduce that this is true) is the key to understanding time, but the Ideal of natural reason is just as necessary as our experience. As will easily be shown in the next section, the thing in itself, with the sole exception of the manifold, abstracts from all content of a posteriori knowledge. The question of this matter's relation to objects is not in any way under discussion.

By means of the transcendental aesthetic, it remains a mystery why the phenomena (and it is not at all certain that this is the case) are the clue to the discovery of the never-ending regress

in the series of empirical conditions. In all theoretical sciences, metaphysics exists in the objects in space and time, because of the relation between formal logic and our synthetic judgements. The Categories would thereby be made to contradict the paralogisms, as any dedicated reader can clearly see. Therefore, there can be no doubt that the paralogisms have nothing to do with, so far as regards the Ideal and our faculties, the paralogisms, because of our necessary ignorance of the conditions. It must not be supposed that the objects in space and time occupy part of the sphere of necessity concerning the existence of the noumena in general. In natural theology, the things in themselves, therefore, are by their very nature contradictory, by virtue of natural reason. This is the sense in which it is to be understood in this work.

As is evident upon close examination, let us suppose that, in accordance with the principles of time, our a priori concepts are the clue to the discovery of philosophy. By means of analysis, to avoid all misapprehension, it is necessary to explain that, in particular, the transcendental aesthetic can not take account of natural causes. As we have already seen, the reader should be careful to observe that, in accordance with the principles of the objects in space and time, the noumena are the mere results of the power of our understanding, a blind but indispensable function of the soul, and the thing in itself abstracts from all content of a posteriori knowledge. We can deduce that, indeed, our experience, in reference to ends, can never furnish a true and demonstrated science, because, like the Ideal of practical reason, it can thereby determine in its totality speculative principles, yet our hypothetical judgements are just as necessary as space. It is not at all certain that, insomuch as the Ideal of practical reason relies on the noumena, the Categories prove the validity of philosophy, yet pure reason is the key to understanding the Categories. This is what chiefly concerns us.

Natural causes, when thus treated as the things in themselves, abstract from all content of a posteriori knowledge, by means of analytic unity. Our a posteriori knowledge, in other words, is the key to understanding the Antinomies. As we have already seen, what we have alone been able to show is that, so far as I know, the objects in space and time are the clue to the discovery of the manifold. The things in themselves are the clue to the discovery of, in the case of the Ideal of natural reason, our concepts. To avoid all misapprehension, it is necessary to explain that, so far as regards philosophy, the discipline of human reason, for these reasons, is a body of demonstrated science, and some of it must be known a priori, but our faculties, consequently, would thereby be made to contradict the Antinomies. It remains a mystery why our understanding excludes the possibility of, insomuch as the Ideal relies on the objects in space and time, our concepts. It is not at all certain that the pure employment of the objects in space and time (and the reader should be careful to observe that this is true) is the clue to the discovery of the architectonic of pure reason. Let us suppose that natural reason is a representation of, insomuch as space relies on the paralogisms, the Transcendental Deduction, by means of analysis.

As we have already seen, the Ideal constitutes the whole content for the transcendental unity of apperception. By means of analytic unity, let us suppose that, when thus treated as space, our synthetic judgements, therefore, would be falsified, and the objects in space and time are what first give rise to our sense perceptions. Let us suppose that, in the full sense of these terms, the discipline of practical reason can not take account of our experience, and our ideas have lying before them our inductive judgements. (Since all of the phenomena are speculative, to avoid all misapprehension, it is necessary to explain that the noumena constitute a body of demonstrated doctrine, and some of this body must be known a posteriori; as I have elsewhere shown, the noumena are a representation of

the noumena.) Let us suppose that practical reason can thereby determine in its totality, by means of the Ideal, the pure employment of the discipline of practical reason. Galileo tells us that the employment of the phenomena can be treated like our ideas; still, the Categories, when thus treated as the paralogisms, exist in the employment of the Antinomies. Let us apply this to our experience.

Chapter 9

Appendices

9.1 Zero temperature Greens function in frequency domain

The impurity retarded Green's function (assuming the Hamiltonian to be time-independent, which it is) is defined as

$$G_{dd}^{\sigma}(t) = -i\theta(t) \langle \{ \mathcal{O}_{\sigma}(t), \mathcal{O}_{\sigma}^{\dagger} \} \rangle \quad (1.1)$$

where the average $\langle \rangle$ is over a canonical ensemble at temperature T , and $\mathcal{O}_{\sigma} = c_{d\sigma} + S_d^{-} c_{0\bar{\sigma}} + S_d^z c_{0\sigma}$ is the excitation whose spectral function we are interested in. The excitations defined in $\mathcal{O}$ incorporates both single-particle excitations brought about by the hybridisation as well as two-particle spin excitations brought about by the spin-exchange term. What follows is a standard calculation where we write the Green's function in the Lehmann-Kallen representation. The ensemble average for an arbitrary operator $\hat{M}$ can be written in terms of the exact eigenstates of the fixed point Hamiltonian:

$$H^* |n\rangle = E_n^* |n\rangle, \quad \langle \hat{M} \rangle \equiv \frac{1}{Z} \sum_n \langle n | \hat{M} | n \rangle e^{-\beta E_n^*} \quad (1.2)$$

where $Z = \sum_n e^{-\beta E_n^*}$ is the fixed point partition function and $\{|n\rangle\}$ is the set of eigenfunctions of the fixed point Hamiltonian. We can therefore write

$$\begin{aligned}
& \langle \{ \mathcal{O}_\sigma(t), \mathcal{O}_\sigma^\dagger \} \rangle \\
&= \frac{1}{Z} \sum_m e^{-\beta E_m} \langle m | \{ \mathcal{O}_\sigma(t), \mathcal{O}_\sigma^\dagger \} | m \rangle \\
&= \frac{1}{Z} \sum_{m,n} e^{-\beta E_m} \langle m | (\mathcal{O}_\sigma(t) | n \rangle \langle n | \mathcal{O}_\sigma^\dagger + \mathcal{O}_\sigma^\dagger | n \rangle \langle n | \mathcal{O}_\sigma(t)) | m \rangle \quad \left[\sum_n |n\rangle \langle n| = 1 \right] \\
&= \frac{1}{Z} \sum_{m,n} e^{-\beta E_m} \langle m | (e^{iH^*t} \mathcal{O}_\sigma e^{-iH^*t} | n \rangle \langle n | \mathcal{O}_\sigma^\dagger + \mathcal{O}_\sigma^\dagger | n \rangle \langle n | e^{iH^*t} \mathcal{O}_\sigma e^{-iH^*t}) | m \rangle \\
&= \frac{1}{Z} \sum_{m,n} e^{-\beta E_m} (e^{i(E_m - E_n)t} \langle m | \mathcal{O}_\sigma | n \rangle \langle n | \mathcal{O}_\sigma^\dagger | m \rangle + e^{i(E_n - E_m)t} \langle m | \mathcal{O}_\sigma^\dagger | n \rangle \langle n | \mathcal{O}_\sigma | m \rangle) \\
&= \frac{1}{Z} \sum_{m,n} e^{i(E_m - E_n)t} || \langle m | \mathcal{O}_\sigma | n \rangle ||^2 (e^{-\beta E_m} + e^{-\beta E_n})
\end{aligned} \tag{1.3}$$

The time-domain impurity Green's function can thus be written as (this is the so-called Lehmann-Kallen representation)

$$G_{dd}^\sigma = -i\theta(t) \frac{1}{Z} \sum_{m,n} e^{i(E_m - E_n)t} || \langle m | \mathcal{O}_\sigma | n \rangle ||^2 (e^{-\beta E_m} + e^{-\beta E_n}) \tag{1.4}$$

We are interested in the frequency domain form.

$$\begin{aligned}
G_{dd}^\sigma(\omega) &= \int_{-\infty}^{\infty} dt e^{i\omega t} G_{dd}^\sigma(t) \\
&= \frac{1}{Z} \sum_{m,n} || \langle m | \mathcal{O}_\sigma | n \rangle ||^2 (e^{-\beta E_m} + e^{-\beta E_n}) (-i) \int_{-\infty}^{\infty} dt \theta(t) e^{i(\omega + E_m - E_n)t}
\end{aligned} \tag{1.5}$$

To evaluate the time-integral, we will use the integral representation of the Heaviside function:

$$\theta(t) = \frac{1}{2\pi i} \lim_{\eta \rightarrow 0^+} \int_{-\infty}^{\infty} \frac{1}{x - i\eta} e^{ixt} dx \tag{1.6}$$

With this definition, the integral in $G_{dd}^\sigma(\omega)$ becomes

$$\begin{aligned}
(-i) \int_{-\infty}^{\infty} dt \theta(t) e^{i(\omega + E_m - E_n)t} &= (-i) \frac{1}{2\pi i} \lim_{\eta \rightarrow 0^+} \int_{-\infty}^{\infty} dx \frac{1}{x - i\eta} \int_{-\infty}^{\infty} dt e^{i(\omega + E_m - E_n + x)t} \\
&= (-i) \frac{1}{2\pi i} \lim_{\eta \rightarrow 0^+} \int_{-\infty}^{\infty} dx \frac{1}{x - i\eta} 2\pi \delta(\omega + E_m - E_n + x) \\
&= (-i) \frac{1}{i} \lim_{\eta \rightarrow 0^+} \frac{-1}{\omega + E_m - E_n - i\eta} \\
&= \frac{1}{\omega + E_m - E_n}
\end{aligned} \tag{1.7}$$

The frequency-domain Green's function is thus

$$G_{dd}^{\sigma}(\omega) = \frac{1}{Z} \sum_{m,n} ||\langle m | \mathcal{O}_{\sigma} | n \rangle||^2 (e^{-\beta E_m} + e^{-\beta E_n}) \frac{1}{\omega + E_m - E_n} \quad (1.8)$$

The zero temperature Green's function is obtained by taking the limit of $\beta \rightarrow \infty$. In both the partition function as well as inside the summation, the only term that will survive is the exponential of the ground state energy E_0 .

$$Z \equiv \sum_m e^{-\beta E_m} \Rightarrow \lim_{\beta \rightarrow \infty} Z = d_0 e^{-\beta E_0}, \quad E_0 \equiv \min \{E_n\}$$

where d_0 is the degeneracy of the ground state. The Greens function then simplifies to

$$\begin{aligned} G_{dd}^{\sigma}(\omega, \beta \rightarrow \infty) &= \frac{1}{d_0 e^{-\beta E_0}} \sum_{m,n} ||\langle m | \mathcal{O}_{\sigma} | n \rangle||^2 [e^{-\beta E_m} \delta_{E_m, E_0} + e^{-\beta E_n} \delta_{E_n, E_0}] \frac{1}{\omega + E_m - E_n} \\ &= \frac{1}{d_0} \sum_{n,0} \left[||\langle 0 | \mathcal{O}_{\sigma} | n \rangle||^2 \frac{1}{\omega + E_0 - E_n} + ||\langle n | \mathcal{O}_{\sigma} | 0 \rangle||^2 \frac{1}{\omega - E_0 + E_n} \right] \end{aligned} \quad (1.9)$$

The label 0 sums over all states $|0\rangle$ with energy E_0 . The spectral function is the imaginary part of this Green's function. To extract the imaginary part, we insert an infinitesimal imaginary part in the denominator:

$$G_{dd}^{\sigma}(\omega, \eta) = \frac{1}{d_0} \lim_{\eta \rightarrow 0^-} \sum_{n,0} \left[||\langle 0 | \mathcal{O}_{\sigma} | n \rangle||^2 \frac{1}{\omega + E_0 - E_n + i\eta} + ||\langle n | \mathcal{O}_{\sigma} | 0 \rangle||^2 \frac{1}{\omega - E_0 + E_n + i\eta} \right] \quad (1.10)$$

The spectral function at zero temperature can then be written as

$$\begin{aligned} \mathcal{A}(\omega) &= -\frac{1}{\pi} \text{Im} [G_{dd}^{\sigma}(\omega)] \\ &= \frac{1}{d_0} \frac{1}{\pi} \text{Im} \left[\lim_{\eta \rightarrow 0^-} \sum_{n,0} \left(\frac{-i\eta ||\langle 0 | \mathcal{O}_{\sigma} | n \rangle||^2}{(\omega + E_0 - E_n)^2 + \eta^2} + \frac{-i\eta ||\langle n | \mathcal{O}_{\sigma} | 0 \rangle||^2}{(\omega - E_0 + E_n)^2 + \eta^2} \right) \right] \\ &= \frac{1}{d_0} \frac{1}{\pi} \sum_{n,0} [||\langle 0 | \mathcal{O}_{\sigma} | n \rangle||^2 \pi \delta(\omega + E_0 - E_n) + ||\langle n | \mathcal{O}_{\sigma} | 0 \rangle||^2 \pi \delta(\omega - E_0 + E_n)] \\ &= \frac{1}{d_0} \sum_{n,0} [||\langle 0 | \mathcal{O}_{\sigma} | n \rangle||^2 \delta(\omega + E_0 - E_n) + ||\langle n | \mathcal{O}_{\sigma} | 0 \rangle||^2 \delta(\omega - E_0 + E_n)] \end{aligned} \quad (1.11)$$

9.2 Spectrum of the Hubbard dimer

Here we document the spectrum of the Hubbard dimer Hamiltonian in eqs. ??.

eigenstate	symbol	eigenvalue
$ 0, 0\rangle$	$ 0\rangle$	$\frac{U^H}{2}$
$\frac{1}{\sqrt{2}}(\sigma, 0\rangle \pm 0, \sigma\rangle)$	$ 0\sigma_{\pm}\rangle$	$\mp t^H$
$ \sigma, \sigma\rangle$	$ \sigma\sigma\rangle$	$-\frac{U^H}{2}$
$\frac{1}{\sqrt{2}}(\uparrow, \downarrow\rangle + \downarrow, \uparrow\rangle)$	$ ST\rangle$	$-\frac{U^H}{2}$
$\frac{1}{\sqrt{2}}(2, 0\rangle - 0, 2\rangle)$	$ CS\rangle$	$\frac{U^H}{2}$
$a_1(U^H, t^H)\frac{1}{\sqrt{2}}(\uparrow, \downarrow\rangle - \downarrow, \uparrow\rangle) + a_2(U^H, t^H)\frac{1}{\sqrt{2}}(2, 0\rangle + 0, 2\rangle)$	$ -\rangle$	$-\frac{1}{2}\Delta(U^H, t^H)$
$-a_2(U^H, t^H)\frac{1}{\sqrt{2}}(\uparrow, \downarrow\rangle - \downarrow, \uparrow\rangle) + a_1(U^H, t^H)\frac{1}{\sqrt{2}}(2, 0\rangle + 0, 2\rangle)$	$ +\rangle$	$\frac{1}{2}\Delta(U^H, t^H)$
$\frac{1}{\sqrt{2}}(\sigma, 2\rangle \pm 2, \sigma\rangle)$	$ 2\sigma_{\pm}\rangle$	$\pm t^H$
$ 2, 2\rangle$	$ 4\rangle$	$\frac{U^H}{2}$

Table 9.1: Spectrum of Hubbard dimer at half-filling

9.3 Greens function of Hubbard model in the atomic limit

The atomic limit is described by the Hamiltonian $H = -\frac{U}{2} \sum_i (\hat{n}_{i\uparrow} - \hat{n}_{i\downarrow})^2$. Since the Hamiltonian has decoupled into $\sum_i \equiv N$ single-site Hamiltonians, we can easily write down the retarded Greens function for a single site by identifying the ground state configuration of a site. The ground states are of course $|\Psi\rangle_i = |\uparrow\rangle, |\downarrow\rangle$, in the absence of any symmetry-breaking. The finite temperature retarded Greens function in the time domain is given by

$$G_{i,\sigma}(T, t) = -i\theta(t) \frac{1}{Z} \sum_n e^{-\beta E_n} \langle n | \{c_{i\sigma}(t), c_{i\sigma}^\dagger\} | n \rangle \quad (3.1)$$

where n labels the eigenstates of the H . At $T \rightarrow 0$, only the ground states will survive in the exponential and the partition function Z , such that the formula then reduces to

$$G_{i,\sigma}(T \rightarrow 0, t) = -i\theta(t) \langle GS | \{c_{i\sigma}(t), c_{i\sigma}^\dagger\} | GS \rangle \quad (3.2)$$

Since the Hamiltonian is simple, we can compute the time-dependent operator $c_{i\sigma}(t) = e^{iHt} c_{i\sigma} e^{-iHt}$. Using the BCH lemma, we have

$$e^{iHt} c_{i\sigma} e^{-iHt} = \sum_{n=0}^{\infty} (it)^n \frac{1}{n!} [H, c_{i\sigma}]_n \quad (3.3)$$

where $[H, c_{i\sigma}]_n = [H, [H, c_{i\sigma}]_{n-1}]$, $[H, c_{i\sigma}]_0 = c_{i\sigma}$. The first two non-trivial commutators are

$$[H, c_{i\sigma}]_1 = -U\tau_{i\bar{\sigma}}c_{i\sigma}, \quad \text{and} \quad [H, c_{i\sigma}]_2 = \frac{1}{4}U^2c_{i\sigma}, \quad (3.4)$$

such that

$$\begin{aligned} e^{iHt} c_{i\sigma} e^{-iHt} &= \left[c_{i\sigma} + \frac{i^2 t^2}{2!} \frac{U^2}{2^2} c_{i\sigma} + \frac{i^4 t^4}{4!} \frac{U^4}{2^4} c_{i\sigma} + \dots \right] + \left[-\frac{it}{1!} U\tau_{i\bar{\sigma}}c_{i\sigma} - \frac{i^3 t^3}{3!} \frac{U^3}{4} \tau_{i\bar{\sigma}}c_{i\sigma} + \dots \right] \\ &= c_{i\sigma} [\cos(Ut/2) - 2i\tau_{i\bar{\sigma}} \sin(Ut/2)] \end{aligned} \quad (3.5)$$

Using this, we can write

$$\left\{ c_{i\sigma}(t), c_{i\sigma}^\dagger \right\} = \left\{ c_{i\sigma}, c_{i\sigma}^\dagger \right\} [\cos(Ut/2) - 2i\tau_{i\bar{\sigma}} \sin(Ut/2)] = [\cos(Ut/2) - 2i\tau_{i\bar{\sigma}} \sin(Ut/2)] \quad (3.6)$$

and the Greens function becomes

$$G_{i,\sigma}(T \rightarrow 0, t) = -i\theta(t) \langle GS | [\cos(Ut/2) - 2i\tau_{i\bar{\sigma}} \sin(Ut/2)] | GS \rangle = -i\theta(t) \left[\cos \frac{Ut}{2} - 2i \sin \frac{Ut}{2} \langle \tau_{i\bar{\sigma}} \rangle \right] \quad (3.7)$$

We now Fourier transform to frequency domain:

$$G_{i,\sigma}(\omega) = -i \int_0^\infty dt e^{i\omega t} \left[\cos \frac{Ut}{2} - 2i \sin \frac{Ut}{2} \langle \tau_{i\bar{\sigma}} \rangle \right] = \frac{1 + \langle \tau_{i\bar{\sigma}} \rangle}{\omega - \frac{U}{2}} + \frac{1 - \langle \tau_{i\bar{\sigma}} \rangle}{\omega + \frac{U}{2}} \quad (3.8)$$

The spin-total Greens function for the site i is

$$G_i(\omega) = \sum_\sigma G_{i,\sigma}(\omega) = \frac{1 + \langle \tau_i \rangle}{\omega - \frac{U}{2}} + \frac{1 - \langle \tau_i \rangle}{\omega + \frac{U}{2}} \quad (3.9)$$

At half-filling, we have $\tau_i = 1$.

9.4 Local Greens function for the Hubbard dimer

From the spectral representation, we have the following expression for the local Greens function for the Hubbard dimer at site 0:

$$G_{D,00}^\sigma(\omega) = \frac{1}{Z} \sum_{m,n} ||\langle m | c_{i\sigma} | n \rangle||^2 (e^{-\beta E_m} + e^{-\beta E_n}) \frac{1}{\omega + E_m - E_n} \quad (4.1)$$

m, n sum over the exact eigenstates. E_m, E_n are the corresponding energies. We are interested in the $T \rightarrow 0$ Greens function. In that limit, all exponentials except that for the ground state E_{gs} will die out. The exponential inside the summation will then cancel the exponential in the partition function.

$$\begin{aligned} G_{D,00}^\sigma(\omega, T \rightarrow 0) &= \sum_n \left[||\langle GS | c_{i\sigma} | n \rangle||^2 \frac{1}{\omega + E_{GS} - E_n} + ||\langle n | c_{i\sigma} | GS \rangle||^2 \frac{1}{\omega + E_n - E_{GS}} \right] \\ &= \sum_n \left[||\langle n | c_{i\sigma}^\dagger | GS \rangle||^2 \frac{1}{\omega + E_{GS} - E_n} + ||\langle n | c_{i\sigma} | GS \rangle||^2 \frac{1}{\omega + E_n - E_{GS}} \right] \end{aligned} \quad (4.2)$$

The ground state $|GS\rangle$ is just the state $|-\rangle$ in the table 9.1. We will choose to look at $\sigma = \uparrow$. Then,

$$\begin{aligned} c_{1\uparrow} |-\rangle &= \frac{a_1}{\sqrt{2}} |0, \downarrow\rangle + \frac{a_2}{\sqrt{2}} |\downarrow, 0\rangle \\ c_{1\uparrow}^\dagger |-\rangle &= -\frac{a_1}{\sqrt{2}} |2, \uparrow\rangle + \frac{a_2}{\sqrt{2}} |\uparrow, 2\rangle \end{aligned} \quad (4.3)$$

The set of states $|n\rangle$ that give non-zero inner product $|GS\rangle$ are therefore

$$\begin{aligned}\{|n\rangle\} &= |0 \downarrow_{\pm}\rangle \\ ||\langle n| c_{\uparrow\sigma} |GS\rangle||^2 &= \frac{1}{4} (a_2 \pm a_1)^2 = \frac{1}{4} (1 \pm 2a_1 a_2) \\ \{E_n\} &= \mp t\end{aligned}\tag{4.4}$$

for the second inner product, and

$$\begin{aligned}\{|n\rangle\} &= |2 \uparrow_{\pm}\rangle \\ ||\langle n| c_{\uparrow\sigma}^\dagger |GS\rangle||^2 &= \frac{1}{4} (a_2 \mp a_1)^2 = \frac{1}{4} (1 \mp 2a_1 a_2) \\ \{E_n\} &= \pm t\end{aligned}\tag{4.5}$$

for the first. The Greens function is therefore

$$G_{D,00}^\uparrow(\omega, T \rightarrow 0) = \left(\frac{1}{2} + \frac{2t}{\Delta}\right) \frac{\omega}{\omega^2 - (t - \frac{\Delta}{2})^2} + \left(\frac{1}{2} - \frac{2t}{\Delta}\right) \frac{\omega}{\omega^2 - (t + \frac{\Delta}{2})^2} = G_{D,00}^\downarrow(\omega, T \rightarrow 0) .\tag{4.6}$$

In the atomic limit ($t = 0$), the Greens function simplifies to

$$G_{D,00}^\uparrow(\omega, T \rightarrow 0) \Big|_{\text{atomic}} = \frac{\omega}{\omega^2 - \frac{1}{4}U^2}\tag{4.7}$$

In the atomic limit, the singly-occupied state has zero energy:

$$E_1(t = 0) = \langle 1, 0 | (U\tau_{0\uparrow}\tau_{0\downarrow} + U\tau_{1\uparrow}\tau_{1\downarrow}) | 1, 0 \rangle = 0\tag{4.8}$$

We can write the atomic limit Greens function in terms of this energy and the self energy:

$$G_{D,00}^\uparrow(\omega, T \rightarrow 0) \Big|_{\text{atomic}} = \frac{1}{\omega - E_1(t = 0) - \Sigma(t = 0)} = \frac{1}{\omega - 0 - \frac{U^2}{4\omega}}\tag{4.9}$$

The self energy in the atomic limit can be read off as

$$\Sigma(t = 0) = \frac{U^2}{4\omega}\tag{4.10}$$

The site local spectral function can also be calculated from the local Greens function:

$$\begin{aligned}A(0 \uparrow, \omega) &= -\frac{1}{\pi} \text{Im } G_{D,00}^\uparrow(\omega) \\ &= \left(\frac{1}{4} - \frac{t}{\Delta}\right) \left[\delta(\omega - \frac{1}{2}\Delta - t) + \delta(\omega + \frac{1}{2}\Delta + t) \right] \\ &\quad + \left(\frac{1}{4} + \frac{t}{\Delta}\right) \left[\delta(\omega - \frac{1}{2}\Delta + t) + \delta(\omega + \frac{1}{2}\Delta - t) \right] \\ &= A(0 \downarrow, \omega) .\end{aligned}\tag{4.11}$$

Finally, the inter-site Greens function for the Hubbard dimer is given by

$$G_{D,01}^{\uparrow}(\omega, T \rightarrow 0) = \left(\frac{1}{2} + \frac{2t}{\Delta}\right) \frac{t - \frac{\Delta}{2}}{\omega^2 - (t - \frac{\Delta}{2})^2} + \left(\frac{1}{2} - \frac{2t}{\Delta}\right) \frac{t + \frac{\Delta}{2}}{\omega^2 - (t + \frac{\Delta}{2})^2} = G_{D,01}^{\downarrow}(\omega, T \rightarrow 0). \quad (4.12)$$

Using the diagonal and off-diagonal real space Greens functions, we can now compute the momentum-space Greens functions. The two momentum states are $ka = 0, \pi$. By Fourier transforming, these two Greens functions can be written as

$$\begin{aligned} G(k=0, \sigma) &= \sum_r e^{ikr} G(r, \sigma) = G(r=0, \sigma) + G(r=a, \sigma) = \frac{1/2 + 2t/\Delta}{\omega - t + \Delta/2} + \frac{1/2 - 2t/\Delta}{\omega - t - \Delta/2} \\ G(k=\pi, \sigma) &= \sum_r e^{ikr} G(r, \sigma) = G(r=0, \sigma) - G(r=a, \sigma) = \frac{1/2 + 2t/\Delta}{\omega + t - \Delta/2} + \frac{1/2 - 2t/\Delta}{\omega + t + \Delta/2} \end{aligned} \quad (4.13)$$

9.5 Contributions of various excitations to the site local spectral function

The site local spectral function is

$$A(0 \uparrow, \omega) = \left(\frac{1}{4} - \frac{t}{\Delta}\right) \left[\delta\left(\omega - \frac{\Delta}{2} - t\right) + \delta\left(\omega + \frac{\Delta}{2} + t\right) \right] + \left(\frac{1}{4} + \frac{t}{\Delta}\right) \left[\delta\left(\omega - \frac{\Delta}{2} + t\right) + \delta\left(\omega + \frac{\Delta}{2} - t\right) \right]$$

If the eigenstates of the $N = 1, S^z = -\frac{1}{2}$ sector are $|1\pm \downarrow\rangle$ and those of $N = 3, S^z = \frac{1}{2}$ sector are $|3\pm \uparrow\rangle$, this spectral function originates from the expression:

$$\begin{aligned} A(0 \uparrow, \omega) &= \langle 1 \downarrow_- | c_{0\uparrow} | \text{GS} \rangle \delta\left(\omega + \frac{\Delta}{2} + t\right) + \langle 2 \uparrow_+ | c_{0\uparrow}^\dagger | \text{GS} \rangle \delta\left(\omega - \frac{\Delta}{2} - t\right) \\ &\quad + \langle 1 \downarrow_+ | c_{0\uparrow} | \text{GS} \rangle \delta\left(\omega + \frac{\Delta}{2} - t\right) + \langle 2 \uparrow_- | c_{0\uparrow}^\dagger | \text{GS} \rangle \delta\left(\omega - \frac{\Delta}{2} + t\right) \end{aligned} \quad (5.1)$$

9.6 Relation between single-particle Greens function and the Greens operator ($T = 0$)

The single-particle Greens function is defined as the solution of the equation:

$$(i\partial_t - H(\vec{r})) G(\vec{r}, \vec{r}', t) = \delta(\vec{r} - \vec{r}') \quad (6.1)$$

and is given by the expression

$$G(\vec{r}, \vec{r}', t) = -i\theta(t) \langle \{c(\vec{r}, t) c^\dagger(\vec{r}', 0)\} \rangle \quad (6.2)$$




Figure 9.1: Position, weight and nature of each of the peaks in the Hubbard dimer site local spectral function

This solution can be written in the Lehmann-Kallen representation and at $T = 0$ as

$$G(\vec{r}\sigma, \vec{r}'\sigma, \omega) = \sum_n \left[\frac{\langle GS | c(\vec{r}, \sigma) | n \rangle \langle n | c^\dagger(\vec{r}', \sigma) | GS \rangle}{\omega + E_{GS} - E_n} + \frac{\langle GS | c^\dagger(\vec{r}', \sigma) | n \rangle \langle n | c(\vec{r}, \sigma) | GS \rangle}{\omega + E_n - E_{GS}} \right] \quad (6.3)$$

The sum is over the exact eigenstates of the Hamiltonian. In what follows, we will represent $\vec{r}, \sigma \equiv \nu$ and $\vec{r}', \sigma \equiv \nu'$.

$$\begin{aligned} G(\nu, \nu', \omega) &= \sum_n \left[\frac{\langle GS | c(\nu) | n \rangle \langle n | c^\dagger(\nu') | GS \rangle}{\omega + E_{GS} - E_n} + \frac{\langle GS | c^\dagger(\nu') | n \rangle \langle n | c(\nu) | GS \rangle}{\omega + E_n - E_{GS}} \right] \\ &= \langle GS | c(\nu) \frac{1}{\omega + E_{GS} - H} c^\dagger(\nu') | GS \rangle + \langle GS | c^\dagger(\nu') \frac{1}{\omega + H - E_{GS}} c(\nu) | GS \rangle \end{aligned} \quad (6.4)$$

If we now define a Greens operator

$$\mathcal{G}(\omega, H) = \frac{1}{\omega - (H - E_{GS})} \quad (6.5)$$

we can write the single-particle Greens function as a sum of the matrix elements of this operator:

$$G(\nu, \nu', \omega) = \langle \nu | \mathcal{G}(\omega, H) | \nu' \rangle - \langle \bar{\nu}' | \mathcal{G}(-\omega, H) | \bar{\nu} \rangle = \mathcal{G}(\omega, H)_{\nu, \nu'} - \mathcal{G}(-\omega, H)_{\bar{\nu}', \bar{\nu}} \quad (6.6)$$

where we have defined the states $|\nu\rangle \equiv c^\dagger(\nu) |GS\rangle$ and $|\bar{\nu}\rangle \equiv c(\nu) |GS\rangle$. The two matrix elements can also be represented in their individual spectral representations:

$$\begin{aligned}\mathcal{G}(\omega, H)_{\nu, \nu'} &= \sum_n \frac{\langle GS | c(\nu) | n \rangle \langle n | c^\dagger(\nu') | GS \rangle}{\omega + E_{GS} - E_n} \\ \mathcal{G}(\omega, H)_{\bar{\nu}', \bar{\nu}} &= \sum_n \frac{\langle GS | c^\dagger(\nu') | n \rangle \langle n | c(\nu) | GS \rangle}{\omega + E_{GS} - E_n}\end{aligned}\tag{6.7}$$

9.7 Writing single-particle excitations of ground state in terms of $N = 3, S^z = \frac{1}{2}$ eigenstates

The excited state $c_{0\uparrow}^\dagger |GS\rangle$ can actually be written in terms of the $N = 3, S^z = +\frac{1}{2}$ eigenstates $|3\pm \uparrow\rangle$ defined in table 9.1.

$$|3\pm \uparrow\rangle = \frac{1}{\sqrt{2}} (|\uparrow, 2\rangle \pm |2, \uparrow\rangle), \quad H^D |3\pm \uparrow\rangle = \pm t |3\pm \uparrow\rangle\tag{7.1}$$

In terms of these eigenstates, we can write

$$\begin{aligned}c_{0\uparrow}^\dagger |GS\rangle &= c_{0\uparrow}^\dagger [a_1 |SS\rangle + a_2 |CT\rangle] \\ &= a_2 \frac{1}{\sqrt{2}} |\uparrow, 2\rangle - a_1 \frac{1}{\sqrt{2}} |2, \uparrow\rangle \\ &= (x + y) \frac{1}{\sqrt{2}} |\uparrow, 2\rangle + (x - y) \frac{1}{\sqrt{2}} |2, \uparrow\rangle \\ &= x |3+ \uparrow\rangle + y |3- \uparrow\rangle\end{aligned}\tag{7.2}$$

where $x + y \equiv a_2$ and $x - y \equiv -a_1$. Similarly, for the other site excitation, we can write

$$\begin{aligned}c_{1\uparrow}^\dagger |GS\rangle &= c_{1\uparrow}^\dagger [a_1 |SS\rangle + a_2 |CT\rangle] \\ &= a_2 \frac{1}{\sqrt{2}} |2, \uparrow\rangle - a_1 \frac{1}{\sqrt{2}} |\uparrow, 2\rangle \\ &= (x + y) \frac{1}{\sqrt{2}} |2, \uparrow\rangle + (x - y) \frac{1}{\sqrt{2}} |\uparrow, 2\rangle \\ &= x |3+ \uparrow\rangle - y |3- \uparrow\rangle\end{aligned}\tag{7.3}$$

Solving for x and y gives

$$x = \frac{a_2 - a_1}{2}, \quad y = \frac{a_2 + a_1}{2}\tag{7.4}$$

Similarly, we can also write the single-hole excitation $c_{0\uparrow} |GS\rangle$ in terms of the $N = 1, S^z = -\frac{1}{2}$ eigenstates, $|1\pm \downarrow\rangle$:

$$|1\pm \downarrow\rangle = \frac{1}{\sqrt{2}} (|\downarrow, 0\rangle \pm |0, \downarrow\rangle), \quad H^D |1\pm \downarrow\rangle = \mp t |1\pm \downarrow\rangle\tag{7.5}$$

$$\begin{aligned}
c_{0\uparrow}|GS\rangle &= a_1 \frac{1}{\sqrt{2}} |0, \downarrow\rangle + a_2 \frac{1}{\sqrt{2}} |\downarrow, 0\rangle = y |1+ \downarrow\rangle + x |1- \downarrow\rangle \\
c_{1\uparrow}|GS\rangle &= a_1 \frac{1}{\sqrt{2}} |\downarrow, 0\rangle + a_2 \frac{1}{\sqrt{2}} |0, \downarrow\rangle = y |1+ \downarrow\rangle - x |1- \downarrow\rangle
\end{aligned} \tag{7.6}$$

9.8 Matrix elements of G^{-1} between single-particle momentum excitations, for the Hubbard dimer

$$G^{-1} \equiv \omega + E_{\text{GS}} - H_D \tag{8.1}$$

The particle excitation momentum space kets are $|k_0\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$, $|k_\pi\rangle = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$. Therefore,

$$\begin{aligned}
(G^{-1})_{k_0 k_0} &= \frac{1}{2} (\langle 0| + \langle 1|) (\omega + E_{\text{GS}} - H_D) (|0\rangle + |1\rangle) \\
&= \frac{1}{2} (2x \langle +|) (\omega + E_{\text{GS}} - H_D) (2x |+\rangle) \\
&= 2x^2 (\omega + E_{\text{GS}} - t)
\end{aligned} \tag{8.2}$$

At the final step, we used $\langle +, + \rangle = 1$ and $\langle +|H_D|+\rangle = t$.

9.9 Topological interpretation of the Wilson ratio

From the Friedel sum rule [Langer and Ambegaokar(1961)], we can relate the phase shift $\delta(0)$ due to scattering (at the Fermi surface) off a local impurity to the number of electrons bound in the potential well produced by that impurity:

$$\tilde{N} = \frac{1}{2\pi i} \text{Tr} \ln S(0) = \int_{\Gamma} dz \partial_z \frac{1}{2\pi i} \text{Tr} \ln S(0) \tag{9.1}$$

From the optical theorem, we can write

$$S = 1 + TG_0 = \frac{G}{G_0} \quad [G = G_0 + G_0 T G_0] \tag{9.2}$$

This allows us to write [Mukherjee and Lal(2020a)]

$$\tilde{N} = \int_{\Gamma} dz \partial_z \frac{1}{2\pi i} \text{Tr} \ln \frac{G}{G_0} \tag{9.3}$$

Since $\text{Tr} \ln \hat{O} = \sum_{\lambda} \ln O_{\lambda} = \ln \prod_{\lambda} O_{\lambda} = \ln \text{Det} \hat{O}$, we get

$$\begin{aligned}
 \tilde{N} &= \int_{\Gamma} dz \partial_z \frac{1}{2\pi i} \ln \text{Det} \frac{G}{G_0} \\
 &= - \int_{\Gamma} dz \partial_z \frac{1}{2\pi i} \ln \frac{\text{Det} G_0}{\text{Det} G} \\
 &\equiv - \int_{\Gamma} dz \partial_z \frac{1}{2\pi i} \ln D \\
 &= - \int_{\Gamma(D)} \frac{dD}{D}
 \end{aligned} \tag{9.4}$$

From the work of Seki and Yunoki [Seki and Yunoki(2017)], we know that this quantity is essentially the winding number of the curve $\Gamma(D)$ in the complex plane spanned by the real and imaginary parts of D , and is equal to the change in Luttinger's volume V_L at $T = 0$.

$$\tilde{N} = - \int_{\Gamma(D)} \frac{dD}{D} = -\Delta V_L \tag{9.5}$$

The incoming electrons can have $\sigma = \uparrow, \downarrow$. Since the impurity singlet ground state is rotationally invariant, we have $\delta_{\uparrow} = \delta_{\downarrow} = \delta(0)$.

$$\begin{aligned}
 \tilde{N} &= \frac{1}{\pi} \sum_{\sigma} \delta_{\sigma}(0) \\
 \Rightarrow \delta(0) &= \frac{\pi}{2} \tilde{N} = -\frac{\pi}{2} \Delta V_L
 \end{aligned} \tag{9.6}$$

$$\begin{aligned}
 R &= 1 + \sin^2 \left(\frac{\pi}{2} \tilde{N} \right) \\
 &= 1 + \sin^2 \left(\frac{\pi}{2} \Delta V_L \right)
 \end{aligned} \tag{9.7}$$

We note that this connection between R and ΔV_L has not been obtained in the existing literature thus far. In the unitary limit, $\delta(0) = \frac{\pi}{2}$, giving $\Delta V_L = -1 = -\tilde{N}$ [Martin(1982b)] (i.e., one electronic state from the impurity has been absorbed into the Luttinger volume of the conduction bath), such that $R = 2$ in this limit. In this way, we see that a change in the topological quantum number $\tilde{N}$ causes the well known renormalisation of the Wilson ratio R from its non-interacting value (1) to the value (2) obtained for the local Fermi liquid [Nozieres(1974)].