

# INSIGHTS ON MOTT TRANSITION AND THE PSEUDOGAP THROUGH THE VEIL OF A QUANTUM IMPURITY MODEL

**ABHIRUP MUKHERJEE**

DPS DAY '25

DEPARTMENT OF PHYSICAL SCIENCES, IISER KOLKATA

MARCH 19, 2025





**Prof. Siddhartha Lal**



**Debraj Debata**



**Siddhartha Patra**  
(Multiverse Computing)



~~~~~  
**Financial support by IISER K and SERB is gratefully  
acknowledged.**  
~~~~~



**Prof. Anamitra Mukherjee**  
(NISER Bhubaneswar)



**Prof. Arghya Taraphder**  
(IIT Kharagpur)



**Prof. N. S. Vidhyadhiraja**  
(JNCASR Bangalore)

# THE ADVENT OF QUANTUM MATERIALS

- Display exotic properties arising due to strong electronic/spin **interactions**.
- **Examples:** heavy fermion compounds ( $\text{CePd}_2\text{Si}_2$ ), copper oxides ( $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ ), iron pnictides ( $\text{BaFe}_2(\text{As}_{1-x}\text{P}_x)_2$ ), twisted bilayer graphene.

# THE ADVENT OF QUANTUM MATERIALS

- Display exotic properties arising due to strong electronic/spin **interactions**.
- **Examples:** heavy fermion compounds ( $\text{CePd}_2\text{Si}_2$ ), copper oxides ( $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ ), iron pnictides ( $\text{BaFe}_2(\text{As}_{1-x}\text{P}_x)_2$ ), twisted bilayer graphene.


## What Makes Them Special - Phenomenology of The Cu-O Superconductors

# THE ADVENT OF QUANTUM MATERIALS

- Display exotic properties arising due to strong electronic/spin **interactions**.
- **Examples:** heavy fermion compounds ( $\text{CePd}_2\text{Si}_2$ ), copper oxides ( $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ ), iron pnictides ( $\text{BaFe}_2(\text{As}_{1-x}\text{P}_x)_2$ ), twisted bilayer graphene.

## What Makes Them Special - Phenomenology of The Cu-O Superconductors

- Cu-O planes separated by insulating layers that can be doped.







# THE ADVENT OF QUANTUM MATERIALS

- Display exotic properties arising due to strong electronic/spin **interactions**.
- **Examples:** heavy fermion compounds ( $\text{CePd}_2\text{Si}_2$ ), copper oxides ( $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ ), iron pnictides ( $\text{BaFe}_2(\text{As}_{1-x}\text{P}_x)_2$ ), twisted bilayer graphene.

## What Makes Them Special - Phenomenology of The Cu-O Superconductors

- Cu-O planes separated by insulating layers that can be doped.
- Strange metal phase beyond 1-particle description (**breakdown of perturbation theory**)







# THE ADVENT OF QUANTUM MATERIALS

- Display exotic properties arising due to strong electronic/spin **interactions**.
- **Examples:** heavy fermion compounds ( $\text{CePd}_2\text{Si}_2$ ), copper oxides ( $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ ), iron pnictides ( $\text{BaFe}_2(\text{As}_{1-x}\text{P}_x)_2$ ), twisted bilayer graphene.

## What Makes Them Special - Phenomenology of The Cu-O Superconductors

- Cu-O planes separated by insulating layers that can be doped.
- Strange metal phase beyond 1-particle description (**breakdown of perturbation theory**)
- Pseudogap phase has Fermi arcs and competing fluctuations (**strong correlations**)





# THE ADVENT OF QUANTUM MATERIALS


- Display exotic properties arising due to strong electronic/spin **interactions**.
- **Examples:** heavy fermion compounds ( $\text{CePd}_2\text{Si}_2$ ), copper oxides ( $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ ), iron pnictides ( $\text{BaFe}_2(\text{As}_{1-x}\text{P}_x)_2$ ), twisted bilayer graphene.

## What Makes Them Special - Phenomenology of The Cu-O Superconductors

- Cu-O planes separated by insulating layers that can be doped.
- Strange metal phase beyond 1-particle description (**breakdown of perturbation theory**)
- Pseudogap phase has Fermi arcs and competing fluctuations (**strong correlations**)
- Superconductivity is **unconventional** (gap vanishes at points on Fermi surface)



s-wave SC gap    d-wave SC gap






# OPEN QUESTIONS

The “normal phases” remain **ill-understood**:

- simple and universal mechanism for strange metals (akin to Fermi liquid theory)
- Microscopic understanding of the pseudogap phase
- How to connect  $T = 0$  results to those at  $T > 0$ ?



# OPEN QUESTIONS



The “normal phases” remain **ill-understood**:

- simple and universal mechanism for strange metals (akin to Fermi liquid theory)
- Microscopic understanding of the pseudogap phase
- How to connect  $T = 0$  results to those at  $T > 0$ ?

Simplest model for realising these phases

- the 2D **Hubbard model** on square lattice

$$H = -t \sum_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger c_{j,\sigma} + \text{h.c.}) + U \sum_i n_{i\uparrow} n_{i\downarrow}$$



# OPEN QUESTIONS

The “normal phases” remain **ill-understood**:

- simple and universal mechanism for strange metals (akin to Fermi liquid theory)
- Microscopic understanding of the pseudogap phase
- How to connect  $T = 0$  results to those at  $T > 0$ ?



Simplest model for realising these phases

- the 2D **Hubbard model** on square lattice

$$H = -t \sum_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger c_{j,\sigma} + \text{h.c.}) + U \sum_i n_{i\uparrow} n_{i\downarrow}$$

**Existing methods** suffer from a variety of problems:


- Quantum Monte Carlo suffers from fermion **sign problem** at low temperatures
- Dynamical mean-field theory: low lack of  **$k$ -space resolution** & large computation cost
- DMRG only works in **quasi-1D** systems



# A BOTTOM-UP APPROACH: STARTING FROM AN IMPURITY MODEL

## What is an Impurity Model?


- Single **correlated site**, hybridising with conduction **bath**



# A BOTTOM-UP APPROACH: STARTING FROM AN IMPURITY MODEL

## What is an Impurity Model?


- Single **correlated site**, hybridising with conduction **bath**
- Impurity-bath hybridisation respects  $C_4$  **symmetry** of 2D square lattice (important!)



# A BOTTOM-UP APPROACH: STARTING FROM AN IMPURITY MODEL

## What is an Impurity Model?

- Single **correlated site**, hybridising with conduction **bath**
- Impurity-bath hybridisation respects  $C_4$  **symmetry** of 2D square lattice (important!)
- (Much!) **Simpler to solve** than Hubbard model, because of non-interacting conduction bath




# A BOTTOM-UP APPROACH: STARTING FROM AN IMPURITY MODEL

## What is an Impurity Model?

- Single **correlated site**, hybridising with conduction **bath**
- Impurity-bath hybridisation respects  $C_4$  **symmetry** of 2D square lattice (important!)
- (Much!) **Simpler to solve** than Hubbard model, because of non-interacting conduction bath

## Mapping to the Lattice Model


- Impurity model describes local environment of each site.



# A BOTTOM-UP APPROACH: STARTING FROM AN IMPURITY MODEL



## What is an Impurity Model?

- Single **correlated site**, hybridising with conduction **bath**
- Impurity-bath hybridisation respects  $C_4$  **symmetry** of 2D square lattice (important!)
- (Much!) **Simpler to solve** than Hubbard model, because of non-interacting conduction bath



## Mapping to the Lattice Model


- Impurity model describes local environment of each site.
- **Translation** operator maps impurity model quantities to those on the lattice model





# MOMENTUM-SPACE RESOLVED IMPURITY PHASE TRANSITION





**Raising bath correlation**  $W$  tunes the impurity model through three phases



# MOMENTUM-SPACE RESOLVED IMPURITY PHASE TRANSITION

**Raising bath correlation  $W$**  tunes the impurity model through three phases





- Impurity **strongly coupled** to Fermi surface



# MOMENTUM-SPACE RESOLVED IMPURITY PHASE TRANSITION

**Raising bath correlation  $W$**  tunes the impurity model through three phases





- Impurity **strongly coupled** to Fermi surface
- Impurity coupled **only to parts** of Fermi surface



# MOMENTUM-SPACE RESOLVED IMPURITY PHASE TRANSITION





**Raising bath correlation  $W$**  tunes the impurity model through three phases

- Impurity **strongly coupled** to Fermi surface
- Impurity coupled **only to parts** of Fermi surface
- Impurity **decoupled** from the Fermi surface




# PSEUDOGAPPING TRANSITION ON THE LATTICE MODEL

- Map Greens functions from impurity model to **lattice model**.
- Momentum-space DOS reveals **partially gapped** Fermi surface.




# MAIN TAKEAWAYS

**Simplifies** the study of lattice models through appropriate **impurity models**.




# MAIN TAKEAWAYS

**Simplifies** the study of lattice models through appropriate **impurity models**.




Our impurity model realises a **pseudo-gapping transition** in a correlated model.




# MAIN TAKEAWAYS

**Simplifies** the study of lattice models through appropriate **impurity models**.



Our impurity model realises a **pseudo-gapping transition** in a correlated model.



## Generalisations and Extensions

- Tune impurity filling - simulate **doping**!
- Multiple impurities - symmetry-broken or **spin liquid** phases