

INSIGHTS ON MOTT TRANSITION AND THE PSEUDOGAP THROUGH THE VEIL OF A QUANTUM IMPURITY MODEL

ABHIRUP MUKHERJEE

DPS DAY '25

DEPARTMENT OF PHYSICAL SCIENCES, IISER KOLKATA

MARCH 19, 2025





Prof. Siddhartha Lal



Debraj Debata



Siddhartha Patra
(Multiverse Computing)



~~~~~  
**Financial support by IISER K and SERB is gratefully  
acknowledged.**  
~~~~~



Prof. Anamitra Mukherjee
(NISER Bhubaneswar)




Prof. Arghya Taraphder
(IIT Kharagpur)



Prof. N. S. Vidhyadhiraja
(JNCASR Bangalore)



THE ADVENT OF QUANTUM MATERIALS: CUPRATE SUPERCONDUCTORS

- Cu-O planes separated by insulating layers that can be doped.




THE ADVENT OF QUANTUM MATERIALS: CUPRATE SUPERCONDUCTORS

- Cu-O planes separated by insulating layers that can be doped.
- **Strange metal** phase beyond 1-particle description



1-electron
picture





many-electron
picture




THE ADVENT OF QUANTUM MATERIALS: CUPRATE SUPERCONDUCTORS

- Cu-O planes separated by insulating layers that can be doped.
- **Strange metal** phase beyond 1-particle description
- * **Missing:** Simple and universal mechanism for strange metals



1-electron
picture






many-electron
picture







THE ADVENT OF QUANTUM MATERIALS: CUPRATE SUPERCONDUCTORS

- Cu-O planes separated by insulating layers that can be doped.
- **Strange metal** phase beyond 1-particle description
 - * **Missing:** Simple and universal mechanism for strange metals
- **Pseudogap** phase has Fermi arcs and competing fluctuations






THE ADVENT OF QUANTUM MATERIALS: CUPRATE SUPERCONDUCTORS

- Cu-O planes separated by insulating layers that can be doped.
- **Strange metal** phase beyond 1-particle description
 - * **Missing:** Simple and universal mechanism for strange metals
- **Pseudogap** phase has Fermi arcs and competing fluctuations
 - * **Missing:** Microscopic understanding of the pseudogap phase




THE ADVENT OF QUANTUM MATERIALS: CUPRATE SUPERCONDUCTORS



Simplest model for realising these phases
- the 2D **Hubbard model** on square lattice

$$H = -t \sum_{\langle i,j \rangle, \sigma} (c_{i,\sigma}^\dagger c_{j,\sigma} + \text{h.c.}) + U \sum_i n_{i\uparrow} n_{i\downarrow}$$


Too hard to solve! Alternative approaches needed.



A BOTTOM-UP APPROACH: STARTING FROM AN IMPURITY MODEL

What is an Impurity Model?


- Single **correlated site**, hybridising with conduction **bath**



A BOTTOM-UP APPROACH: STARTING FROM AN IMPURITY MODEL

What is an Impurity Model?


- Single **correlated site**, hybridising with conduction **bath**
- Impurity-bath hybridisation respects C_4 **symmetry** of 2D square lattice (important!)



A BOTTOM-UP APPROACH: STARTING FROM AN IMPURITY MODEL

What is an Impurity Model?

- Single **correlated site**, hybridising with conduction **bath**
- Impurity-bath hybridisation respects C_4 **symmetry** of 2D square lattice (important!)
- (Much!) **Simpler to solve** than Hubbard model, because of non-interacting conduction bath




A BOTTOM-UP APPROACH: STARTING FROM AN IMPURITY MODEL

What is an Impurity Model?

- Single **correlated site**, hybridising with conduction **bath**
- Impurity-bath hybridisation respects C_4 **symmetry** of 2D square lattice (important!)
- (Much!) **Simpler to solve** than Hubbard model, because of non-interacting conduction bath

Mapping to the Lattice Model


- Impurity model describes local environment of each site.



A BOTTOM-UP APPROACH: STARTING FROM AN IMPURITY MODEL



What is an Impurity Model?

- Single **correlated site**, hybridising with conduction **bath**
- Impurity-bath hybridisation respects C_4 **symmetry** of 2D square lattice (important!)
- (Much!) **Simpler to solve** than Hubbard model, because of non-interacting conduction bath




Mapping to the Lattice Model

- Impurity model describes local environment of each site.
- **Translation** operator maps impurity model quantities to those on the lattice model



MOMENTUM-SPACE RESOLVED IMPURITY PHASE TRANSITION





Raising bath correlation W tunes the impurity model through three phases



MOMENTUM-SPACE RESOLVED IMPURITY PHASE TRANSITION

Raising bath correlation W tunes the impurity model through three phases





- Impurity **strongly coupled** to Fermi surface



MOMENTUM-SPACE RESOLVED IMPURITY PHASE TRANSITION

Raising bath correlation W tunes the impurity model through three phases





- Impurity **strongly coupled** to Fermi surface
- Impurity coupled **only to parts** of Fermi surface



MOMENTUM-SPACE RESOLVED IMPURITY PHASE TRANSITION





Raising bath correlation W tunes the impurity model through three phases

- Impurity **strongly coupled** to Fermi surface
- Impurity coupled **only to parts** of Fermi surface
- Impurity **decoupled** from the Fermi surface




PSEUDOGAPPING TRANSITION ON THE LATTICE MODEL

- Map Greens functions from impurity model to **lattice model**.
- Momentum-space DOS reveals **partially gapped** Fermi surface.




MAIN TAKEAWAYS

Simplifies the study of lattice models through appropriate **impurity models**.




MAIN TAKEAWAYS

Simplifies the study of lattice models through appropriate **impurity models**.




Our impurity model realises a **pseudo-gapping transition** in a correlated model.




MAIN TAKEAWAYS

Simplifies the study of lattice models through appropriate **impurity models**.



Our impurity model realises a **pseudo-gapping transition** in a correlated model.



Generalisations and Extensions

- Tune impurity filling - simulate **doping**!
- Multiple impurities - symmetry-broken or **spin liquid** phases