

LOCAL METAL-INSULATOR TRANSITION IN AN EXTENDED ANDERSON IMPURITY MODEL

JRF-to-SRF Upgradation Presentation

Abhirup Mukherjee

August 14, 2022

Supervisor: Dr. Siddhartha Lal

Department of Physical Sciences, IISER Kolkata, Mohanpur



Summary of Work

Summary of Work

Completed Projects

- ✓ Unveiling the Kondo cloud: Unitary renormalization-group study of the Kondo model

Phys. Rev. B 105, 085119, arXiv:2111.10580v3

A. Mukherjee, Abhirup Mukherjee, N. S. Vidhyadhiraja, A. Taraphder, and S. Lal

- ✓ Frustration shapes multi-channel Kondo physics: A star graph perspective

under review at PRB, arXiv:2205.00790

S. Patra, Abhirup Mukherjee, A. Mukherjee, N. S. Vidhyadhiraja, A. Taraphder, S. Lal



Ongoing Projects

- ✓ Metal-insulator transition in an extended Anderson impurity model (*manuscript in preparation*)
-

- ✓ Holography and topology of entanglement scaling in free fermions (*manuscript in preparation*)

- ✓ URG-based auxiliary model approach to correlated systems (*ongoing*)


Local MIT in an extended Anderson impurity model




Introducing the extended Anderson impurity model

DMFT on the Bethe lattice: Exact in $d = \infty$


- ✓ shows **metal-insulator transition** on the Bethe lattice with ∞ coordination number



- ✓ Conduction bath obtained by imposing self-consistency shows **non-trivial correlations**




- ✓ Spectral function develops three peaks and then **gaps out**



Introducing the extended Anderson impurity model

Standard Anderson impurity model

- ✓ no local-moment phase, $A(\omega)$ gapless
- ✓ cannot explain insulating phase of DMFT




Gap in spectral function requires additional physics!



Extended Anderson impurity model

- ✓ impurity-bath spin correlation: J
- ✓ bath zeroth site local correlation: U_b



Phase Diagram & Ground-States



Nature of RG flows

- ✓ URG Equations reveal **critical** point at $r = -U_b/J = 1/4$



$$\Delta J = J(J + 4U_b)/|d_3|$$

- ✓ allows averting strong-coupling behaviour
- ✓ U_b always marginal




RG Phase Diagram

✓ blue phase $\rightarrow U_b < -J/4$: V, J are **irrelevant** \rightarrow local moment flows

✓ yellow phase: $J \ll D_0$: involves **V, U, U_b**
vanishes for large systems

✓ gray phase: $J \sim D_0$: **all** couplings irrelevant
vanishes for large systems





✓ red phase $\rightarrow U_b > -J/4$: V, J are **relevant** \rightarrow strong-coupling flows

Low-energy effective Hamiltonians and ground-states

Regime 1: $|U_b| < J/4$

- ✓ J relevant,
- ✓ V subdominant,
- ✓ U irrelevant

$$H = J\vec{S}_d \cdot \vec{S}_0 - U_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2 + \sum_{k < \Lambda^*} \epsilon_k \hat{n}_{k\sigma}$$



$$H = J\vec{S}_d \cdot \vec{S}_0 - U_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2$$

Zero-bandwidth limit



- ✓ two-spin Heisenberg, attractive zeroth site
- ✓ **singlet** ground state


$$|\Psi\rangle_{GS} = \frac{1}{\sqrt{2}} [|\uparrow_d \downarrow_0\rangle - |\downarrow_d \uparrow_0\rangle]$$

Low-energy effective Hamiltonians and ground-states


Regime 2: $|U_b| \sim J/4$

- ✓ J relevant,
- ✓ V relevant,
- ✓ U irrelevant

$$H = J\vec{S}_d \cdot \vec{S}_0 + V \sum_{\sigma} (c_{d\sigma}^\dagger c_{0\sigma} + \text{h.c.}) - U_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2 + \sum_{k < \Lambda^*} \epsilon_k \hat{n}_{k\sigma}$$



Zero-bandwidth limit



$$H = J\vec{S}_d \cdot \vec{S}_0 + V \sum_{\sigma} (c_{d\sigma}^\dagger c_{0\sigma} + \text{h.c.}) - U_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2$$

- ✓ **spin+charge** dimer with attractive zeroth site
- ✓ spin-singlet + charge-triplet-zero in gr-state


$$|\Psi\rangle_{GS} = \frac{1}{2\sqrt{2}} [|\uparrow_d \downarrow_0\rangle - |\downarrow_d \uparrow_0\rangle] + \frac{1}{2\sqrt{2}} [|\downarrow_d, 0_0\rangle - |0_d, 2_0\rangle]$$

Low-energy effective Hamiltonians and ground-states

Regime 3: $|U_b| > J/4$

- ✓ J, V irrelevant,
- ✓ U relevant,

$$H = -U/2(\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2 - U_b(\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2 + \sum_{k<\Lambda^*} \epsilon_k \hat{n}_{k\sigma}$$



impurity site



zeroth site

$$H = -U/2(\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2 - U_b(\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2$$

Zero-bandwidth limit




- ✓ impurity site detaches from bath
- ✓ **local moment** ground-state

$$|\Psi\rangle_{GS} = |\uparrow, \downarrow\rangle_d \otimes |0, 2\rangle_0$$

Low-energy effective Hamiltonians and ground-states


Ground-state overlaps with spin singlet and charge triplet zero



Nature of the Transition

Gapping of the impurity spectral function

- ✓ Broad central peak at $|U_b| \ll J/4$




- ✓ Correlated **three peak** structure at $|U_b| \lesssim J/4$

- ✓ hard central **gap** for $|U_b| > J/4$


Destruction of the Kondo cloud

The Kondo cloud **weakens, and is destroyed** at the transition.

✓ vanishing of impurity-bath correlations




✓ transfer of entanglement into the bath




Growth of pairing fluctuations in the bath

Subdominant pairing fluctuations, near the transition...

- ✓ growth of fluctuations in Cooper channel, at the cost of spin-flip fluctuations



- ✓ mutual information within the bath maximised after transition




Universal Theory near the Transition

Minimal effective model for the transition


- ✓ For $|U_b| \leq J/4$, central peak and side peaks are **well-separated**
 - ✓ **Integrate out** charge fluctuations through Schrieffer-Wolff transformation

$$H_{\text{eff}} = \tilde{J} \vec{S}_d \cdot \vec{S}_0 - U_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2 + H_{\text{KE}}$$

RG equation for \tilde{J} : $\Delta\tilde{J} \sim \tilde{J}(\tilde{J} + 4U_h)$




- ✓ **captures** the criticality, and the strong-coupling and local moment phases



Suggests that J and U_h are the minimal & universal ingredients for transition!

Capturing the level crossing at the transition from a two-site model



- ✓ Obtain two-site model by taking **zero bandwidth** limit
- ✓ spectrum shows **level crossing** between singlet and local moment states



Insights into DMFT


Extended SIAM in the context of DMFT

DMFT represents excursions from a site into the correlated bath in the form of self-energies for the bath and the impurity. This results in a three-peak DOS of the bath.



Equivalence of the impurity site and the bath zeroth site

- ✓ Integrate out impurity site from fixed point Hamiltonian via a single URG transformation
- ✓ Generates additional  correlation U_0 on zeroth site




Essence of self-consistency: Equivalence of impurity and zeroth sites! 

Equivalence of the impurity site and the bath zeroth site

- ✓ J is relevant and the largest scale → **repulsive correlation**:

$$U_0 + U_b \approx J > 0$$

- ✓ J acts a **symmetrisation mechanism** between impurity and zeroth sites
- ✓ **Coherent** spin-flip scatterings ensure similarity of spectral functions




Essence of self-consistency: Equivalence of impurity and zeroth sites! 

Observation of a coexistence region


- ✓ DMFT observes a **coexistence region** near the critical point, for $U_{c1} < U < U_{c2}$

- ✓ Insulating when coming in from the insulator, metallic when coming in from the metal



- ✓ spectral function shows **preformed gap** from metallic side


- ✓ **True** transition believed to occur at U_{c2}



Observation of a coexistence region

Can be explained heuristically using the two site spectrum


- ✓ Initial point is when the side peaks get separated (near-zeroes in the spectral function)



Observation of a coexistence region

Can be explained heuristically using the two site spectrum


- ✓ U_{c2} is the point where the levels cross



Observation of a coexistence region

Can be explained heuristically using the two site spectrum


- ✓ Coming from $U > U_{c2}$, **adiabatic continuity** allows DMFT to stay on the local moment state...



Observation of a coexistence region

Can be explained heuristically using the two site spectrum

- ✓ For $U < U_{c1}$, local moment state is too unstable, **relaxes** to the true ground state.




Comparison against NRG-DMFT correlation functions

Poor Man's scaling of the effective Kondo model

[K. Held, R. Peters, and A. Toschi. PRL 2013]

- shows **quantitative** agreement with NRG-DMFT (crossover scale and kinks in self-energy)
- Suggests that the minimal model can capture **spin susc.**

[K. Held, R. Peters, and A. Toschi.
PRL 110, 246402 (2013)]



- Our $J - U_b$ model **goes further** by capturing physics beyond the transition


Comparison against NRG-DMFT correlation functions

Doublon-holon correlators of the Hubbard model



[S. B. Lee, J. v Delft, and A. Weichselbaum. PRL 2017]



Lee et. al show **peaks** in doublon-holon correlators near zero energy within the central peak.



- We find support for this in the form of **increasing ground-state charge correlations and overlap**.



Low-energy excitations of the bath

effective Hamiltonian for the excitations of the bath near the transition

emergence of NFL terms at the critical point

Broad conclusions

- The extended SIAM appears to capture the DMFT transition and **self-consistency**.
- **Minimal ingredients** are the Kondo physics J and the attractive local correlation U_b .
- Crucial feature of the journey is the enhancement of **pairing fluctuations** in the bath.
- SOMETHING ABOUT THE FINAL CALCULATIONS (MAYBE NON-FERMI LIQUID PHYSICS)



Future Prospects

Future Prospects

- ✓ The extended SIAM can be improved by considering **multiple impurities** and general impurity **filling**.
- ✓ We are developing a new **tiling-based auxiliary model method** can used for studying other models of strong-correlations as well as topologically active or flat band systems.
- ✓ The URG can be applied to **heavy-fermion materials** towards a study of phase diagram and unconventional superconductivity, as well as Kondo insulators.
- ✓ Interacting systems in a magnetic field is also a potential area of study, specifically **fractional Chern insulators** (e.g. the fractional quantum hall effects).

Acknowledgements

Gracious thanks to

- ✓ my collaborators **S. Patra, A. Mukherjee, Prof. A. Taraphdar** and **Prof. N. S. Vidhyadhiraja**,
- ✓ my RPC members **Prof. Ritesh Singh** and **Prof. Anandamohan Ghosh** for instructive feedback,
- ✓ **Prof. H. Casini, Prof. N. Banerjee** and **Shibendu G. Chowdhury** for very fruitful discussions, and
- ✓ IISER Kolkata for funding.

References

- Affleck, Ian (1995). "Conformal field theory approach to the Kondo effect". In: *Acta Phys.Polon.B* 26.1869.
- Affleck, Ian and Andreas W.W. Ludwig (1991). "Critical theory of overscreened Kondo fixed points". In: *Nuclear Physics B* 360.2, pp. 641–696. ISSN: 0550-3213.
- Affleck, Ian and Andreas WW Ludwig (1993). "Exact conformal-field-theory results on the multichannel Kondo effect: Single-fermion Green's function, self-energy, and resistivity". In: *Physical Review B* 48.10, p. 7297.
- Anderson, P. W. (1961). "Localized Magnetic States in Metals". In: *Phys. Rev.* 124 (1), pp. 41–53.
- — (1978). "Local moments and localized states". In: *Rev. Mod. Phys.* 50 (2), pp. 191–201.
- Anderson, PW (1970). "A poor man's derivation of scaling laws for the Kondo problem". In: *Journal of Physics C: Solid State Physics* 3.12, p. 2436.
- Andrei, N. (1980). "Diagonalization of the Kondo Hamiltonian". In: *Phys. Rev. Lett.* 45 (5), pp. 379–382.
- Andrei, N, K Furuya, and J H Lowenstein (1983). "Solution of the Kondo problem". In: *Rev. Mod. Phys.* 55, p. 331.
- Arias, Raúl E, David D Blanco, and Horacio Casini (2015). "Entanglement entropy as a witness of the Aharonov–Bohm effect in QFT". In: *Journal of Physics A: Mathematical and Theoretical* 48.14, p. 145401.
- Bulla, R, A C Hewson, and Th Pruschke (1998). "Numerical renormalization group calculations for the self-energy of the impurity Anderson model". In: *Journal of Physics: Condensed Matter* 10.37, pp. 8365–8380.
- Calabrese, Pasquale and John Cardy (2004). "Entanglement entropy and quantum field theory". In: *Journal of Statistical Mechanics: Theory and Experiment* 2004.06, P06002.
- Casini, H, C D Fosco, and M Huerta (2005). "Entanglement and alpha entropies for a massive Dirac field in two dimensions". In: *Journal of Statistical Mechanics: Theory and Experiment* 2005.07, P07007–P07007.
- Chen, Xiao et al. (2017). "Two-cylinder entanglement entropy under a twist". In: *Journal of Statistical Mechanics: Theory and Experiment* 2017.4, p. 043104.
- Costi, T. A. (2000). "Kondo Effect in a Magnetic Field and the Magnetoresistivity of Kondo Alloys". In: *Phys. Rev. Lett.* 85 (7), pp. 1504–1507.
- Costi, T.A. and A.C. Hewson (1990). "A new approach to the calculation of spectra for strongly correlated systems". In: *Physica B: Condensed Matter* 163.1, pp. 179–181. ISSN: 0921-4526.
- Cox, D. L. and N. Grewe (1988). "Transport properties of the Anderson lattice". In: *Zeitschrift für Physik B Condensed Matter* 71.3, pp. 321–340.
- Emery, V. J. and S. Kivelson (1992). "Mapping of the two-channel Kondo problem to a resonant-level model". In: *Phys. Rev. B* 46 (17), pp. 10812–10817.
- Gan, J (1994). "On the multichannel Kondo model". In: 6.24, pp. 4547–4568.
- Georges, Antoine and Gabriel Kotliar (1992). "Hubbard model in infinite dimensions". In: *Physical Review B* 45.12, p. 6479.
- Goldhaber-Gordon, D. et al. (1998). "Kondo effect in a single-electron transistor". In: *Nature* 391.6663, pp. 156–159. ISSN: 1476-4687.
- Heath, Joshua T and Kevin S Bedell (2020). "Necessary and sufficient conditions for the validity of Luttinger's theorem". In: *New Journal of Physics* 22.6, p. 063011.
- Held, K., R. Peters, and A. Toschi (2013). "Poor Man's Understanding of Kinks Originating from Strong Electronic Correlations". In: *Phys. Rev. Lett.* 110 (24), p. 246402.
- Hewson, A. C. (1993). *The Kondo Problem to Heavy Fermions*. Cambridge University Press.
- Holzhey, Christoph, Finn Larsen, and Frank Wilczek (1994). "Geometric and renormalized entropy in conformal field theory". In: *Nuclear Physics B* 424.3, pp. 443–467. ISSN: 0550-3213.
- Kondo, Jun (1964). "Resistance minimum in dilute magnetic alloys". In: *Progress of theoretical physics* 32.1, pp. 37–49.
- Kotliar, G. et al. (2006). "Electronic structure calculations with dynamical mean-field theory". In: *Rev. Mod. Phys.* 78 (3), pp. 865–951.

References

- ▶ Krishna-murthy, H. R., J. W. Wilkins, and K. G. Wilson (1980). "Renormalization-group approach to the Anderson model of dilute magnetic alloys. I. Static properties for the symmetric case". In: *Phys. Rev. B* 21 (3), pp. 1003–1043.
- ▶ Kuramoto, Y and T Watanabe (1987). "Theory of momentum-dependent magnetic response in heavy-fermion systems". In: *Proceedings of the Yamada Conference XVIII on Superconductivity in Highly Correlated Fermion Systems*. Elsevier, pp. 80–83.
- ▶ Lee, Ching Hua and Xiao-Liang Qi (2016). "Exact holographic mapping in free fermion systems". In: *Physical Review B* 93.3, p. 035112.
- ▶ Lee, Sung-Sik (2010). "Holographic description of quantum field theory". In: *Nuclear Physics B* 832.3, pp. 567–585.
- ▶ — (2014). "Quantum renormalization group and holography". In: *Journal of High Energy Physics* 2014.1, p. 76.
- ▶ Luttinger, JM (1960). "Fermi surface and some simple equilibrium properties of a system of interacting fermions". In: *Physical Review* 119.4, p. 1153.
- ▶ Luttinger, Joaquin Mazdak and John Clive Ward (1960). "Ground-state energy of a many-fermion system. II". In: *Physical Review* 118.5, p. 1417.
- ▶ Maier, Thomas et al. (2005). "Quantum cluster theories". In: *Rev. Mod. Phys.* 77 (3), pp. 1027–1080.
- ▶ Maldacena, Juan (1999). "The large-N limit of superconformal field theories and supergravity". In: *International journal of theoretical physics* 38.4, pp. 1113–1133.
- ▶ Metzner, Walter and Dieter Vollhardt (1989). "Correlated Lattice Fermions in $d = \infty$ Dimensions". In: *Phys. Rev. Lett.* 62 (3), pp. 324–327.
- ▶ Mukherjee, Anirban and Siddhartha Lal (2020a). "Unitary renormalisation group for correlated electrons-I: a tensor network approach". In: *Nuclear Physics B* 960, p. 115170.
- ▶ — (2020b). "Unitary renormalisation group for correlated electrons-II: insights on fermionic criticality". In: *Nuclear Physics B* 960, p. 115163.
- ▶ — (2022). "Superconductivity from repulsion in the doped 2D electronic Hubbard model: an entanglement perspective". In: *Journal of Physics: Condensed Matter* 34.27, p. 275601.
- ▶ Murciano, Sara, Paola Ruggiero, and Pasquale Calabrese (2020). "Symmetry resolved entanglement in two-dimensional systems via dimensional reduction". In: *Journal of Statistical Mechanics: Theory and Experiment* 2020.8, p. 083102.
- ▶ Nozaki, Masahiro, Shinsei Ryu, and Tadashi Takayanagi (2012). "Holographic geometry of entanglement renormalization in quantum field theories". In: *Journal of High Energy Physics* 2012.10, p. 193.
- ▶ Nozieres, P (1974). "A "Fermi-liquid" description of the Kondo problem at low temperatures". In: *Journal of Low Temperature Physics* 17, p. 31.
- ▶ Nozières, Ph. and Blandin, A. (1980). "Kondo effect in real metals". In: *J. Phys. France* 41.3, pp. 193–211.
- ▶ Ohashi, Takuma et al. (2008). "Finite Temperature Mott Transition in Hubbard Model on Anisotropic Triangular Lattice". In: *Phys. Rev. Lett.* 100 (7), p. 076402.
- ▶ Oshikawa, Masaki (2000). "Topological approach to Luttinger's theorem and the Fermi surface of a Kondo lattice". In: *Physical Review Letters* 84.15, p. 3370.
- ▶ Parcollet, O., G. Biroli, and G. Kotliar (2004). "Cluster Dynamical Mean Field Analysis of the Mott Transition". In: *Phys. Rev. Lett.* 92 (22), p. 226402.
- ▶ Qi, Xiao-Liang (2013). "Exact holographic mapping and emergent space-time geometry". In: *arXiv preprint arXiv:1309.6282*.
- ▶ Ryu, Shinsei and Tadashi Takayanagi (2006a). "Aspects of holographic entanglement entropy". In: *Journal of High Energy Physics* 2006.08, p. 045.
- ▶ — (2006b). "Holographic derivation of entanglement entropy from the anti-de sitter space/conformal field theory correspondence". In: *Physical review letters* 96.18, p. 181602.
- ▶ Sakai, Osamu, Yukihiko Shimizu, and Tadao Kasuya (1989). "Single-Particle and Magnetic Excitation Spectra of Degenerate Anderson Model with Finite f-f Coulomb Interaction". In: *Journal of the Physical Society of Japan* 58.10, pp. 3666–3678. eprint: <https://doi.org/10.1143/JPSJ.58.3666>.
- ▶ Seki, Kazuhiro and Seiji Yunoki (2017). "Topological interpretation of the Luttinger theorem". In: *Physical Review B* 96.8, p. 085124.

References


- ▶ Sen, Sudeshna, Patrick J. Wong, and Andrew K. Mitchell (2020). "The Mott transition as a topological phase transition". In: *Phys. Rev. B* 102 (8), p. 081110.
- ▶ Tsvelick, A M and P B Wiegmann (1983). "Exact results in the theory of magnetic alloys". In: *Adv. in Phys.* 32, p. 453.
- ▶ Tsvelick, A. M. and P. B. Wiegmann (1985). "Exact solution of the multichannel Kondo problem, scaling, and integrability". In: *Journal of Statistical Physics* 38.1, pp. 125–147. ISSN: 1572-9613.
- ▶ V. Borzenets, Ivan et al. (2020). "Observation of the Kondo screening cloud". In: *Nature* 579.7798, pp. 210–213. ISSN: 1476-4687.
- ▶ Van Raamsdonk, Mark (2010). "Building up spacetime with quantum entanglement". In: *General Relativity and Gravitation* 42.10, pp. 2323–2329.
- ▶ Wilson, Kenneth G. (1975). "The renormalization group: Critical phenomena and the Kondo problem". In: *Rev. Mod. Phys.* 47 (4), pp. 773–840.
- ▶ Wilson, Kenneth G. and J. Kogut (1974). "The renormalization group and the ϵ -expansion". In: *Physics Reports* 12.2, pp. 75 –199.

Other Projects

Theory for the single-channel Kondo cloud



Phys. Rev. B 105, 085119

Anirban Mukherjee, **Abhirup Mukherjee**, N. S. Vidhyadhiraja, A. Taraphder, and Siddhartha Lal



Theory for the single-channel Kondo cloud

- ✓ spectral function & magnetic susceptibility




- ✓ local Fermi liq. & orthogonality catastrophe
- ✓ thermal entropy

Role of degeneracy in the multi-channel Kondo problem



arXiv:2205.00790

Siddhartha Patra, **Abhirup Mukherjee**, Anirban Mukherjee, N. S. Vidhyadhiraja, A. Taraphder, Siddhartha Lal




Role of degeneracy in the multi-channel Kondo problem

- ✓ Intermediate-coupling RG fixed point Hamiltonian and **degenerate** ground states
- ✓ Degree of compensation, magnetization and susceptibility show **incomplete screening**





Role of degeneracy in the multi-channel Kondo problem

- ✓ Local **marginal Fermi liquid** within the low-energy excitations of the bath
- ✓ **Duality** relations constrain the RG flows of the MCK model




Entanglement scaling in free fermions: holography & topology



Entanglement scaling in free fermions: holography & topology - Summary

- Under coarse-graining or fine-graining in k -space, entanglement of free Dirac fermions shows hierarchical onion-like structure.
- Entanglement scaling can be used to define distances, leads to additional spatial dimension \rightarrow holography.
- Emergent distances and curvature can be related to RG beta function; the sign of the curvature is topological.
- Pole structure of the entanglement tracks the Luttinger volume - invariant under the scaling transformations.



Creating subsystems

Free Dirac fermions on torus: $k_x^n = \frac{2\pi}{L_x} n, \quad n \in \mathbb{Z}; \quad$ define **sparsity** = $\Delta n = 1$



Simplest choice: the entire set

sparsity = 1 $\longrightarrow n \in \{-N, -(N - 1), -(N - 2), \dots, -1, 0, 1, \dots, N - 2, N - 1, N\}$

Coarser choices: increase sparsity

sparsity = 2 $\longrightarrow n \in \{-N, -(N - 2), -(N - 4), \dots, -2, 0, 2, \dots, N - 4, N - 2, N\}$

sparsity = 4 $\longrightarrow n \in \{-N, -(N - 4), -(N - 8), \dots, -4, 0, 4, \dots, N - 8, N - 4, N\}$



Subsystem entanglement entropy: Entanglement hierarchy

$$S_{A_z(j)} = f_z(j) c \alpha L_x - c \log |2 \sin(\pi f_z(j)\phi)|$$

$$i < j, \quad S_{i \cup j} = \begin{cases} S_i, & z > 0 \\ S_j, & z < 0 \end{cases}$$



- ✓ presents a **hierarchy** of entanglement → EE distributed across RG steps
RG transformation → reveals entanglement
- ✓ distribution of entanglement also present in **multipartite** entanglement

Mutual information = distance


Mutual information: $I^2(A : B) \equiv S(A) + S(B) - S(A \cup B)$ (non-negative)

Define distances using mut. info.

$$x_z(j) = \log t_z(j), \quad y_z(j) = \log t_z(j \pm 1)$$

$$v_z(j) \equiv \Delta y_z(j)/\Delta x_z(j), \quad v' = \Delta v_z(j)/\Delta x_z(j)$$

Curvature as well: $\kappa_z(j) = \frac{v'_z(j)}{\left[1 + v_z(j)^2\right]^{\frac{3}{2}}}$



RG evolution = emergent distance

- ✓ Distances and curvature can be related to an RG **beta function**
- ✓ Amounts to an **explicit demonstration** of the holographic principle
- ✓ Sign of curvature is **topological**, can be written in terms of winding numbers

Topological nature of geometry-independent term

$$S_{A_z(j)} = f_z(j)caL_x - \underbrace{c \log |2 \sin(\pi f_z(j)\phi)|}_{=Q(\phi), \text{geometry-independent term}}$$

- ✓ $Q(\phi)$ is periodic in the flux ϕ , $\phi = 1$ transports a charge across Fermi surface
- ✓ pole structure of $(\sin \frac{\pi}{4} - |\sin(\pi f_z(j)\phi)|)^{-1}$ counts number of states → tracks Luttinger volume
- ✓ Luttinger volume is topological, so is $Q(\phi)$; $Q(\phi)$ can be expressed in terms of winding numbers